

Early Life History of *Lefua costata* (Cypriniformes : Balitoridae) from Korea

Jae-Min Park¹, Seong-Jang Cho², and †Kyeong-Ho Han³

¹Gyeongsangbuk-Do Native Fish Business Center, Uiseong 37366, Korea

²Boryeong Freshwater Eco Center, Boryeong 33416, Korea

³Fishery Science, Chonnam National University, Yeosu 59626, Korea



Received: October 18, 2020

Revised: October 24, 2020

Accepted: November 1, 2020

†Corresponding author

Kyeong-Ho Han

Fishery Science, Chonnam National University, Yeosu 59626, Korea.

Tel: +82-61-659-7163

E-mail: aqua05@jnu.ac.kr

Copyright © 2020 The Korean Society of Developmental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Park Jae Min

<https://orcid.org/0000-0001-6398-1748>

Cho Seong Jang

<https://orcid.org/0000-0002-7347-1706>

Han Kyeong Ho

<https://orcid.org/0000-0001-5918-3196>

Conflict of interests

The authors declare no potential conflict of interest.

Acknowledgements

Not applicable.

Authors' contributions

Conceptualization: Park JM

Data curation: Park JM

Methodology: Cho SJ, Han KH

Software: Park JM, Cho SJ

Validation: Cho SJ, Han KH

Investigation: Cho SJ

Writing-original draft: Park JM

Writing-review & editing: Han KH

Ethics approval

This article does not require IRB/IACUC

Abstract

The purpose of this study is to observe the early life history of Korean *Lefua costata* and use the result as basic taxonomic research data for balitorid fish. The fertilized eggs were light green color with completely circle shape and mean size was 1.21 ± 0.06 mm ($n=30$). Immediately after hatching, the size of the larvae was 2.81 ± 0.11 mm ($n=5$) in mean length, with egg yolk. On the 3rd day after hatching, the preflexion larvae had a mean length of 4.64 ± 0.09 mm ($n=5$), and their mouth was opened to start feeding. On the 8th day after hatching, a mean length of the postflexion larvae was 9.43 ± 0.46 mm ($n=5$), the distal part of the notochord was bent to 45° , and 16 fin rays were developed on the caudal fin. On the 31st day after hatching, a mean length of juveniles was 22.3 ± 0.85 mm ($n=5$), and the number of fin rays was the same as that of adult fish with (iv8) dorsal fins and (iii8) anal fins.

Keywords: Egg development, Juveniles, Larvae, *Lefua costata*, Early life history

INTRODUCTION

In particular, many traits revealed during the development of fish provide important basic data for clarifying the taxonomic relationships of allied species (Blaxter, 1974; Balon, 1985). In the study of the systematics traits of fish of the Cypriniformes during the development, the characteristics of the fertilized eggs were found as demersal egg and epipelagic eggs even within the *Squalidus* same genus. Melanophore vesicles were formed after hatching (AH) for short fishes with a hatching time of 1–2 days. At the end of the egg development, melanophore vesicles are developed in the eye of the embryo. This will be used as a systematics trait to grasp the taxonomic relations of the genus *Squalidus* (Song et al., 2017).

Fish belonging to Cypriniformes, Cobitoidei, and balitorid are distributed in Europe and Asia (Kottelat, 2012; Nelson et al., 2016). Balitorid fish living in Korea includes *Orthobias toni*, *O. nudus*, and *Lefua costata* (Kim, 1997; Kim et al., 2005; Kim & Park, 2007), of which *L. costata* is distributed in Korea, Japan, and the Maritime Province of Siberia in Russia (Kim, 1997; Kim et al., 2005).

The genus *Lefua* fish in Korea are morphologically similar to the fish of the same genus in Japan. They show distinguishable characteristics that appeared in early life history. This study is to investigate the differences in the characteristics of genus *Lefua* fish between Korea and Japan.

approval because there are no human and animal participants.

Genus *Lefua* fish living in Japan are 5 species including *L. costata*, *L. nikkonis*, *L. echigonia*, *L. torrentis*, and *L. tokaiensis* (Hosoya, 2019). In case of *L. torrentis*, previous research by Aoyama & Doi (2011) distinguished the morphological differences of the head and the distribution location of melanocyte appearing in the early life period. In addition, it was found that *L. sp.* lives in the same water stream as *L. echigonia* but *L. echigonia* lives in wetland areas and *L. sp.* in mountainous valley areas. Later, *L. sp.* was reported as a new species *L. torrentis* by Hosoya et al. (2018).

Record of early life history for *L. costata* only been recorded Uchida (1939). It is necessary to clearly observe the characteristics of the early life history that appear during the development.

Therefore, this study aims to observe the early life history, such as egg development and fish larvae morphology of *L. costata* and compare the results with those of related species to use them as basic data for taxonomic studies.

MATERIALS AND METHODS

1. Collecting broodstocks

The broodstocks used in the study were collected using a stake net at Sanguncheon located in Sonyang-myeon, Yangyang-gun, Gangwon-do in June 2018, and were transported to the laboratory.

Housed in a square water tank (1×1×1 m), the broodstocks were bred in a raceway culture, and frozen mosquito larvae were supplied twice/day as feed (Hikari, Japan), and the water temperature was 22°C–24°C (mean 23±1°C).

2. Induction of spawning

For broodstocks, females had a total length of 7–9 cm and a weight of 4–5 g (n=10), and males had a total length of 5–7 cm and a weight of 2–3 g (n=20). In order to induce natural spawning, males and females were anesthetized with an anesthetic (MS-222, Ethyl 3-aminobenzoate methanesulfonate, Sigma-Aldrich, St. Louis, MO, USA). Then a sexual hormone for maturity (LHRH, Syndel, Nanaimo, Canada) was injected into the muscle on the basis of 0.5 mL per kg. In order to stimulate the water temperature, the breeding of water temperature in the tank was raised from 23°C–25°C (mean 24±1°C).

3. Egg development and fish larvae morphology

The fertilized eggs, the rearing of water temperature was maintained at 24°C–26°C (mean 25±1°C). In order to observe the egg development, 30 eggs were randomly selected and observed with a stereomicroscope (JP SMZ800, Nikon, Tokyo, Japan), and up to 0.01 mm was measured using a universal projector (JP V-12BM, Nikon). For the food of fish larvae, nauplius larvae of *Artemia* sp. were supplied from 15 days after hatching (DAH). From then on, the dry feed (350 µm, Jeilfeed, Daejeon, Korea) was kneaded and supplied. In order to describe the morphological characteristics from juvenile fish which were selected, the external morphology of five was observed with a stereoscopic microscope immediately AH, and the size was measured up to 0.01 mm with a universal projector. We followed the study of Okiyama (1988) for the criteria for each developmental stage of fish larvae. Ontogenetic patterns of the melanophore was observed by selecting 8 sites for each growth stage (Fig. 1).




Fig. 1. The parts for observation of melanophore of larvae and juveniles of *Lefua costata*.

RESULTS

1. Number and characteristics of spawned eggs

The female spawning amount was about 900–1,200 eggs (mean $1,050 \pm 150$). The eggs were a light green color with completely circle shape, and were adherent demersal eggs with adhesiveness on the surface. The egg sizes were 1.15–1.27 mm (mean 1.21 ± 0.06 mm, $n=30$).

2. Egg development

At 30 minutes after fertilization (AF), blastodisc was formed on the upper part of the egg yolk (Fig. 2A), and it reached the 2-cell stage as the cleavage occurred in the animal pole in 1 hour AF




Fig. 2. Eggs development of *Lefua costata*. Scale bars=1.00 mm. Time required for each development stage is shown in Table 1.

Table 1. Eggs development of *Lefua costata* at water temperature 25±1°C

Stage	Elapsed time	Characters	Fig. 2
Cell cleave period			
Blastodisc	0 h 30 min	Blastodisc is formed	A
Two cell	1 h 00 min	2–1 array of blastomeres	B
Four cell	1 h 30 min	2–2 array of blastomeres	C
Eight cell	2 h 00 min	2–4 array of blastomeres	D
Sixteen cell	2 h 30 min	4–4 array of blastomeres	E
Thirty-two cell	3 h 00 min	4–8 array of blastomeres	F
Sixty-four cell	3 h 30 min	8–8 array of blastomeres	G
Morula	4 h 00 min	The size of the blastomere is getting smaller	H
Blastula	6 h 30 min	The surface of the blastomere coincides with the egg yolk	I
Gastrulation 1/3	9 h 00 min	Covered 1/3 of egg yolk	J
Gastrulation 2/3	10 h 30 min	Covered 2/3 of egg yolk	K
Gastrulation 3/3	12 h 00 min	Covered 3/3 of egg yolk	L
Embryonic period			
	14 h 00 min	Development of embryo	M
	19 h 00 min	Development of Kuffer's vesicle	N
	30 h 00 min	Appearance of otocyst and long tail	O
Embryo just before hatching	33 h 00 min	The movement of the embryonic body was fastered	P
Hatching period			
	42 h 00 min	Hatching started	Q
	48 h 00 min	Hatching rate 50%	R

(Fig. 2B). At 1 hour 30 minutes AF, it reached the 4-cell stage as the meridional cleavage occurred in the same size (Fig. 2C). At 2 hours AF, it reached the 8-cell stage in the same manner (Fig. 2D), and reached the 16-cell stage at 2 hours 30 minutes AF (Fig. 2E). At 3 hours AF, it reached the 32-cell stage as the number of cell divisions increased (Fig. 2F), and at 3 hours 30 minutes AF, it reached the 64-cell stage (Fig. 2G). At 4 hours AF, it reached the loss stage as the number of cells increased beyond count (Fig. 2H). At 6 hours 30 minutes AF, it reached the blastula stage (Fig. 2I), and at 9 hours AF, it reached the early gastrula stage as covering down from the animal pole to the vegetal pole (Fig. 2J). At 10 hours 30 minutes AF, half of the 50% was covered and it reached the middle gastrula stage (Fig. 2K), and at 12 hours AF, 90% was covered and it reached the late gastrula stage (Fig. 2L). At 14 hours AF, the blastopore was closed and an embryonic body was formed (Fig. 2M). At 19 hours AF, the optic vesicle was formed, 10 myotomes were formed, and Kuffer's vesicle was formed (Fig. 2N). At 30 hours AF, Kuffer's vesicle was lost. As the tail became longer, it began to separate from the egg yolk, and blood flow was observed from the heart along the egg yolk (Fig. 2O). At 33 hours AF, the tail lengthened to the upper part of the head, and the movement of the embryonic body became active (Fig. 2P). At 42 hours AF, melanocyte began to deposit in the eyes, and hatching started as they broke through the egg membrane from the head (Fig. 2Q). At 48 hours AF, more than 50% of the fertilized eggs were hatched (Fig. 2R).

3. Fish larval morphology

The larvae immediately AH are 2.70–2.93 mm in total length (mean 2.81±0.11 mm, n=5), their mouth and anus were not open, had egg yolk, and the entire body was covered with membrane fins. Melanocytes were deposited in the eyes, and the body was lying sideways on the floor with no swimming ability, and the movement of the tail was observed (Fig. 3A).

On the 1st day AH, the egg yolk-sac larvae was 3.71–3.87 mm in total length (mean 3.79±0.08




Fig. 3. Morphological development of yolk-sac larvae and postflexion larvae and juveniles of *Lefua costata*. (A) Newly hatched yolk-sac larvae, mean 2.81 mm TL (total length), (B) 1 days after hatching (DAH), mean 3.79 mm TL, (C) 8 DAH, mean 9.43 mm TL, (D) 15 DAH, mean 13.0 mm TL, (E) 20 DAH, mean 15.5 mm TL, (F) 26 DAH, mean 18.5 mm TL, (G) 31 DAH, mean 22.3 mm TL. Scale bars=1.00 mm.

mm, n=5), and twig-shaped melanocyte was deposited on the upper part of the head, body, and the upper part of the egg yolk. The melanocyte deposited on the eyes became darker, and the anus began to open (Fig. 3B).

On the 8th day AH, the postflexion larvae are 8.97–9.90 mm in total length (mean 9.43 ± 0.46 mm, n=5), 7 fin rays were developed in the differentiated dorsal fins, and 4 fin rays were developed as the anal fins begins to differentiate. The distal part of the notochord was bent 45° to the upper part, and 16 fin rays were developed in the caudal fins. The medial fin was in the form of a non-differentiated membrane fin, and the dorsal, anal and caudal fins had developed girdle but were connected by a membrane. The melanocyte deposited on the body became spot-shaped and mostly disappeared in the body part, and some remained in the head and the front part of the body (Fig. 3C).

On the 15th day AH, the postflexion larvae are 12.2–13.8 mm in total length (mean 13.0 ± 0.80 mm, n=5), and the number of fin rays for each part increased to 9 dorsal fins and 6 anal fins, and 3 fin rays were formed in ventral fins. Membranous dorsal, anal, caudal, and ventral fins were all separated, and the caudal fins were made of a membrane from the caudal part to the middle part of fin rays. In the abdomen, one air bladder in the shape of a cylinder was divided into two, which

repeated staying stationary and swimming in the middle layer. Two pairs of barbels were formed around the mouth (Fig. 3D).

On the 20th day AH, the postflexion larvae are 14.8–16.3 mm in total length (mean 15.5 ± 0.75 mm, $n=5$), and the number of fin rays for each part increased to 12 dorsal fins, 10 anal fins and 17 caudal fins, and 3 fin rays were formed in ventral fins. The melanocyte was scattered in the shape of small spots on the upper part around the upper part of the gill and the side line (Fig. 3E).

On the 26th day AH, juveniles are 17.7–19.4 mm in total length (mean 18.5 ± 0.85 mm, $n=5$), and the number of fin rays for each part reached a whole number, and the number of barbels increased to 4 pairs, and the length of the barbels was shorter than the ones located in the middle between the top and bottom (Fig. 3F).

On the 31st day AH, juveniles are 21.5–23.2 mm in total length (mean 22.3 ± 0.85 mm, $n=5$), and the fin rays for each part were IV8 dorsal fins and III8 anal fins, and the melanocyte in the shape of irregular small spots along the middle of the lateral line of the body formed a horizontal band, and the outer shape was the same as that of broodstocks (Fig. 3G).

DISCUSSION

The eggs of *L. costata* are demersal eggs with adherent properties and exhibited the same properties as *M. mizolepis*, *Kichulchoia multifasciata*, *Koreocobitis naktongensis* belonging to Cobitidae (Kim et al., 1987; Kim & Lee, 1995; Song et al., 2009). The size of the eggs is 1.15–1.27 mm (mean 1.21 ± 0.06 mm, $n=30$), which is similar to mean 1.12 mm of *M. mizolepis* (Kim et al., 1987), Cobitid fish and smaller than mean 2.50 mm of *K. multifasciata* (Kim & Lee, 1995). Immediately AH, the larvae were 2.70–2.93 mm in total length (mean 2.81 ± 0.11 mm, $n=5$), and Uchida (1939) recorded the total length as 3.90 mm. Given that the back of the head is connected to the end of egg yolk with the membrane fin, and melanocyte is not deposited on other areas except the eyes, the morphological characteristics were not significantly different from the results of this study.

In this study, we compared the morphological characteristics of the larvae of *L. costata* with its related species (eg. Japanese *L. echigonia* and *L. torrentis*). The body lengths of hatching larvae of *L. echigonia* and *L. torrentis* are 2.56–2.60 mm and 3.28–4.28 mm, respectively, which are larger than *L. echigonia* but smaller than *L. torrentis*, showing the middle size of the two species (Table 2). The melanocyte of hatching larvae of *L. costata* was generally deposited in the eyes, but that of *L. echigonia*, and *L. torrentis* were not deposited. The egg yolk larvae of *L. costata* was 3.71–3.87 mm,

Table 2. Total length and body length (mm) of each developmental stage of *Lefua* genus fishes

Stage	Species			
	<i>Lefua costata</i> ¹⁾ in Sanguncheon, Korea (TL, mm)	<i>L. costata</i> ²⁾ (TL, mm)	<i>L. echigonia</i> ³⁾ in Kako river, Japan (BL, mm)	<i>L. sp. (L. torrentis)</i> ³⁾ in Kako river, Japan (BL, mm)
Hatched larvae	2.70–2.93 (2.81)	3.9	2.56–2.60*	3.28–4.28
Yolk-sac larvae	3.71–3.87 (3.79)	-	-	5.16–5.56
Preflexion larvae	4.55–4.74 (4.64)	4.1	4.21–4.92	5.78–6.19
Flexion larvae	5.94–6.23 (6.08)	-	5.46–7.64	6.17–8.75
Postflexion larvae	8.97–9.90 (9.43)	5.5	7.23–11.3	8.74–11.0
Juveniles	17.7–19.4 (18.5)	17.5	12.3–13.0	11.4–12.9

* Total length of *L. echigonia*, ¹⁾ Present study, ²⁾ Uchida (1939), ³⁾ Aoyama & Doi (2011).

TL, total length; BL, body length.

and twig-shaped melanocyte was deposited in the upper part of the head, back and even the top of the egg yolk and tails. *L. torrentis* was 5.16–5.56 mm in body length, and twig-shaped melanocyte was deposited in the upper part of the head, back and even the top of the egg yolk and tails, and the distribution position was similar to that of *L. costata* and *L. torrentis* had an egg yolk, an open mouth, and developed pectoral fins, whereas *L. costata* during a similar period did not have an open mouth nor developed pectoral fins, showing a morphological difference during the egg yolk larvae period.

The preflexion larvae of *L. costata* were 4.55–4.74 mm in total length, and the melanocyte deposited in the upper part of the head was deposited more widely. During this period, their mouth was open and the pectoral fins developed. The preflexion larvae of *L. echiogonia* are 4.21–4.92 mm in body length, and the upper part of the head had the spot-shaped melanocyte which is smaller than that of the twig-shaped melanocyte deposited in the entire body. The preflexion larvae of *L. torrentis* were 5.78–6.19 mm in body length, and both small spot-shaped and twig-shaped melanocytes were deposited in the upper part of the head, so the distribution location of the melanocyte during the preflexion larvae period showed an intermediate form between *L. costata* and *L. echiogonia*.

The flexion larvae of *L. costata* are 5.94–6.23 mm in total length, and the size of the melanocyte deposited in the trunk and the upper part of the egg yolk decreased, and the notochord distal end of the tail part began to bend to 45°, and fin rays were developed in the caudal fin. The flexion larvae of *L. echiogonia* are 5.46–7.64 mm in body length, and the size of small spot-shaped melanocyte on the upper part of the head increased, and it was widely deposited along the upper part of the gills and the middle of the body. The flexion larvae of *L. torrentis* are 6.17–8.75 mm in body length, and the melanocyte deposited on the upper part of the head was deposited to a similar size, and the distribution range of the melanocyte in the middle and anterior part of the body became wider. The flexion larvae of *L. costata* had a similar form of the melanocyte deposited on the upper part of the gills to that of *L. torrentis*, and the flexion larvae of *L. echiogonia* were widely distributed, showing a difference.

The postflexion larvae of *L. costata* are 8.97–9.90 mm in body length, and the size of the twig-shaped melanocyte deposited on the upper part of the head decreased, and the fins of each part were developed, and fin rays were formed as the dorsal and anal fins developed. The postflexion larvae of *L. echiogonia* are 7.23–11.3 mm in body length, and the melanocyte deposited on the upper part of the gills, the middle of the body, and the dorsal side was deposited more widely, and fin rays were formed on the dorsal and anal fins. The postflexion larvae of *L. torrentis* are 8.74–11.0 mm in body length, and the melanocyte deposited on the upper part of the head, the middle of the body, and the dorsal side became smaller in the shape of small spots, and was deposited all over the body except for the back of the snout and the upper part of the gill. The developmental pattern of melanocytes during the postflexion larvae period was similar to that of *L. torrentis*. During this period, the membranes connected to the dorsal, anal and caudal fins did not disappear, which was the same as that of closely related species.

Juveniles of *L. costata* are 17.7–19.4 mm in total length, and the shape of the melanocyte was similar to that of adult fish. One pair of barbels was developed in the nostrils and three pairs of barbels around the mouth. Juveniles of *L. echiogonia* are 12.3–13.0 mm in body length, and most of the melanocyte was irregularly deposited on the upper half of the body. One pair of barbels was developed in the nostrils and three pairs of barbels around the mouth. Juveniles of *L. torrentis* are 11.4–12.9 mm in body length, and small spot-shaped melanocyte was evenly deposited all over the body. One pair of barbels was developed in the nostrils and three pairs of barbels around the mouth. In the case of *L. costata* and closely related species, the fins connected by the membrane were all separated into the dorsal, anal and caudal fins as migrating to the juvenile period, and the number of barbels was the same (Fig. 4).


















Stages	Species		
	<i>L. costata</i> ¹⁾	<i>L. echigonia</i> ²⁾	<i>L. torrentis</i> ²⁾
Hatched larvae	 * 2.70-2.93 mm	 * 2.56-2.60 mm	 3.28-4.28 mm
Yolk-sac larvae	 * 3.71-3.87 mm	-	 5.16-5.56 mm
Preflexion larvae	 * 4.55-4.74 mm	 4.21-4.92 mm	 5.78-6.19 mm
Flexion larvae	 * 5.94-6.23 mm	 5.46-7.64 mm	 6.17-8.75 mm
Postflexion larvae	 * 8.97-9.90 mm	 7.23-11.3 mm	 8.74-11.0 mm
Juveniles	 * 17.7-19.4 mm	 12.3-13.0 mm	 11.4-12.9 mm

Fig. 4. Comparison of larvae and juveniles melanophore distribution in *Lefua* genus fishes. **L. costata* and *L. echigonia* in newly hatched larvae : total length; *L. echigonia* and *L. torrentis* : body length. ¹⁾ Present study, ²⁾ Aoyama & Doi (2011) with CC-BY.

Aoyama & Doi (2011) suggested the external morphological differences between the two species because *L. echigonia*'s eye size is larger than the length from mouth to nostrils, and *L. torrentis* has an eye size smaller than the length from mouth to nostrils. *L. costata*'s eye size (0.003–0.005 mm) is similar to the length from mouth to nostrils (0.003–0.005 mm), showing a difference from *L. echigonia*, and *L. torrentis*.

When observed from the back side, the cornea of the eye is off the head in the *L. costata*'s head, and the same shape was also seen when observed from the bottom. When observed from the back and the bottom, the *L. echigonia*'s cornea was outside the head, while the cornea was located inside the head in *L. torrentis*, and the head shape of *L. costata* was similar to that of *L. echigonia*. Ito et al. (2019) compared the distribution of melanocytes in the dorsal and caudal fins including the head shape with the closely related species and reported *L. tokaiensis* as a new species. When comparing the result with that of *L. costata*, *L. tokaiensis* has the melanocyte deposited on its dorsal and caudal fin, and the dark band is formed at the starting point of the caudal fin. *L. echigonia* has the distribution pattern of melanocytes which is similar to that of *L. tokaiensis*, but showed a difference because there was no dark band, and melanocytes and dark bands were not observed on the dorsal and caudal fins. *L. costata* was similar to *L. torrentis* among the closely related species as no melanocyte and dark bands were observed on the dorsal and caudal fins (Fig. 5).

According to the results of the study, *L. costata*'s head shape, size of hatching larvae showed the intermediate shape between Japanese *L. echigonia* and *L. torrentis*, and the distribution locations of




Fig. 5. Pigmentation and shapes of the dorsal and ventral view of head and dorsal and caudal fins in *Lefua costata*. (A) dorsal view of head, (B) ventral view of head, (C) dorsal fin, (D) caudal fin. Scale bars=1.0 mm.

melanocytes were found to vary according to the growth stage of fish larvae, showing differences from the closely related species, thus providing basic data for taxonomic studies.

REFERENCES

- Aoyama S, Doi T (2011) Morphological comparison of early stages of two Japanese species of eight-barbel loaches: *Lefua echigonia* and *Lefua* sp. (Nemacheilidae). *Folia Zool* 60:355-361.
- Balon EK (1985) *Early Life Histories of Fishes: New Developmental, Ecological and Evolutionary Perspectives*. Dr W Junk, Dordrecht, Nederland, p 280.
- Blaxter JHS (1974) *The Early Life History of Fish*. Springer-Verlag Berlin, Heidelberg, Germany, p 765.
- Hosoya K (2019) *Japanese Freshwater Fish, Supplemented and Revised Compiled*. Mountain and Gyeoksha Shrine, Tokyo, Japan, p 560.
- Hosoya K, Ito T, Miyazaki JI (2018) *Lefua torrentis*, a new species of loach from western Japan (Teleostei: Nemacheilidae). *Ichthyol Expl Freshwater* 2:1-9.
- Ito TK, Hosoya K, Miyazaki JI (2019) *Lefua tokaiensis*, a new species of nemacheilid loach from central Japan (Teleostei: Nemacheilidae). *Ichthyol Res* 66:479-487.
- Kottelat M (2012) *Conspectus Cobitidum: an inventory of the loaches of the world (Teleostei: Cypriniformes: Cobitoidei)*. *Raffles Bull Zool* 26:1-199.
- Kim IS (1997) *Illustrated Encyclopedia of Fauna and Flora of Korea, Vol. 37, Freshwater Fishes*. Ministry of Education, Yeongi, Korea, p 629.
- Kim IS, Lee EH (1995) Studies on early embryonic development of *Nirwaella multifasciata* (Pisces: Cobitidae). *Korean J Limnol* 28:455-462.
- Kim IS, Park JY (2007) *Freshwater Fishes of Korea*. Kyohak, Seoul, Korea, p 467.
- Kim IS, Choi Y, Lee CL, Lee YJ, Kim BJ, Kim JH (2005) *Illustrated Book of Korean Fishes*. Kyohak, Seoul, Korea, p 134.

- Kim YU, Kim DS, Park YS (1987) Development of eggs, larvae and juveniles of loach, *Misgurnus mizolepis* Güther. Bull Korean Fish Soc 20:16-23.
- Nelson JS, Grande TC, Wilson MVH (2016) Fishes of the World. 5th ed. John Wiley & Sons, Hoboken, NJ, p 192.
- Okiyama M (2015) An Atlas of the Early Stage Fishes in Japan. Tokai Univ Press, Tokyo, Japan, p 25-43.
- Song HY, Ko MH, Seo IY, Moon SJ, Bang IC (2017) Morphological development of egg and larvae of *Squalidus multimaculatus* (Gobioninae). Korean J Ichthyol 29:52-61.
- Song HY, Yang HY, Cho EM, Shin HC, Bang IC (2009) Morphological development of egg and larvae of *Koreocobitis naktongensis* (Cobitidae). Korean J Ichthyol 21: 247-252.
- Uchida K (1939) The fishes of Tyosen (Korea) part I. nematognathi, eventognathi. Bull Fish Exp St Gov Gen Tyosen 6:399-458.