



Preoperative Magnetic Resonance Imaging Features Associated with Positive Resection Margins in Patients with Invasive Lobular Carcinoma

Jiyoung Yoon, MD, Eun-Kyung Kim, MD, PhD, Min Jung Kim, MD, PhD, Hee Jung Moon, MD, PhD, Jung Hyun Yoon, MD, PhD, Vivian Y. Park, MD, PhD

All authors: Department of Radiology, Severance Hospital, Research Institute of Radiological Science, Yonsei University College of Medicine, Seoul, Korea

Objective: To investigate preoperative magnetic resonance imaging (MRI) findings associated with resection margin status in patients with invasive lobular carcinoma (ILC) who underwent breast-conserving surgery.

Materials and Methods: One hundred and one patients with ILC who underwent preoperative MRI were included. MRI (tumor size, multifocality, type of enhancing lesion, distribution of non-mass enhancement [NME], and degree of background parenchymal enhancement) and clinicopathological features (age, pathologic tumor size, presence of ductal carcinoma *in situ* [DCIS] or lobular carcinoma *in situ*, presence of lymph node metastases, and estrogen receptor/progesterone receptor/human epidermal growth factor receptor type 2 status) were analyzed. A positive resection margin was defined as the presence of invasive cancer or DCIS at the inked surface. Logistic regression analysis was performed to determine pre- and postoperative variables associated with positive resection margins.

Results: Among the 101 patients, 21 (20.8%) showed positive resection margins. In the univariable analysis, NME, multifocality, axillary lymph node metastasis, and pathologic tumor size were associated with positive resection margins. With respect to preoperative MRI findings, multifocality (odds ratio [OR] = 3.977, $p = 0.009$) and NME (OR = 2.741, $p = 0.063$) were associated with positive resection margins in the multivariable analysis, although NME showed borderline significance.

Conclusion: In patients with ILC, multifocality and the presence of NME on preoperative breast MRI were associated with positive resection margins.

Keywords: Breast; Carcinoma, lobular; Magnetic resonance imaging; Margins of excision

INTRODUCTION

Breast-conserving surgery (BCS) with radiation therapy is

the standard treatment for early-stage invasive breast cancer, and there is no significant difference in survival outcome between patients who undergo mastectomy and those who undergo BCS after 20 years of follow-up (1, 2). However, positive resection margins after BCS still pose difficulties in patient management. Positive resection margins are associated with a two-fold increase in the risk of ipsilateral breast tumor recurrence compared with negative margins (3). Consequently, almost 25% of patients who undergo BCS require additional re-excisions (4, 5).

Magnetic resonance imaging (MRI) has been reported to be more sensitive than mammography in detecting invasive cancer or ductal carcinoma *in situ* (DCIS), and is known to be more accurate than mammography or ultrasonography (US) in determining tumor diameters and margins and detecting multifocal, multicentric, and synchronous contralateral

Received: September 6, 2019 **Revised:** March 2, 2020

Accepted: March 3, 2020

This study was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2017R1D1A1B03035995).

Corresponding author: Vivian Y. Park, MD, PhD, Department of Radiology, Research Institute of Radiological Science, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea.

• E-mail: VIVIANPARK0116@yuhs.ac

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

breast cancers (6-8). Several studies have investigated preoperative MRI features associated with resection margin status in invasive breast cancer, and reported that non-mass enhancement (NME) and less convex margins are associated with positive resection margins (9, 10).

However, these results were based on study populations composed mostly of patients with invasive ductal carcinoma (IDC) (9, 10). Furthermore, the majority of malignant NME lesions depicted by MRI in previous studies were confirmed as DCIS, so the sensitivity advantage of MRI over conventional imaging in depicting local tumor extent was reported to increase as DCIS component size increased, which would be less frequently detected in invasive lobular carcinoma (ILC) (11, 12). Therefore, it is unclear whether similar MRI features are associated with resection margin status in this subgroup.

Although the association between preoperative MRI and surgical outcomes in patients with breast cancer is still under debate, multiple studies have shown that preoperative MRI can significantly reduce re-excision rates after initial surgery in patients with ILC (13-16). Such results have been incorporated into recommendations from the European Society of Breast Cancer Specialists working group and the U.S. National Comprehensive Cancer Network guidelines (17, 18), in which preoperative MRI is recommended for patients with ILC and several other subgroups of patients with breast cancer.

However, to the best of our knowledge, no studies have reviewed specific preoperative MRI features related to surgical outcomes in patients with ILC. Despite the improved accuracy of MRI in demonstrating tumor extent, the increasing incidence of BCS is less clear in patients with ILC compared with those with IDC (19). Further identification of imaging features related to resection margin status would aid in patient management by enhancing confidence in performing BCS for ILC patients at low risk for positive margins, and identifying patients at high risk for re-excision who may require more aggressive surgery.

Therefore, the purpose of this study was to investigate preoperative MRI findings associated with resection margin status in patients with ILC.

MATERIALS AND METHODS

Patients

Our Institutional Review Board approved this retrospective study and waived the requirement to obtain informed

consent. We conducted a retrospective chart review and identified 197 consecutive patients with newly diagnosed ILC who underwent preoperative MRI and surgery from May 2012 to August 2017. Patients who underwent vacuum-assisted biopsy (n = 4), surgical excisional biopsy (n = 2), or neoadjuvant chemotherapy before MRI (n = 17) were excluded. Among the remaining 174 patients, 73 patients who underwent total mastectomy were excluded. Finally, 101 patients (mean age, 52.1 years; range, 34–76 years) who underwent BCS were included in this study (Fig. 1).

Histopathologic Analysis

Pathologic data including pathologic tumor size, histologic grade, estrogen receptor (ER)/progesterone receptor (PR)/human epidermal growth factor receptor type 2 (HER2) status, and presence of metastasis in surgically resected axillary lymph nodes were obtained from the final pathologic reports. A tumor was considered ER/PR-positive when 10% of tumor cells in the testing sample had positively stained nuclei (20). The immunohistochemical staining results for HER2 were scored as 0, 1+, 2+, or 3+ according to the number of cells with positively stained membranes. Tumors with a score of 0 or 1+ were classified as HER2-negative, whereas tumors with a score of 3+ were classified as HER2-positive. In tumors with a score of 2+, additional gene amplification using silver *in situ* hybridization was performed to determine HER2 status. HER2 positivity was defined as a HER2/chromosome 17 ratio greater than 2.0 (21).

Lesion Localization and Assessment of the Resection Margin

Preoperative wire localization of non-palpable lesions was performed with mammogram or US guidance. In all cases, intraoperative margin excision and frozen section analysis were performed. According to the Society of Surgical Oncology-American Society for Radiation Oncology-American Society of Clinical Oncology consensus guideline (22), a “positive resection margin” was defined as the presence of invasive cancer or DCIS at the inked surface of the resected specimen, in either the frozen section or permanent specimen. The presence of lobular carcinoma *in situ* (LCIS) at the inked surface was considered to indicate a negative resection margin.

MRI Technique

Breast MRI examinations were performed using one of two 3T MRI systems: Discover 750 (GE Healthcare,

Milwaukee, WI, USA) or Ingenia (Philips Healthcare, Best, The Netherlands). Patients lay in the prone position, and imaging was performed with a dedicated phased array breast coil (8-channel [GE Healthcare] or 16-channel [Philips Healthcare]). Before the contrast agent was injected, a three-plane localizing sequence, axial T2-weighted fast spin echo and T2-stimulated inversion recovery sequence, and axial T1-weighted sequence with/without fat suppression were obtained. A bolus injection of a gadolinium-based contrast agent (Dotarem [gadoterate meglumine], Guerbet, Paris, France; Magnevist [gadopentetate dimeglumine], Berlex Laboratories, Wayne, NJ, USA; or Gadovist [gadobutrol], Bayer AG, Berlin, Germany) at a dose of 0.2 mmol/kg was administered at a rate of 2 mL/s, followed by a 20 mL saline flush. After the contrast agent was injected, T1-weighted three-dimensional (3D) dynamic contrast-enhanced images were acquired in the axial plane. This included one pre-contrast acquisition and six post-contrast bilateral axial acquisitions. Post-contrast images were obtained immediately after the contrast material was injected with no time delay. The length of each dynamic series was 63 (Discover 750) or 64 seconds (Ingenia). T1-weighted 3D delayed post-contrast images were later obtained in the sagittal plane. Image subtraction was performed after the dynamic series, including subtraction

of the pre-contrast images from the 1st, 2nd, and 6th post-contrast images.

Image Analysis

One board certified, breast-dedicated radiologist with 5 years of subspecialty experience in breast imaging reviewed the breast MR images according to the American College of Radiology Breast Imaging-Reporting and Data System (23). Tumor size was defined as the maximal diameter of the enhancing tumor on the early post-contrast subtraction images, which were obtained by subtracting the pre-contrast images from the 2nd post-contrast images acquired approximately 2 minutes after contrast material injection. Multifocal disease was defined as findings suspicious of additional sites of malignancy within the same quadrant as the index cancer. Enhancing lesions were divided into masses or NME. We further categorized the 101 cases into masses without NME and NME with or without a mass. In patients with NME, we analyzed NME distribution (focal, linear, or segmental). Background parenchymal enhancement (BPE) was categorized as minimal, mild, moderate, or marked based on a combination of pre-contrast, 2nd post-contrast T1-weighted images and subtraction images (23). After categorization, we dichotomized BPE into weak (minimal or mild) and strong (moderate or marked).




Fig. 1. Patient selection diagram. DCIS = ductal carcinoma *in situ*, MRI = magnetic resonance imaging

Statistical Analysis

The patients were divided into two groups according to resection margin status. Clinicopathological variables and imaging features were compared between the groups using the chi-square test or Fisher's exact test for categorical variables and the independent *t* test for continuous variables. Univariable logistic regression analysis was used to identify variables associated with positive resection margins. Multivariable logistic regression analysis was used to identify preoperative variables significantly associated with positive resection margins, using only preoperative variables that showed a significant association ($p < 0.05$) in the univariate analysis. Additional multivariable logistic regression analysis was performed using preoperative and postoperative variables that showed a significant association ($p < 0.05$) in the univariate analysis. All statistical analyses were performed using SPSS version 23.0.0 (IBM Corp., Armonk, NY, USA), and a p value < 0.05 was considered statistically significant.

RESULTS

The mean pathologic tumor size was 15.2 mm (standard deviation, 8.9 mm; range, 3–43 mm). Among the 101 patients, 11 (10.8%) had combined DCIS and 62 (61.4%) had combined LCIS. Negative resection margins were achieved in all of the patients with combined DCIS. Of the 101 patients, 15 (14.9%) had axillary lymph node metastasis.

Among the 101 patients, 21 (20.8%) showed positive resection margins. Among the 21 patients with positive resection margins, 9 (42.9%) were converted to mastectomy

at the time of the initial operation, 11 (52.4%) underwent additional local excision at the time of the initial operation, and 1 (4.8%) underwent re-excision later. Of the 11 patients who underwent immediate additional local excision, 3 (27.3%) later underwent mastectomy due to a positive margin on the re-excision specimen. Therefore, 12 (57.1%) of the 21 patients with positive resection margins were eventually converted to mastectomy (Figs. 2, 3).




Fig. 3. 48 year-old female with invasive lobular carcinoma at right upper outer breast. Preoperative axial early contrast-enhanced T1-weighted subtraction MR image show suspicious non-mass enhancement at right upper outer breast (arrow). At breast-conserving surgery, positive resection margin was confirmed by intraoperative frozen sectional analysis and additional local excision was performed immediately.




Fig. 2. 71-year-old female with invasive lobular carcinoma at left upper breast.

A, B. Preoperative axial early contrast-enhanced T1-weighted subtraction MR images show malignant mass at left upper medial breast (arrow in **A**) with adjacent suspicious non-mass enhancement (long arrow in **B**) and additionally detected suspicious focus at left subareolar breast (short arrow in **B**). At breast-conserving surgery, positive resection margin was confirmed on intraoperative frozen sectional analysis and additional local excision was performed immediately.

When comparing MRI features between the positive and negative resection margin groups, the presence of NME (NME with/without mass) showed significant difference between the two groups (47.6% vs. 21.3%, $p = 0.015$). The incidence of multifocality on preoperative breast MRI also showed significant difference between the two groups (57.1% vs. 22.5%, $p = 0.002$). The mean pathologic tumor size was significantly larger in the positive resection margin group (22.3 ± 10.5 mm vs. 13.4 ± 7.5 mm, $p < 0.001$), and

patients with positive resection margins had a higher rate of pathologic lymph node metastasis (38.1% vs. 8.8%, $p = 0.003$) (Table 1).

In the univariable logistic regression analysis, NME on preoperative breast MRI (odds ratio [OR] = 3.369; 95% confidence interval [CI] = 1.227–9.251; $p = 0.018$), multifocality on preoperative breast MRI (OR = 4.593; 95% CI = 1.671–12.623; $p = 0.003$), pathologic tumor size (OR = 1.117; 95% CI = 1.052–1.186; $p < 0.001$), and

Table 1. Comparison of Variables according to Status of Resection Margin

Characteristics	Positive Resection Margin (n = 21)	Negative Resection Margin (n = 80)	P
Age (years)	49.6 ± 8.9	52.8 ± 10.0	0.193
Pathologic tumor size (mm)	22.3 ± 10.5	13.4 ± 7.5	< 0.001
DCIS present			0.114
Yes	0 (0.0)	11 (13.8)	
No	21 (100.0)	69 (86.3)	
LCIS present			0.288
Yes	15 (71.4)	47 (58.8)	
No	6 (28.6)	33 (41.3)	
Lymph node metastases			0.003
Yes	8 (38.1)	7 (8.8)	
No	13 (61.9)	73 (91.3)	
Estrogen receptor			> 0.999
Positive	20 (95.2)	75 (93.8)	
Negative	1 (4.8)	5 (6.3)	
Progesterone receptor			> 0.999
Positive	17 (81.0)	64 (80.0)	
Negative	4 (19.0)	16 (20.0)	
HER2			0.635
Positive	2 (9.5)	5 (6.3)	
Negative	19 (90.5)	75 (93.8)	
Wire localization			> 0.999
Yes	17 (81.0)	64 (80.0)	
No	4 (19.0)	16 (20.0)	
Size on MRI (mm)	20.1 ± 9.4	17.6 ± 8.5	0.234
BPE			0.368
Weak	15 (71.4)	65 (81.3)	
Strong	6 (28.6)	15 (18.8)	
Presence of NME			0.015
Yes	10 (47.6)	17 (21.3)	
No	11 (52.4)	63 (78.8)	
Distribution of NME			0.290
Focal	5 (50.0)	12 (70.6)	
Linear/segmental	5 (50.0)	5 (29.4)	
Multifocality			0.002
Yes	12 (57.1)	18 (22.5)	
No	9 (42.9)	62 (77.5)	

Unless otherwise indicated, data in parentheses are percentages. BPE = background parenchymal enhancement, DCIS = ductal carcinoma *in situ*, HER2 = human epidermal growth factor receptor type 2, LCIS = lobular carcinoma *in situ*, MRI = magnetic resonance imaging, NME = non-mass enhancement

axillary lymph node metastasis (OR = 6.418; 95% CI = 1.985–20.750; $p = 0.002$) were significantly associated with positive resection margins (Table 2).

In the multivariable logistic regression analysis of preoperative variables, multifocality on preoperative breast MRI was independently associated with positive resection margins (OR = 3.977; 95% CI = 1.408–11.231; $p = 0.009$). NME on preoperative breast MRI tended to be associated with positive resection margins but showed borderline significance (OR = 2.741; 95% CI = 0.948–7.925; $p = 0.063$)

Table 2. Univariate Logistic Regression Analysis of Variables for Predicting Positive Resection Margins

Variables	OR	95% CI	P
Pathologic tumor size (mm)	1.117	1.052–1.186	< 0.001
DCIS present			
Yes	0.000	0.000–0.000	0.999
No	Ref		
LCIS present			
Yes	1.755	0.617–4.997	0.292
No	Ref		
Lymph node metastases			
Yes	6.418	1.985–20.750	0.002
No	Ref		
Estrogen receptor			
Positive	1.333	0.147–12.069	0.798
Negative	Ref		
Progesterone receptor			
Positive	1.062	0.314–3.596	0.922
Negative	Ref		
HER2			
Positive	1.558	0.280–8.661	0.613
Negative	Ref		
Wire localization			
Yes	1.062	0.314–3.596	0.922
No	Ref		
Size on MRI (mm)	1.033	0.979–1.089	0.234
BPE			
Weak	Ref		
Strong	1.733	0.577–5.211	0.327
Presence of NME			
Yes	3.369	1.227–9.251	0.018
No	Ref		
Distribution of NME			0.290
Focal	Ref		
Linear/segmental	2.400	0.475–12.130	
Multifocality			
Yes	4.593	1.671–12.623	0.003
No	Ref		

CI = confidence interval, OR = odds ratio, Ref = reference

(Table 3).

In the multivariable logistic regression analysis of pre- and postoperative variables, only multifocality on preoperative breast MRI (OR = 3.263; 95% CI = 1.034–10.304; $p = 0.044$) and pathologic tumor size (OR = 1.092; 95% CI = 1.023–1.166; $p = 0.008$) were independently associated with positive resection margins (Table 3).

DISCUSSION

In our study, we found that multifocality on preoperative breast MRI was independently associated with positive resection margins in patients with ILC, whereas NME showed borderline significance. The rate of positive resection margins in our study was 20.8% (21/101), which was within the previously reported range of 20–70% (24, 25). As preoperative MRI is recommended for patients with ILC (17, 18), our study results could provide assurance in cases which are unlikely to yield positive margins at BCS, help identify high-risk cases in which the extent of resection should be increased, and allow patients to be warned about the higher likelihood of additional procedures being needed to attain negative margins.

Although ILC is the second most common histological subtype of breast cancer, it is more challenging clinically and radiologically to detect than IDC (26). Due to its infiltrative nature, both mammography and US show lower sensitivity in depicting ILC, and MRI has been shown to be particularly useful in the assessment of tumor extent in this subgroup (26). However, surgeons may still prefer to adopt a more aggressive local approach for these patients, mainly due to the higher risk of incomplete resection with BCS (19). In the past five years, several studies have reported that preoperative MRI can reduce re-excision rates in patients with ILC, but none has reviewed the MRI features of ILC related to surgical outcomes (13–16).

In past studies with study populations composed mostly of patients with IDC, NME on preoperative MRI, tumor size exceeding 5 cm, and multifocal lesions were factors associated with positive resection margins after BCS (9, 27, 28). As DCIS lesions commonly appear as NME on MRI (29, 30), previous studies suggested an association between DCIS components and positive resection margins (12). In one study, 55% (22/40) of the NME lesions around index cancer masses were malignant, with 81.8% (18/22) confirmed as DCIS (11). In addition, as DCIS commonly presents as microcalcifications on mammography (31), previous studies

Table 3. Multivariate Analysis of Pre/Postoperative Variables for Predicting Positive Resection Margins

Preoperative Variables				Pre/Postoperative Variables			
Variables	OR	95% CI	P	Variables	OR	95% CI	P
Presence of NME				Presence of NME			
Yes	2.741	0.948–7.925	0.063	Yes	1.805	0.543–6.000	0.335
No	Ref			No	Ref		
Multifocality				Multifocality			
Yes	3.977	1.408–11.231	0.009	Yes	3.263	1.034–10.304	0.044
No	Ref			No	Ref		
				Pathologic tumor size (mm)	1.092	1.023–1.166	0.008
				Lymph node metastases			
				Yes	2.735	0.676–11.065	0.158
				No	Ref		

have reported that the presence of microcalcifications on mammography and presence of DCIS were both associated with resection margin status in patient populations composed mostly of patients with IDC (32).

In this study, we questioned whether similar MRI features were also associated with resection margin status in patients with ILC. Of the 101 patients included in our study, 11 (10.8%) had combined DCIS, but negative resection margins were achieved in all of these patients. However, our study results showed that multifocality and NME on preoperative MRI were associated with positive resection margins in ILC patients, although NME showed borderline significance. Tumors with higher cellularity usually present as masses, whereas scattered tumor cells and tumors with lower cellularity present as NME, which may underlie the association between NME and resection margin status (33). Among the enhancing patterns of ILC on MRI, NME has been correlated with malignant cells streaming in a single-file fashion in the breast stroma or small tumor aggregates separated by normal tissue, which would likely cause underestimation of tumor extent (34, 35). Our results are also consistent with previous studies, which have reported that NME lesions show higher discordance than mass lesions in estimating tumor size (36).

In addition, the lack of MRI-guided lesion localization may have also attributed to the higher positive resection margin rate in NME lesions. Although previous trials investigating the benefit of breast MRI for treatment planning in women with invasive breast cancer have failed to demonstrate a positive effect on surgical outcome, one recent study that utilized MRI-guided biopsy and MRI-guided lesion bracketing demonstrated a low overall positive margin rate of 3.7% (12). Therefore, we can infer

that in real-world clinical practice where the use of MRI-guided lesion localization is often lacking, the improved assessment of tumor extent by MRI may not be fully applicable to surgery. This would be especially related to NME, which is often less discernible on conventional imaging (37).

ILC has been associated with higher rates of positive resection margins after BCS compared to IDC (38, 39). In a previous study that focused on the associations between clinicopathologic variables and positive resection margins, tumor size exceeding 2 cm and lymph node involvement were associated with positive resection margins in ILC (40). Our study showed similar results, with multifocality at MRI and tumor size being independently associated with positive resection margins. Axillary lymph node metastasis was also associated with positive resection margins, although this was not statistically significant in the multivariable analysis. Our study results imply that multifocality on MRI and tumor size are most strongly associated with positive margins in patients with ILC, for whom more aggressive surgery such as resection of shave margins, wide excision, or mastectomy should be considered. Patients with none of these risk factors and without NME on MRI have a lower risk of positive margins and may more confidently receive BCS.

There are several limitations of this study. First, this was a retrospective study and patients were recruited from a single tertiary referral center, so selection bias was inevitable. Second, one breast-dedicated radiologist reviewed the breast MR images. Although a high agreement rate among readers has been reported when differentiating NME from mass lesions, interobserver variability would be present if another radiologist were to review the same images (41). Third, our study had a small number of events

(21 cases with positive resection margins), which may have caused overfitting of the model. An event per variable ≥ 10 would be preferable, and larger studies are required to confirm our results (42).

In conclusion, in patients with ILC, multifocality and NME on preoperative breast MRI were associated with positive resection margins. These findings may aid in determining the extent of resection required prior to BCS.

Conflicts of Interest

The authors have no potential conflicts of interest to disclose.

ORCID iDs

Vivian Y. Park

<https://orcid.org/0000-0002-5135-4058>

Jiyoung Yoon

<https://orcid.org/0000-0003-2266-0803>

Eun-Kyung Kim

<https://orcid.org/0000-0002-3368-5013>

Min Jung Kim

<https://orcid.org/0000-0003-4949-1237>

Hee Jung Moon

<https://orcid.org/0000-0002-5643-5885>

Jung Hyun Yoon

<https://orcid.org/0000-0002-2100-3513>

REFERENCES

1. Fisher B, Anderson S, Bryant J, Margolese RG, Deutsch M, Fisher ER, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med* 2002;347:1233-1241
2. Veronesi U, Cascinelli N, Mariani L, Greco M, Saccozzi R, Luini A, et al. Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med* 2002;347:1227-1232
3. Moran MS, Schnitt SJ, Giuliano AE, Harris JR, Khan SA, Horton J, et al. Society of Surgical Oncology-American Society for Radiation Oncology consensus guideline on margins for breast-conserving surgery with whole-breast irradiation in stages I and II invasive breast cancer. *Int J Radiat Oncol Biol Phys* 2014;88:553-564
4. McCahill LE, Single RM, Aiello Bowles EJ, Feigelson HS, James TA, Barney T, et al. Variability in reexcision following breast conservation surgery. *JAMA* 2012;307:467-475
5. Morrow M, Jagsi R, Alderman AK, Griggs JJ, Hawley ST, Hamilton AS, et al. Surgeon recommendations and receipt of mastectomy for treatment of breast cancer. *JAMA* 2009;302:1551-1556
6. Van Goethem M, Tjalma W, Schelfhout K, Verslegers I, Biltjes I, Parizel P. Magnetic resonance imaging in breast cancer. *Eur J Surg Oncol* 2006;32:901-910
7. Gruber IV, Rueckert M, Kagan KO, Staebler A, Siegmann KC, Hartkopf A, et al. Measurement of tumour size with mammography, sonography and magnetic resonance imaging as compared to histological tumour size in primary breast cancer. *BMC Cancer* 2013;13:328
8. Houssami N, Ciatto S, Macaskill P, Lord SJ, Warren RM, Dixon JM, et al. Accuracy and surgical impact of magnetic resonance imaging in breast cancer staging: systematic review and meta-analysis in detection of multifocal and multicentric cancer. *J Clin Oncol* 2008;26:3248-3258
9. Kang JH, Youk JH, Kim JA, Gweon HM, Eun NL, Ko KH, et al. Identification of preoperative magnetic resonance imaging features associated with positive resection margins in breast cancer: a retrospective study. *Korean J Radiol* 2018;19:897-904
10. Dashevsky BZ, Oh JH, Apte AP, Bernard-Davila B, Morris EA, Deasy JO, et al. MRI features predictive of negative surgical margins in patients with HER2 overexpressing breast cancer undergoing breast conservation. *Sci Rep* 2018;8:315
11. Gweon HM, Jeong J, Son EJ, Youk JH, Kim JA, Ko KH. The clinical significance of accompanying NME on preoperative MR imaging in breast cancer patients. *PLoS One* 2017;12:e0178445
12. Kuhl CK, Strobel K, Bieling H, Wardelmann E, Kuhn W, Maass N, et al. Impact of preoperative breast MR imaging and MR-guided surgery on diagnosis and surgical outcome of women with invasive breast cancer with and without DCIS component. *Radiology* 2017;284:645-655
13. Houssami N, Turner R, Morrow M. Preoperative magnetic resonance imaging in breast cancer: meta-analysis of surgical outcomes. *Ann Surg* 2013;257:249-255
14. Fortune-Greeley AK, Wheeler SB, Meyer AM, Reeder-Hayes KE, Biddle AK, Muss HB, et al. Preoperative breast MRI and surgical outcomes in elderly women with invasive ductal and lobular carcinoma: a population-based study. *Breast Cancer Res Treat* 2014;143:203-212
15. McGhan LJ, Wasif N, Gray RJ, Giurescu ME, Pizzitola VJ, Lorans R, et al. Use of preoperative magnetic resonance imaging for invasive lobular cancer: good, better, but maybe not the best? *Ann Surg Oncol* 2010;17 Suppl 3:255-262
16. Mann RM, Loo CE, Wobbes T, Bult P, Barentsz JO, Gilhuijs KG, et al. The impact of preoperative breast MRI on the re-excision rate in invasive lobular carcinoma of the breast. *Breast Cancer Res Treat* 2010;119:415-422
17. Sardanelli F, Boetes C, Borisch B, Decker T, Federico M, Gilbert FJ, et al. Magnetic resonance imaging of the breast: recommendations from the EUSOMA working group. *Eur J Cancer* 2010;46:1296-1316
18. National Comprehensive Cancer Network.® *Breast cancer*

- screening and diagnosis. Version 1.2015. NCCN clinical practice guidelines in oncology. Plymouth Meeting: NCCN, 2015
19. Truin W, Roumen RM, Siesling S, van der Heiden-van der Loo M, Duijm LE, Tjan-Heijnen VC, et al. Patients with invasive lobular breast cancer are less likely to undergo breast-conserving surgery: a population based study in the Netherlands. *Ann Surg Oncol* 2015;22:1471-1478
 20. Hammond ME, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *J Clin Oncol* 2010;28:2784-2795
 21. Wolff AC, Hammond ME, Schwartz JN, Hagerty KL, Allred DC, Cote RJ, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *Arch Pathol Lab Med* 2007;131:18-43
 22. Morrow M, Van Zee KJ, Solin LJ, Houssami N, Chavez-MacGregor M, Harris JR, et al. Society of Surgical Oncology-American Society for Radiation Oncology-American Society of Clinical Oncology consensus guideline on margins for breast-conserving surgery with whole-breast irradiation in ductal carcinoma in situ. *Ann Surg Oncol* 2016;23:3801-3810
 23. D'Orsi CJ, Sickles EA, Mendelson EB, Morris EA. *ACR BI-RADS atlas: breast imaging reporting and data system, 5th ed.* Reston: American College of Radiology, 2013
 24. Lai HW, Huang RH, Wu YT, Chen CJ, Chen ST, Lin YJ, et al. Clinicopathologic factors related to surgical margin involvement, reoperation, and residual cancer in primary operable breast cancer-An analysis of 2050 patients. *Eur J Surg Oncol* 2018;44:1725-1735
 25. Singletary SE. Surgical margins in patients with early-stage breast cancer treated with breast conservation therapy. *Am J Surg* 2002;184:383-393
 26. Berg WA, Gutierrez L, NessAiver MS, Carter WB, Bhargavan M, Lewis RS, et al. Diagnostic accuracy of mammography, clinical examination, US, and MR imaging in preoperative assessment of breast cancer. *Radiology* 2004;233:830-849
 27. Park SY, Kang DK, Kim TH. Does background parenchymal enhancement on MRI affect the rate of positive resection margin in breast cancer patients? *Br J Radiol* 2015;88:20140638
 28. Jang M, Kim SM, Yun BL, Kim SW, Kang EY, Park SY, et al. Magnetic resonance imaging factors predicting re-excision in breast cancer patients having undergone conserving therapy. *J Korean Soc Magn Reson Med* 2014;18:133-143
 29. Tozaki M, Fukuda K. High-spatial-resolution MRI of non-masslike breast lesions: interpretation model based on BI-RADS MRI descriptors. *AJR Am J Roentgenol* 2006;187:330-337
 30. Liberman L, Morris EA, Lee MJ, Kaplan JB, LaTrenta LR, Menell JH, et al. Breast lesions detected on MR imaging: features and positive predictive value. *AJR Am J Roentgenol* 2002;179:171-178
 31. Yamada T, Mori N, Watanabe M, Kimijima I, Okumoto T, Seiji K, et al. Radiologic-pathologic correlation of ductal carcinoma in situ. *Radiographics* 2010;30:1183-1198
 32. Kurniawan ED, Wong MH, Windle I, Rose A, Mou A, Buchanan M, et al. Predictors of surgical margin status in breast-conserving surgery within a breast screening program. *Ann Surg Oncol* 2008;15:2542-2549
 33. Zhou J, Li G, Sheng F, Qiao P, Zhang H, Xing X. Magnetic resonance imaging evaluation of residual tumors in breast cancer after neoadjuvant chemotherapy: surgical implications. *Acta Radiol* 2016;57:529-537
 34. Qayyum A, Birdwell RL, Daniel BL, Nowels KW, Jeffrey SS, Agoston TA, et al. MR imaging features of infiltrating lobular carcinoma of the breast: histopathologic correlation. *AJR Am J Roentgenol* 2002;178:1227-1232
 35. Shin SU, Cho N, Lee HB, Kim SY, Yi A, Kim SY, et al. Neoadjuvant chemotherapy and surgery for breast cancer: preoperative MRI features associated with local recurrence. *Radiology* 2018;289:30-38
 36. Mann RM, Bult P, van Laarhoven HW, Span PN, Schlooz M, Veltman J, et al. Breast cancer size estimation with MRI in BRCA mutation carriers and other high risk patients. *Eur J Radiol* 2013;82:1416-1422
 37. Spick C, Baltzer PA. Diagnostic utility of second-look US for breast lesions identified at MR imaging: systematic review and meta-analysis. *Radiology* 2014;273:401-409
 38. Raje D, Bollard R, Wilson A. Invasive lobular cancer of the breast-Is breast conservation surgery a good option? *Breast J* 2006;12:574-575
 39. Hussien M, Lioe TF, Finnegan J, Spence RA. Surgical treatment for invasive lobular carcinoma of the breast. *Breast* 2003;12:23-35
 40. Wanis ML, Wong JA, Rodriguez S, Wong JM, Jabo B, Ashok A, et al. Rate of re-excision after breast-conserving surgery for invasive lobular carcinoma. *Am Surg* 2013;79:1119-1122
 41. Lunkiewicz M, Forte S, Freiwald B, Singer G, Leo C, Kubik-Huch RA. Interobserver variability and likelihood of malignancy for fifth edition BI-RADS MRI descriptors in non-mass breast lesions. *Eur Radiol* 2020;30:77-86
 42. Cen D, Hu W, Wang X, Wu X. Re: identification of preoperative magnetic resonance imaging features associated with positive resection margins in breast cancer? *Korean J Radiol* 2019;20:999-1000