

# *Bowlesia incana* Ruiz & Pav. (Apiaceae), a New Invasive Alien Plant in Korea

Eun Su Kang<sup>1</sup>, Yoon-Young Kim<sup>2</sup>, Myoung Ja Nam<sup>3</sup>, Nak Yong Kim<sup>4</sup>,  
Seong-Jin Ji<sup>5</sup> and Dong Chan Son<sup>6\*</sup>

<sup>1</sup>Master's Degree Researcher and <sup>6</sup>Researcher, Division of Forest Biodiversity,  
Korea National Arboretum, Pocheon 11186, Korea

<sup>2</sup>Researcher, Division of Variety Examination, National Forest Seed Variety Center, Chungju 27495, Korea

<sup>3</sup>Teacher, Onhye Elementary School, Andong 36604, Korea

<sup>4</sup>Researcher, Subalpine Plant Research Group, Seoul 01776, Korea

<sup>5</sup>Researcher, General Exhibition Division, National Science Museum, Daejeon 34143, Korea

**Abstract** - *Bowlesia incana* Ruiz & Pav., a new invasive alien plant in Korea, was found in Ulju-gun, Ulsan Metropolitan City. This species is characterized as being similar to the genera *Hydrocotyle* Tourn. ex L. and *Centella* L. However, it differs morphologically from these two genera in leaves, flowers, and fruits, as well as stellate pubescence growing on the whole plant. In Korea, *Bowlesia incana* Ruiz & Pav. is recorded for the first time in this study. A morphological description, distribution map, and illustrations based on the Korean materials collected are presented.

**Key words** – Apiaceae, *Bowlesia*, Flora, New record, Republic of Korea

## Introduction

*Bowlesia* Ruiz & Pav. is a genus of flowering plant in the family Apiaceae, and about 15 species are known worldwide (Mathias and Constance, 1965). This genus can be distinguished from the related genera by its prostrate herbaceous habit, simple umbels, short carpophores, and stellate pubescence (Reiche, 1902).

Initially, *Bowlesia* was classified as belonging to the subfamily Hydrocotyloideae (Drude, 1898; Pimenov and Leonov, 1993), however, it has recently been placed in Azorelloideae by studies based on molecular data (Plunkett *et al.*, 2004). Representative genera of Azorelloideae are *Azorella* Lam., *Asteriscium* Cham. & Schtdl. and *Diposis* DC., however, none of the taxa in subfamily have been recorded in Korea.

During extensive field surveys for floristic research in Korea in 2018, we collected an unidentifiable plant characterized by leaves similar to those of *Hydrocotyle* and flowers and fruits similar to those of *Centella* (Figs. 1, 2).

After we examined floras of various regions, herbarium specimens of Korea and adjacent countries, and the relevant literature, we concluded that the collected specimens represented an Apiaceae species hitherto unrecorded in Korea. We have formally described it below as *Bowlesia incana* Ruiz & Pav.




Fig. 1. Distribution of *Bowlesia incana* Ruiz & Pav. in South Korea. Illustration by Eun Su Kang.

\*Corresponding author. E-mail : sdclym@korea.kr  
Tel. +82-31-540-8813

## Taxonomic Treatment

*Bowlesia* Ruiz & Pav., Fl. Peruv. Prodr. 44 (1794).

**Vernacular (Korean) name:** Byeong-pul-a-jae-bi-sok (병풀아재비속; new Korean name).

*Bowlesia incana* Ruiz & Pav., Fl. Peruv. 3: 28 (1802) (Figs. 2, 3)

**Type:** (neotype, designated by Mathias and Constance, 1965):

—*P*: ‘Habitat in Peruviae collibus ad Huanuci, Rondos et Pillao tractus.’ (present address: Peru: Huanuco Dist. & Rondos Dist.), s.d., *Dombey* s.n. (F, digital photograph)

**Annual** creeping, small herb, covered with stellate hairs. **Stem** slender, glabrate to slightly stellate pubescent; hairs 8-rayed, unstalked. **Leaf** opposite, with fimbriate stipule and usually long petiole; blade palmate, 0.5–3 cm long, rounded-reniform; lobes 5–9, widely lanceolate to round, generally obtuse with deep sinus, stellate pubescent; stalked hairs



Fig. 2. Photographs of *Bowlesia incana* Ruiz & Pav. A: Habit, B: Stem, C: Petiole, D: Leaf (lateral view), E–G: Flower, H: Fruit. Photographs by Yoon-Young Kim.




Fig. 3. Illustrations of *Bowlesia incana* Ruiz & Pav. A–B: Habit, C: Shoot, D: Inflorescence position, E: Stem cross-section, F: Petiole hair (stellate), G–K: Leaf (H: ventral surface, I: margin, J: apex, K: dorsal surface), L: Bract, M: Bractlet, N: Inflorescence, O: Flower longitudinal section, P: Flower (top view), Q: Calyx, R: Stamen, S: Fruit, T: Seed. Illustrations by Hyeryun Jo.

8-rayed on both surfaces. **Inflorescence** umbel, simple, axillary, few-flowered. **Bract** of a few scarious, white to purplish white, lacerate bractlet. **Flower** small; calyx teeth prominent; petal oblong to ovate, greenish white to purple; stamen inserted, anther purple. **Stylopodium** conical; style short; carpel densely stellate pubescent. **Fruit** small, broadly ellipsoid to globose ridge inconspicuous, ca. 2 mm long, 1.5 mm wide, covered with stellate hairs.

**Phenology:** Flowering Mar. to Apr.

**Distribution:** Korea (Ulju-gun), Japan (Honshu), South America, southern USA, Europe, western Pakistan, New Zealand.

**Vernacular (Korean) name:** Byeong-pul-a-jae-bi (병풀아재비); new Korean name).

**Specimens examined:** KOREA. Ulsan Metropolitan City, Ulju-gun, March 20, 2018, *Y.Y. Kim 180320-001* (KH)

**Introduction route:** *Bowlesia incana* was found on the slopes next to the protective walls installed along the road. Thus, we assume that their seeds unintentionally transported by vehicles along with construction materials in the construction sites. Eventually they germinate and produce seeds which dispersed throughout area. Although *Bowlesia incana* was found in the forest areas, recently recorded other invasive

alien plants, for instance, *Ludwigia peploides* (Kunth) P.H.Raven subsp. *montevidensis* (Spreng.) P.H. Raven and *Geranium purpureum* Vill., have been found in lakes and railroads (Kim *et al.*, 2019a; 2019b). This indicated that invasive alien plants can grow in every possible environment, thus it is important to evaluate their invasiveness through regular monitoring and distribution survey. The final results should be provided as information for establishing the management of invasive alien plants.

**Taxonomic note:** While *Bowlesia incana*, native to Peru of South America, has been naturalized in countries in from tropic to temperate region including North America, Europe, Asia and New Zealand (Fig. 4). In Korea, this species is similar to *Hydrocotyle sibthorpioides* Lam. in terms of the appearance of young leaves and white-colored early blooms, and *Centella asiatica* (L.) Urb. with respect to the shape of the flower and fruit. However, *B. incana* can be distinguished by several features from these two species; in the former case, the white flower color changes into reddish as it grows, ridges on the fruit surface, which are not seen in *H. sibthorpioides*. Furthermore, *B. incana* has palmate leaves and hairy fruits, whereas *C. asiatica* has heart-shaped or reniform leaves and glabrous fruits. Above all, stellate pubescence growing all




Fig. 4. Distribution of *Bowlesia incana* Ruiz & Pav. in the world. Illustration by Eun Su Kang.

over the surface of *B. incana* is the most important characteristic that is not found in the two similar species (Figs. 2, 3). According to our investigation, this genus is recorded for Korean flora as the species *B. incana*. This is the first record of this species in the Korea area. The number of genera of Apiaceae in the Flora of Korea (Korea National Arboretum, 2017) has increased to 41 with the addition of *Bowlesia*. Meanwhile, *Drusa* DC. and *Homalocarpus* Hook. & Arn. are known to be related with *Bowlesia* (Plunkett *et al.*, 2018). However, these genera were not found in Korea, thus we described *Bowlesia incana* comparing apparently similar genera, *Hydrocotyle* and *Centella* in Apiaceae. The identification key of *Bowlesia* and related genera of the family Apiaceae has been recorded and is given as follows.

**Key to *Bowlesia* and morphologically similar genera of Apiaceae in Korea**

- 1. Fruit without ridges .....  
..... *Hydrocotyle* Tourn. ex L. 피막이속
- 1. Fruit with reticulate ridges.
  - 2. Herbs perennial, glabrous. Adventitious root development at node. Leaf cordate or reniform. Calyx and petals glabrous. Fruits reticulate, ridged .....  
..... *Centella* L. 병풀속
  - 2. Herbs annual, with stellate trichomes. Adventitious roots absent. Leaf palmate. Calyx and petals covered with stellate hairs on dorsal surface. Fruits not ridged .....  
..... *Bowlesia* Ruiz & Pav. 병풀아재미속

\* Although *Hydrocotyle* is recently placed in Araliaceae by some authors based on molecular phylogenetic studies (Downie *et al.*, 2000; Plunkett *et al.*, 2004; Erbar *et al.*, 2004; Angiosperm Phylogeny Group, 2009), we described it as belonging to Apiaceae in accordance with Flora of Korea (Korea National Arboretum, 2017; Park *et al.*, 2017)

**Acknowledgments**

This study was supported by the Korea National Arboretum, project No. “KNA1-2-27, 16-5.” We would like to thank Hyeryun Jo for helping with the line drawing.

**References**

Angiosperm Phylogeny Group 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. Bot. J. Linnean Soc. 161(2):105-121.

Downie, S.R., D.S. Katz-Downie and M.F. Watson. 2000. A phylogeny of the flowering plant family Apiaceae based on chloroplast DNA *rpl16* and *rpoC1* intron sequences: Towards a suprageneric classification of subfamily Apioideae. Am. J. Bot. 87(2):273-292.

Drude, C.G.O. 1898. Umbelliferae: In Engler, A. and K. Prantl (eds.), Die Natürlichen Pflanzenfamilien, Wilhelm Engelmann, Leipzig, Germany. Vol. 3(8). pp. 63-250.

Erbar, C., P. Leins, B.E. van Wyk and P.M. Tilney. 2004. Sympetaly in Apiales (Apiaceae, Araliaceae, Pittosporaceae). S. Afr. J. Bot. 70(3):458-467.

Kim, H.Y., D.C. Son, S.H. Park, C.S. Jang, E.M. Sun, H. Jo, S.M. Yun and K.S. Chang. 2019a. Unrecorded alien plant in south Korea: *Ludwigia peploides* subsp. *montevidensis* (Spreng.) P.H. Raven. Korean J. Plant Res. 32(2):201-206 .

Kim, H.Y., E.M. Sun, S.Y. Jung and D.C. Son. 2019b. *Geranium purpureum* Vill.: A new causal alien plant in Korea. Korean J. Pl. Taxon. 49(3):209-214.

Korea National Arboretum 2017. Checklist of Vascular Plants in Korea. Korea National Arboretum, Pocheon, Korea. p. 1000.

Mathias, M.E. and L. Constance. 1965. A revision of the Genus *Bowlesia* Ruiz & Pav. (Umbelliferae-Hydrocotyloideae) and its relatives. Univ. Calif. Publ. Bot. 38:1-73.

Park, C.W., B.Y. Lee, J.H. Song and K. Kim. 2017. Apiaceae. In Flora of Korea Editorial Committee (eds.) Flora of Korea, vol. 5c. Junghaengsa, Seoul, Korea. pp. 77-149.

Pimenov, M.G. and M.V.E. Leonov. 1993. The Genera of the Umbelliferae: A Nomenclator. Royal Botanic Gardens, Kew and London Botanical Garden of Moscow University, Russia. p. 156.

Plunkett, G.M., G.T. Chandler, P.P. Lowry II, S.M. Pinney and T.S. Sprenkle. 2004. Recent advances in understanding apiales and a revised classification. S. Afr. J. Bot. 70(3): 371-381.

Plunkett, G.M., M.G. Pimenov, J.P. Reduron, E.V. Kljuykov, B.E. van Wyk, T.A. Ostroumova, M.J. Henwood, P.M. Tilney, K. Spalik, M.F. Watson, B.Y. Lee, F.D. Pu, C.J.

Webb, J.M. Hart, A.D. Mitchell and B. Muckensturm. 2018. Apiaceae: *In* Kadereit, J.W. and C. Jeffrey (eds.), Flowering Plants. Eudicots. The Families and Genera of Vascular Plants. Cham, Springer, Berlin, Germany. 15:9-206.

Reiche, K. 1902. Estudios Críticos Sobre la Flora de Chile: *In* Faride, Z.C. (ed.), Anales de la Universidad de Chile, Santiago, Chile. Vol. 3. pp. 56-57.

(Received 25 February 2020 ; Revised 10 April 2020 ; Accepted 13 April 2020)