

A Pediatric Case of Long-term Untreated Distal Renal Tubular Acidosis

Philavanh Kedsatha, M.D.¹
Hee Young Shin, M.D., Ph.D.²
Yong Choi, M.D., Ph.D.²
Hae Il Cheong, M.D., Ph.D.²
Tae-Joon Cho, M.D., Ph.D.³
Eunsang Yi, M.D., Ph.D.⁴
Mayfong Maisai, M.D., Ph.D.^{5,6,7}

Department of Pediatrics¹, Children Hospital, Vientiane, Lao PDR, Department of Pediatrics², Orthopedic Surgery³, Seoul National University Children's Hospital, Seoul, Korea, Department of Pediatrics⁴, Korea University Guro Hospital, Seoul, Korea, Institute of Research and Education Development⁵, University of Health Sciences, Ministry of Health, Vientiane, Lao PDR, Lao-Oxford-Mahosol Hospital Welcome Trust Research Unit⁶, Mahosol Hospital, Vientiane, Lao PDR, Centre for Tropical Medicine and Global Health⁷, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

Corresponding author:

Hae Il Cheong, M.D., Ph.D.
Department of Pediatrics, Seoul National University Children's Hospital 101 Daehak-ro, Jongno-gu, Seoul 03080, Korea
Tel: +82-2-2072-2810
Fax: +82-2-743-3455
E-mail: cheonghi@snu.ac.kr

Received: 30 June 2020
Revised: 12 August 2020
Accepted: 8 September 2020

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Distal renal tubular acidosis (dRTA) is a rare renal tubular disorder characterized by normal anion gap metabolic acidosis, hypokalemia, and high urine pH. It can be inherited or acquired. In untreated pediatric patients with dRTA, rickets and growth retardation are common. We report the case of a 12-year-old Lao girl who presented with typical clinical features of dRTA with severe bone deformities that developed after a bed-ridden state due to a bicycle accident at the age of 8 years. Initial laboratory tests revealed metabolic acidosis with a normal anion gap, hypokalemia, and alkali urine. Renal ultrasonography revealed bilateral medullary nephrocalcinosis. Whole exome sequencing revealed no pathogenic mutations. After treatment with oral alkali, potassium, and vitamin D, she could walk and run. Later, she underwent corrective orthopedic surgeries for bony deformities. Thus, in pediatric dRTA patients, despite severe symptoms remaining untreated, accurate diagnosis and proper management can improve quality of life.

Key words: Distal renal tubular acidosis, Failure to thrive, Bony deformity, Muscle paralysis

Introduction

Renal tubular acidosis (RTA) is a group of diseases characterized by the development of hypokalemic metabolic acidosis with a normal anion gap due to a defect in the ability of the renal tubules to either reabsorb bicarbonate or excrete hydrogen ions in response to acidemia^{1,2}. Distal tubular acidosis (dRTA), which was first described in 1946³, is also referred to as type 1 RTA and is characterized by the inability to secrete hydrogen ions from the distal tubules. The etiology of dRTA is diverse and can be inherited or acquired. Common clinical presentations of dRTA in the pediatric age group include failure to thrive, polyuria, constipation, rickets, and nephrocalcinosis. We report the case of a 12-year-old Lao girl with long-term untreated dRTA, who presented with severe complications of progressive bony deformities and muscle paralysis.

Case report

A 12-year-old girl visited the Children's Hospital in Vientiane Capital, Lao

PDR with severe bony deformities and progressive paralysis of the extremity muscles. The patient was born via normal vaginal delivery without any perinatal complications. Her birth weight was 2.9 kg. She was breastfed since birth and started solid food at 6 months of age, and she completed vaccination as scheduled. She is the second child of the family, and her growth and development were normal until the age of 8 years. (Fig. 1A) At the age of 8 years, she had a bicycle accident, resulting in dislocation of the right hip and wrist joints. After that, she was bedridden and stayed at home without treatment because of the very challenging socioeconomic status of the family. The range of motion of her upper and lower extremities gradually decreased, accompanied by progressive limb bone deformities. She also lost appetite and weight. At the age of 12 years, the patient visited the outpatient clinic at the Children's Hospital in Vientiane Capital, Lao PDR once but only received a multi-vitamin prescription without an accurate diagnosis. However, her symptoms gradually worsened resulting in severe contracture of extremity joints and whole-body muscle weakness. Therefore, she visited the Children's Hospital again.

Her father had been healthy until the age of 30 years, when he had progressive paralysis of both arms after working hard, cutting trees. After some treatment and physical therapy at a military hospital, the symptoms of the left arm improved, but the weakness in the right arm continued to this day. Her mother, sister, and cousins were healthy, without any medical condition.

At admission, her body weight was 10 kg (<3rd percentile, she weighed 20 kg at the age of 8 years just before the bicycle accident, according to her father), and her height was 87 cm (<3rd percentile). She was bedridden and looked cachectic. She had a rachitic rosary on the thoracic wall and severe bony deformities with tenderness in the upper and lower extremities. (Fig. 1B–F) A blood test showed hypokalemia (2.3–2.8 mEq/L), metabolic acidosis (serum HCO₃⁻ 15 mEq/L) with normal anion gap (13 mEq/L). Serum calcium level was 8.0 mg/dL, phosphorus 1.0 mmol/L, sodium 135 mEq/L, and chloride 107 mEq/L. Urine pH was high (6.6–9.0 by dipstick) and the specific gravity was 1005. There was no proteinuria or hematuria. The urine anion gap was 78 mEq/L. Kidney ultrasonography showed normal sized kidneys with bilateral medullary nephrocalcinosis. Other labora-

tory tests revealed the following: WBC, 5,500; hematocrit, 38%; hemoglobin, 13 g/dL; platelet, 407,000/mm³; serum creatinine, 0.4 mg/dL; alkaline phosphatase, 1,340 IU/L; and blood sugar, 116 mg/dL. The patient was clinically diagnosed with dRTA and was treated with oral sodium bicarbonate (started from 2.4 g/day and increased to 9.4 g/day), potassium phosphate (4 g/day), vitamin D (400 IU/day), and high calcium diet for 4 months, and then sodium bicarbonate was changed to potassium citrate (Urocitra®, Pharmbio Korea, Inc., 3–5 mEq/kg/day). Her general condition began to improve after one month of Urocitra treatment, and the follow-up laboratory tests revealed serum HCO₃⁻, 21–24 mEq/L; potassium, 4.1–4.6 mEq/L; and pho-



Fig. 1. Serial medical photographs and bone radiographs of the patient. A: A photograph of the patient (right) with her older sister (left). The patient was 4 years old and healthy at that time. B–D: Medical photographs taken at the age of 12 years show a rachitic rosary in the chest wall (B) and severe bony deformities in the upper (C) and lower (D) extremities. E and F: Bone radiographs show severe bony deformities in the upper (E) and lower (F) extremities. G and H: Medical photographs taken 9 months after alkali treatment show disappearance of the rachitic rosary (G) and increased muscle bulk in the lower extremities (H). I: A bone radiograph of the right foot taken after the 2nd orthopedic surgery. K: A currently taken photograph.

sphorus, 6–6.5 mmol/L. She could stand alone, ride a bike, and attend school after 2 months of treatment. After 6 months of treatment, her body weight and height increased to 15 kg and 98 cm, respectively. After 2.5 years of treatment, she underwent two orthopedic surgeries for correction of the limb bone deformities in the Department of Pediatric Orthopedic Surgery, Seoul National University Children's Hospital, Seoul, Korea with financial support from the Seoul National University Children's Hospital Fund. Two years after the corrective surgery, plate removal from both legs was performed in the Mittaphab Hospital, Lao PDR. (Fig. 1G–K) Currently, she is 17 years old with a height of 133 cm and taking potassium citrate (Urocitra, 1 mEq/kg/day) only.

Sanger sequencing of the *ATP6V1B1*, *ATP6V0A4*, and *SLC4A1* genes as well as whole exome sequencing, which was performed in the Department of Pediatrics, Seoul National University Children's Hospital, revealed no pathogenic mutations.

Discussion

The patient had typical biochemical findings of RTA, including metabolic acidosis with a normal anion gap and hypokalemia. Additional findings of alkaline urine, positive urine anion gap, nephrocalcinosis, and rickets suggested that the patient had dRTA. The patient also had growth retardation with severe bony deformities and muscle weakness.

In children, dRTA is almost always observed as a primary entity with underlying genetic causes, and at least three genes are known to cause primary dRTA: *ATP6V1B1*, *ATP6V0A4*, and *SLC4A1*^{4,5}. *ATP6V1B1* and *ATP6V0A4* encode the B1 and A4 subunits of the apical H⁺-ATPase pump, that is, the proton pump, in α -intercalated cells, respectively, and mutations in these genes cause autosomal recessive (AR) forms of dRTA with or without sensorineural hearing loss^{6,7}. In a recent large multicenter study by the European dRTA Consortium, the prevalence of hearing loss was 88% in patients with *ATP6V1B1* mutations and 36% in patients with *ATP6V0A4* mutations⁸. *SLC4A1* encodes the basolateral anion (Cl⁻/HCO₃⁻) exchanger 1 (AE1)⁹. *SLC4A1* mutations can cause dRTA and/or hemolytic

anemia with abnormal red cell morphology because AE1 is expressed not only in α -intercalated cells but also in erythrocytes^{9,10}. In non-tropical areas, the majority of *SLC4A1* mutations have been reported in association with the autosomal dominant (AD) form of dRTA⁹. However, AR form of dRTA caused by *SLC4A1* mutations has been reported mostly from tropical Southeast Asia and other tropical areas, and is therefore called tropical recessive dRTA¹¹. We previously reported three Lao patients from two unrelated families with AR dRTA in association with *SLC4A1* mutations¹².

Genotype–phenotype analysis in the European dRTA Consortium revealed that patients with AD *SLC4A1* mutations had a milder phenotype and older onset age compared with those with mutations in the proton pump subunits⁸. While 77% of patients with proton pump subunit mutations presented in the first year of life and all before 10 years of age, 12% of patients with AD *SLC4A1* mutations presented in adulthood⁸. However, phenotypes of AR *SLC4A1* mutations are quite different from the phenotypes of AD *SLC4A1* mutations, but are similar in many respects to the AR dRTA caused by proton pump subunit mutations, such as younger onset age and more severe biochemical changes¹¹. The present case showed unusual phenotypes, that is, later disease onset at the age of 8 years (compatible with AD *SLC4A1* mutations) with severe biochemical and bone involvement (compatible with the AR form of dRTA). Genetic studies, including whole exome sequencing, in the patient revealed no pathogenic mutations. However, it is difficult to determine whether the patient has a hereditary or acquired form of dRTA because she may have a genetic defect that cannot be detected by Sanger sequencing and whole exome sequencing. In the European dRTA Consortium study⁸, 36 of 206 (17.5%) patients who underwent complete genetic tests had no pathogenic mutations. Otherwise, this patient may have a secondary form of dRTA due to prolonged immobilization-induced renal medullary nephrocalcinosis.

Growth failure and/or rickets, resulting from defective bone mineralization in association with chronic metabolic acidosis, are other characteristic features of untreated dRTA patients^{13–15}. However, catch-up growth is observed in most patients after the initiation of alkaline treatment¹⁵. In this patient, bone involvement was unusually severe because

she was left in a bed-ridden state for a long time without adequate treatment. The patient also complained of progressive muscle weakness, which may be due to chronic persistent hypokalemia and malnutrition¹³. Biochemical abnormalities and muscle weakness in the patient were normalized by supplementation therapy with alkali, vitamin D, and potassium. However, bony deformities required corrective surgical treatment.

In conclusion, dRTA in children is a controllable disease with favorable long-term prognosis, and accurate diagnosis and proper management are very important. Although this is a single case report of dRTA, the severe phenotypes in this long-term untreated case are of some educational value.

Funding

Dr. Philavanh Kedsatha was a fellow of Lee Jong-Wook Seoul Project in 2019. Lee Jong-Wook Seoul Project is a cooperative project between University of Health Sciences, Lao PDR and Seoul National University College of Medicine, Korea (2011-2020) funded by Korea Foundation for International Healthcare (KOFIH).

This research was supported by a grant of the Korea Health Technology Research & Development Project through the Korea Health Industry Development Institute, funded by the Ministry of Health & Welfare, Republic of Korea (grant number: HI18C0013).

Patient consent

This study was approved by the Institutional Review Board of Seoul National University Hospital, and the consent was waived due to the nature of the retrospective study (IRB number 2004-027-1114).

ORCID IDs

Philavanh Kedsatha <https://orcid.org/0000-0003-3962-7313>

Hee Young Shin <https://orcid.org/0000-0003-2091-1947>
Yong Choi <https://orcid.org/0000-0003-2287-3745>

Hae Il Cheong <https://orcid.org/0000-0001-7556-1265>

Tae-Joon Cho <https://orcid.org/0000-0001-8514-377X>

Eunsang Yi <https://orcid.org/0000-0001-8214-2106>

Mayfong Maisai <https://orcid.org/0000-0002-6056-4516>

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

References

- Rodríguez-Soriano J. Renal tubular acidosis: the clinical entity. *J Am Soc Nephrol* 2002;13:2160-70.
- Fry AC, Karet FE. Inherited renal acidosis. *Physiology (Bethesda)* 2007;22:20211.
- Albright F, Burnett CH, Parson W, Reifenstein EC Jr, Roos A. Osteomalacia and late rickets; the various etiologies met in the United States with emphasis on that resulting from a specific form of renal acidosis, the therapeutic indications for each etiological sub-group, and the relationship between osteomalacia and Milkman's syndrome. *Medicine* 1946;25:399-479.
- Karet FE. Inherited distal renal tubular acidosis. *J Am Soc Nephrol* 2002;13:2178-84.
- Battle D, Haque SK. Genetic causes and mechanisms of distal renal tubular acidosis. *Nephrol Dial Transplant* 2012;27:3691-704.
- Karet FE, Finberg KE, Nelson RD, Nayir A, Mocan H, Sanjad SA, et al. Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. *Nat Genet* 1999;21:84-90.
- Smith AN, Skaug J, Choate KA, Nayir A, Bakkaloglu A, Ozen S, et al. Mutations in ATP6N1B, encoding a new kidney vacuolar proton pump 116-kD subunit, cause recessive distal renal tubular acidosis with preserved hearing. *Nat Genet* 2000;26:71-5.
- Lopez-Garcia SC, Emma F, Walsh SB, Fila M, Hooman N, Zaniew M, et al. Treatment and long-term outcome in primary distal renal tubular acidosis. *Nephrol Dial Transplant* 2019;34:981-91.
- Karet FE, Gainza FJ, Györy AZ, Unwin RJ, Wrong O, Tanner MJ, et al. Mutations in the chloride-bicarbonate exchanger gene AE1 cause autosomal dominant but not autosomal recessive distal renal tubular acidosis. *Proc Natl Acad Sci U S A* 1998;95:6337-42.
- Tanphaichitr VS, Sumboonnanon A, Ideguchi H, Shayakul C, Brugnara C, Takao M, et al. Novel AE1 mutations in recessive distal renal tubular acidosis: loss-of-function is rescued by glycoporphin. *J Clin Invest* 1998;102:2173-9.
- Khositseth S, Bruce LJ, Walsh SB, Bawazir WM, Ogle GD, Unwin RJ, et al. Tropical distal renal tubular acidosis: clinical and epide-

- miological studies in 78 patients. *QJM* 2012;105:861-77.
12. Park E, Phaymany V, Yi ES, Phangmanixay S, Cheong HI, Choi Y. Primary autosomal recessive distal renal tubular acidosis caused by a common homozygous SLC4A1 mutation in two Lao families. *J Korean Med Sci* 2018;33:e95.
 13. Ranawaka R, Dayasiri K, Gamage M. A child with distal (type 1) renal tubular acidosis presenting with progressive gross motor developmental regression and acute paralysis. *BMC Res Notes* 2017;10:618.
 14. Soares SBM, de Menezes Silva LAW, de Carvalho Mrad FC, Simões E Silva AC. Distal renal tubular acidosis: genetic causes and management. *World J Pediatr* 2019;15:422-31.
 15. Rodriguez-Soriano J, Vallo A, Castillo G, Oliveros R. Natural history of primary distal renal tubular acidosis treated since infancy. *J Pediatr* 1982;101:669-76.