Original Research Article

## Anti-skin Aging Potential of Alcoholic Extract of Phragmites communis Rhizome

Chang Woo Ha<sup>1†</sup>, Sung Hyeok Kim<sup>1†</sup>, Sung Ryul Lee<sup>2†</sup>, Sohee Jang<sup>3</sup>, Seung Namkoong<sup>4</sup>, Sungsil Hong<sup>4</sup>, Hyosun Lim<sup>5</sup>, Youn Kyu Kim<sup>6</sup> and Eun-Hwa Sohn<sup>4,7</sup>\*

<sup>1</sup>Graduate Student, <sup>3</sup>Undergraduate Student and <sup>4</sup>Professor, College of Health Sciences, Kangwon National University, Samcheok 25949, Korea

<sup>2</sup>Professor, Department of Convergence Biomedical Science, Cardiovascular and Metabolic Disease Center, College of Medicine, Inje University, Busan 47392, Korea

<sup>5</sup>Master's Degree Researcher and <sup>6</sup>Post-doc, Korea Research Institute Bio Science Co., Ltd., Yeoju 12668, Korea <sup>7</sup>Professor, Department of Bio-Health Convergence, Kangwon National University, Chuncheon 24341, Korea

Abstract - Chronological aging and photoaging affect appearance, causing wrinkles, pigmentation, texture changes, and loss of elasticity in the skin. *Phragmites communis* is a tall perennial herb used for its high nutritional value and for medicinal purposes, such as relief from fever and vomiting and facilitation of diuresis. In this study, we investigated the effects of ethanol extract of *P. communis* rhizome (PCE) on skin aging. The total flavonoid and total phenolic content in PCE were  $2.92 \pm 0.007 \ \mu g$  of quercetin equivalents (QE) and  $231.8 \pm 0.001 \ \mu g$  of gallic acid equivalents (GAE) per 100 mg of dried extract (n = 3). The half-maximal inhibitory concentration (IC50) values of PCE for 2,2-diphenyl-1-picrylhydrazyl radical (DPPH) and hydrogen peroxide scavenging activities were 0.96 and 0.97 mg/mL, respectively. PCE showed inhibitory effects on tyrosinase when L-tyrosine (IC50 = 1.25 mg/mL) and L-3,4-dihydroxyphenylalanine (IC50 = 0.92 mg/mL) were used as substrates. PCE treatment up to 200 µg/mL for 24 h did not cause any significant cytotoxicity in B16F10 melanocytes, human dermal fibroblasts (HDFs), and HaCaT keratinocytes. In B16F10 melanocytes, PCE (25 and 50  $\mu$ g /mL) inhibited melanin production and cellular tyrosinase activity after challenge with  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH; p < 0.05). In HDFs, PCE suppressed the mRNA expression of matrix metalloproteinase-1 (MMP-1) and reduced the activity of elastase ( $p \le 0.05$ ). In addition, ultraviolet B (UVB)-mediated downregulation of hyaluronic acid synthase-2 gene expression in HaCaT keratinocytes was also effectively suppressed by PCE treatment. Overall, our results showed that PCE has potential anti-skin aging activity associated with the suppression of hyperpigmentation, wrinkle formation, and reduction in dryness. PCE is a promising candidate for the development of an anti-skin aging cosmetic ingredient.

Key words - Anti-aging, Elastase, Hyaluronic acid synthase-2, Matrix metalloproteinase-1, Phragmites communis

## Introduction

The skin is a physical barrier to external stimuli, infection, and moisture loss (Lee *et al.*, 2006). Keratinocytes form the major cell population in the epidermis and are involved in the development of the body's barrier against environmental insults and water loss. Fibroblasts contribute to building the architecture of the extracellular matrix (ECM) in the dermis and produce various types of matrix metalloproteinases

\*Corresponding author. E-mail : ehson@kangwon.ac.kr Tel. +82-33-540-3322

<sup>†</sup> These authors contributed equally to this work.

(MMPs), including MMP-1 and elastase (Freitas-Rodríguez *et al.*, 2017). Skin aging is caused by chronological aging and/or photoaging, which is a premature type of skin aging caused by repeated and prolonged exposure to sunlight (Fisher *et al.*, 2002). It has been postulated that reactive oxygen species (ROS) play a central role in both chronological aging and photoaging (Fisher *et al.*, 2002). Excessive ROS production due to intrinsic or extrinsic factors, such as sunlight, increases MMP expression and transforming growth factor- $\beta$  (TGF- $\beta$ ) signaling, which controls collagen homeostasis and affects wrinkle formation (Shin *et al.*, 2019). ROS stimulate the  $\alpha$ -melanin-stimulating hormone ( $\alpha$ -MSH)-

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

tyrosinase signaling pathway and consequently, increase melanin production from keratinocytes (Costin and Hearing, 2007). The cumulative structural and physiological alterations in the ECM caused by the aging process result in unattractive changes in skin appearance, such as folds and wrinkles (Ganceviciene *et al.*, 2012). Wrinkles are a sign of loss of flexibility and dryness in the skin. Good skin health and beauty are considered to be representative of an overall high quality-of-life and a perception of good health in humans. In addition to invasive procedures, including chemical peelings and injection of botulinum toxin or filler, extensive efforts have been made to develop topical pharmacological agents (Shin *et al.*, 2019). The development and standardization of plant-derived agents to inhibit skin aging is a key challenge for the cosmetics industry.

Phragmites communis (common reed) is a hydrophytic species inhabiting fresh swamps, riverbanks, and lakes; it grows well in temperate regions such as China, Korea, and Japan. Fresh and/or dried Phragmites rhizome has long been prescribed in Korea to relieve fever and vomiting and to facilitate diuresis (Park and Choi, 2016). A recent toxicological study showed that the aqueous extract of Phragmites rhizome had neither genotoxic nor mutagenic activities (Kim et al., 2019). Polysaccharides isolated from P. communis exerted anti-aging effects in a mouse model of aging (Miao et al., 2004). In addition, young leaves of P. communis played a role in the suppression of melanogenesis in B16F10 melanocytes (Sim et al., 2017). However, there are few reports on whether the ethanol extract of P. communis rhizomes (PCE) has antiskin aging effects. In this study, we evaluated the total flavonoid and phenolic contents as well as the antioxidative potential of PCE based on ROS scavenging activities. The anti-melanogenic effects of PCE were evaluated through the inhibitory effects on tyrosinase in vitro and the suppression of melanin production in B16F10 melanocytes stimulated with a-MSH. In addition to the acceleration of melanogenesis, repeated ultraviolet (UV) irradiation damages the skin, resulting in a visible "solar scar" or wrinkles associated with photoaging (Helfrich et al., 2008). The anti-wrinkle potential of PCE was evaluated based on changes in the mRNA expression of MMP-1 and the elastase activity in human dermal fibroblasts (HDFs). In addition, to evaluate skin

moisturization, changes in hyaluronic acid synthase-2 (HAS-2) gene expression in the presence or absence of UVB radiation in HaCaT keratinocytes were evaluated using qRT-PCR.

## Materials and Methods

## Chemicals

Ascorbic acid, arbutin, dimethylsulfoxide (DMSO), 2,2-diphenyl-1-picryl-hydrazyl radical (DPPH), mushroom tyrosinase,  $\alpha$ -MSH, L-3,4-dihydroxyphenylalanine (L-DOPA), L-tyrosine, phenylmethylsulfonyl fluoride, phosphoramidon, retinol, N-succinyl-tri-alanyl-p-nitroanilide (STANA), TGF- $\beta$ , and deionized distilled water were purchased from Sigma-Aldrich (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM) was obtained from Welgene Inc. (Daegu, Korea). Fetal bovine serum was obtained from GenDEPOT (Katy, TX, USA). Absolute ethanol was obtained from DUKSAN (Seoul, Korea), penicillin/streptomycin was obtained from Gibco (Grand Island, NY, USA), and Trizol reagent was obtained from Invitrogen (Carlsbad, CA, USA). Unless otherwise indicated, all other chemicals were purchased from Sigma-Aldrich.

### Preparation of P. communis Trinius root extract

The dried roots of *P. communis* Trinius were purchased from a local market in Suncheon-si (Korea) in January 2018. The material was identified by Dr. Y. K. Kim of the Korea Research Institute of Bio Science Inc. and stored in the dark at  $4^{\circ}$ C until use. The dried roots (10 g) were mixed with 200 mL of 100% EtOH and extracted in a water bath at 60 °C for 4 h. The extract was filtered to remove any particulate matter, and evaporated under reduced pressure to obtain a powder, which was stored at -20 °C for further use in experiments. The yield of PCE was 4.2% (weight/weight).

## Determination of total phenolic content and total flavonoid content

The total phenolic content was determined using the Folin-Ciocalteu method, as previously reported (So *et al.*, 2019). Briefly, the extract was mixed with Folin–Ciocalteu's phenol reagent (50 g/100 mL) and sodium carbonate (2 g/100 mL). After adjusting the final volume to 5 mL with deionized water, the reaction mixture was incubated at 25 °C for 30 min. The absorbance at 750 nm was measured using a VICTOR X3 microplate reader (PerkinElmer, Waltham, MA, USA). The total phenolic content of PCE was calculated from a standard curve and expressed as  $\mu g$  of gallic acid equivalent (GAE) /100 mg of dried extract.

The total flavonoid content of PCE was determined using an aluminum chloride colorimetric method (So *et al.*, 2019). Briefly, the extract was mixed with aluminum chloride hexahydrate (10 g/100 mL), 95% EtOH, and 1 M potassium acetate. After adjusting the final volume to 5 mL with deionized water, the reaction mixture solution was incubated at 25°C for 40 min. The absorbance at 415 nm was measured using a VICTOR X3 microplate reader. The total flavonoid content in the sample was calculated from the standard curve and expressed as  $\mu$ g of quercetin equivalent (QE)/100 mg of dried extract. All experiments were performed at least in triplicate.

### Determination of DPPH radical scavenging activity

DPPH radical scavenging activity was determined as previously reported (So *et al.*, 2019). Briefly, 100  $\mu$ L reaction mixture containing different concentrations of PCE was reacted with 100  $\mu$ L of 0.4 mM DPPH solution in 99% EtOH at 25 °C for 30 min. Decolorization of DPPH was measured at 540 nm using a VICTOR X3 microplate reader. The halfmaximal inhibitory concentration (IC<sub>50</sub>) for DPPH radical scavenging activity was calculated graphically by plotting the inhibition percentage against the tested PCE concentrations.

#### Determination of hydrogen peroxide scavenging activity

Hydrogen peroxide scavenging activity was determined as previously reported (So *et al.*, 2019). Briefly, reaction mixtures containing different concentrations of PCE were reacted with an equal volume of 1.0 M H<sub>2</sub>O<sub>2</sub> and 0.1 M phosphate buffer at  $37^{\circ}$ C for 5 min. Then, 3-ethylbenzothiazoline-6-sulfonic acid and peroxidase were added. After the mixture was incubated at  $37^{\circ}$ C for 10 min, changes in the absorbance at 405 nm were measured using a VICTOR X3 microplate reader. The IC<sub>50</sub> value for hydrogen peroxide scavenging activity was calculated graphically by plotting the percentage inhibition against the tested PCE concentrations.

## Cell culture

Mouse melanoma B16F10 cells were obtained from the Korean Cell Line Bank (Seoul, Korea). Human dermal fibroblasts (HDFs) and human keratinocyte HaCaT cells were purchased from Gibco and CLS Cell Lines Service (Eppelheim, Baden-Württemberg, Germany), respectively. Cells were maintained in DMEM containing 10% fetal bovine serum and 1% penicillin/streptomycin in a humidified atmosphere of 5% CO<sub>2</sub> at  $37^{\circ}$ C.

### Cell viability

Cell viability was measured using the quantitative colorimetric 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, as previously reported (Koo *et al.*, 2017). B16F10 melanocytes in the exponential growth phase were seeded ( $1 \times 10^4$  cells/well) into 96-well tissue culture plates and treated with different concentrations (25-400 µg /mL) of PCE for 24 and 48 h. HDFs and HaCaT keratinocytes were seeded ( $1 \times 10^4$  cells/well) into 96-well tissue culture plates and treated with different concentrations (25-400 µg /mL) of PCE for 24 h. After incubation with MTT (150 µg /mL) of PCE for 24 h. After incubation with MTT (150 µg /mL) for 4 h, the formed formazan crystals were dissolved in DMSO, and the optical density at 540 nm was measured using a VICTOR X3 microplate reader. Cytotoxicity was expressed as a percentage of the untreated control cells.

## Measurement of melanin production in B16F10 melanocytes

B16F10 melanocytes were seeded (2 × 10<sup>5</sup> cells/well) into 6-well cell culture plates and allowed to adhere overnight. The cells were then treated with  $\alpha$ -MSH (100 nM) in the presence or absence of PCE (25-200 µg/mL) for 48 h. The cells were washed with ice-cold phosphate-buffered saline (PBS) and harvested using a cell scraper. After centrifugation (2500 × g, 25 °C, 1 min), the pellets were dissolved in 1 N NaOH for 1 h at 60°C. The absorbance at 490 nm was measured using a VICTOR X3 microplate reader.

#### Measurement of tyrosinase inhibitory activity

As previously reported (So *et al.*, 2019), the inhibitory activity of PCE on tyrosinase was determined in a cell-free system using a colorimetric method with L-DOPA and L-tyrosine as the substrates. Different concentrations of PCE were added to the reaction buffer containing mushroom tyrosinase, and the enzymatic reaction was initiated after the addition of L-tyrosine or L-DOPA. After incubation at 37°C for 15 min, the absorbance at 490 nm was measured using a VICTOR X3 microplate reader. The IC<sub>50</sub> value for tyrosinase activity was calculated from the graph by plotting the inhibition percentage against the tested PCE concentrations. To determine the inhibitory effect of PCE on cellular tyrosinase, B16F10 melanocytes were seeded ( $2 \times 10^5$  cells/ well) into 6-well cell culture plates, and the cells were treated with  $\alpha$ -MSH (100 nM) in the presence or absence of PCE (25-200  $\mu$ g/mL) for 48 h. The cells were harvested, washed with PBS, and then lysed with lysis buffer containing 50 mM sodium phosphate buffer (pH 6.8), 1% Triton X-100, and 0.1 mM phenylmethylsulfonyl fluoride. After centrifugation (15,000  $\times$  g, 4°C, 30 min), tyrosinase activity of the cell supernatant (40  $\mu$ g) was determined in a reaction mixture containing 40 µL of 100 mM sodium phosphate buffer (pH 6.8) and 160 µL of 10 mM L-DOPA for 1 h. Then, the absorbance at 490 nm was measured using a VICTOR X3 microplate reader.

### Measurement of cellular elastase activity

HDFs were seeded (2 × 10<sup>6</sup> cells/well) into 100-mm tissue culture dishes. Confluent cells were lysed with PRO-PREP<sup>TM</sup> protein extraction solution (Invitrogen, Seongnam, Korea), and the cell lysates were collected by centrifugation (15,000 × g, 4°C, 30 min). Equal volumes of cell lysates (each containing 100 µg protein) were added to a 96-well plate. Then, PCE (25-200 µg/mL) and STANA (50 mM) were added, successively. After incubation at 37 °C for 90 min, the release of p-nitroaniline was measured at 405 nm using a VICTOR X3 microplate reader. Elastase activity was expressed as a percentage of the untreated control.

#### **UVB** irradiation

UVC 500 UltraViolet Crosslinker (Amersham, Bucks, UK) was used for UVB irradiation, as previously described (Park *et al.*, 2018). Briefly, HaCaT keratinocytes at approximately 90% confluence were switched to serum-free DMEM for 4 h. The cells were then covered with PBS and exposed to UVB light (34.06 s) for a total dose of 30 mJ/cm<sup>2</sup>. After washing with PBS, the cells were further cultured in DMEM containing

10% fetal bovine serum with or without PCE for 24 h. The control group treated in the same manner but did not receive UVB irradiation.

## Quantitative RT-PCR (qRT-PCR) analysis of MMP-1 and HAS-2 mRNA expression

The mRNA expression of MMP-1 and HAS-2 was detected in HDFs and HaCaT keratinocytes, respectively. Total cellular RNA was extracted using Trizol<sup>TM</sup> reagent by the phenol-chloroform extraction method and reverse-transcribed to synthesize cDNA using a cDNA synthesis kit (ECDNA100, NanoHelix, Daejeon, Korea). Subsequently, PCR amplification was performed using a premier qPCR kit (PQL-S500, NanoHelix). The following reverse transcription conditions were used:  $65^{\circ}$  for 5 min,  $42^{\circ}$  for 60 min, and  $95^{\circ}$  for 5 min. The two-step PCR amplification conditions were 95°C for 30 s, followed by 40 cycles of 95  $^{\circ}$ C for 5 s and 60  $^{\circ}$ C for 30 s to produce a melt curve. The primer sequences used were: 5'-AGT GGC CCA GTG GTT GAA AA-3' (forward) and 5'-CCA CAT CAG GCA CTC CAC AT-3' (reverse) for MMP-1, 5'-TGG ATC TCA TTC CTC AGC AG-3' (forward) and 5'-TCC ACA AAC TCA TGC AAC AA-3' (reverse) for HAS-2; and 5'-GTG GCA AAG TGG AGA TTG CC-3' (forward) and 5'-GAT GAT GAC CCG TTT GGC TCC-3' (reverse) for GAPDH.

#### Statistical analysis

Each experiment was repeated at least three times. The results were expressed as the mean  $\pm$  standard error of the mean (SEM) and were analyzed using one-way ANOVA followed by Tukey's test (Systat Software Inc., San Jose, CA, USA). A statistical probability value of p < 0.05 was considered significant.

## Results

#### Total phenolic content and total flavonoid content

Phenolic acids and flavonoids are the major antioxidants found in plants (Cai *et al.*, 2004). In addition to their broad biological functions, such as antioxidant and anti-inflammation activity, both types of compounds possess photoprotective activity because of their structural similarity to chemical filters (Saewan and Jimtaisong, 2015). The total phenolic content in PCE was determined to be 231.8  $\pm$  0.001 µg GAE/100 mg of dried extract (n = 3) and that of flavonoid was 2.92  $\pm$  0.007 µg QE/100 mg of dried extract (n = 3).

### **Radical scavenging activity of PCE**

The antioxidant activities of up to 200  $\mu$ g/mL PCE were evaluated in terms of the scavenging potential of DPPH radicals (Fig. 1A) and hydrogen peroxide (Fig. 1B). The IC<sub>50</sub> values of PCE for the DPPH radical and hydrogen peroxide scavenging were 0.96 and 0.97 mg/mL, respectively.

#### Inhibitory effect of PCE on tyrosinase activity in vitro

Suppression of tyrosinase activity is a key event in antimelanogenesis. Polyphenols belong to the largest group of tyrosinase inhibitors and are potential skin-whitening agents (Chang, 2009). To determine the anti-melanogenic effect of PCE, we determined its inhibitory effect on tyrosinase in a cell-free system with different substrates, L-tyrosine and L-DOPA. The IC<sub>50</sub> values of PCE on tyrosinase were 1.25 and



Fig. 1. Radical scavenging activity of PCE. The antioxidant activities of PCE were determined in terms of DPPH (A) and hydrogen peroxide (B) scavenging activities. Ascorbic acid was used as the control.  $IC_{50}$  values were determined using the regression equation. Data are presented as mean  $\pm$  SEM (n = 4). DPPH, 2,2-diphenyl-1-picryl-hydrazyl-hydrate;  $IC_{50}$ , half-maximal inhibitory concentration; PCE, ethanol extract of *P. communis* rhizome.



Fig. 2. Inhibitory effects of PCE on tyrosinase in a cell-free system. Tyrosinase regulatory activities were determined using a colorimetric method. L-tyrosine (A) and L-DOPA (B) were used as substrates for tyrosinase.  $IC_{50}$  values were determined using the regression equation. Data are means  $\pm$  SEM (n = 4) and are expressed as a percentage of the untreated control.  $IC_{50}$ , half-maximal inhibitory concentration; PCE, ethanol extract of *P. communis* rhizome.

0.92 mg/mL for the substrates L-tyrosine (Fig. 2A) and L-DOPA (Fig. 2B), respectively.

## Effect of PCE on the viability of B16F10 melanocytes, HDFs, and HaCaT keratinocytes

The MTT assay was performed to determine the cytotoxicity of PCE treatment in B16F10 melanocytes, HDFs, and HaCaT keratinocytes. Unlike the subsequent experiments performed in HDFs and HaCaT keratinocytes, the measurement of melanin production in B16F10 melanocytes requires treatment for 48 h of treatment. Thus, the effect of PCE on the viability of B16F10 melanocytes was also determined at 48 h. No significant cytotoxicity of PCE was observed up to 200  $\mu$ g /mL in B16F10 melanocytes (Fig. 3A), HDFs (Fig. 3C), and HaCaT keratinocytes (Fig. 3D) at 24 h. However, at 400  $\mu$ g /mL, PCE resulted in significant cytotoxicity in B16F10 melanocytes at 48 h (Fig. 3B). Thus, a PCE concentration below 200  $\mu$ g/mL was used for the subsequent experiments.

## Inhibitory effect of PCE on melanin production in B16F10 melanocytes

 $\alpha$ -MSH is a strong inducer of tyrosinase expression and is involved in the promotion of melanogenesis. The effect of PCE on melanin production in B16F10 melanocytes was determined in the presence of  $\alpha$ -MSH (100 nM). Arbutin strongly inhibited tyrosinase and was thus used as a positive control.  $\alpha$ -MSH stimulation increased melanin production in B16F10 melanocytes by 147 ± 1.49% compared with



Fig. 3. Cytotoxicity of PCE. Melanocyte B16F10 cells were incubated with PCE for 24 h (A) and 48 h (B). Human dermal fibroblasts (HDFs) (C) and keratocytes (HaCaT) (D) were incubated in the presence of PCE for 24 h. The treatment dose of PCE ranged from 12.5 to 400  $\mu$ g/mL. Cell cytotoxicity was determined using the MTT assay. Data are means  $\pm$  SEM (n = 5) and are expressed as percentages of the untreated control. MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; PCE, ethanol extract of *P. communis* rhizome.



Fig. 4. Effects of PCE on melanin production and tyrosinase activity in B16F10 cells. B16F10 cells were exposed to  $\alpha$ -MSH (100 nM) in the presence or absence of PCE (25–200  $\mu$ g/mL) for 48 h. The content of melanin is expressed as a percentage of the untreated control cells (A). Cellular tyrosinase activity was determined by a colorimetric method and expressed as a percentage of the untreated control (B). Data are presented as mean ± SEM (n = 5). ##p < 0.01 vs. untreated control and \*\*p < 0.01 vs.  $\alpha$ -MSH-treated control.  $\alpha$ -MSH,  $\alpha$ -melanin-stimulating hormone; PCE, ethanol extract of *P. communis* rhizome.

untreated control (p < 0.05).  $\alpha$ -MSH-mediated melanin production was significantly suppressed in the presence 25 and 50  $\mu$ g/mL PCE. However, the inhibitory effect of PCE on melanin production not observed at PCE concentrations above 100  $\mu$ g/mL (Fig. 4A).

# Inhibitory effect of PCE on cellular tyrosinase activity stimulated by $\alpha$ -MSH in B16F10 melanocytes

PCE suppressed melanin production in  $\alpha$ -MSH-stimulated B16F10 melanocytes (Fig. 4A). As shown in Fig. 3, PCE showed a direct inhibitory effect on tyrosinase activity. Therefore, we subsequently examined the effect of PCE on tyrosinase activity in  $\alpha$ -MSH-stimulated B16F10 melanocytes (Fig. 4B). Consistent with the results of changes in the melanin content (Fig. 4A), significant tyrosinase inhibitory activity of PCE was observed at 25 and 50  $\mu$ g/mL (p < 0.05).

## Suppressive effect of PCE on the mRNA expression of MMP-1 in HDFs

Upon exposure to UVB, activator protein 1 (AP-1) is activated through various signaling pathways. The activation of AP-1 involves the upregulation of collagen-degrading enzymes (e.g., MMPs) and inhibition of collagen synthesis by counteracting the TGF- $\beta$  signaling pathway (Fisher *et al.*, 2002). MMP-1 is involved in the initiation of collagen cleavage (Fisher *et al.*, 2009) and is thus a major contributor to degradation of the ECM. When the mRNA expression of MMP-1 in HDFs was determined via qRT-PCR, 25 and 50  $\mu$ g /mL PCE was found to significantly downregulate MMP-1 expression (Fig. 5A, p < 0.05).

### Suppressive effect of PCE on elastase activity in HDFs

Skin fibroblasts secrete elastase, which belongs to the family of metalloproteinases. Skin fibroblast elastase plays a crucial role in the degradation of elastic fibers associated with wrinkle formation (Tsuji *et al.*, 2001). Here, phosphoramidon, an inhibitor of skin fibroblast-derived elastase isolated from *Pseudomonas aersa* (Morihara and Tsuzuki, 1978), was used as a positive control. As shown in Fig. 5B, PCE significantly inhibited elastase activity in HDFs (p < 0.05).

### Effect of PCE on the mRNA expression of HAS-2

Skin moisture is closely associated with water retention in the skin. Skin hyaluronan, or hyaluronic acid (HA), which is synthesized by the membrane-bound enzyme HAS, is a major contributor to water retention (Papakonstantinou *et al.*, 2012). During skin aging, a significant reduction in the content of HA and the downregulation of HAS-2 have been identified (Papakonstantinou *et al.*, 2012). To determine the moisturizing effect of PCE, we evaluated the mRNA expression of HAS-2



Fig. 5. Effects of PCE on mRNA expression levels of MMP-1 and elastase activity in HDFs cells. HDFs were treated with different doses of PCE for 24 h. The mRNA expression levels of the MMP-1 gene were determined using qRT-PCR and normalized to mRNA levels of GAPDH. Results are expressed as an index of the untreated control (A). Elastase activity was measured using STANA as a substrate and expressed as a percentage of the untreated control (B). TGF- $\beta$  and PPR were used as positive controls for mRNA expression of MMP-1 and elastase activity, respectively. Data are presented as mean ± SEM (n = 5). ##p < 0.01 vs. CON and \*\*p < 0.01 vs.  $\alpha$ -MSH-treated control.  $\alpha$ -MSH,  $\alpha$ -melanin stimulating hormone; CON, untreated control; MMP-1, matrix metal-loprotease-1; PCE, ethanol extract of *P. communis* rhizome; PPR, phosphoramidon; STANA, N-succinyl-tri-alanyl-p-nitroanilide; TGF- $\beta$ , transforming growth factor- $\beta$ .



Fig. 6. Effect of PCE on the mRNA expression levels of HAS-2. UVB irradiation strength was set at 30 mJ/cm<sup>2</sup>. One group of HaCaT cells was treated with different concentrations of PCE (A), whereas the other group was treated with different concentrations of PCE followed by UVB irradiation (B). The mRNA expression levels of the HAS-2 gene were determined using qRT-PCR and normalized to mRNA levels of the GAPDH gene. ROL (10  $\mu$ g/mL) was used as a stimulator of HAS-2 gene expression. Data are means ± SEM (n = 5) and expressed as an index of the respective control. ##p < 0.01 vs. CON and \*\*p < 0.01 vs. UVB-treated control. CON, untreated control; HAS-2, hyaluronan synthase-2; PCE, ethanol extract of *P. communis* rhizome; ROL, retinol; UVB, ultraviolet B irradiation.

in HaCaT keratinocytes in the presence or absence of UVB. Retinol is a strong inducer of HAS gene expression in skin tissue (Li *et al.*, 2017) and was used as the positive control. In basal conditions, the mRNA expression of HAS-2 did not increase at the applied PCE concentrations, except for 200  $\mu$ g /mL (Fig. 6A). However, PCE treatment reversed the decrease in HAS-2 gene expression that followed UVB exposure (Fig. 6B). This result suggested that PCE did not directly upre-

gulate the expression of HAS-2, but suppressed the UVBmediated downregulation of HAS-2 gene expression.

## Discussion

The results of our study showed the value of PCE in the alleviation of skin aging. PCE possessed tyrosinase inhibitory activity and suppressed the  $\alpha$ -MSH-mediated increase in melanin production in B16F10 melanocytes. PCE could not only suppress the mRNA expression of MMP-1, but also reduced the activity of elastase in HDFs. In addition, the UVB-mediated downregulation of HAS-2 gene expression in HaCaT keratinocytes was also effectively suppressed in the presence of PCE.

The conservation of smooth skin and youthful complexion is considered a very important part of an attractive appearance and beauty in Asia (Smit et al., 2009). Similar to other organs, the skin will age over time. Numerous approaches have been undertaken in the cosmetic industry to maintain healthy skin or prevent skin aging (Jadoon et al., 2015). Skin aging leads to the occurrence of folds and wrinkles due to atrophy of the dermis and deterioration of ECM organization (Helfrich et al., 2008). Prolonged and repeated exposure to sunlight, especially UVB, is a major cause of skin photoaging. Sunscreen provides a physical barrier against sunlight (e.g., UVB); however, simple protection against sunlight is not sufficient to maintain healthy skin. In some cases, synthetic skincare cosmetics may cause allergic reactions in the skin; however, natural skincare products may less frequently induce such reactions (Mukherjee et al., 2011). Alternatively, herbal extracts are good sources of bioactive compounds with anti-skin aging potential (Mukherjee et al., 2011). The suggested mechanisms of action of herbal extracts are associated with: 1) suppression of MMP activity; 2) antioxidant and free radical scavenging; or 3) increased hyaluronic acid content (Mukherjee et al., 2011).

In addition to the established use of Phragmitis rhizome as a traditional medicine (Park and Choi, 2016), we explored the anti-skin aging potential of PCE in this study. Antioxidants are recognized as important functional ingredients of cosmetic materials, with the ability to inhibit aging and skin pigmentation by ROS scavenging. PCE exhibits radical scavenging

activities (Fig. 1) and contains phenolic acids and flavonoids. In addition to their various biological functions, phenolic acids present in PCE may function as a UVB blocker and attenuate skin damage caused by sunlight. Tyrosinase is the key enzyme in melanogenesis and is increased in the skin during UVB exposure (Gillbro and Olsson, 2011). Thus, tyrosinase inhibitors are the most characterized cellular targets of skin-whitening agents (Chang, 2009). At 25-200  $\mu$ g /mL, PCE displayed tyrosinase inhibitory capacities in vitro and was more potent when L-DOPA was used as a substrate (Fig. 2). PCE possessed tyrosinase inhibitory activity and suppressed the a-MSH-mediated increase in melanin production in B16F10 melanocytes (Fig. 4). Collagen and elastin are the major constituents of the ECM that maintain skin architecture. The accelerated breakdown of the ECM impairs the dermal structure, resulting in the formation of wrinkles and the loss of skin elasticity. MMP-1 and elastase are involved in the degradation of ECM in the skin. The topical treatment of elastase inhibitors participates in the suppression of wrinkle formation caused by UVB exposure (Tsuji et al., 2001). The mRNA expression of MMP-1 and the activity of elastase were both reduced in the presence of PCE (Fig. 5); thus, these effects of PCE may contribute to the suppression of wrinkle formation. Finally, the maintenance of skin moisture, which is associated with hyaluronic acid content, is necessary to preserve the firmness and springiness of skin (Li et al., 2017). The levels of hyaluronic acid in the skin determine skin dryness and are frequently decreased by UVB exposure or normal aging. UVB exposure caused the downregulation of HAS-2 expression; however, this decrease in HAS-2 expression did not occur in the presence of PCE (Fig. 6). Therefore, PCE can contribute to the maintenance of hyaluronic acid in the skin. Interestingly, the suppressive effects of PCE on both cellular tyrosinase activity and the reduction of a-MSH-mediated melanin production were only observed with 25 and 50  $\mu$ g/mL PCE (Fig. 4). The disappearance of the inhibitory activity of PCE above 50  $\mu$ g/mL was also shown by the gene expression of MMP-1 in HDFs (Fig. 5A). However, the suppressive effect of PCE on the gene expression of HAS-2 was dose dependent. It should be noted that PCE shows differences in concentration or efficacy according to the type of skin cells used. The reason for this is unclear, and

requires further examination.

In conclusion, PCE is a potential cosmetic ingredient owing to its suppressive effects on melanogenesis, wrinkle formation, and water loss in the skin. Further studies are warranted to identify the active compounds involved in the individual anti-aging pathways.

## Acknowledgements

This research was conducted as part of the project entitled "Development of multi-functional cosmetic materials derived from halophytes" funded by the Ministry of Oceans and Fisheries, Korea.

## **Conflict of Interest**

The authors declare that they have no conflict of interest.

## References

- Cai, Y., Q. Luo, M. Sun and H. Corke. 2004. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. Life Sci. 74:2157-2184.
- Chang, T.S. 2009. An updated review of tyrosinase inhibitors. Int. J. Mol. Sci. 10:2440-2475.
- Costin, G.E. and V.J. Hearing. 2007. Human skin pigmentation: Melanocytes modulate skin color in response to stress. FASEB J. 21:976-994.
- Fisher, G.J., S. Kang, J. Varani, Z. Bata-Csorgo, Y. Wan, S. Datta and J.J. Voorhees. 2002. Mechanisms of photoaging and chronological skin aging. Arch. Dermatol. 138:1462-1470.
- Fisher, G.J., T. Quan, T. Purohit, Y. Shao, M.K. Cho, T. He, J. Varani, S. Kang and J.J. Voorhees. 2009. Collagen fragmentation promotes oxidative stress and elevates matrix metalloproteinase-1 in fibroblasts in aged human skin. Am. J. Pathol. 174:101-114.
- Freitas-Rodríguez, S., A.R. Folgueras and C. López-Otín. 2017. The role of matrix metalloproteinases in aging: Tissue remodeling and beyond. Biochim. Biophys. Acta. Mol. Cell. Res. 1864: 2015-2025.
- Ganceviciene, R., A.I. Liakou, A. Theodoridis, E. Makrantonaki and C.C. Zouboulis. 2012. Skin anti-aging strategies.

Dermato-endocrinology 4:308-319.

- Gillbro, J.M. and M.J. Olsson. 2011. The melanogenesis and mechanisms of skin-lightening agents--existing and new approaches. Int. J. Cosmet. Sci. 33: 210-221.
- Helfrich, Y.R., D.L. Sachs and J.J. Voorhees. 2008. Overview of skin aging and photoaging. Dermatol. Nurs. 20:177-183.
- Jadoon, S., S. Karim, M.H. Bin Asad, M.R. Akram, A.K. Khan, A. Malik, C. Chen and G. Murtaza. 2015. Anti-aging potential of phytoextract loaded-pharmaceutical creams for human skin cell longetivity. Oxid. Med. Cell. Longev. 2015:709628.
- Kim, N.S., S. Shin, G.G. Shin and O.S. Bang. 2019. Genotoxicity evaluation of a *Phragmitis* rhizoma extract using a standard battery of *in vitro* and *in vivo* assays. J. Ethnopharmacol. 241:112025.
- Koo, H.J., S.R. Lee, S.C. Kang, J.E. Kwon, D.E. Lee, E.S. Choung, J.S. Lee, J.W. Lee, Y. Park, D.S. Sim and E.H. Sohn. 2017. The suppressive effect of *Pueraria lobata* root extract and its biotransformed preparation against skin wrinkle formation. Korean J. Plant Res. 30:272-279.
- Lee, S.H., S.K. Jeong and S.K. Ahn. 2006. An update of the defensive barrier function of skin. Yonsei Med. J. 47:293-306.
- Li, W.H., H.K. Wong, J. Serrano, M. Randhawa, S. Kaur, M.D. Southall and R. Parsa. 2017. Topical stabilized retinol treatment induces the expression of HAS genes and HA production in human skin *in vitro* and *in vivo*. Arch. Dermatol. Res. 309:275-283.
- Miao, M.S., L.Y. Gu, X.Y. Fang and Y.Y. Miao. 2004. Effect of *Phragmites communis* polysaccharide on the aged-model mice. Zhongguo Zhong Yao Za Zhi 29:673-675.
- Morihara, K. and H. Tsuzuki. 1978. Phosphoramidon as an inhibitor of elastase from *Pseudomonas aeruginosa*. Jpn. J. Exp. Med. 48:81-84.
- Mukherjee, P.K., N. Maity, N.K. Nema and B.K. Sarkar. 2011. Bioactive compounds from natural resources against skin aging. Phytomedicine 19:64-73.
- Papakonstantinou, E., M. Roth and G. Karakiulakis. 2012. Hyaluronic acid: A key molecule in skin aging. Dermatoendocrinology 4:253-258.
- Park, J.C. and G. Choi. 2016. Review on herbal medicinal materials in the Korean pharmacopoeia and the Korean herbal pharmacopoeia. Korean Herb. Med. Inf. 4:9-35 (in Korean).
- Park, Y.A., S.R. Lee, J.W. Lee, H.J. Koo, S.A. Jang, S.W. Yun, H.J. Kim, J.S. Woo, M.R. Park, S.C. Kang, Y.K. Kim and E.H. Sohn. 2018. Suppressive effect of fermented *Angelica*

*tenuissima* root extract against photoaging: Possible involvement of hemeoxygenase-1. J. Microbiol. Biotechnol. 28:1391-1400.

- Saewan, N. and A. Jimtaisong. 2015. Natural products as photoprotection. J. Cosme. Dermatol. 14:47-63.
- Shin, J.W., S.H. Kwon, J.Y. Choi, J.I. Na, C.H. Huh, H.R. Choi and K.C. Park. 2019. Molecular mechanisms of dermal aging and antiaging approaches. Int. J. Mol. Sci. 20:2126.
- Sim, M.O., J.R. Ham and M.K. Lee. 2017. Young leaves of reed (*Phragmites communis*) suppress melanogenesis and oxidative stress in B16F10 melanoma cells. Biomed. Pharmacother.

93:165-171.

- Smit, N., J. Vicanova and S. Pavel. 2009. The hunt for natural skin whitening agents. Int. J. Mol. Sci. 10:5326-5349.
- So, G.S., S.R. Lee, S.H. Kim, C.W. Ha, Y. Park, S. Jang, J.P. Bak, H.J. Koo and E.H. Sohn. 2019. Antioxidant and antimelanogenic effects of *Stevia rebaudiana* flower extract. Korean J. Plant Res. 32:220-227.
- Tsuji, N., S. Moriwaki, Y. Suzuki, Y. Takema and G. Imokawa. 2001. The role of elastases secreted by fibroblasts in wrinkle formation: Implication through selective inhibition of elastase activity. Photochem. Photobiol. 74:283-290.

(Received 24 June 2020; Revised 30 July 2020; Accepted 3 August 2020)