


Complete genome sequence of *Clostridium perfringens* B20, a bacteriocin-producing pathogen

Axel G. Elnar and Geun-Bae Kim*

Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea



Received: Sep 10, 2021
Revised: Oct 8, 2021
Accepted: Oct 8, 2021

*Corresponding author

Geun-Bae Kim
Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea.
Tel: +82-31-670-3027
E-mail: kimgeun@cau.ac.kr

Copyright © 2021 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Axel G. Elnar
<https://orcid.org/0000-0002-2716-4924>
Geun-Bae Kim
<https://orcid.org/0000-0001-8531-1104>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

Not applicable.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Kim GB.
Data curation: Elnar AG.
Formal analysis: Elnar AG, Kim GB.

Abstract

Clostridium perfringens B20 was isolated from chicken feces collected from a local farm associated with Chung-Ang University (Anseong, Korea). The whole genome of *C. perfringens* B20 was sequenced using the PacBio RS II platform and assembled *de novo*. The genome is 2,982,563 bp long and assembled in two contigs. Annotation analyses revealed 2,668 protein-coding sequences, 30 rRNA genes, and 94 tRNA genes, with 28.2% G + C (guanine + cytosine) content. *In silico* genomic analysis revealed the presence of genes encoding a class IId bacteriocin, lactococcin A, and associated ABC transporter and immunity proteins, as well as a putative bacteriocin gene.

Keywords: *Clostridium perfringens*, Whole genome sequence, Pathogen, Virulence, Bacteriocin, Lactococcin A

INTRODUCTION

Bacteriocin production among lactic acid bacteria is currently a rapidly expanding field of research, which has been stimulated to a large extent by issues regarding the safety of using live probiotics and has, thus, driven a shift toward the development and elaboration of postbiotics [1]. Postbiotics are currently defined as bioactive compounds that are not probiotic, prebiotic, or paraprobiotic, and can include soluble factors, cell compounds, and metabolic products and/or by-products. Compared with live probiotics, some of the advantages of using postbiotics, including bacteriocins, include safer administration, longer shelf-life, and lower probability of the acquisition and spread of resistance genes and other virulence factors [2]. However, although bacteriocin production is considered a favorable trait for potential probiotic strains, the same may not hold true for pathogenic microbes.

In order to successfully colonize a host and establish a suitable niche, pathogens require diverse mechanisms that would enable them to evade possible immune responses, replicate, and ensure disease transmission. *Clostridium perfringens*, a Gram-positive bacterium, is a known causative agent for several disorders in humans (clostridial myonecrosis and food poisoning) and animals (necrotic enteritis in poultry) [3]. The pathogenicity of *C. perfringens* is associated with the production of major toxins (α -toxin, β -toxin, ι -toxin, and ϵ -toxin), subsidiary toxins (collagenase, enterotoxin, and perfringolysin O), and antimicrobial peptide resistance genes (e.g., tetracycline resistance efflux protein and specific aminoglycoside resistance genes) [4]. Here, based on genetic analysis, we report evidence of bacteriocin production by a *C. perfringens* isolate, provisionally designated as strain B20. Bacteriocin production by

Methodology: Elnar AG, Kim GB.
 Software: Elnar AG, Kim GB.
 Validation: Kim GB.
 Investigation: Elnar AG.
 Writing - original draft: Elnar AG.
 Writing - review & editing: Elnar AG, Kim GB.

Ethics approval and consent to participate

This article does not require IRB/ACUC approval because there are no human and animal participants.

a pathogen may enhance its pathogenicity through the same mechanisms employed by probiotic strains to become more beneficial; specifically, through direct inhibition of competing strains and niche establishment, as well as through modulating the local microbiota [2]. Thus, it is important to deduce how pathogens acquire genes for bacteriocin production and thereby gain a better understanding of how these traits may affect their pathogenicity.

C. perfringens B20 was isolated from the feces of chickens maintained in a Chung-Ang University-affiliated farm (Anseong, Korea). Strain B20 was routinely cultured anaerobically in Brain-Heart Infusion (BD Bacto, Sparks, MD, USA) medium at 37°C for 24 h [5]. Genomic DNA was extracted from 12 h cultures using a QIAamp PowerFecal DNA Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The genomic DNA obtained was sequenced commercially at ChunLab (Korea) using the Pacific Biosciences (PacBio, Menlo Park, CA, USA) RSII Single Molecule Real-Time (SMRT) platform with a 20-kb SMRTbell™ template library. *De novo* assembly of the PacBio reads was performed using the PacBio SMAR Analysis ver. 2.3.0 program, and genome annotation was performed based on Rapid Annotation using Subsystem Technology (RAST) with default parameters (<https://rast.nmpdr.org/>) and CLgenomics™ ver. 1.55 software. Transfer RNAs (tRNAs) were identified using tRNAscan-SE ver. 1.3.1 [6], whereas non-coding ribosomal RNAs (rRNAs) were identified using INFERNAL ver. 1.1.3 software in conjunction with the Rfam 12.0 database [7]. The complete genome of *C. perfringens* B20 (Fig. 1) is 2,982,563 bp long with a G + C (guanine + cytosine) content of 28.2% and is assembled into two




Fig. 1. Circular genome map of *Clostridium perfringens* B20. Circles represent the following characteristics from the outermost circle to the center: (1) contig information, (2) coding sequences on forward strand, (3) coding sequences on reverse strand, (4) transfer RNAs (tRNAs) and ribosomal RNAs (rRNAs), (5) GC skew, and (6) GC ratio. G, guanine; C, cytosine.

Table 1. Genome features of *Clostridium perfringens* B20

Attribute	Value
Genome size (bp)	2,982,563
GC content (%)	28.2
No. of contigs	2
Total genes	2,792
Protein-coding gene	2,668
tRNA	94
rRNA	30
Plasmids	0
GenBank accession No.	JAGKIL000000000

G, guanine; C, cytosine.




Fig. 2. Subsystem category distribution by KEGG annotation (A) and cluster of orthologous groups (B) based on the functional classification of the whole genome of *Clostridium perfringens* B20. KEGG, Kyoto Encyclopedia of Genes and Genomes.




Fig. 3. Predicted bacteriocin gene cluster in *Clostridium perfringens* B20 genome showing a single ORF for lactococcin A (green) using BAGEL4 software. ORF, open reading frame.

contigs with an N_{50} value of 2,925,822 bp. The genome contains 2,668 protein-coding sequences, 30 rRNA, and 94 tRNA genes (Table 1). Functional annotation of protein-coding sequences (Fig. 2) was performed using PRODIGAL ver. 2.6.2 software [8] and compared with the selected protein databases (SwissProt, KEGG, SEED, EggNOG).

Functional annotation revealed the presence of several multidrug resistance efflux pumps and tetracycline resistance genes. Putative bacteriocin-encoding gene clusters were determined *in silico* using the BAGEL4 online tool (<http://bagel4.molgenrug.nl/>) which revealed two areas of interest in contig 53CPB201_2, corresponding to the core bacteriocin, transport, and immunity genes (Fig. 3). One open reading frame encoding the core peptide of lactococcin A (E value, 4.37×10^{-49}) was also detected. The amino acid sequence of the core peptide, deduced from the putative bacteriocin gene, showed a single peptide, non-pediocin-like bacteriocin, thereby indicating that it could be classified as a class IIId bacteriocin [9]. Lactococcin A was initially isolated from *Lactococcus lactis* subsp. *cremoris* LMG 2130, in which it was located on a plasmid, and showed a narrow spectrum of activity against lactococci [10]. Additionally, *LanT* and *HylD*, encoding a bacteriocin ATP binding cassette (ABC) transporter and bacteriocin binding protein, respectively, were detected downstream of the core peptide gene, and they conceivably play functional roles in membrane-associated immunity. Moreover, a putative 63-amino acid-long bacteriocin, which was not identified using BAGEL4, was detected using RAST, with the following amino acid sequence: MENLNLNQLENINGGSTLGWATATATQLGLAAAATAEVPPLSAGLAAAGAV LAGGVAVYEAWH. A BLASTP search revealed identical protein sequences associated with *C. perfringens*, indicating that the putative bacteriocin gene could be novel. Genomic analyses provide evidence on bacteriocin production by *C. perfringens* B20 and could potentially enhance its pathogenicity by eliminating the *Lactococcus* community and other sensitive microorganisms in the chicken gut, thereby making disease control more challenging.

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The sequence obtained in this Whole Genome Shotgun project has been deposited in DDBJ/ENA/GenBank under the accession number JAGKIL000000000. The version described in this paper is version JAGKIL010000000.

REFERENCES

1. Cuevas-González PF, Liceaga AM, Aguilar-Toalá JE. Postbiotics and paraprobiotics:

- from concepts to applications. *Food Res Int.* 2020;136:109502. <https://doi.org/10.1016/j.foodres.2020.109502>
2. Moradi M, Kousheh SA, Almasi H, Alizadeh A, Guimarães JT, Yılmaz N, et al. Postbiotics produced by lactic acid bacteria: the next frontier in food safety. *Compr Rev Food Sci Food Saf.* 2020;19:3390-415. <https://doi.org/10.1111/1541-4337.12613>
 3. Villagrán-de la Mora Z, Macías-Rodríguez ME, Arratia-Quijada J, Gonzalez-Torres YS, Nuño K, Villarruel-López A. *Clostridium perfringens* as foodborne pathogen in broiler production: pathophysiology and potential strategies for controlling necrotic enteritis. *Animals.* 2020;10:1718. <https://doi.org/10.3390/ani10091718>
 4. Kiu R, Caim S, Alexander S, Pachori P, Hall LJ. Probing genomic aspects of the multi-host pathogen *Clostridium perfringens* reveals significant pangenome diversity, and a diverse array of virulence factors. *Front Microbiol.* 2017;8:2485. <https://doi.org/10.3389/fmicb.2017.02485>
 5. Heo S, Kim MG, Kwon M, Lee HS, Kim GB. Inhibition of *Clostridium perfringens* using bacteriophages and bacteriocin producing strains. *Korean J Food Sci Anim Resour.* 2018;38:88-98. <https://doi.org/10.5851/kosfa.2018.38.1.88>
 6. Chan PP, Lowe TM. tRNAscan-SE: searching for tRNA genes in genomic sequences. *Methods Mol Biol.* 2019;1962:1-14. https://doi.org/10.1007/978-1-4939-9173-0_1
 7. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics.* 2013;29:2933-5. <https://doi.org/10.1093/bioinformatics/btt509>
 8. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics.* 2010;11:119. <https://doi.org/10.1186/1471-2105-11-119>
 9. Acedo JZ, Chiorean S, Vederas JC, van Belkum MJ. The expanding structural variety among bacteriocins from Gram-positive bacteria. *FEMS Microbiol Rev.* 2018;42:805-28. <https://doi.org/10.1093/femsre/fuy033>
 10. Holo H, Nilssen O, Nes IF. Lactococcin A, a new bacteriocin from *Lactococcus lactis* subsp. *cremoris*: isolation and characterization of the protein and its gene. *J Bacteriol.* 1991;173:3879-87. <https://doi.org/10.1128/jb.173.12.3879-3887.1991>