

Brachial Plexus Injuries in Adults with Traumatic Brain Injury : A Retrospective Study

Nihal Tezel,¹ Aslı Can,¹ Damla Cankurtaran,¹ Ece Ünlü Akyüz,¹ Aytül Çakıcı²

Department of Physical Medicine and Rehabilitation,¹ University of Health Sciences Diskapi Yıldırım Beyazıt Training and Research Hospital, Ankara, Turkey

Department of Physical Medicine and Rehabilitation,² Losante Hospital, Ankara, Turkey

Objective : We aimed to investigate the presence of brachial plexus injury (BPI) in traumatic brain injury (TBI) patients and to draw attention to BPI, which can be overlooked by physicians in TBI patients.

Methods : The study was designed retrospectively by examining the files of 58 patients with moderate to severe TBI to investigate coexistence of TBI and BPI.

Results : BPI was detected in six of 58 TBI patients (10.3%). BPI was detected after an average 116 days from the initial injury. Three patients had lower trunk BPI and three patients had panplexopathy.

Conclusion : Diagnosis of BPI in patients with TBI is delayed in the acute period of injury. The clinicians should keep in mind that BPIs may occur and remain undiagnosed in patients with TBI.

Key Words : Brachial plexus · Brain injuries, Traumatic · Electromyography.

INTRODUCTION

Traumatic brain injury (TBI) is a brain dysfunction caused by an external source⁸⁾. Trauma to the brain can lead to a wide variety of long-term pathological changes. Although TBI affects all ages, it is mostly seen in children, adolescents, and over 75 years of age¹²⁾.

Both the central nervous system (CNS) and the peripheral nervous system may be deteriorated as a result of TBI³⁾. As muscle weakness and sensory deficits are often attributed to the CNS injuries, peripheral nerve injuries (PNI) may be

missed in patients with TBI¹³⁾. Life-threatening injuries need to be treated in the acute phase and a detailed orthopedic examination may be lacking sometimes. These factors may delay the diagnosis of concomitant PNIs such as brachial plexus injuries (BPIs) in patients with TBI.

BPIs are PNIs that may lead to motor and sensory deficits, pain, functional limitation, and impaired quality of life⁴⁾. They are most commonly caused by direct trauma, especially motor vehicle accidents. It has been known that BPIs also may accompany TBI⁴⁾. However, in the literature, there are only a few studies to evaluate BPI in patients with TBI¹⁴⁾. The aims of

• Received : June 27, 2020 • Revised : July 27, 2020 • Accepted : August 29, 2020

• Address for reprints : **Nihal Tezel**

Department of Physical Medicine and Rehabilitation, University of Health Sciences Diskapi Yıldırım Beyazıt Training and Research Hospital, Şehit Ömer Halisdemir Street, No:1, Altındag, Ankara 06110, Turkey

Tel : +90 0312 596 20 00, Fax : +90 0312 318 66 90, E-mail : nihaltezel@gmail.com, ORCID : <https://orcid.org/0000-0002-7623-7827>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

the study were to investigate the presence of BPI in TBI patients and to draw attention to BPI, which can be overlooked by physicians in TBI patients.

MATERIALS AND METHODS

The study protocol was approved by the local Ethics Committee (approve date : March 2, 2020, approve number : 83/08). The study was conducted in accordance with the principles of the Declaration of Helsinki. The study was designed retrospectively by examining the files of 58 patients with moderate to severe TBI. All patients received inpatient treatment in a physical medicine and rehabilitation clinic between

Table 1. Demographic features and disease characteristics of the TBI patients

Characteristic	TBI patients (n=58)
Age (years)	32.84±14.07
Gender	
Female	14 (24.1)
Male	44 (75.9)
Educational status	
Illiterate	0 (0.0)
Literate	0 (0.0)
Primary school	25 (43.1)
Secondary school	18 (31.0)
High school	10 (17.3)
University or higher	5 (8.6)
Tracheostomy	33 (56.9)
Gastrostomy	35 (60.3)
Mechanical ventilation	0 (0.0)
Speech disorder	44 (75.9)
Bladder dysfunction	45 (77.6)
Bowel dysfunction	44 (75.9)
DRS (0–29)	15.48±7.87
FIM	
Motor	24.81±16.72
Cognitive	15.60±10.57
Total score	41.20±25.72

Values are presented as mean±standard deviation or number (%). TBI : traumatic brain injury, DRS : disability rating scale, FIM : functional independent measurement

Table 2. Demographic features and disease characteristics of the TBI patients with brachial plexopathy

Characteristic	TBI patients with BPI (n=6)
Age (years)	27.6±10.2
Gender	
Female	1
Male	5
Educational status	
Illiterate	0
Literate	1
Primary school	1
Secondary school	2
High school	2
University or higher	
Hand dominancy	
Right	6
Left	
Tracheostomy	2
Gastrostomy	1
Mechanical ventilation	0
Speech disorder	4
Bladder dysfunction	1
Bowel dysfunction	1
DRS (0–29)	13.59±4.48
FIM	
Motor	31±18
Cognitive	29.3±9.8
Total score	
BPI side	
Right	1
Left	5
BPI detection time after injury (days)	116±3
Types of BPI	
Lower trunk injury	3
Panplexopathy	3
Etiology of Injury	
Traffic accident	4
Work accident	1
Gunshot injury	1
Mechanism of Injury	
Traction	4
Penetration	1
Avulsion	1
Proximal humerus fracture	3
Proximal humerus and clavícula fracture	1

Values are presented as mean±standard deviation or number. TBI : traumatic brain injury, BPI : brachial plexus injury, DRS : disability rating scale, FIM : functional independent measurement

January 2017 and December 2019.

The patients aged between 18 and 60 and diagnosed with BPI according to electroneuromyographic findings were included in the study. In our rehabilitation clinic, BPI is suspect-

ed in patients with upper extremity atrophy, hyporeflexia/areflexia, flaccid muscle tone, do not have expected motor recovery, and shoulder girdle fracture. To confirm the diagnosis, electroneuromyography (ENMG) is performed. Patients

Table 3. MRC* grades of affected muscles of patients with BPI

	No. 1 patient	No. 2 patient	No. 3 patient	No. 4 patient	No. 5 patient	No. 6 patient
Brachial plexus lesion	Panplexopathy	Panplexopathy	Panplexopathy	Lower trunk injury (severe)	Lower trunk injury (severe)	Lower trunk injury (severe)
Time of evaluation (days)	92	104	110	90	134	168
Deltoid	1	0	5/5	5/5	4/5	3/5
Supraspinatus	1	0	3/5	5/5	3/5	3/5
Pectoralis	1	0	5/5	5/5	4/5	3/5
Biceps	0	0	4/5	5/5	3/5	3/5
Triceps	0	0	4/5	5/5	4/5	3/5
ECRL-ECRB	0	0	4/5	5/5	3/5	3/5
FDS-FDP	4	0	4/5	0/5	1/5	3/5

*0 : no contraction, 1 : flicker of contraction, 2 : active movement, with gravity eliminated, 3 : active movement against gravity, 4 : active movement against gravity and resistance, 5 : muscle contracts normally against full resistance. MRC : Medical Research Council, BPI : brachial plexus injury, ECRL : extensor carpi radialis longus, ECRB : extensor carpi radialis brevis, FDS : flexor digitorum superficialis, FDP : flexor digitorum profundus

Table 4. Electrophysiologic findings of the patients

Patient	Localisation, root/trunk	SNAP	CMAP	Needle EMG
No. 1 patient	C5-6/upper	A	A	Reduced activity/SA : +
	C7/middle	A	A	Reduced activity/SA : +
	C8-T1/lower	A	A	Reduced activity/SA : +
No. 2 patient	C5-6/upper	A	A	No activity/SA : none
	C7/middle	A	A	No activity/SA : none
	C8-T1/lower	A	A	No activity/SA : +
No. 3 patient	C5-6/upper	A	D	Reduced activity/SA : none
	C7/middle	A	D	Reduced activity/SA : none
	C8-T1/lower	A	D	Reduced activity/SA : +
No. 4 patient	C5-6/upper	N	N	N
	C7/middle	N	N	N
	C8-T1/lower	A	D	No activity/SA : +
No. 5 patient	C5-6/upper	N	N	N
	C7/middle	N	N	N
	C8-T1/lower	A	A	No activity/SA : +
No. 6 patient	C5-6/upper	N	N	N
	C7/middle	N	N	N
	C8-T1/lower	A	D	Reduced activity/SA : +

SNAP : sensory nerve action potential, CMAP : compound motor action potential, EMG : electromyography, A : absent, SA : spontaneous activity, D : diminished, N : normal

with a history of upper extremity PNI were excluded from the study. Demographic, clinical, and electromyographic data of the patients with TBI and BPI were recorded.

Statistics

Descriptive statistics were performed for demographic and clinical characteristics of the patients. Continuous variables were presented as mean±standard deviation. Categorical variables were presented as frequencies. Statistical analysis was performed using the PASW version 17.0 software (SPSS Inc, Chicago, IL, USA).

RESULTS

Fifty-eight patients with TBI were screened for the presence of BPI. The demographic and clinic characteristics of all patients with TBI were summarized in Table 1.

Brachial plexus injury was detected in six of 58 TBI patients

(10.3%). Five patients were male and one patient was female. All patients were right-handed. BPI was on the right side in one patient and on the left side in five patients. BPI was detected after an average 116 days from the initial injury. Three patients had lower trunk brachial plexus injury and three patients had panplexopathy.

The etiology of injury was a traffic accidents in four patients, work-related accident in one patient, and gunshot injury in one patient. The mechanism of injury was traction in four patients, avulsion in one patient, and laceration in one patient. Four patients had proximal humerus fracture and one patient had proximal humerus and clavicle fractures in affected upper extremity. The demographic and clinic characteristics of TBI patients with BPI were summarized in Table 2. ENMG findings and Medial Research Council grades of affected muscles of patients with BPI has been shown in Tables 3 and 4.




Fig. 1. The flow-sheet for diagnosis fo BPI and other PNIs of upper extremity in TBI patients. TBI : traumatic brain injury, BPI : brachial plexus injury, PNIs : peripheral nerve injuries, ENMG : electroneuromyography.

DISCUSSION

TBI is a serious, well recognized public health problem around the worldwide. It is an important cause of mortality and morbidity⁷⁾. Aside from associated CNS trauma, there may be injuries to peripheral nerves. Concomitant PNIs such as BPIs contribute to functional disability and can lead to unemployment and economic burden in patients with TBI.

In a previous study, Stone and Keenan¹⁴⁾ detected PNI in 34% of 50 TBI patients. Ten percent of these patients had BPI. They also reported that none of the BPI patients was detected at the time of TBI¹⁴⁾. Ciaramitaro et al.²⁾ investigated the epidemiological data of traumatic nerve injuries. They found that the concomitance of traumatic nerve lesion and TBI was approximately 15%. They also reported that the diagnosis may delay due to changes in consciousness²⁾. Similarly, in the present study, 10% of patients with TBI had a concomitant BPI. The diagnosis of BPI was made after average of 116 days from the head trauma. According to these results, it is possible to obtain an idea that BPI is commonly accompanied to TBI, and recognition of BPI is often overlooked at the time of initial injury.

Traumatic BPIs are most commonly secondary to high-energy trauma, especially motor vehicle accidents and, less commonly, secondary to falls, bullet injuries, and penetrating injuries with a sharp object. Suggested mechanisms of the injury are traction, avulsion, laceration, compression, and contusion^{6,11)}. In this study, the major etiology was motor vehicle accidents. The other causes were bullet injury and work-related accidents. The most common injury mechanism was traction and the others were avulsion and laceration. Similar to our results, Stone and Keenan¹⁴⁾ observed that major etiology and mechanism were traction-induced motor vehicle accidents in patients with TBI and concomitant BPI.

The association between proximal humerus or clavicle fractures and BPI has been previously reported^{5,11)}. Fractures of proximal humerus or clavicle may lead to the traction of brachial plexus. In addition, brachial plexus may be compressed by fractures of these bones¹¹⁾. In the present study, three BPI patients had proximal humerus fracture, and one patient had proximal humerus and clavicle fractures together. In contrast to our results, Stone and Keenan¹⁴⁾ did not observe any upper extremity fracture in their TBI patients with concomitant BPI.

It seems that the prevalence of BPI in TBI patients is higher than previously suspected. It is often overlooked by clinicians because motor deficits are often attributed to the CNS injury. In addition, unstable medical condition, impaired cognitive functioning, or inadequately treated proximal humerus and/or clavicle fractures may lead to unrecognized BPI. Even though the patients do not report any complaints, the clinician should keep in mind that BPI can be seen concomitant in TBI patients, especially those with the proximal humerus or clavicle fractures^{1,9,10)}. Detailed neurological examination is crucial for the diagnosis of BPI and should be performed periodically after stabilization of medical condition. Patients with TBI typically present with upper motor neuron symptoms of spasticity, weakness, clonus, pathologic reflexes, and hyperreflexia in the affected extremities. In the presence of BPI, lower motor neuron symptoms such as flaccid muscle tone, areflexia/hyporeflexia, and muscle atrophy are detected. In addition, patients who do not have expected motor recovery in the affected extremity should be assessed for BPI. ENMG should be performed in TBI patients with clinical suspicion of BPI. ENMG is the most important diagnostic tool for assessing BPI. It gives not only information about the diagnosis, but also about the localization, severity, and prognosis of BPI. The flow-sheet for diagnosis of BPI and other PNIs of the upper extremity in TBI patients has been shown in Fig. 1.

There were some limitations in our study. First, the number of patients analyzed was limited. Second, this study was a single-center study. Future studies with a larger sample size are needed to confirm the results of this study.

CONCLUSION

The diagnosis of BPI in patients with TBI is delayed in the acute period of injury because of unstable medical condition, impaired cognitive functioning or inadequately treated shoulder girdle fractures. The clinicians should keep in mind that BPIs may occur and remain undiagnosed in patients with TBI. Detailed neurological examination should be performed in all TBI patients after stabilization of medical condition and in patients with clinical suspicion of BPI, ENMG should be considered.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

INFORMED CONSENT

This type of study does not require informed consent.

AUTHOR CONTRIBUTIONS

Conceptualization : NT, AC

Data curation : NT, AC, AÇ

Formal analysis : DC

Methodology : EÜA

Project administration : NT, AC

Visualization : NT, AC

Writing - original draft : NT, AC

Writing - review & editing : EÜA, AÇ

ORCID

Nihal Tezel <https://orcid.org/0000-0002-7623-7827>

Aslı Can <https://orcid.org/0000-0002-4189-7323>

Damla Cankurtaran <https://orcid.org/0000-0002-6208-3345>

Ece Ünlü Akyüz <https://orcid.org/0000-0003-4718-5981>

Aytül Çakıcı <https://orcid.org/0000-0002-2121-9330>

References

1. Brien WW, Gellman H, Becker V, Garland DE, Waters RL, Wiss DA : Management of fractures of the humerus in patients who have an injury of the ipsilateral brachial plexus. **J Bone Joint Surg Am** 72 : 1208-1210, 1990
2. Ciaramitaro P, Mondelli M, Logullo F, Grimaldi S, Battiston B, Sard A, et al. : Traumatic peripheral nerve injuries: epidemiological findings, neuropathic pain and quality of life in 158 patients. **J Peripher Nerv Syst** 15 : 120-127, 2010
3. Dang B, Chen W, He W, Chen G : Rehabilitation treatment and progress of traumatic brain injury dysfunction. **Neural Plast** 2017 : 1582182, 2017
4. de Moraes FB, Kwae MY, da Silva RP, Porto CC, Magalhães Dde P, Paulino MV : Clinical aspects of patients with traumatic lesions of the brachial plexus following surgical treatment. **Rev Bras Ortop** 50 : 556-561, 2015
5. Garland DE, Keenan MA : Orthopedic strategies in the management of the adult head-injured patient. **Phys Ther** 63 : 2004-2009, 1983
6. Kim DH, Murovic JA, Kline DG : Brachial plexus injury: mechanisms, surgical treatment and outcomes. **Korean Neurosurg Soc** 36 : 177-185, 2004
7. Mansukhani KA : Electrodiagnosis in traumatic brachial plexus injury. **Ann Indian Acad Neurol** 16 : 19-25, 2013
8. Menon DK, Schwab K, Wright DW, Maas AI; Demographics and Clinical Assessment Working Group of the International and Interagency Initiative toward Common Data Elements for Research on Traumatic Brain Injury and Psychological Health : Position statement: definition of traumatic brain injury. **Arch Phys Med Rehabil** 91 : 1637-1640, 2010
9. Midha R : Epidemiology of brachial plexus injuries in a multitrauma population. **Neurosurgery** 40 : 1182-1189, 1997
10. Noble J, Munro CA, Prasad VS, Midha R : Analysis of upper and lower extremity peripheral nerve injuries in a population of patients with multiple injuries. **J Trauma** 45 : 116-122, 1998
11. Park HR, Lee GS, Kim IS, Chang JC : Brachial plexus injury in adults. **The Nerve** 3 : 1-11, 2017
12. Robbins W, Cifu D : Geriatric Traumatic Brain Injury in Eapen BC, Cifu DX (eds) : **Rehabilitation After Traumatic Brain Injury**, ed 1. St. Louis : Elsevier, 2018, pp109-114
13. Robinson LR : Traumatic injury to peripheral nerves. **Muscle Nerve** 23 : 863-873, 2000
14. Stone L, Keenan MA : Peripheral nerve injuries in the adult with traumatic brain injury. **Clin Orthop Relat Res** (233) : 136-144, 1988