



## SHORT COMMUNICATION

# A Preliminary Study: Antibiotic Resistance of *Escherichia coli* and *Staphylococcus aureus* from the Meat and Feces of Various South African Wildlife Species

Michaela Sannettha van den Honert<sup>1,2\*</sup>, Pieter Andries Gouws<sup>1</sup>, and Louwrens Christiaan Hoffman<sup>2,3</sup>

## OPEN ACCESS

Received June 22, 2020

Revised July 24, 2020

Accepted July 30, 2020

### \*Corresponding author :

Michaela Sannettha van den Honert  
Centre for Food Safety, Department of Food  
Science, University of Stellenbosch, Private  
Bag X1, Matieland 7602, South Africa  
Tel: +27-21-808-2609  
E-mail: [michaelavdh@sun.ac.za](mailto:michaelavdh@sun.ac.za)

### \*ORCID

Michaela Sannettha van den Honert  
<https://orcid.org/0000-0002-1886-8516>  
Pieter Andries Gouws  
<https://orcid.org/0000-0002-7061-2403>  
Louwrens Christiaan Hoffman  
<https://orcid.org/0000-0003-2736-1933>

<sup>1</sup>Centre for Food Safety, Department of Food Science, University of Stellenbosch, Private Bag X1, Matieland 7602, South Africa

<sup>2</sup>Department of Animal Sciences, University of Stellenbosch, Private Bag X1, Matieland 7602, South Africa

<sup>3</sup>Centre for Nutrition and Food Sciences, Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Coopers Plains, QLD 4108, Australia

**Abstract** This study determined the antibiotic resistance patterns of *Escherichia coli* and *Staphylococcus aureus* from the raw meat and feces of three game species from three different farms across South Africa. The Kirby-Bauer disk diffusion method was used according to the Clinical and Laboratory Standards Institute 2018 guidelines. *E. coli* was tested against ampicillin, ceftazidime, chloramphenicol, streptomycin, sulphafurazole and tetracycline. *S. aureus* was tested against tetracycline, erythromycin, vancomycin, penicillin, oxacillin and ceftazidime. There were no significant differences in the *E. coli* antibiotic resistance profiles between the meat and fecal samples (except towards ceftazidime where 5% of the meat isolates were resistant and 0% of the fecal isolates). The *S. aureus* meat isolates showed high (75%) resistance towards penicillin and on average, 13% were resistant to oxacillin/ceftazidime, indicating methicillin resistance. The results from this study indicate that there is incidence of antibiotic resistant bacteria from the feces and meat of wildlife species across South Africa, suggesting that cross contamination of the meat occurred during slaughter by antibiotic resistant bacteria from the abattoir personnel or equipment and or from carcass fecal matter. In addition, the results highlight the importance of food safety and hygiene procedures during slaughter to prevent cross-contamination of antibiotic resistant bacteria, as well as pathogens, onto raw meat.

**Keywords** antimicrobial resistance, game, bacteria, pathogen

## Introduction

Various studies have demonstrated that wild animals and their surrounding environments are significant reservoirs of antibiotic resistance genes and antibiotic

resistant bacteria (Cantas et al., 2013; Costa et al., 2008; Karesh et al., 2012). This is a growing public health issue, due to increased wildlife contact with humans, as well as livestock and domestic animals due to co-habitation on the same farm (van den Honert et al., 2018). In addition, there is a rising trend in the consumption of game meat over recent years (Cantas et al., 2013; Dias et al., 2015). Majority of these wild animals are slaughtered in field abattoirs (van Shalkwyk and Hoffman, 2010). This can potentially increase the risk that bacteria from the gut and skin of animals can cross-contaminate meat during the slaughter process, typically due to inadequate hygiene conditions and handling (Bakhtiary et al., 2016; van Shalkwyk and Hoffman, 2010). Schlegelová et al. (2004) has stated that bacteria which contaminate meat can be the source of foodborne diseases and also possibly a cause of drug resistant human pathogenic bacteria (Schlegelová et al., 2004).

Although studies are limited, especially on bacteria from wildlife, several studies have been conducted on the antibiotic resistance patterns within a meat processing facility and conclude that cross-contamination of antibiotic resistant bacteria onto the meat does occur and originates from various sources such as the animal feces, staff hands or facility equipment and machinery (Amir et al., 2017; Aslam et al., 2003; Schlegelová et al., 2004).

This study evaluated the antibiotic resistance patterns of *Escherichia coli* and *Staphylococcus aureus* isolated from the meat and feces of three commonly hunted/harvested game species from different regions in South Africa.

## Methods

### Ethics number

All animals were hunted and sampled according to the standard operating procedure approved by the Stellenbosch University Animal Care and Use Committee (ethics number: SU-ACUM14-001SOP).

### Study area and sample collection

All three farms are private game farms where the wildlife are free to roam with the other wildlife species on the farm. The wildlife graze and drink on the farm's natural resources and are given supplementary feed when necessary, especially in times of drought when the grass has become depleted.

Meat (the *infraspinatus* muscle) and fecal (from the ileum of the small intestine) samples were collected from the same animal in order to directly compare the antibiotic resistance profiles of the meat and feces of each animal. All samples from the same farm were collected on the same day. After sample collection, all samples were stored frozen at  $-20^{\circ}\text{C}$  until analysis commenced.

Impala (*Aepyceros melampus*) meat (n=5) and fecal samples (n=5) were collected from five different impala from a farm in Modimolle, Limpopo, South Africa. Bontebok (*Damaliscus pygargus*) meat (n=5) and fecal samples (n=5) were collected from five different bontebok from a farm in Wellington, Western Cape, South Africa. Springbok (*Antidorcas marsupialis*) meat (n=5) and fecal samples (n=5) were collected from five different springbok from a farm in Witsand, Western Cape, South Africa.

### Isolation and species identification of bacteria

Samples were defrosted at room temperature until thawed before analysis commenced. The bacteria were isolated from the samples using a series of plating on selective agar media. The inoculated agar plates were incubated overnight at  $35^{\circ}\text{C}$ . For *E. coli*, Violet Red Bile Dextrose agar (Merck Bioloab, Modderfontein, South Africa) and then Eosin Methylene Blue agar (Oxoid, Hampshire, UK) was used to isolate the bacteria from the fecal and meat samples. Baird Parker agar (Oxoid,

Hampshire, UK) was used to isolate *S. aureus* from the meat samples. Characteristic growth of *E. coli* on Violet Red Bile Dextrose agar is purple/red colonies surrounded by a halo and characteristic growth on Eosin Methylene Blue agar is colonies with a dark purple center with a green metallic sheen. *S. aureus* characteristic growth on Baird Parker agar is black colonies with a clear halo.

Gram's stain and various biochemical tests, the citrate test, catalase test and Staphylase test (Oxoid, Hampshire, UK) were used to confirm colony identity.

After pure cultures were obtained, colonies from the selective agar plates were streaked onto Nutrient agar (Merck Biolab) plates and incubated overnight at 35°C, which were then used for the antibiotic susceptibility test (AST).

### Antibiotic susceptibility testing

Antibiotic susceptibility of all samples were tested in triplicate as each animal was sub-sampled three times in the isolation step. Fresh cultures grown overnight on Nutrient agar (Merck Biolab) were used for the antibiotic susceptibility analysis. The disk diffusion method was used according to the Clinical & Laboratory Standards Institute (CLSI, 2018) 2018 guidelines (M100S) using Mueller-Hinton agar (Merck Biolab). For *E. coli*, the antibiotic discs (Oxoid, Johannesburg, South Africa) ampicillin 10 µg, chloramphenicol 30 µg, ceftazidime 30 µg, streptomycin 10 µg, sulphafurazole 300 µg and tetracycline 30 µg were tested. For *S. aureus*, the antibiotic discs (Oxoid, Johannesburg, South Africa) cefoxitin 30 µg, erythromycin 15 µg, oxacillin 1 µg, penicillin 10 U, tetracycline 30 µg and vancomycin 30 µg were tested. *E. coli* ATCC 25922 and *S. aureus* ATCC 25923 (Thermo Fisher Scientific, Lake Charles, LA, USA) were used as quality controls and an uninoculated agar plate was used as a negative control. The results of the disc diffusion test classified the isolates as resistant, intermediately resistant or susceptible to the selection of antibiotics, according to the zone diameter specifications as listed by the CLSI 2018 guidelines.

### Antibiotic resistant gene detection

A crude extraction method using lysis buffer and boiling was used to extract DNA from fresh overnight broth cultures of the isolated *E. coli* meat and fecal samples. The ZymoBiomics DNA kit (Inqaba Biotec, Muckleneuk, South Africa) was used according to the manufacturer's instructions to extract DNA from fresh overnight broth cultures of the isolated *S. aureus* meat samples.

Polymerase chain reaction (PCR) was used to detect various antibiotic resistant genes which are commonly associated with phenotypic antibiotic resistance. The antibiotic resistant genes detected in the *E. coli* isolates were: *tetA* at 502 bp and *tetB* at 930 bp (tetracycline resistance), *sul1* at 433 bp and *sul2* at 721 bp (sulphonamide resistance), *bla*CMY at 1,000 bp (ampicillin resistance) and *aadA* at 525 bp (streptomycin resistance). The antibiotic resistant genes detected in the *S. aureus* isolates were: *tetK* at 1,515 bp, *tetL* at 229 bp and *tetM* at 406 bp (tetracycline resistance), *vanA* at 732 bp and *vanB* at 647 bp (vancomycin resistance) and *blaZ* at 498 bp (penicillin resistance). All reactions were performed in duplicate. The primers and reaction conditions used in this study are described by van den Honert et al. (2020), except the *blaZ* gene, which is the most common gene encoding production of beta-lactamases to hydrolyse penicillin. The *blaZ* gene was detected using the following primers and reaction conditions: forward primer sequence: 5'-AAGAGATTTGCCTATGCTTC-3', reverse primer sequence: 5'-GCTTGACCACTTTTATCAGC-3'; 5 min initial denaturation at 94°C followed by 35 cycles of 94°C for 30s, 55°C for 30s and 72°C for 10 min and a final extension step of 72°C for 10 min (Russi et al., 2015).

Gel electrophoresis was performed using 1.2% agarose gel (Lonza SeaKem, Rockland, ME, USA) stained with EZ-Vision®

in-gel solution DNA dye (Amresco, Solon, OH, USA). Gel visualisation was performed using the Bio-Rad Gel Doc XR+ System (Bio-Rad, Hercules, CA, USA) in combination with Image Lab Software V5.2.1.

### Statistical analysis


Statistical analysis was performed on the *E. coli* isolate results to determine if there were any significant differences between the meat and faecal sample antibiotic resistant levels. The statistical analysis was performed using Statistica 13.2 software (Tibco, Palo Alto, CA, USA). The data was analysed using one-way analysis of variance (ANOVA). The main effect was meat versus faecal sample. Significant results were identified by least significant means (LSM) by using a 95% confidence interval ( $p \leq 0.05$ ).

## Results and Discussion


Antibiotic resistant bacteria and some of the associated antibiotic resistance genes were found in both the faecal and meat samples of the wildlife species from all three of the farms (Figs. 1 and 2). Table 1 shows the phenotype-genotype correlations of the *E. coli* and *S. aureus* isolates. Antibiotic resistant genes were detected in all the isolates which were classified as phenotypically resistant to the various antibiotics.

*Escherichia coli* was isolated from 22 (49%) and 45 (100%) of the meat and faecal samples, respectively. A summary of the AST results for the *E. coli* isolates is shown in Fig. 1. There were no significant differences in the *E. coli* antibiotic resistance patterns between the meat and faecal samples, except for ceftazidime. Thus it can be speculated that contamination of the game meat occurred from *E. coli* both from the carcass feces and from the surrounding environment, equipment and/ or the slaughter personnel (Gouws et al., 2017).

The presence of *E. coli* (49%) on the meat samples indicates that faecal contamination from the carcass occurred on the meat of these animals during the slaughter process (Aslam et al., 2003). The presence of *S. aureus* on the meat samples (36%) indicates that unhygienic practices and thus cross-contamination occurred onto the meat from the hide and/ or from the meat handlers. Ultimately, the presence of *S. aureus* on raw meat is indicative of poor hygiene conditions in the food chain, mainly



**Fig. 1.** Antibiotic susceptibility profiles of *E. coli* isolates from meat (n=22) and faecal samples (n=45) from springbok, bontebok and impala to ampicillin (AMP)  $p > 0.05$ , ceftazidime (CAZ)  $p \leq 0.05$ , chloramphenicol (C)  $p > 0.05$ , streptomycin (ST)  $p > 0.05$ , sulphafurazole (SF)  $p > 0.05$  and tetracycline (TE)  $p > 0.05$ .



**Fig. 2.** Antibiotic susceptibility profile of *S. aureus* isolates (n=16) from meat samples from springbok and bontebok to tetracycline (TE), erythromycin (E), vancomycin (VA), penicillin (P) and oxacillin (OX) / ceftiofur (FOX).

**Table 1.** Phenotype-genotype antibiotic resistance patterns of *E. coli* and *S. aureus*

Location	Animal	Phenotypic resistance <sup>1)</sup>	Genotypic resistance					
			<i>bla</i> CMY	<i>sul1</i>	<i>sul2</i>	<i>aadA</i>	<i>tetA</i>	<i>tetB</i>
<i>Escherichia coli</i>								
Witsand	Springbok	AMP(S), SF(R), ST(R), TE(R)	–	+	+	+	–	+
Witsand	Springbok	AMP(I), SF(I), ST(I), TE(S)	–	–	–	+	–	–
Witsand	Springbok	AMP(I), SF(I), ST(I), TE(S)	–	–	–	+	–	–
Witsand	Springbok <sup>2)</sup>	AMP(S), SF(S), ST(R), TE(S)	+	–	–	+	–	–
Witsand	Springbok <sup>2)</sup>	AMP(S), SF(S), ST(I), TE(S)	+	–	–	+	–	–
Modimolle	Impala	AMP(I), SF(I), ST(R), TE(S)	–	–	–	+	–	–
Modimolle	Impala	AMP(S), SF(S), ST(I), TE(S)	+	–	–	+	–	–
Modimolle	Impala <sup>2)</sup>	AMP(R), SF(S), ST(S), TE(S)	+	–	–	–	–	–
Modimolle	Impala <sup>2)</sup>	AMP(S), SF(S), ST(S), TE(S)	–	–	–	–	–	–
Modimolle	Impala <sup>2)</sup>	AMP(S), SF(S), ST(I), TE(S)	–	–	–	+	–	–
Wellington	Bontebok	AMP(R), SF(I), ST(I), TE(S)	+	–	–	+	–	–
Wellington	Bontebok	AMP(R), SF(S), ST(I), TE(S)	+	–	–	+	–	–
Wellington	Bontebok	AMP(R), SF(S), ST(I), TE(S)	+	–	–	+	–	–
Wellington	Bontebok	AMP(S), SF(S), ST(I), TE(S)	+	–	–	+	–	–
Wellington	Bontebok <sup>2)</sup>	AMP(R), SF(S), ST(S), TE(S)	+	–	–	–	–	–
<i>Staphylococcus aureus</i>								
			<i>tetL</i>	<i>tetK</i>	<i>tetM</i>	<i>vanA</i>	<i>vanB</i>	<i>blaZ</i>
Witsand	Springbok <sup>2)</sup>	TE(S), VA(S), P(R)	+	–	–	+	–	+
Witsand	Springbok <sup>2)</sup>	TE(S), VA(S), P(R)	+	–	–	–	–	+
Witsand	Springbok <sup>2)</sup>	TE(S), VA(S), P(R)	–	–	–	–	–	+
Witsand	Springbok <sup>2)</sup>	TE(S), VA(S), P(R)	–	–	–	–	–	+
Wellington	Bontebok <sup>2)</sup>	TE(S), VA(S), P(R)	–	–	–	–	–	+

<sup>1)</sup> AMP, ampicillin (*bla*CMY gene); SF, sulphonamide (*sul1* and *sul2* genes); ST, streptomycin (*aadA* gene); TE, tetracycline (*tetA* and *tetB* genes); TE, tetracycline (*tetL*, *tetK* and *tetM* genes); VA, vancomycin (*vanA* and *vanB* genes); P, penicillin (*blaZ* gene); S, susceptible; I, intermediate; R, resistant.

<sup>2)</sup> Meat samples (all others are fecal samples).

due to contamination by food handlers and equipment (Naas et al., 2019). The detection of *E. coli* and *S. aureus* from the meat samples was expected and the frequency of isolation is similar to those found by Schlegelová et al. (2004), who detected *E. coli* and *S. aureus* in meat samples of beef in a slaughterhouse in 67% and 24% of samples, respectively.

The *E. coli* isolates were most frequently resistant to streptomycin, followed by ampicillin, sulphafurazole, tetracycline and then ceftazidime. No *E. coli* isolates were resistant to chloramphenicol.

The *E. coli* isolates were resistant to streptomycin (average 9% meat; 5% fecal) where a high percentage were intermediately resistant (average 59% meat; 68% fecal) (Fig. 1). Other studies have reported that streptomycin resistance is common in food animals due to its extensive use in both agricultural and clinical settings (Boerlin et al., 2005; Bryan et al., 2004; Kozak et al., 2009; Wilkerson et al., 2004). In addition, streptomycin is present in the soil, produced by organisms such as *Streptomyces griseus*, and could confer a natural low-level resistance to the grazing wildlife (van Overbeek et al., 2002; Wegst-Uhrich et al., 2014).

Moreover, the fecal and meat *E. coli* isolates were notably resistant to ampicillin (average 18% meat; 20% fecal). More specifically, the high average of ampicillin resistance was mainly attributed to the fecal (100% resistant) and meat (73% resistant) samples from the bontebok species from the Wellington farm (data not shown). The higher ampicillin resistance seen in the bontebok *E. coli* isolates can be attributed to the fact that this farm which hosts the bontebok was previously a dairy and sheep farm about thirty years ago. The penicillin antibiotic class, which includes ampicillin, is the most widely used antibiotic class in sheep farming (Wegst-Uhrich et al., 2014). The application of antibiotics during the dairy and sheep farming period could have stimulated the development of antibiotic resistant bacteria within the soil which could be transferred to the grazing wildlife (Wegst-Uhrich et al., 2014).

Furthermore, the *E. coli* isolates had low levels of resistance towards sulphafurazole (average 9% meat; 2% fecal) and tetracycline (average 9% meat; 4% fecal). This is consistent with Li et al. (2007) study which reported similar resistant levels of *E. coli* isolates from game meat, with sulphafurazole resistance at 7.9% and tetracycline resistance at 13%.

No *E. coli* meat or fecal isolates were resistant to chloramphenicol. Low resistance was expected as chloramphenicol is not permitted for food animal use in South Africa and many other countries (Rawat and Nair, 2010).

The only significant difference in antibiotic resistance levels between the *E. coli* isolates from the meat and fecal samples was towards ceftazidime, where 5% of the meat isolates and 0% of the fecal isolates were classified as resistant. Resistance to ceftazidime indicates suspicion for extended-spectrum  $\beta$ -lactamase (ESBL) production (Overdevest et al., 2011). However, additional phenotypic confirmatory tests would still need to be performed to confirm ESBL-production (Dahms et al., 2015; Henton et al., 2011; Overdevest et al., 2011). Other studies have also speculated that environmental ESBL *E. coli* is a result of human influence, as the majority of ESBLs are reported from human clinical isolates due to the direct use of novel sub-classes of  $\beta$ -lactam antibiotics (Guenther et al., 2011; Skurnik et al., 2006). Thus it can be speculated that contamination of the meat occurred predominantly from human influence, most likely during the skinning and evisceration steps, as also found by Schlegelová et al. (2004), via resistant genotype and phenotype analysis.

*S. aureus* was isolated from 16 of meat samples (36%). A summary of the AST results for the *S. aureus* isolates is shown in Fig. 2. The *S. aureus* isolates were most frequently resistant to penicillin, followed by oxacillin/ cefoxitin, tetracycline and then erythromycin. No isolates were classified as resistant to vancomycin.

There were 12 *S. aureus* isolates from the game meat which were resistant to penicillin (75%) (Fig. 2). This was anticipated, as resistance to penicillin is now widespread in humans and animals since the 1960s, in both community, hospital and meat staphylococcal isolates (Appelbaum, 2007; Chambers and DeLeo, 2009; Lowy, 2003; Schlegelová et al., 2004).

Methicillin resistant *S. aureus* (MRSA) was only detected in two of the bontebok meat isolates (13% average) (indicated by oxacillin and confirmed by cefoxitin). Although genetic confirmation to confirm methicillin resistance should be performed by detection of the *mecA* gene. Other studies have concluded that contamination of meat with MRSA can result from cross contamination of the carcass from the animal itself or from the people involved in the meat handling during slaughter and processing (Gilmore et al., 2008).

The *S. aureus* meat isolates were classified as tetracycline resistant in 2 (12%) of the samples. Furthermore, 1 (6%) of the *S. aureus* isolates from the game meat were classified as resistant to erythromycin, although 5 (31%) were classified as intermediately resistant. Other studies have found varying frequencies of resistance to erythromycin (4.3%–30%) and tetracycline (~50%) of *S. aureus* from meat (retail non-game meat), where tetracycline resistance is generally more common than erythromycin resistance (Kelman et al., 2011).

None of the *S. aureus* meat isolates were classified as resistant to vancomycin. Other studies have also reported negligible to very low levels (0%–3%) of vancomycin resistant *S. aureus* from raw commercial meat samples (Das and Mazumder, 2016; Jackson et al., 2013; Pesavento et al., 2007).

At least one of the selected antibiotic resistant genes were detected in all samples which showed to have a corresponding phenotypic antibiotic resistance pattern. There were seven samples where the antibiotic resistant gene was detected but the phenotypic method classified the isolates as susceptible. This occurred for ampicillin (4) from the *E. coli* isolates and for tetracycline (2) and vancomycin (1) from the *S. aureus* isolates. Some possible explanations for these resistance genes being detected in these samples could be that they are inactive genes, meaning that they are present but are not active because there is no antibiotic resistance selective pressure to phenotypically express the gene. Alternatively, PCR can be considered a more sensitive method to the disc diffusion phenotypic method, as resistance is dependent on the size of the zone of inhibition, which is determined by the CLSI committee on an annual basis, whereas resistance in PCR is determined simply by the detection of a resistance gene (Gilmore et al., 2008).

## Conclusion

Antibiotic resistant bacteria were detected in the feces and on the raw meat of the wildlife species, with ampicillin and streptomycin resistance being the most prevalent in *E. coli* from both sample types. The *S. aureus* isolates from the game meat showed high resistance to penicillin but fairly low resistance to the other five antibiotics.

Although it seems unlikely that antibiotic resistant bacteria would be found in wildlife, movement of antibiotic resistance genes and resistant bacteria can reach these more isolated environments from pollution of human and farm animal environments as well as via supplementary feed and water sources. Contamination via humans during the slaughter and processing steps can also be a source of antibiotic resistant bacteria onto the raw meat.

The *E. coli* isolated from the meat and feces of the same animal showed to have similar antibiotic resistance patterns except towards ceftazidime, where there was a significant difference in resistance frequencies between the meat and fecal samples. These results indicate that cross contamination of the meat occurred from bacteria both from the carcass and from human origin.

In order to prevent cross-contamination of harmful and/ or antibiotic resistant bacteria from the hides or feces onto raw meat, various precautionary steps can be put in place. For example, exposed muscle must avoid contact with workers hands and the animal's skin as best as possible. Workers hands are important sources of contamination during processing and thus



hand washing is essential in preventing contamination of the carcass. Furthermore, it is important to clean all equipment and meat processing machinery to reduce the effect of cross-contamination.

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This research is supported by the South African Research Chairs Initiative (SARChI) and partly funded by the South African Department of Science and Technology (UID number: 84633), as administered by the National Research Foundation (NRF) of South Africa, and partly by the Department of Trade and Industry's THRIP program (THRIP/64/19/04/2017) with Wildlife Ranching South Africa as partner and by Stellenbosch University. Any opinions, findings and conclusion or recommendations expressed in this material are that of the author(s) and the National Research Foundation does not accept any liability in this regard.

## Author Contributions

Conceptualization: van den Honert MS, Gouws PA, Hoffman LC. Data curation: van den Honert MS, Gouws PA, Hoffman LC. Formal analysis: van den Honert MS. Methodology: van den Honert MS, Gouws PA, Hoffman LC. Software: van den Honert MS. Validation: van den Honert MS. Investigation: van den Honert MS. Writing - original draft: van den Honert MS. Writing - review & editing: van den Honert MS, Gouws PA, Hoffman LC.

## Ethics Approval

All animals were sampled according to the standard operating procedure approved by the Stellenbosch University Animal Care and Use Committee (ethics number: SU-ACUM14-001SOP).

## References

- Amir M, Muhammad R, Chang Y-F, Akhtar S, Yoo SH, Sheikh AS, Kashif M. 2017. Impact of unhygienic conditions during slaughtering and processing on spread of antibiotic resistant *Escherichia coli* from poultry. *Microbiol Res* 8:35-40.
- Appelbaum PC. 2007. Microbiology of antibiotic resistance in *Staphylococcus aureus*. *Clin Infect Dis* 45:S165-S170.
- Aslam M, Nattress F, Greer G, Yost C, Gill C, McMullen L. 2003. Origin of contamination and genetic diversity of *Escherichia coli* in beef cattle. *Appl Environ Microbiol* 69:2794-2799.
- Bakhtiary F, Sayevand HR, Remely M, Hippe B, Hosseini H, Haslberger AG. 2016. Evaluation of bacterial contamination sources in meat production line. *J Food Qual* 39:750-756.
- Boerlin P, Travis R, Gyles CL, Reid-Smith R, Janecko N, Lim H, Nicholson V, McEwen SA, Friendship R, Archambault M. 2005. Antimicrobial resistance and virulence genes of *Escherichia coli* isolates from swine in Ontario. *Appl Environ Microbiol* 71:6753-6761.



- Bryan A, Shapir, N, Sadowsky MJ. 2004. Frequency and distribution of tetracycline resistance genes in genetically diverse, nonselected, and nonclinical *Escherichia coli* strains isolated from diverse human and animal sources. *Appl Environ Microbiol* 70:2503-2507.
- Cantas L, Shah SQA, Cavaco LM, Manaia CM, Walsh F, Popowska M, Garelick H, Bürgmann H, Sørsum H. 2013. A brief multi-disciplinary review on antimicrobial resistance in medicine and its linkage to the global environmental microbiota. *Front Microbiol* 4:96.
- Chambers HF, DeLeo FR. 2009. Waves of resistance: *Staphylococcus aureus* in the antibiotic era. *Nat Rev Microbiol* 7:629-641.
- Clinical and Laboratory Standards Institute. 2018. M100 Performance standards for antimicrobial susceptibility testing. 28<sup>th</sup> ed. CLSI, Waune, PA, USA.
- Costa D, Poeta P, Sáenz Y, Vinué L, Coelho AC, Matos M, Rojo-Bezares B, Rodrigues J, Torres C. 2008. Mechanisms of antibiotic resistance in *Escherichia coli* isolates recovered from wild animals. *Microb Drug Resist* 14:71-77.
- Dahms C, Hübner N-O, Kossow A, Mellmann A, Dittmann K, Kramer A. 2015. Occurrence of ESBL-producing *Escherichia coli* in livestock and farm workers in Mecklenburg-Western Pomerania, Germany. *PLOS ONE* 10:e0143326.
- Das P, Mazumder PB. 2016. Prevalence of *Staphylococcus* in raw meat samples in Southern Assam, India. *IOSR J Agric Vet Sci* 9:23-29.
- Dias D, Torres RT, Kronvall G, Fonseca C, Mendo S, Caetano T. 2015. Assessment of antibiotic resistance of *Escherichia coli* isolates and screening of *Salmonella* spp. in wild ungulates from Portugal. *Res Microbiol* 166:584-593.
- Gilmore KS, Gilmore MS, Sahn DF. 2008. Methicillin resistance in *Staphylococcus aureus*. In *Bacterial resistance to antimicrobials*. 2<sup>nd</sup> ed. Wax RG, Lewis K, Salyers A, Taber H (ed). CRC Press, Florida, FL, USA. pp 291-296.
- Gouws PA, Shange N, Hoffman LC. 2017. Microbial quality of springbok (*Antidorcas marsupialis*) meat in relation to harvesting and production process. In *Game meat hygiene: Food safety and security*. Paulsen P, Bauer A, Smulders FJM (ed). Wageningen Academic Publishers, Gelderland, The Netherlands. pp 223-228.
- Guenther S, Ewers C, Wieler LH. 2011. Extended-spectrum beta-lactamases producing *E. coli* in wildlife, yet another form of environmental pollution? *Front Microbiol* 2:246.
- Henton MM, Eagar HA, Swan GE, van Vuuren M. 2011. Antibiotic management and resistance in livestock production. *S Afr Med J* 101:583-586.
- Jackson CR, Davis JA, Barrett JB. 2013. Prevalence and characterization of methicillin-resistant *Staphylococcus aureus* isolates from retail meat and humans in Georgia. *J Clin Microbiol* 51:1199-1207.
- Karesh WB, Loh E, Machalaba C. 2012. Food safety: A view from the wild side. In *Improving food safety through a one health approach*. National Academies Press, Washington, DC, USA. pp 207-211.
- Kelman A, Soong YA, Dupuy N, Shafer D, Richbourg W, Johnson K, Brown T, Kestler E, Li Y, Zheng J, McDermott P, Meng J. 2011. Antimicrobial susceptibility of *Staphylococcus aureus* from retail ground meats. *J Food Prot* 74:1625-1629.
- Kozak GK, Boerlin P, Janecko N, Reid-Smith RJ, Jardine C. 2009. Antimicrobial resistance in *Escherichia coli* isolates from swine and wild small mammals in the proximity of swine farms and in natural environments in Ontario, Canada. *Appl Environ Microbiol* 75:559-566.
- Li Q, Sherwood JS, Loque CM. 2007. Characterization of antimicrobial resistant *Escherichia coli* isolated from processed bison carcasses. *J Appl Microbiol* 103:2361-2369.

- Lowy F. 2003. Antimicrobial resistance: The example of *Staphylococcus aureus*. J Clin Invest 111:1265-1273.
- Naas HT, Edarhoby RA, Garbaj AM, Azwai SM, Abolghait SK, Gammoudi FT, Moawad AA, Barbieri I, Eldaghayes IM. 2019. Occurrence, characterization, and antibiogram of *Staphylococcus aureus* in meat, meat products, and some seafood from Libyan retail markets. Vet World 12:925-931.
- Overdeest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, Heck M, Savelkoul P, Vandenbroucke-Grauls C, van der Zwaluw K, Huijsdens X, Kluytmans J. 2011. Extended-spectrum  $\beta$ -lactamase genes of *Escherichia coli* in chicken meat and humans, the Netherlands. Emerg Infect Dis 17:1216-1222.
- Pesavento G, Ducci B, Comodo N, Lo Nostro A. 2007. Antimicrobial resistance profile of *Staphylococcus aureus* isolated from raw meat: A research for methicillin resistant *Staphylococcus aureus* (MRSA). Food Control 18:196-200.
- Rawat D, Nair D. 2010. Extended-spectrum  $\beta$ -lactamases in gram negative bacteria. J Glob Infect Dis 2:263-274.
- Russi NB, Maito J, Dieser SA, Renna MS, Signorini ML, Camussone C, Neder VE, Pol M, Tirante L, Odierno LM, Calvino LF. 2015. Comparison of phenotypic tests for detecting penicillin G resistance with presence of *blaZ* gene in *Staphylococcus aureus* isolated from bovine intramammary infections. J Dairy Res 82:317-321.
- Schlegelová J, Nápravníková E, Dendis M, Horváth R, Benedik J, Babák V, Klimová E, Navrátilová P, Šustáčková A. 2004. Beef carcass contamination in a slaughterhouse and prevalence of resistance to antimicrobial drugs in isolates of selected microbial species. Meat Sci 66:557-565.
- Skurnik D, Ruimy R, Andremont A, Amarin C, Rouquet P, Picard B, Denamur E. 2006. Effect of human vicinity on antimicrobial resistance and integrons in animal faecal *Escherichia coli*. J Antimicrob Chemother 57:1215-1219.
- van den Honert MS, Gouws PA, Hoffman LC. 2018. Importance and implications of antibiotic resistance development in livestock and wildlife farming in South Africa: A review. S Afr J Anim Sci 48:401-412.
- van den Honert MS, Gouws PA, Hoffman LC. 2020. A preliminary study: Antibiotic resistance patterns of *Escherichia coli* and *Enterococcus* species from wildlife species subjected to supplementary feeding on various South African farms. Animals 10:396.
- van Overbeek LS, Wellington EMH, Egan S, Smalla K, Heuer H, Collard JM, Guillaume G, Karagouni AD, Nikolakopoulou TL, van Elsas JD. 2002. Prevalence of streptomycin-resistance genes in bacterial populations in European habitats. FEMS Microbiol Ecol 42:277-288.
- van Shalkwyk DL, Hoffman LC. 2010. Guidelines for the harvesting of game for meat export. AgriPublishers, Windhoek, Namibia.
- Wegst-Uhrich SR, Navarro DAG, Zimmerman L, Aga DS. 2014. Assessing antibiotics sorption in soil: A literature review and new case studies on sulphonamides and macrolides. Chem Cent J 8:5.
- Wilkerson C, Samadpour M. 2004. Antibiotic resistance and distribution of tetracycline resistance genes in *Escherichia coli* O157:H7 isolates from humans and bovines. Antimicrob Agents Chemother 48:1066-1067.