



T-Cell Death-Associated Gene 51 Is a Novel Negative Regulator of PPAR γ That Inhibits PPAR γ -RXR α Heterodimer Formation in Adipogenesis

Sumi Kim^{1,4}, Nari Lee^{1,4}, Eui-Soon Park¹, Hyeongseok Yun¹, Tae-Uk Ha¹, Hyeon Jeon¹, Jiyeon Yu¹, Seunga Choi¹, Bongjin Shin¹, Jungeun Yu¹, Sang Dal Rhee², Yongwon Choi³, and Jaerang Rho^{1,*}

¹Department of Microbiology and Molecular Biology, Chungnam National University, Daejeon 34134, Korea, ²Therapeutics and Biotechnology Division, Korea Research Institute of Chemical Technology, Daejeon 34114, Korea, ³Department of Pathology and Laboratory Medicine, University of Pennsylvania School of Medicine, Philadelphia, PA 19104, USA, ⁴These authors contributed equally to this work.

*Correspondence: jrrho@cnu.ac.kr

<https://doi.org/10.14348/molcells.2020.0143>

www.molcells.org

The nuclear receptor peroxisome proliferator-activated receptor γ (PPAR γ) is the master transcriptional regulator in adipogenesis. PPAR γ forms a heterodimer with another nuclear receptor, retinoid X receptor (RXR), to form an active transcriptional complex, and their transcriptional activity is tightly regulated by the association with either coactivators or corepressors. In this study, we identified T-cell death-associated gene 51 (TDAG51) as a novel corepressor of PPAR γ -mediated transcriptional regulation. We showed that TDAG51 expression is abundantly maintained in the early stage of adipogenic differentiation. Forced expression of TDAG51 inhibited adipocyte differentiation in 3T3-L1 cells. We found that TDAG51 physically interacts with PPAR γ in a ligand-independent manner. In deletion mutant analyses, large portions of the TDAG51 domains, including the pleckstrin homology-like, glutamine repeat and proline-glutamine repeat domains but not the proline-histidine repeat domain, are involved in the interaction with the region between residues 140 and 506, including the DNA binding domain, hinge, ligand binding domain and activation function-2 domain, in PPAR γ . The heterodimer formation of PPAR γ -RXR α was competitively inhibited in a ligand-independent manner by TDAG51 binding to PPAR γ . Thus,

our data suggest that TDAG51, which could determine adipogenic cell fate, acts as a novel negative regulator of PPAR γ by blocking RXR α recruitment to the PPAR γ -RXR α heterodimer complex in adipogenesis.

Keywords: adipocyte differentiation, adipogenesis, pleckstrin homology-like domain A family, peroxisome proliferator-activated receptor gamma, retinoid X receptor alpha, T-cell death-associated gene 51

INTRODUCTION

Adipogenesis is the process of mature adipocyte formation from adipocyte precursors, and adipocytes play a vital role in lipid metabolism, energy homeostasis and insulin sensitivity (Sarjeant and Stephens, 2012). Adipogenesis is mainly induced by the coordinated interplay among transcriptional factors, such as CCAAT/enhancer binding proteins (C/EBPs) and peroxisome proliferator-activated receptor γ (PPAR γ) (Rosen et al., 2002; 2009).

C/EBPs, such as C/EBP α , C/EBP β , and C/EBP δ , are members of the basic leucine zipper transcription factor family

Received 3 July, 2020; revised 6 November, 2020; accepted 27 November, 2020; published online 18 December, 2020

eISSN: 0219-1032

©The Korean Society for Molecular and Cellular Biology. All rights reserved.

©This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>.

(Rosen et al., 2009). C/EBP β and C/EBP δ are expressed at the early stage of adipocyte differentiation, but their expression is not sufficient for complete adipogenic differentiation (Rosen et al., 2002; 2009; Tanaka et al., 1997). C/EBP β and C/EBP δ are required for the early stage of adipogenic differentiation by inducing C/EBP α and PPAR γ expression (Rosen et al., 2002; 2009). The expression of C/EBP α is enhanced late in adipogenesis, and C/EBP α promotes adipogenesis by inducing PPAR γ expression through direct binding of C/EBP recognition elements on the PPAR γ promoter (Elberg et al., 2000; Wu et al., 1996). Consequently, C/EBPs are considered to be important adipogenic transcription factors together with PPAR γ in adipogenesis. Interestingly, ectopic expression of PPAR γ in C/EBP α -deficient fibroblasts was shown to induce adipocyte differentiation, whereas C/EBP α expression in PPAR γ -deficient fibroblasts was not observed (Rosen et al., 2002; Wu et al., 1999). Thus, PPAR γ , a member of the nuclear receptor gene family, acts as the master transcriptional regulator in adipogenesis (Lefterova and Lazar, 2009).

PPAR γ is expressed as two major different isoforms, PPAR γ 1 and PPAR γ 2, as a result of alternative promoter usage; PPAR γ 2 has an extra 30 residues in its N-terminal end (Kroker and Bruning, 2015; Tontonoz et al., 1994). PPAR γ 2 is predominantly expressed in adipose tissue, where it regulates adipogenesis, while PPAR γ 1 is broadly expressed in adipose tissue, colon, retina, liver, spleen and cardiac muscles (Saladin et al., 1999). Upon PPAR γ activation by PPAR γ ligand binding, PPAR γ forms a heterodimer with another nuclear receptor, such as retinoid X receptors (RXRs) (Chandra et al., 2008; Tontonoz et al., 1994). The formation of PPAR γ -RXR α heterodimers is crucial to induce the expression of adipogenic markers, such as adipocyte fatty acid-binding protein 2 (aP2) and adiponectin, through direct binding to PPAR responsive elements (PPREs) in their promoters during adipogenesis (Juge-Aubry et al., 1997; Lefterova and Lazar, 2009). Interestingly, the transcriptional activity of PPAR γ -RXR α heterodimers is tightly regulated by the association of either coactivators or corepressors in many different tissues and cell types (Lefterova and Lazar, 2009). Thus, elucidating a functional role of the novel PPAR γ regulator may improve our understanding of the molecular regulatory mechanism of PPAR γ in various tissues and cells and in their relevance to human metabolic diseases, such as obesity and diabetes.

T-cell death-associated gene 51 (TDAG51), also known as the pleckstrin homology-like (PHL) domain A family 1 (PHL-DA1) in humans, possesses an N-terminal PHL domain and a glutamine repeat (QQ) domain (Park et al., 1996). In the C-terminus, TDAG51 contains a proline-glutamine repeat (PQ) domain and a proline-histidine repeat (PH) domain (Park et al., 1996; 2013). In eukaryotic systems, PHL domains are mainly involved in protein-protein interactions, and the QQ, PQ, and PH domains are predominantly present in transcriptional activators (Gehring et al., 1994; Scheffzek and Welti, 2012; Xiao and Jeang, 1998). Thus, TDAG51 was proposed to be a putative transcriptional regulator involved in many biological processes, such as cell proliferation, migration, apoptosis and differentiation (Chen et al., 2018; Nagai, 2016; Park et al., 1996). TDAG51 expression is modulated by various cellular stimuli, such as heat shock, oxidative stress, inflam-

mation, tumor induction and endoplasmic reticulum stress (Hayashida et al., 2006; Hossain et al., 2013; Jiao et al., 2016; Nagai, 2016; Park et al., 2013). In addition, microarray analysis showed that TDAG51 expression is only maintained at the early stage of adipogenic lineage and is rapidly decreased during the process of adipogenesis (Burton et al., 2004). Moreover, TDAG51-deficient (TDAG51^{-/-}) mice showed mature-onset obesity, hepatic steatosis and insulin resistance by regulating lipogenesis (Basseri et al., 2013). Thus, TDAG51 may negatively contribute to the regulation of adipogenesis. However, the exact role of TDAG51 in adipogenesis remains unclear.

In this study, we identified TDAG51 as a novel PPAR γ regulator. Adipogenesis is reduced by TDAG51 expression. The heterodimer formation of PPAR γ -RXR α is competitively inhibited by TDAG51 binding to PPAR γ . Thus, our data suggest that TDAG51 is a crucial negative regulator of PPAR γ -induced adipogenesis.

MATERIALS AND METHODS

Antibodies and cells

Specific antibodies were purchased from the following commercial sources: anti-Flag, anti-Myc, and anti- β -Actin from Sigma-Aldrich (USA); anti-Xpress (Xp) from Invitrogen (USA); and anti-GST, anti-TDAG51, and anti-PPAR γ antibodies from Santa Cruz Biotechnology (USA). Murine preadipocyte 3T3-L1 and human embryonic kidney 293T and PlatE cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Welgene, Korea) supplemented with 10% fetal bovine serum (FBS; Invitrogen) and 1 \times antibiotic/antimycotic solution (Welgene).

Plasmids

The luciferase reporter harboring the TDAG51 promoter fragments containing nt -2,450 to +551 (P_{T51}-3K-Luc) was described previously (Park et al., 2013). For the construction of aP2 luciferase reporter (aP2-Luc), the aP2 promoter fragments from nt -5,379 to +21 were amplified by polymerase chain reaction (PCR) from the murine genomic DNA and subcloned into the pGL3-basic vector (Promega, USA). The eukaryotic expression plasmid of Flag-tagged TDAG51 (Flag-TDAG51) was described previously (Park et al., 2013). Flag-tagged RXR α (Flag-RXR α) was kindly provided by Yong-Ho Ahn (Yonsei University College of Medicine, Korea) (Kim et al., 2009). The epitope-tagged (Flag-, Myc-, Xp-, or GST-tagged) eukaryotic expression plasmids for TDAG51, PPAR γ , RXR α , and their corresponding deletion mutants were generated by PCR amplification and subcloned into pFLAG-CMV2 (Sigma-Aldrich), pcDNA3.1/myc-His/lacZ (Invitrogen), pcDNA3.1-His (Invitrogen), pEBG (Park et al., 2015), and pMXS-puro (Park et al., 2015).

Adipogenic differentiation and analysis

Differentiation of adipocytes was performed as described previously (Basseri et al., 2013). Briefly, 3T3-L1 preadipocytes (1 \times 10⁵ cells/ml) were cultured in 10% FBS/DMEM supplemented with differential cocktail (1 μ M dexamethasone, 0.5 mM 3-isobutyl-1-methylxanthine, 100 μ M indomethacin, and

10 μ g/ml insulin [Sigma-Aldrich]) for 6 days. For the ectopic expression of TDAG51, retroviral supernatants harboring the Flag-tagged TDAG51 transgene were generated from PlatE cells as described previously (Yu et al., 2016). 3T3-L1 cells (1×10^5 cells/ml) were infected by retroviral supernatants with 8 μ g/ml polybrene for 6 h. At 24 h postinfection, infected 3T3-L1 cells were selected by puromycin (2 μ g/ml) for 2 days. After puromycin selection, 3T3-L1 cells were seeded at a density of 1×10^5 cells/ml in adipocyte differentiation media and differentiated into mature adipocytes for 6 days. Finally, adipocyte differentiation was analyzed by oil red O staining and solution assays as described previously (Basseri et al., 2013; Hossain et al., 2013).

Real-time PCR analysis

Real-time PCR was performed as described previously (Park et al., 2015). Briefly, differentiated 3T3-L1 cells were harvested, and total RNA was isolated using TRI reagent (MRC, USA) according to the manufacturers' instructions. Reverse transcription was performed with 1 μ g of total RNA and M-MLV reverse transcriptase (USB, USA) for 1 h at 42°C. The synthesized cDNAs were then subjected to real-time PCR analysis using a CFX Connect Real-time PCR Detection System (Bio-Rad, USA) with the appropriate primers: PPAR γ , 5'-ACC AGG GAG TTC CTC AAA AG-3' and 5'-TTG TCT TGG ATG TCC TCG AT-3'; adiponectin, 5'-ACT CCT GGA GAG AAG GGA GA-3' and 5'-GAA TGG GTA CAT TGG GAA CA-3'; and β -actin, 5'-ATG AAG ATC CTC CTG ACC GAG CG-3' and 5'-TAC TTG CGC TGA GGA GC-3'. β -Actin was used as an internal normalization control.

Luciferase reporter assay

A luciferase reporter assay was performed as described previously (Kim et al., 2019; Yu et al., 2016). Briefly, 293T (3.5×10^5 cells/ml) or 3T3-L1 (1×10^5 cells/ml) cells cultured in 24-well plates were cotransfected with reporters and an appropriate combination of epitope-tagged eukaryotic expression plasmids for TDAG51, PPAR γ or RXR α in triplicate using TurboFect reagent (Fermentas, USA) according to the manufacturers' instructions. Plasmids P_{T51}-3K-Luc (0.1 μ g), PPAR γ -Luc (0.1 μ g), aP2-Luc (0.1 μ g), and pcDNA3.1/His/LacZ (0.1 μ g, Invitrogen) were used as reporters. The reporters were cotransfected with the epitope-tagged expression plasmids (0.1-0.5 μ g). The β -galactosidase activity derived from the expression of pcDNA3.1/His/LacZ was used as an internal normalization control for transfection. For the reporter assay in 3T3-L1 cells, P_{T51}-3K-Luc-transfected 3T3-L1 cells were differentiated with adipocyte differentiation media for 2 days. For the treatment of rosiglitazone, the transfected cells were treated with 5 μ M rosiglitazone for 36 h, as described previously (Dowell et al., 2003). At 24 h posttransfection, reporter activities were measured using a luciferase assay kit (Promega) and β -galactosidase assay kit (Applied Biosystems, USA) according to the manufacturers' instructions.

Protein interaction and immunoblotting

Protein-protein interaction and immunoprecipitation (IP) assays were performed as described previously (Yu et al., 2016). Briefly, 293T cells seeded at 3.5×10^5 cells/ml in 6-well plates

were cotransfected with an appropriate combination of epitope-tagged eukaryotic expression plasmids for TDAG51 (0.2-2.5 μ g), PPAR γ (1.5 μ g), or RXR α (0.25-1.0 μ g). For the treatment of rosiglitazone, the transfected cells were treated with 5 μ M rosiglitazone for 36 h as described previously (Dowell et al., 2003). At 24 h posttransfection, the cells were lysed in cold lysis buffer (250 mM Tris-HCl [pH 7.5], 150 mM NaCl, 1 mM EDTA, 1 mM NaF, 1 mM sodium orthovanadate, 1 mM phenylmethylsulfonyl fluoride, 5% glycerol, and 0.5% Triton X-100). The lysates were pulled down (PD) by glutathione S-transferase (GST) beads or immunoprecipitated by anti-Flag beads. The bead-bound proteins were washed three times with lysis buffer and finally analyzed by immunoblotting (IB) with antibodies against epitopes, as described previously (Park et al., 2015; Son et al., 2020). For the endogenous protein-protein interaction assay, 3T3-L1 cells seeded at 1×10^5 cells/ml were cultured with adipocyte differentiation media for 2 days. The differentiated cells were lysed in cold lysis buffer, and the lysates were incubated with anti-TDAG51 or anti-PPAR γ antibody overnight at 4°C with rocking. Then, the lysates were precipitated with protein G beads (50% slurry) for 2 h at 4°C. The beads were washed three times with lysis buffer, and the interacting proteins were analyzed by IB with antibodies against anti-TDAG51 or anti-PPAR γ antibody.

Statistical analysis

Student's *t*-test was used to determine the significance of differences between experimental samples ($n = 3$ per group) using IBM SPSS Statistics (ver. 24.0; IBM, USA). The data represent the mean \pm SD. *P* values < 0.05 were considered statistically significant.

RESULTS

TDAG51 is negatively involved in the regulation of PPAR γ -induced adipocyte formation

Because TDAG51 expression is decreased in adipogenesis, as revealed by a microarray study (Burton et al., 2004), we determined whether the expression of TDAG51 is downregulated in adipogenesis using 3T3-L1 preadipocytes. We observed that TDAG51 expression was abundantly maintained at day 0 and was then rapidly reduced after day 2 of adipogenic differentiation, while PPAR γ expression was induced after day 2 (Fig. 1A). In a TDAG51 reporter assay, TDAG51 promoter activities were reduced (41.6%-79.3% reduction, $P < 0.05$ or $P < 0.01$) in a differentiation-dependent manner in 3T3-L1 cells (Fig. 1B). In addition, adipocyte formation was significantly inhibited (25.0% or 45.9% reduction at day 6, $P < 0.05$) by the forced expression of TDAG51 in 3T3-L1 preadipocytes (Fig. 1C). Consistent with these results, we observed that the mRNA levels of the adipogenic markers PPAR γ and adiponectin were inversely correlated with the levels of TDAG51 expression during adipogenic differentiation in 3T3-L1 preadipocytes (Fig. 1D). These results indicate that the downregulation of TDAG51 expression during adipogenesis is crucial for adipocyte formation and adipogenic marker expression.

As shown in Fig. 1, we observed that the mRNA levels of PPAR γ were downregulated by forced TDAG51 expression. Furthermore, Basseri et al. (2013) demonstrated that




Fig. 1. TDAG51 inhibits adipogenesis. (A) Downregulation of TDAG51 expression in 3T3-L1 preadipocyte differentiation. The 3T3-L1 preadipocytes were cultured in adipocyte differentiation media for 6 days (d). The expressed proteins were analyzed by immunoblotting against anti-PPAR γ or anti-TDAG51 antibody indicated by an arrow. β -Actin was used as the loading control. (B) Downregulation of TDAG51 transcription in 3T3-L1 preadipocytes. A schematic diagram of the luciferase reporter of the TDAG51 promoter (P_{T51}-3K-Luc) is shown in the left panel. The putative transcription factor binding sites are illustrated, and the nucleotide positions were numbered based on the transcriptional start site. NF- κ B, nuclear factor kappa B; CREB, cAMP response element-binding protein; AP-1, activator protein 1. TDAG51 promoter activities during 3T3-L1 preadipocyte differentiation were analyzed in P_{T51}-3K-Luc-transfected 3T3-L1 preadipocytes using reporter assay kits (right panel). (C) Inhibition of 3T3-L1 preadipocyte differentiation by TDAG51 expression. The Flag-tagged TDAG51-transduced 3T3-L1 preadipocytes were cultured in adipocyte differentiation media for 6 days. The differentiated cells were stained with oil red O. The oil red O levels in the differentiated cells were monitored at 490 nm (left panel). The number of stained cells was counted (right panel). Mock, gray box; Flag-TDAG51 expression, open box. (D) Downregulation of adipogenic markers in 3T3-L1 preadipocyte differentiation by TDAG51 expression. The expression levels of the adipogenic markers PPAR γ and adiponectin were analyzed by real-time PCR. * P < 0.05; ** P < 0.01.

TDAG51^{-/-} mice exhibited greater lipogenic potential and mature-onset obesity. Thus, we postulated that TDAG51 is negatively involved in the regulation of the master transcription factor PPAR γ in adipogenesis. To address this possibility, we constructed a luciferase reporter of the PPAR γ 2 promoter (PPAR γ -Luc) harboring two putative PPREs in the promoter fragment from nt -2,983 to +68, and the functionality of these putative PPREs was confirmed by site-directed mutagenesis and reporter assays (Supplementary Fig. S1). We next analyzed PPAR γ -induced PPAR γ -Luc activation by forced TDAG51 expression using reporter assays. Our results showed that PPAR γ -induced PPAR γ -Luc activation was dose-dependently inhibited (13.6%-29.0% reduction, P < 0.01 or P < 0.001) by forced TDAG51 expression (Supplementary Fig. S2A). Contrary to the TDAG51 inhibitory effect

on PPAR γ promoter activity, however, TDAG51 promoter activities were not inhibited by the forced expression of PPAR γ (Supplementary Fig. S2B). Taken together, these results indicate that TDAG51 expression is negatively involved in the regulation of PPAR γ -induced transcriptional activation.

TDAG51 interacts with PPAR γ

We next analyzed whether TDAG51 physically interacts with PPAR γ . In the GST bead pulldown (PD) assay with GST-tagged TDAG51 (GST-TDAG51), we observed an interaction between TDAG51 and PPAR γ (Fig. 2A). Consistent with these results, the anti-Flag IP assay with Flag-tagged PPAR γ (Flag-PPAR γ) revealed that Flag-PPAR γ interacted with Myc-TDAG51 (Fig. 2B). To further confirm this observation, we examined their endogenous interaction using an IP assay in





Fig. 2. TDAG51 interacts with PPAR γ . (A) TDAG51 binds to PPAR γ . 293T cells were transfected with GST-TDAG51 (1.5 μ g) with or without Flag-PPAR γ (2.5 μ g). GST-TDAG51 was pulled down (PD) using GST beads. The bound proteins were visualized via anti-Flag (top) or anti-GST (middle) immunoblotting (IB). The level of PPAR γ expression in whole cell lysates was detected via anti-Flag IB (bottom). GST alone (mock) was used as a control. Protein expression is indicated by an arrow. (B) PPAR γ binds to TDAG51. 293T cells were transfected with Flag-PPAR γ (1.5 μ g) with or without Myc-TDAG51 (2.5 μ g). The bound proteins by anti-Flag IP were visualized via anti-Myc (top) or anti-Flag IB (middle). The level of TDAG51 expression in whole cell lysates was detected via anti-Myc IB (bottom). Flag alone (mock) was used as a control. (C) Identification of the endogenous TDAG51-PPAR γ interaction. Endogenous TDAG51 was immunoprecipitated (IP) with an anti-(α -)TDAG51 antibody and visualized via anti-TDAG51 IB (second). Proteins bound to TDAG51 were visualized via anti-PPAR γ IB (top). The levels of endogenous TDAG51 and PPAR γ in whole cell lysates were detected via anti-TDAG51 (third) and anti-PPAR γ IB (bottom), respectively. (D) Identification of the endogenous PPAR γ -TDAG51 interaction. Endogenous PPAR γ was immunoprecipitated with an anti-(α -)PPAR γ antibody and visualized via anti-PPAR γ IB (second). Proteins bound to PPAR γ were visualized via anti-TDAG51 IB (top). The levels of endogenous PPAR γ and TDAG51 in whole cell lysates were detected via anti-PPAR γ (third) and anti-TDAG51 IB (bottom), respectively. (E) Schematic diagram of TDAG51 mutants. PHL, pleckstrin homology-like domain; QQ, glutamine repeat; PQ, proline-glutamine repeat; PH, proline-histidine repeat. The number of amino acid residues is shown. The interactions of deletion mutants are summarized at the right panel. (F) Analysis of TDAG51 deletion mutants interacting with PPAR γ . The bound proteins by GST-PD were visualized via anti-Flag (top) or anti-GST IB (middle). The level of PPAR γ expression in whole cell lysates was detected via anti-Flag IB (bottom). (G) Schematic diagram of PPAR γ mutants. AF-1, activation function-1 domain; DBD, DNA binding domain; Hinge, hinge domain; AF-2, activation function-2 domain; LBD, ligand binding domain. The interactions of deletion mutants are summarized at the right panel. (H) Analysis of PPAR γ deletion mutants interacting with TDAG51. The bound proteins by GST-PD were visualized via anti-Flag (top) or anti-GST IB (middle). The PPAR γ expression level in whole cell lysates was detected via anti-Flag IB (bottom).

G



H




Fig. 2. Continued.

3T3-L1 cells. Endogenous TDAG51 was immunoprecipitated by anti-TDAG51 antibodies, and the bound proteins were analyzed by IB with anti-PPAR γ antibodies. Our results showed that endogenous TDAG51 was associated with endogenous PPAR γ in 3T3-L1 cells (Fig. 2C). In an IP assay using anti-PPAR γ antibodies, we also observed that endogenous PPAR γ was associated with endogenous TDAG51 in 3T3-L1 cells (Fig. 2D). We next constructed deletion mutants of TDAG51, as shown in Fig. 2E, and analyzed particular domains involved in interacting with PPAR γ using a GST-PD assay. All TDAG51 deletion mutants, except for TDAG51 [202-262] containing the PH domain only, could interact with PPAR γ , indicating that the PHL, QQ, and PQ domains in TDAG51 are involved in the PPAR γ interaction (Fig. 2F). To determine the precise domain of PPAR γ that interacts with TDAG51, we constructed expression plasmids for PPAR γ deletion mutants lacking the N-terminal, C-terminal, and internal regions (Fig. 2G). GST-PD assays for the N- and C-terminal deletion mutants of PPAR γ revealed that only the PPAR γ [1-139] mutant was not bound to TDAG51, indicating that the activation function-1 (AF-1) domain in PPAR γ is not involved in the TDAG51 interaction (Fig. 2H). In the GST-PD assay with the internal deletion mutants, all PPAR γ internal deletion mutants could interact with TDAG51, although the PPAR γ Δ [281-435] mutant lacking the ligand binding domain (LBD) showed a markedly reduced interaction with TDAG51 (Fig. 2H). Collectively, we observed that the region between residues 140 and 506, including the DNA binding domain (DBD), hinge, LBD, and AF-2 domain, in PPAR γ is required for the TDAG51 interaction (Fig. 2H). Taken together, these results indicate that TDAG51 is a binding partner of PPAR γ .

The interaction of TDAG51 and PPAR γ inhibits PPAR γ -induced aP2 promoter activity

PPAR γ is a crucial transcriptional regulator of the induction of adipogenic markers, such as aP2 and adiponectin (Juge-Aubry et al., 1997; Lefterova and Lazar, 2009). Thus, we investigated whether the interaction of TDAG51-PPAR γ could regulate PPAR γ -induced aP2 transcriptional activation using a luciferase reporter of the aP2 promoter (aP2-Luc) harboring transcriptional regulatory elements, such as PPREs and C/EBPs, in the promoter fragment from nt -5,379 to +21 (Fig. 3A). In a PPAR γ -induced aP2-Luc reporter assay, PPAR γ -induced aP2-Luc activities were inhibited (19.5%-45.5% reduction, $P < 0.05$ or $P < 0.01$) in a dose-dependent manner by forced TDAG51 expression (Fig. 3B). Consistent with the results shown in Fig. 2F, the inhibitory effect of TDAG51 in the PPAR γ -induced aP2-Luc reporter assay was not observed by the PPAR γ -nonbinding mutant of TDAG51 (GST-TDAG51[202-262]), while the wild-type TDAG51 (GST-TDAG51) or its PPAR γ -binding mutant (GST-TDAG51[1-201]) showed inhibitory effects (39.0% or 34.1% reduction, $P < 0.01$) (Fig. 3C). Taken together, these results indicate that the binding of TDAG51 to PPAR γ is negatively involved in the regulation of PPAR γ -induced adipogenic marker expression.

TDAG51 binding to PPAR γ inhibits the formation of PPAR γ -RXR α heterodimers

The formation of PPAR γ -RXR α heterodimers by PPAR γ ligand stimulation is crucial to induce the expression of adipogenic markers through binding to PPREs in their promoters in adipogenesis (Chandra et al., 2008; Lefterova and Lazar, 2009; Tontonoz et al., 1994). Thus, we examined whether TDAG51 acts as a corepressor of the PPAR γ -RXR α heterodimer. In a GST-PD assay, we observed a direct interaction of




Fig. 3. TDAG51 inhibits PPAR γ -induced aP2 promoter activity. (A) Schematic diagram of the aP2 promoter (aP2-Luc). The putative transcription factor binding sites are illustrated, and the nucleotide positions are numbered based on the transcriptional start site. PPRE, PPAR γ responsive element; C/EBP, CCAAT/enhancer binding protein; SP1, specificity protein 1; AP-2, activator protein 2. (B) The inhibitory effect of TDAG51 on PPAR γ -induced aP2-Luc reporter activity. Reporter plasmids (aP2-Luc [0.1 μ g] and pcDNA3.1/His/LacZ [0.1 μ g]) were cotransfected with epitope-tagged expression plasmids (Flag-PPAR γ [0.3 μ g] and/or Myc-TDAG51 [0.2 and 0.5 μ g]) into 293T cells. The expression of transfected plasmids was analyzed by immunoblotting against anti-epitope antibodies. β -Actin was used as the loading control. Protein expression is indicated by an arrow. (C) The inhibitory effects of TDAG51 deletion mutants on PPAR γ -induced aP2-Luc reporter activity. Reporter plasmids were cotransfected with epitope-tagged expression plasmids (Flag-PPAR γ [0.3 μ g] and/or GST-TDAG51 [0.5 μ g]) into 293T cells. GST alone (mock) was used as a control. * P < 0.05; ** P < 0.01; n.s., not significant.

TDAG51 with PPAR γ but not with RXR α (Fig. 4A). Moreover, the interaction of TDAG51 and PPAR γ was not blocked by treatment with the PPAR γ ligand rosiglitazone (Fig. 4B). Thus, TDAG51 binding to PPAR γ *per se* may negatively affect the formation of PPAR γ -RXR α heterodimers. To address this possibility, we performed competitive IP assays with Flag-PPAR γ by dose-dependent expression of TDAG51 or RXR α (Figs. 4C-4E). Our results showed that the formation of PPAR γ -RXR α heterodimers was inhibited in a dose-dependent manner by forced TDAG51 expression (Fig. 4C), whereas the interaction between PPAR γ and TDAG51 was dose-dependently reduced by forced RXR α expression (Fig. 4D). Consistent with the results shown in Figs. 2F and 3C, in IP assays with TDAG51 deletion mutants, the formation of the PPAR γ -RXR α heterodimer was not inhibited by the forced expression of the PPAR γ -non-binding mutant (GST-TDAG51[202-262]), while wild-type TDAG51 (GST-TDAG51) or its PPAR γ -binding mutant

(GST-TDAG51[1-201]) showed competitive inhibitory effects on the formation of the PPAR γ -RXR α heterodimer (Fig. 4E). Taken together, these results indicate that TDAG51 inhibits the formation of PPAR γ -RXR α heterodimers through direct interaction with PPAR γ .

TDAG51 inhibits aP2 promoter activity induced by the PPAR γ -RXR α heterodimer

We next examined whether TDAG51 could inhibit aP2 promoter activity induced by the PPAR γ -RXR α heterodimer using an aP2-Luc reporter assay (Fig. 5A). In the aP2-Luc reporter assay, we observed enhanced promoter activities of PPAR γ -induced aP2-Luc (3.4-fold induction, P < 0.01) by coexpressing RXR α compared to PPAR γ alone (Fig. 5A). These enhanced aP2 promoter activities induced by the PPAR γ -RXR α heterodimer were dose-dependently inhibited (14.6%-28.9% reduction, P < 0.01) by forced TDAG51 ex-




Fig. 4. The interaction of TDAG51 and PPAR γ inhibits the formation of the PPAR γ -RXR α heterodimer. (A) No interaction between TDAG51 and RXR α . First, 293T cells were cotransfected with GST-TDAG51 (1.5 μ g) with or without Flag-PPAR γ (1.5 μ g) or Flag-RXR α (0.1 μ g). The bound proteins by GST-PD were visualized via anti-Flag (top) or anti-GST IB (middle). The PPAR γ and RXR α expression levels in whole cell lysates were detected via anti-Flag IB (bottom). GST alone (mock) was used as a control. Protein expression is indicated by an arrow. (B) No effect of rosiglitazone treatment on the TDAG51-PPAR γ interaction. The 293T cells cotransfected with epitope-tagged expression plasmids were treated with 5 μ M rosiglitazone for 36 h. The protein interaction was monitored by GST-PD and IB analysis. (C) The inhibitory effect of TDAG51 on the formation of PPAR γ -RXR α heterodimers. The Xp-tagged TDAG51 plasmid (Xp-TDAG51 [0.2, 0.5, and 2.0 μ g]) was dose-dependently cotransfected with Flag-PPAR γ (1.5 μ g) and Myc-RXR α (0.1 μ g) expression plasmids into 293T cells. The protein interactions were monitored by anti-Flag IP and IB analysis (top three panels). The TDAG51 and RXR α expression levels in whole cell lysates were detected via anti-Xp and anti-Myc IB, respectively (bottom two panels). (D) The inhibitory effect of RXR α on the interaction of PPAR γ and TDAG51. The Myc-tagged RXR α plasmid (Myc-RXR α [0.25, 0.5, and 1.0 μ g]) was dose-dependently cotransfected with Flag-PPAR γ (1.5 μ g) and Myc-TDAG51 (1.5 μ g) expression plasmids into 293T cells. (E) The inhibitory effects of TDAG51 deletion mutants on the formation of PPAR γ -RXR α heterodimers. The GST-tagged TDAG51 deletion mutant plasmids (0.5 and 2.0 μ g) were dose-dependently cotransfected with Flag-PPAR γ (1.5 μ g) and Myc-RXR α (0.1 μ g) expression plasmids into 293T cells.

pression (Fig. 5A). Consistent with the results shown in Fig. 4B, treatment with rosiglitazone did not affect the inhibitory effect of TDAG51 on the activation of PPAR γ -induced aP2 promoter activity (Fig. 5B). Finally, we analyzed the inhibitory effects of TDAG51 using TDAG51 deletion mutants. Similar to the results shown in Figs. 2F, 3C, and 4E, the inhibitory effects of TDAG51 on the activation of aP2 promoter activity induced by the PPAR γ -RXR α heterodimer were observed (20.8% or 29.4% reduction, $P < 0.01$) with wild type or PPAR γ -binding mutant (GST-TDAG51[1-201]) expression but not with the PPAR γ -nonbinding mutant (GST-TDAG51[202-262]) (Fig. 5C). Taken together, these results indicate that TDAG51 functions as a corepressor of PPAR γ in adipogenic marker expression by inhibiting the transcriptional activity of the PPAR γ -RXR α heterodimer.

DISCUSSION

PPAR γ plays a central role in regulating the expression of a wide range of genes involved in adipogenesis, lipid metabolism, insulin sensitivity, energy storage, inflammation and differentiation (Kroger and Bruning, 2015; Lefterova and Lazar,

2009). In particular, the crucial role of PPAR γ in adipogenesis has been extensively studied by many researchers (Lefterova and Lazar, 2009). Although PPAR γ functions as the master transcriptional regulator of adipocyte differentiation, few genes have been validated as direct PPAR γ regulators or targets (Lefterova and Lazar, 2009). Thus, the identification of novel regulators of PPAR γ is important for understanding the transcriptional regulatory mechanism of PPAR γ in various cell types or tissues. PPAR γ contains AF-1 and DBD in the N-terminal region, the flexible hinge domain in the middle, and LBD and AF-2 in the C-terminal region (Kroger and Bruning, 2015). Upon the binding of the PPAR γ ligand, PPAR γ forms a heterodimer with RXR α through their LBD domain in the C-terminal region (Chandra et al., 2008). In this study, we identified TDAG51 as a novel negative regulator of PPAR γ that blocks PPAR γ -RXR α heterodimer formation in adipogenesis. We showed that the region between residues 140 and 506, including the DBD, hinge, LBD and AF-2 domain, of the PPAR γ protein is involved in direct interactions with the PHL, QQ, and PQ domains, except for the PH domain, in TDAG51 (Fig. 2). Furthermore, we observed that the TDAG51-PPAR γ interaction is not inhibited by the binding of




Fig. 5. TDAG51 inhibits aP2 promoter activity induced by the PPAR γ -RXR α heterodimer. (A) The inhibitory effects of TDAG51 on the promoter activity of aP2-Luc induced by PPAR γ -RXR α coexpression. Reporter plasmids (aP2-Luc [0.1 μ g] and pcDNA3.1/His/LacZ [0.1 μ g]) were cotransfected with epitope-tagged expression plasmids (Flag-PPAR γ [0.25 μ g], Myc-RXR α [0.1 μ g], and/or Xp-TDAG51 [0.25-0.5 μ g]) into 293T cells. The expression of transfected plasmids was analyzed by immunoblotting against anti-epitope antibodies. β -Actin was used as the loading control. (B) The inhibitory effects of TDAG51 on the promoter activity of aP2-Luc induced by PPAR γ expression with rosiglitazone treatment. (C) The inhibitory effects of TDAG51 deletion mutants on the promoter activity of aP2-Luc induced by PPAR γ -RXR α coexpression. Reporter plasmids were cotransfected with epitope-tagged expression plasmids (Flag-PPAR γ [0.25 μ g], Myc-RXR α [0.1 μ g], and/or GST-TDAG51 [0.5 μ g]) into 293T cells. GST alone (mock) was used as a control. ** P < 0.01; *** P < 0.001; n.s., not significant.

synthetic PPAR γ ligand and that TDAG51 is more strongly associated with PPAR γ alone than the PPAR γ -RXR α heterodimer in protein-protein interaction assays (Fig. 4). To date, no DNA binding motif in TDAG51 has been found. TDAG51, which has evolutionarily conserved PHL, PQ, and PH domains, is considered a putative transcriptional regulator (Nagai, 2016). Thus, based on our results and previous studies, TDAG51 can function as a ligand-independent competitive inhibitor of RXR α recruitment toward assembly as a PPAR γ -RXR α heterodimer complex in adipogenesis. Interestingly, we showed that TDAG51 expression is downregulated in a differentiation-dependent manner during adipogenic differentiation and that this downregulation is not directly mediated by the adipogenic transcription factor PPAR γ (Fig. 1, Supplementary Fig. S2). Hence, further exploration of the transcriptional regulatory pathways involved in the downregulation of TDAG51 at the early stage of adipogenesis would be of interest.

PPAR γ activity is modulated by several posttranslational modifications, including phosphorylation, sumoylation and ubiquitination (Ahmadian et al., 2013). Interestingly, the phosphorylation of PPAR γ can repress or enhance its activity depending on the phosphorylation site or kinase activation, while PPAR γ sumoylation has been shown to repress PPAR γ activity by the recruitment of corepressors with histone deacetylases (Hu et al., 1996; Iankova et al., 2006; Pascual et al., 2005). Notably, TDAG51 was reported to be involved in regulating several kinase pathways, such as aurora A kinase, ErbB2 tyrosine kinase, Akt and receptor tyrosine kinase, in cancer cells (Fearon et al., 2018; Johnson et al., 2011; Li et al., 2014). Hence, based on previous studies, the inhibitory effect of TDAG51 on PPAR γ activity is indirectly mediated by possible recruitment of another corepressor or kinase. Thus, further studies are needed to elucidate whether any other regulators or kinases are involved in indirect regulation of the TDAG51-mediated negative effects on PPAR γ -induced adipogenesis. Ubiquitination and proteasome-dependent degradation are also linked to the regulation of PPAR γ activity by ligand binding, IFN γ stimulation or recruitment of mediators, including Wnt1-inducible signaling pathway protein-1 (Hausser et al., 2000; Waite et al., 2001). However, we assumed that TDAG51 function is not linked to the proteasomal degradation of PPAR γ because the levels of PPAR γ were not altered by the TDAG51 interaction (Figs. 2 and 4).

Several transcription factors have been shown to regulate PPAR γ expression and activity in adipogenesis (Miard and Fajas, 2005). Of the six GATA family transcription factors, GATA-2 and GATA-3 were shown to downregulate PPAR γ expression by directly interacting with the PPAR γ 2 promoter and C/EBP family, thereby inhibiting adipogenic differentiation (Tong et al., 2000). The expression of GATA-2/3 is abundant in preadipocytes and progressively decreases during adipocyte differentiation (Tong et al., 2000). Thus, GATA-2/3 have a potential role in commitment to the adipogenic lineage (Kamata et al., 2014). Similar to the pattern of GATA-2/3 expression shown in the work of Tong et al. (2000), the most abundant expression of TDAG51 was observed only in the early stage of adipogenic differentiation (Fig. 1). Although we have not yet experimentally determined the preadipocyte restriction by TDAG51 expression in the early stage of adi-

pogenic differentiation, we propose that TDAG51 may also have a potential role in commitment to the adipogenic lineage. FoxO1, a member of the forkhead transcription factor family, is known to be a negative regulator of adipogenesis (Nakae et al., 2003). This molecule represses both PPAR γ expression through direct binding to the FoxO1 binding consensus sequences on the PPAR γ 2 promoter and PPAR γ activity by inhibiting the DNA binding activity of the PPAR γ -RXR α heterodimer through direct interaction with PPAR γ (Armoni et al., 2006; Dowell et al., 2003; Fan et al., 2009). Furthermore, the physical interaction among FoxO1, C/EBP α and small heterodimer partner (SHP) has been reported (Park et al., 2007; Qiao and Shao, 2006; Yamagata et al., 2004). Interestingly, SHP, an orphan nuclear receptor that lacks a DBD, is also known to repress PPAR γ activity by directly interacting with RXR α , but not with PPAR γ , which blocks RXR α recruitment to the PPAR γ -RXR α heterodimer, although the precise role of SHP in PPAR γ activity modulation remains controversial (Kim et al., 2009; Nishizawa et al., 2002). Consequently, the mechanism of FoxO1 or SHP that represses PPAR γ activity is quite distinct from the inhibitory mechanism of TDAG51 that blocks the recruitment of RXR α to the PPAR γ -RXR α heterodimer complex by directly interacting with PPAR γ .

In conclusion, we demonstrated that TDAG51 is a novel negative regulator of PPAR γ in adipogenesis. In this study, we showed that TDAG51 expressed abundantly in the early stage of adipocyte differentiation inhibits adipogenesis. TDAG51 interacts with PPAR γ in a ligand-independent manner, thereby inhibiting PPAR γ activity by blocking the formation of the PPAR γ -RXR α heterodimer complex. Thus, these results indicate that TDAG51, which could determine adipogenic cell fate, is a ligand-independent competitive inhibitor of PPAR γ in adipogenesis, expanding the importance of TDAG51 in the regulation in metabolic diseases, including obesity and diabetes.

Note: Supplementary information is available on the Molecules and Cells website (www.molcells.org).

ACKNOWLEDGMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) and was funded by the Ministry of Education (NRF-2017R1A2B4007327, 2019R1A2C1084311, 2019M3F6A1109486) and by the research fund of Chungnam National University.

AUTHOR CONTRIBUTIONS

S.K. and N.L. performed experiments and wrote the manuscript. E.S.P., H.Y., T.U.H., H.J., J.Y., S.C., B.S., and J.Y. performed experiments or analyzed the data. S.D.R. and Y.C. contributed essential reagents or tools and designed the research study. J.R. supervised the research and edited the manuscript.

CONFLICT OF INTEREST

The authors have no potential conflicts of interest to disclose.

ORCID

Sumi Kim <https://orcid.org/0000-0002-6524-6170>
Nari Lee <https://orcid.org/0000-0003-3104-4877>
Eui-Soon Park <https://orcid.org/0000-0002-1375-8027>
Hyeongseok Yun <https://orcid.org/0000-0001-9959-7254>
Tae-Uk Ha <https://orcid.org/0000-0002-1434-2365>
Hyoen Jeon <https://orcid.org/0000-0002-2141-6123>
Jiyeon Yu <https://orcid.org/0000-0001-5368-3093>
Seunga Choi <https://orcid.org/0000-0002-6688-8239>
Bongjin Shin <https://orcid.org/0000-0002-3763-2613>
Jungeun Yu <https://orcid.org/0000-0002-4036-4867>
Sang Dal Rhee <https://orcid.org/0000-0003-0620-1893>
Yongwon Choi <https://orcid.org/0000-0002-1007-7998>
Jaerang Rho <https://orcid.org/0000-0002-0019-4939>

REFERENCES

- Ahmadian, M., Suh, J.M., Hah, N., Liddle, C., Atkins, A.R., Downes, M., and Evans, R.M. (2013). PPAR γ signaling and metabolism: the good, the bad and the future. *Nat. Med.* 19, 557-566.
- Armoni, M., Harel, C., Karni, S., Chen, H., Bar-Yoseph, F., Ver, M.R., Quon, M.J., and Karnieli, E. (2006). FOXO1 represses peroxisome proliferator-activated receptor- γ 1 and - γ 2 gene promoters in primary adipocytes. A novel paradigm to increase insulin sensitivity. *J. Biol. Chem.* 281, 19881-19891.
- Basseri, S., Lhoták, Š., Fullerton, M.D., Palanivel, R., Jiang, H., Lynn, E.G., Ford, R.J., Maclean, K.N., Steinberg, G.R., and Austin, R.C. (2013). Loss of TDAG51 results in mature-onset obesity, hepatic steatosis, and insulin resistance by regulating lipogenesis. *Diabetes* 62, 158-169.
- Burton, G.R., Nagarajan, R., Peterson, C.A., and McGehee, R.E., Jr. (2004). Microarray analysis of differentiation-specific gene expression during 3T3-L1 adipogenesis. *Gene* 329, 167-185.
- Chandra, V., Huang, P., Hamuro, Y., Raghuram, S., Wang, Y., Burris, T.P., and Rastinejad, F. (2008). Structure of the intact PPAR- γ -RXR- nuclear receptor complex on DNA. *Nature* 456, 350-356.
- Chen, Y., Takikawa, M., Tsutsumi, S., Yamaguchi, Y., Okabe, A., Shimada, M., Kawase, T., Sada, A., Ezawa, I., Takano, Y., et al. (2018). PHLDA1, another PHLDA family protein that inhibits Akt. *Cancer Sci.* 109, 3532-3542.
- Dowell, P., Otto, T.C., Adi, S., and Lane, M.D. (2003). Convergence of peroxisome proliferator-activated receptor γ and Foxo1 signaling pathways. *J. Biol. Chem.* 278, 45485-45491.
- Elberg, G., Gimble, J.M., and Tsai, S.Y. (2000). Modulation of the murine peroxisome proliferator-activated receptor γ 2 promoter activity by CCAAT/enhancer-binding proteins. *J. Biol. Chem.* 275, 27815-27822.
- Fan, W., Imamura, T., Sonoda, N., Sears, D.D., Patsouris, D., Kim, J.J., and Olefsky, J.M. (2009). FOXO1 transrepresses peroxisome proliferator-activated receptor γ transactivation, coordinating an insulin-induced feed-forward response in adipocytes. *J. Biol. Chem.* 284, 12188-12197.
- Fearon, A.E., Carter, E.P., Clayton, N.S., Wilkes, E.H., Baker, A.M., Kapitonova, E., Bakhouch, B.A., Tanner, Y., Wang, J., Gadaleta, E., et al. (2018). PHLDA1 mediates drug resistance in receptor tyrosine kinase-driven cancer. *Cell Rep.* 22, 2469-2481.
- Gehring, W.J., Affolter, M., and Burglin, T. (1994). Homeodomain proteins. *Annu. Rev. Biochem.* 63, 487-526.
- Hauser, S., Adelmant, G., Sarraf, P., Wright, H.M., Mueller, E., and Spiegelman, B.M. (2000). Degradation of the peroxisome proliferator-activated receptor γ is linked to ligand-dependent activation. *J. Biol. Chem.* 275, 18527-18533.
- Hayashida, N., Inouye, S., Fujimoto, M., Tanaka, Y., Izu, H., Takaki, E., Ichikawa, H., Rho, J., and Nakai, A. (2006). A novel HSF1-mediated death

- pathway that is suppressed by heat shock proteins. *EMBO J.* **25**, 4773-4783.
- Hossain, G.S., Lynn, E.G., Maclean, K.N., Zhou, J., Dickhout, J.G., Lhotak, S., Trigatti, B., Capone, J., Rho, J., Tang, D., et al. (2013). Deficiency of TDAG51 protects against atherosclerosis by modulating apoptosis, cholesterol efflux, and peroxiredoxin-1 expression. *J. Am. Heart Assoc.* **2**, e000134.
- Hu, E., Kim, J.B., Sarraf, P., and Spiegelman, B.M. (1996). Inhibition of adipogenesis through MAP kinase-mediated phosphorylation of PPAR γ . *Science* **274**, 2100-2103.
- Iankova, I., Petersen, R.K., Annicotte, J.S., Chavey, C., Hansen, J.B., Kratchmarova, I., Sarraf, D., Benkirane, M., Kristiansen, K., and Fajas, L. (2006). Peroxisome proliferator-activated receptor gamma recruits the positive transcription elongation factor b complex to activate transcription and promote adipogenesis. *Mol. Endocrinol.* **20**, 1494-1505.
- Jiao, H.W., Jia, X.X., Zhao, T.J., Rong, H., Zhang, J.N., Cheng, Y., Zhu, H.P., Xu, K.L., Guo, S.Y., Shi, Q.Y., et al. (2016). Up-regulation of TDAG51 is a dependent factor of LPS-induced RAW264.7 macrophages proliferation and cell cycle progression. *Immunopharmacol. Immunotoxicol.* **38**, 124-130.
- Johnson, E.O., Chang, K.H., de Pablo, Y., Ghosh, S., Mehta, R., Badve, S., and Shah, K. (2011). PHLDA1 is a crucial negative regulator and effector of Aurora A kinase in breast cancer. *J. Cell Sci.* **124**, 2711-2722.
- Juge-Aubry, C., Pernin, A., Favez, T., Burger, A.G., Wahli, W., Meier, C.A., and Desvergne, B. (1997). DNA binding properties of peroxisome proliferator-activated receptor subtypes on various natural peroxisome proliferator response elements. Importance of the 5'-flanking region. *J. Biol. Chem.* **272**, 25252-25259.
- Kamata, M., Okitsu, Y., Fujiwara, T., Kanehira, M., Nakajima, S., Takahashi, T., Inoue, A., Fukuhara, N., Onishi, Y., Ishizawa, K., et al. (2014). GATA2 regulates differentiation of bone marrow-derived mesenchymal stem cells. *Haematologica* **99**, 1686-1696.
- Kim, J.L., Kaufman, R.J., Back, S.H., and Moon, J.Y. (2019). Development of a reporter system monitoring regulated intramembrane proteolysis of the transmembrane bZIP transcription factor ATF6 α . *Mol. Cells* **42**, 783-793.
- Kim, T.H., Kim, H., Park, J.M., Im, S.S., Bae, J.S., Kim, M.Y., Yoon, H.G., Cha, J.Y., Kim, K.S., and Ahn, Y.H. (2009). Interrelationship between liver X receptor alpha, sterol regulatory element-binding protein-1c, peroxisome proliferator-activated receptor gamma, and small heterodimer partner in the transcriptional regulation of glucokinase gene expression in liver. *J. Biol. Chem.* **284**, 15071-15083.
- Kroker, A.J. and Bruning, J.B. (2015). Review of the structural and dynamic mechanisms of PPAR γ partial agonism. *PPAR Res.* **2015**, 816856.
- Lefterova, M.I. and Lazar, M.A. (2009). New developments in adipogenesis. *Trends Endocrinol. Metab.* **20**, 107-114.
- Li, G., Wang, X., Hibshoosh, H., Jin, C., and Halmos, B. (2014). Modulation of ErbB2 blockade in ErbB2-positive cancers: the role of ErbB2 Mutations and PHLDA1. *PLoS One* **9**, e106349.
- Miard, S. and Fajas, L. (2005). Atypical transcriptional regulators and cofactors of PPAR γ . *Int. J. Obes. (Lond.)* **29** Suppl 1, S10-S12.
- Nagai, M.A. (2016). Pleckstrin homology-like domain, family A, member 1 (PHLDA1) and cancer. *Biomed. Rep.* **4**, 275-281.
- Nakae, J., Kitamura, T., Kitamura, Y., Biggs, W.H., 3rd, Arden, K.C., and Accili, D. (2003). The forkhead transcription factor Foxo1 regulates adipocyte differentiation. *Dev. Cell* **4**, 119-129.
- Nishizawa, H., Yamagata, K., Shimomura, I., Takahashi, M., Kuriyama, H., Kishida, K., Hotta, K., Nagaretani, H., Maeda, N., Matsuda, M., et al. (2002). Small heterodimer partner, an orphan nuclear receptor, augments peroxisome proliferator-activated receptor gamma transactivation. *J. Biol. Chem.* **277**, 1586-1592.
- Park, C.G., Lee, S.Y., Kandala, G., Lee, S.Y., and Choi, Y. (1996). A novel gene product that couples TCR signaling to Fas(CD95) expression in activation-induced cell death. *Immunity* **4**, 583-591.
- Park, E.S., Choi, S., Shin, B., Yu, J., Hwang, J.M., Yun, H., Chung, Y.H., Choi, J.S., Choi, Y., and Rho, J. (2015). Tumor necrosis factor (TNF) receptor-associated factor (TRAF)-interacting protein (TRIP) negatively regulates the TRAF2 ubiquitin-dependent pathway by suppressing the TRAF2-sphingosine 1-phosphate (S1P) interaction. *J. Biol. Chem.* **290**, 9660-9673.
- Park, E.S., Kim, J., Ha, T.U., Choi, J.S., Soo Hong, K., and Rho, J. (2013). TDAG51 deficiency promotes oxidative stress-induced apoptosis through the generation of reactive oxygen species in mouse embryonic fibroblasts. *Exp. Mol. Med.* **45**, e35.
- Park, M.J., Kong, H.J., Kim, H.Y., Kim, H.H., Kim, J.H., and Cheong, J.H. (2007). Transcriptional repression of the gluconeogenic gene PEPCK by the orphan nuclear receptor SHP through inhibitory interaction with C/EBP α . *Biochem. J.* **402**, 567-574.
- Pascual, G., Fong, A.L., Ogawa, S., Gamliel, A., Li, A.C., Perissi, V., Rose, D.W., Willson, T.M., Rosenfeld, M.G., and Glass, C.K. (2005). A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR- γ . *Nature* **437**, 759-763.
- Qiao, L. and Shao, J. (2006). SIRT1 regulates adiponectin gene expression through Foxo1-C/enhancer-binding protein alpha transcriptional complex. *J. Biol. Chem.* **281**, 39915-39924.
- Rosen, E., Eguchi, J., and Xu, Z. (2009). Transcriptional targets in adipocyte biology. *Expert Opin. Ther. Targets* **13**, 975-986.
- Rosen, E.D., Hsu, C.H., Wang, X., Sakai, S., Freeman, M.W., Gonzalez, F.J., and Spiegelman, B.M. (2002). C/EBP α induces adipogenesis through PPAR γ : a unified pathway. *Genes Dev.* **16**, 22-26.
- Saladin, R., Fajas, L., Dana, S., Halvorsen, Y.D., Auwerx, J., and Briggs, M. (1999). Differential regulation of peroxisome proliferator activated receptor gamma1 (PPARgamma1) and PPARgamma2 messenger RNA expression in the early stages of adipogenesis. *Cell Growth Differ.* **10**, 43-48.
- Sarjeant, K. and Stephens, J.M. (2012). Adipogenesis. *Cold Spring Harb. Perspect. Biol.* **4**, a008417.
- Scheffzek, K. and Welti, S. (2012). Pleckstrin homology (PH) like domains - versatile modules in protein-protein interaction platforms. *FEBS Lett.* **586**, 2662-2673.
- Son, H.E., Min, H.Y., Kim, E.J., and Jang, W.G. (2020). Fat mass and obesity-associated (FTO) stimulates osteogenic differentiation of C3H10T1/2 cells by inducing mild endoplasmic reticulum stress via a positive feedback loop with p-AMPK. *Mol. Cells* **43**, 58-65.
- Tanaka, T., Yoshida, N., Kishimoto, T., and Akira, S. (1997). Defective adipocyte differentiation in mice lacking the C/EBPbeta and/or C/EBPdelta gene. *EMBO J.* **16**, 7432-7443.
- Tong, Q., Dalgin, G., Xu, H., Ting, C.N., Leiden, J.M., and Hotamisligil, G.S. (2000). Function of GATA transcription factors in preadipocyte-adipocyte transition. *Science* **290**, 134-138.
- Tontonoz, P., Hu, E., Graves, R.A., Budavari, A.I., and Spiegelman, B.M. (1994). mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes Dev.* **8**, 1224-1234.
- Waite, K.J., Floyd, Z.E., Arbour-Reilly, P., and Stephens, J.M. (2001). Interferon-gamma-induced regulation of peroxisome proliferator-activated receptor gamma and STATs in adipocytes. *J. Biol. Chem.* **276**, 7062-7068.
- Wu, Z., Bucher, N.L., and Farmer, S.R. (1996). Induction of peroxisome proliferator-activated receptor gamma during the conversion of 3T3 fibroblasts into adipocytes is mediated by C/EBPbeta, C/EBPdelta, and glucocorticoids. *Mol. Cell. Biol.* **16**, 4128-4136.
- Wu, Z., Rosen, E.D., Brun, R., Hauser, S., Adelmant, G., Troy, A.E., McKeon, C., Darlington, G.J., and Spiegelman, B.M. (1999). Cross-regulation of C/EBP alpha and PPAR gamma controls the transcriptional pathway of

Negative Effect of TDAG51 on PPAR γ Activity
Sumi Kim et al.

adipogenesis and insulin sensitivity. *Mol. Cell* 3, 151-158.

Xiao, H. and Jeang, K.T. (1998). Glutamine-rich domains activate transcription in yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 273, 22873-22876.

Yamagata, K., Daitoku, H., Shimamoto, Y., Matsuzaki, H., Hirota, K., Ishida, J., and Fukamizu, A. (2004). Bile acids regulate gluconeogenic gene expression via small heterodimer partner-mediated repression of

hepatocyte nuclear factor 4 and Foxo1. *J. Biol. Chem.* 279, 23158-23165.

Yu, J., Yun, H., Shin, B., Kim, Y., Park, E.S., Choi, S., Yu, J., Amarasekara, D.S., Kim, S., Inoue, J., et al. (2016). Interaction of tumor necrosis factor receptor-associated factor 6 (TRAF6) and Vav3 in the receptor activator of nuclear factor kappaB (RANK) signaling complex enhances osteoclastogenesis. *J. Biol. Chem.* 291, 20643-20660.