

Second Order Parallel Tensor on Almost Kenmotsu Manifolds

VENKATESHA VENKATESHA*

Department of Mathematics, Kuvempu University, Shivamogga 577-451, India
e-mail: vensmath@gmail.com

DEVARAJA MALLESHA NAIK

Department of Mathematics, Kuvempu University, Shivamogga 577-451, India
Department of Mathematics, CHRIST (Deemed to be University), Bengaluru
560029, Karnataka, India
e-mail: devaraja.mallesha@christuniversity.in

AYSEL-TURGUT VANLI

Department of Mathematics, Gazi University, Ankara, Turkey
e-mail: avanli@gazi.edu.tr

ABSTRACT. Let M be an almost Kenmotsu manifold of dimension $2n + 1$ having non-vanishing ξ -sectional curvature such that $tr\ell > -2n - 2$. We prove that any second order parallel tensor on M is a constant multiple of the associated metric tensor and obtained some consequences of this. Vector fields keeping curvature tensor invariant are characterized on M .

1. Introduction

In 1923, Eisenhart [13] proved that if a positive definite Riemannian manifold (M, g) admits a second order parallel symmetric covariant tensor other than the constant multiple of metric tensor, then it is reducible. In 1926, Levy [18] proved that a second order parallel symmetric non-singular tensor in a space of constant curvature

* Corresponding Author.

Received November 14, 2019; revised July 18, 2020; accepted July 21, 2020.

2010 Mathematics Subject Classification: 53C15, 53C25, 53D10.

Key words and phrases: almost Kenmotsu manifold, second order parallel tensor, nullity distribution, homothetic vector field.

The author D.M. Naik is financially supported by University Grants Commission, New Delhi (Ref. No.:20/12/2015 (ii)EU-V) in the form of Junior Research Fellowship.

The authors V. Venkatesha and D.M. Naik are thankful to Department of Science and Technology, New Delhi for financial assistance to the Department of Mathematics, Kuvempu University under FIST program (Ref. No. SR/FST/MS-I/2018/23(C)).

is a constant multiple of the metric tensor. Using Ricci identities, Sharma in [24] gives a global approach to the Eisenhart problem, and generalized the Levy's theorem. This problem was studied by the same author in contact geometry [25, 26, 27] on different manifolds, for example for K-contact manifolds in [26]. Since then many geometers have investigated the Eisenhart problem on various contact manifolds: nearly Sasakian[30], P -Sasakian [7, 29], f -Kenmotsu manifold [4], $N(\kappa)$ -quasi Einstein manifold [6], 3-dimensional normal paracontact geometry [1], contact manifolds having non-vanishing ξ -sectional curvature [14], (κ, μ) -contact metric manifold [19], almost Kenmotsu manifolds [34] and 3-dimensional non-cosymplectic normal almost contact pseudo-metric manifold of non-vanishing ξ -sectional curvature [32].

In contact geometry, Kenmotsu manifolds, introduced by Kenmotsu in [17], are one of the important classes of manifolds. Such manifolds were observed to be normal. Let $(M, \varphi, \xi, \eta, g)$ be an almost Kenmotsu structure (see Section 2) on a $(2n + 1)$ -dimensional differentiable manifold. The purpose of this paper is to study second order parallel tensors on M under certain conditions. Throughout the paper, we suppose that the almost Kenmotsu manifold is of dimension $2n + 1$. Denoting the Ricci tensor by S , the tensor $\frac{1}{2}\mathcal{L}_\xi\varphi$ by h , where \mathcal{L} denotes the Lie differentiation, and the operator $R(\cdot, \xi)\xi$ by ℓ , we prove the following.

Theorem 1.1. *Let α be a second order symmetric parallel tensor, and A be the $(1, 1)$ -tensor metrically equivalent to α on an almost Kenmotsu manifold M . The following hold*

- (i) $A\xi = \alpha(\xi, \xi)\xi$, if M has non-vanishing ξ -sectional curvature;
- (ii) $trA = \alpha(\xi, \xi) - tr(A(h^2 - 2\varphi h - \varphi(\nabla_\xi h))) + \alpha(\xi, \xi)S(\xi, \xi)$;
- (iii) $tr(A\ell) = \alpha(\xi, \xi)S(\xi, \xi)$,

where tr denotes the trace.

Since the Ricci tensor S is a second order tensor, we have:

Corollary 1.1. *If an almost Kenmotsu manifold M is Ricci symmetric, and if M has non-vanishing ξ -sectional curvature, then*

- (i) $Q\xi = -(2n + trh^2)\xi$;
- (ii) the scalar curvature $r = \|Q\xi\|^2 - 2n - tr(h^2) - tr(Q(h^2 - 2\varphi h - \varphi(\nabla_\xi h)))$;
- (iii) $tr(Q\ell) = \|Q\xi\|^2$,

where Q is the Ricci operator determined by $S(X, Y) = g(QX, Y)$.

Theorem 1.2. *Let M be an almost Kenmotsu manifold having non-vanishing ξ -sectional curvature such that $tr\ell > -2n - 2$. The second order parallel tensor on M is a constant multiple of the associated metric tensor.*

Corollary 1.2. *If M is an almost Kenmotsu manifold having non-vanishing ξ -sectional curvature such that $tr\ell > -2n - 2$ is Ricci symmetric, then it is Einsteinian.*

Since $tr\ell = S(\xi, \xi) = -2n$ and the ξ -sectional curvature $K(\xi, X) = -1$ for Kenmotsu manifold, we have the following.

Corollary 1.3. *Any Ricci symmetric Kenmotsu manifold is Einsteinian.*

Corollary 1.4. *An affine Killing vector field on M which has non-vanishing ξ -sectional curvature such that $tr\ell > -2n - 2$ is homothetic.*

Blair, Koufogiorgos and Papantoniou [3] introduced (κ, μ) -nullity distributions on a contact metric manifold generalizing the notion of κ -nullity distributions by defining

$$N_p(\kappa, \mu) = \{Z \in T_pM : R(U, V)Z = \kappa[g(V, Z)U - g(U, Z)V] + \mu[g(V, Z)hU - g(U, Z)hV]\},$$

for any $p \in M$, where $\kappa, \mu \in \mathbb{R}$.

Corollary 1.5. *A second order parallel tensor on an almost Kenmotsu manifold with $\xi \in N(\kappa, \mu)$ is a constant multiple of the associated metric tensor.*

The above corollary has been proved by Wang and Liu in [34]. Recently, Dileo-Pastore [12] introduced $(\kappa, \mu)'$ -nullity distribution on an almost Kenmotsu manifold which is defined by

$$N_p(\kappa, \mu)' = \{Z \in T_pM : R(U, V)Z = \kappa[g(V, Z)U - g(U, Z)V] + \mu[g(V, Z)h'U - g(U, Z)h'V]\},$$

for any $p \in M$, where $h' = h \circ \varphi$ and $\kappa, \mu \in \mathbb{R}$. Here we recall the following results due to Dileo-Pastore [11, 12].

Lemma 1.1.([12, Proposition 4.1]) *Let M be an almost Kenmotsu manifold such that $h' \neq 0$ with $\xi \in N(\kappa, \mu)'$. Then $\kappa < -1$, $\mu = -2$ and $\text{spect}(h') = \{0, \lambda, -\lambda\}$ with 0 as simple eigenvalue and $\lambda = \sqrt{-1 - k}$.*

Lemma 1.2.([11, Theorem 6]) *Let M be a locally symmetric almost Kenmotsu manifold such that $h' \neq 0$ and $R(X, Y)\xi = 0$ for any $X, Y \in \mathcal{D}$, where $\mathcal{D} = \ker(\eta)$. Then, M is locally isometric to $\mathbb{H}^{n+1}(-4) \times \mathbb{R}^n$.*

Lemma 1.3.([12, Corollary 4.2]) *Let M be an almost Kenmotsu manifold such that $h' \neq 0$ with $\xi \in N(\kappa, \mu)'$. Then M is locally symmetric if and only if $\text{spect}(h') = \{0, 1, -1\}$ that is if and only if $k = -2$.*

Lemma 1.4.([12, Proposition 4.3]) *Let M be an almost Kenmotsu manifold such that $h' \neq 0$ with $\xi \in N(\kappa, \mu)'$. Then the ξ -sectional curvature satisfies*

$$K(\xi, X) = \begin{cases} \kappa - 2\lambda, & \text{if } X \in [\lambda]' \\ \kappa + 2\lambda, & \text{if } X \in [-\lambda]', \end{cases}$$

where $[\lambda]'$ denotes the eigenspace of h' related to the eigenvalue λ .

We use these to prove:

Theorem 1.3. *Let M be an almost Kenmotsu manifold with $\xi \in N(\kappa, \mu)'$ and $h' \neq 0$. If M admits a second order parallel tensor, then either the second order parallel tensor is a constant multiple of the associated metric tensor or M is locally isometric to $\mathbb{H}^{n+1}(-4) \times \mathbb{R}^n$.*

In [34], Wang and Liu proved the above theorem in another way. Now waving the hypothesis non-vanishing ξ -sectional curvature in Theorem 1.2 by $\nabla_{\xi}h = 0$, we prove:

Theorem 1.4. *Let M be an almost Kenmotsu manifold with $\nabla_{\xi}h = 0$ such that $tr\ell > -2n - 2$. Then the second order parallel tensor on M is a constant multiple of the associated metric tensor.*

In [23] Naik et al. proved that every vector field which leaves the curvature tensor invariant are Killing in a (κ, μ) -almost Kenmotsu manifold with $h' \neq 0$ and $\kappa \neq -2$. Here, as an application of Theorem 1.4, we prove the following.

Theorem 1.5. *Let M be an almost Kenmotsu manifold with $\nabla_{\xi}h = 0$ such that $tr\ell > -2n - 2$. Then every vector field keeping curvature tensor invariant are homothetic.*

The m -Bakry-Emery Ricci tensor is a natural extension of the Ricci tensor to smooth metric measure spaces and is given by

$$S_f^m = S + \text{Hess}f - \frac{1}{m}df \otimes df,$$

where f is a smooth function on M and m is an integer such that $0 < m \leq \infty$. If S_f^m is a constant multiple of the metric g , then the Riemannian manifold (M, g) is called m -quasi-Einstein manifold (see [5, 15] and the references therein). Now applying Theorem 1.2, Theorem 1.3 and Theorem 1.4, we deduce the following statement.

Theorem 1.6. *Let M be an almost Kenmotsu manifold either having non-vanishing ξ -sectional curvature such that $tr\ell > -2n - 2$ or $\xi \in N(\kappa, \mu)'$ and $h' \neq 0$ or $\nabla_{\xi}h = 0$ such that $tr\ell > -2n - 2$. Then the m -Bakry-Emery Ricci tensor $S_f^m = S + \text{Hess}f - \frac{1}{m}df \otimes df$ is parallel if and only if M is m -quasi-Einstein manifold.*

A Ricci soliton on a Riemannian manifold (M, g) is defined by

$$\mathcal{L}_V g + 2S + 2\rho g = 0,$$

where V is a smooth vector field and ρ is a constant. In the context of contact geometry and paracontact geometry, Ricci solitons are studied in [8, 9, 10, 20, 21, 22, 28, 31, 33]. In the similar vein as Theorem 1.6. we state the following.

Theorem 1.7. *Let M be an almost Kenmotsu manifold either having non-vanishing ξ -sectional curvature such that $tr\ell > -2n - 2$ or $\xi \in N(\kappa, \mu)'$ and $h' \neq 0$ or $\nabla_{\xi}h = 0$ such that $tr\ell > -2n - 2$. Then the the second order tensor $\mathcal{L}_V g + 2S$ is parallel if*

and only if M admits a Ricci soliton.

2. Preliminaries

An almost contact metric structure on a $(2n + 1)$ -Riemannian manifold M is a quadruple (φ, ξ, η, g) , where φ is an endomorphism, ξ a global vector field, η a 1-form and g a Riemannian metric, such that

$$(2.1) \quad \varphi^2 = -I + \eta \otimes \xi, \quad \eta(\xi) = 1,$$

$$(2.2) \quad g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y).$$

We easily obtain from (2.1) that $\varphi\xi = 0$ and $\eta \circ \varphi = 0$ (see [2]).

A manifold M with (φ, ξ, η, g) structure is said to be an almost contact metric manifold. We define the fundamental 2-form on $(M, \varphi, \xi, \eta, g)$ by $\Phi(X, Y) = g(X, \varphi Y)$. If the 1-form η is closed and $d\Phi = 2\eta \wedge \Phi$, then M is said to be an almost Kenmotsu manifold [16]. It is well known that a normal almost Kenmotsu manifold is a Kenmotsu manifold.

The two tensor fields $\ell := R(\cdot, \xi)\xi$ and $h := \frac{1}{2}\mathcal{L}_\xi\varphi$ are known to be symmetric and satisfy

$$(2.3) \quad \ell\xi = 0 \quad h\xi = 0, \quad trh = 0, \quad h\varphi + \varphi h = 0.$$

Further, one has the following formulas:

$$(2.4) \quad \nabla_\xi\varphi = 0,$$

$$(2.5) \quad \nabla_X\xi = -\varphi^2X - \varphi hX,$$

$$(2.6) \quad (\nabla_\xi h)X = -\varphi X - 2hX - \varphi h^2X - \varphi R(X, \xi)\xi,$$

$$(2.7) \quad R(X, \xi)\xi - \varphi R(\varphi X, \xi)\xi = 2(\varphi^2X - h^2X),$$

$$(2.8) \quad tr\ell = S(\xi, \xi) = -2n - trh^2.$$

3. Proof of Theorems

Now we prove the results stated in Section 1.

Proof of Theorem 1.1. Suppose that α is a symmetric $(0, 2)$ -tensor and A be the $(1, 1)$ -tensor metrically equivalent to α , that is, $g(AX, Y) = \alpha(X, Y)$. Note that $\nabla\alpha = 0$ implies $\nabla A = 0$, and so

$$R(X, Y)AZ = AR(X, Y)Z.$$

Therefore

$$g(R(X, Y)Z, AW) = -g(R(X, Y)AW, Z) = -g(R(X, Y)W, AZ).$$

Thus we get

$$\begin{aligned} g(R(Z, AW)X, Y) &= g(R(X, Y)Z, AW) \\ &= -g(R(X, Y)W, AZ) = -g(R(W, AZ)X, Y). \end{aligned}$$

Setting $X = Z = \xi$ in the above equation, we get

$$(3.1) \quad g(R(\xi, AW)\xi, Y) + g(R(W, A\xi)\xi, Y) = 0.$$

For $W = \xi$ and $Y = A\xi$, equation (3.1) becomes

$$(3.2) \quad g(R(\xi, A\xi)\xi, A\xi) = 0.$$

Since the ξ -sectional curvature $K(\xi, X)$ is non-vanishing, equation (3.2) implies that $A\xi = f\xi$, for some scalar function f on M . As $g(\xi, \xi) = 1$, we have

$$f = g(A\xi, \xi) = \alpha(\xi, \xi).$$

This proves (i). Applying φ to (2.6) shows that

$$(3.3) \quad R(\xi, X)\xi = h^2X - \varphi^2X - 2\varphi hX - \varphi(\nabla_\xi h)X.$$

Using (3.3) in (3.1), we get

$$g(h^2AW + AW - \eta(AW)\xi - 2\varphi hAW - \varphi(\nabla_\xi h)AW, Y) + g(R(W, A\xi)\xi, Y) = 0.$$

If $\{e_i\}$ is a local orthonormal basis, then putting $W = Y = e_i$ in the above equation and summing it over i , leads to

$$tr(h^2A) + trA - \alpha(\xi, \xi) - 2tr(\varphi hA) - tr(\varphi(\nabla_\xi h)A) + \alpha(\xi, \xi)S(\xi, \xi) = 0.$$

This gives (ii). Now plugging X by AX in (3.3) and then contracting with respect to X gives

$$tr(A\ell) = tr(\varphi(\nabla_\xi h)A) - trA + \alpha(\xi, \xi) + 2tr(\varphi hA) - tr(h^2A).$$

Using this in (ii) yields (iii). This finishes the proof. \square

Proof of Corollary 1.1. As the Ricci tensor S is parallel, it follows from Theorem 1.1 that

$$(3.4) \quad Q\xi = S(\xi, \xi)\xi.$$

Now (i) follows from (2.8). Note that from (3.4), we have

$$S(\xi, \xi)S(\xi, \xi) = g(Q\xi, Q\xi) = \|Q\xi\|^2.$$

Hence (ii) and (iii) follows directly from (ii) and (iii) of Theorem 1.1. \square

Proof of Theorem 1.2. Let $\{e_i, \varphi e_i, \xi\}_{i=1}^n$ be a local orthonormal basis such that $he_i = \lambda_i e_i$. Then $h\varphi e_i = -\lambda_i \varphi e_i$. Hence as $h\xi = 0$, we have

$$trh^2 = 2 \sum_{i=1}^n \lambda_i^2.$$

But by hypothesis, $tr\ell = -2n - trh^2 > -2n - 2$, and so $trh^2 < 2$. Therefore $\sum_{i=1}^n \lambda_i^2 < 1$ which means $\lambda_i^2 < 1$ for each i . This fact will be used in the rest of analysis for this theorem.

Let α be a $(0, 2)$ -tensor such that $\nabla\alpha = 0$, and A be the dual $(1, 1)$ -type tensor which is metrically equivalent to α , that is, $\alpha(X, Y) = g(AX, Y)$. We will analyse the symmetric and anti-symmetric cases of second order tensor separately.

First, suppose that α is symmetric. Then from item (i) of Theorem 1.1, we get

$$(3.5) \quad A\xi = \alpha(\xi, \xi)\xi.$$

To show $\alpha(\xi, \xi)$ is constant, we differentiate it along X to obtain

$$\begin{aligned} X(\alpha(\xi, \xi)) &= 2\alpha(\nabla_X \xi, \xi) = 2g(-\varphi^2 X - \varphi hX, A\xi) \\ &= 2\alpha(\xi, \xi)g(-\varphi^2 X - \varphi hX, \xi) = 0. \end{aligned}$$

Now differentiating (3.5) along X yields

$$A(-\varphi^2 X - \varphi hX) = \alpha(\xi, \xi)(-\varphi^2 X - \varphi hX).$$

Replacing X by φX , it follows that

$$(3.6) \quad A(\varphi X - hX) = \alpha(\xi, \xi)(\varphi X - hX).$$

Now putting $X = e_i$ and $X = \varphi e_i$ in (3.6) respectively gives

$$(3.7) \quad A(\varphi e_i) - \lambda_i A(e_i) = \alpha(\xi, \xi)\{\varphi e_i - \lambda_i e_i\},$$

and

$$(3.8) \quad \lambda_i A(\varphi e_i) - A(e_i) = \alpha(\xi, \xi)\{\lambda_i \varphi e_i - e_i\}.$$

Multiplying λ_i to (3.7) and then subtracting it with (3.8) shows

$$(\lambda_i^2 - 1)A(e_i) = \alpha(\xi, \xi)(\lambda_i^2 - 1)e_i.$$

Similarly, we can find

$$(\lambda_i^2 - 1)A(\varphi e_i) = \alpha(\xi, \xi)(\lambda_i^2 - 1)\varphi e_i.$$

Since $\lambda_i^2 < 1$, we have $A(e_i) = \alpha(\xi, \xi)e_i$ and $A(\varphi e_i) = \alpha(\xi, \xi)\varphi e_i$ for every $i = 1, \dots, n$. But $A\xi = \alpha(\xi, \xi)\xi$, and so

$$AX = \alpha(\xi, \xi)X,$$

for any $X \in TM$, that is, α is a constant multiple of g . Now suppose that α is skew-symmetric. Note that $\nabla\alpha = 0$ implies

$$g(R(X, Y)AZ, W) = g(AR(X, Y)Z, W).$$

The skew-symmetry of α then gives

$$(3.9) \quad g(R(X, Y)AZ, W) + g(R(X, Y)Z, AW) = 0.$$

For $X = A^2\xi$, $Y = Z = \xi$ and $W = A\xi$, we find

$$(3.10) \quad g(R(A^2\xi, \xi)\xi, A^2\xi) = 0.$$

From (3.10), the hypothesis $K(\xi, X)$ is non-vanishing imply that $A^2\xi = f\xi$. Clearly,

$$(3.11) \quad f = -g(A\xi, A\xi) = -\|A\xi\|^2.$$

One can easily verify the constancy of f by differentiating (3.11) along any vector field and getting the derivative 0. Now differentiating $A^2\xi = -\|A\xi\|^2\xi$ along φX yields

$$A^2(\varphi X - hX) = -\|A\xi\|^2(\varphi X - hX).$$

Next, we take $X = e_i$ and φe_i successively in the above equation and argue as before in order to obtain $A^2e_i = -\|A\xi\|^2e_i$ and $A^2(\varphi e_i) = -\|A\xi\|^2\varphi e_i$, for every $i = 1, 2, \dots, n$. So, $A^2X = -\|A\xi\|^2X$ for any X orthogonal to ξ . Hence, we obtain

$$A^2 = -\|A\xi\|^2I.$$

If $\|A\xi\| \neq 0$, then $J = \|A\xi\|^{-1}A$ defines a Kaehlerian structure on M leading to a contradiction that M is odd dimensional. Thus $\|A\xi\| = 0$, and so $A\xi = 0$. Differentiating it along φX gives

$$A(\varphi X - hX) = 0.$$

Now, as before we put $X = e_i$ and φe_i successively in the above equation to conclude that $AX = 0$, for any $X \perp \xi$. Since $A\xi = 0$, we obtain $AX = 0$ for any $X \in TM$. This completes the proof. \square

Proof of Corollary 1.4. A vector field V such that $\mathcal{L}_V\nabla = 0$ is called an affine Killing vector field. Note that $\mathcal{L}_V\nabla = 0$ is equivalent to $\nabla(\mathcal{L}_Vg) = 0$. Now the result follows from Theorem 1.2. \square

Proof of Corollary 1.5. Let M be an almost Kenmotsu manifold with $\xi \in N(\kappa, \mu)'$, that is

$$(3.12) \quad R(U, V)\xi = \kappa\{\eta(V)U - \eta(U)V\} + \mu\{\eta(V)hU - \eta(U)hV\},$$

for all $U, V \in TM$. It then follows from Dileo-Pastore [12] that $k = -1$ and $h = 0$. Hence (3.12) gives $tr\ell = -2n$, $K(\xi, X) = -1$ and so the conclusion follows from Theorem 1.2. \square

Proof of Theorem 1.3. Observe that, if X is an eigenvector of h with eigenvalue λ , and thus $h\varphi X = -\lambda\varphi X$, then $X + \varphi X$ is an eigenvector of h' with eigenvalue $-\lambda$, while $X - \varphi X$ is eigenvector with eigenvalue λ . Thus, it follows that h and h' admit the same eigenvalues.

First suppose that $\kappa \neq -2$. Note that, if X is such that $X \perp \xi$ and $hX = \lambda X$, then from Lemma , we see λ is different from $+1$ and -1 . Thus Lemma implies that $K(\xi, X)$ is non-vanishing. If the second order parallel tensor α is symmetric, then part (i) of Theorem 1.1 shows that

$$(3.13) \quad A\xi = \alpha(\xi, \xi)\xi.$$

That $\alpha(\xi, \xi)$ is constant, can be verified by differentiating it and getting the derivative equal to 0. Now differentiating (3.13) along X and φX , where $X \perp \xi$ and $hX = \lambda X$, gives

$$(\lambda^2 - 1)A(X) = \alpha(\xi, \xi)(\lambda^2 - 1)X,$$

and

$$(\lambda^2 - 1)A(\varphi X) = \alpha(\xi, \xi)(\lambda^2 - 1)\varphi X.$$

Since λ is different from $+1$ and -1 , we obtain

$$AX = \alpha(\xi, \xi)X,$$

for any $X \in TM$, that is, α is a constant multiple of g . Now suppose that α is skew-symmetric. Then as in the proof of Theorem 1.2, we obtain $A^2\xi = -\|A\xi\|^2$, where $\|A\xi\|^2$ is constant. Differentiating it along X and φX successively, where $X \perp \xi$ and $hX = \lambda X$, gives

$$(\lambda^2 - 1)A^2(X) = -\|A\xi\|^2(\lambda^2 - 1)X,$$

and

$$(\lambda^2 - 1)A^2(\varphi X) = -\|A\xi\|^2(\lambda^2 - 1)\varphi X.$$

Thus, we obtain

$$A^2 = -\|A\xi\|^2I.$$

Proceeding the similar manner as in the Theorem 1.2 one gets $AX = 0$ for any $X \in TM$.

Now suppose that $\kappa = -2$ and $\xi \in N(\kappa, \mu)'$, that is,

$$R(U, V)\xi = \kappa\{\eta(V)U - \eta(U)V\} + \mu\{\eta(V)h'U - \eta(U)h'V\}.$$

Then Lemma shows M is locally symmetric. Now it follows from Lemma that M is locally isometric to $\mathbb{H}^{n+1}(-4) \times \mathbb{R}^n$, since $R(X, Y)\xi = 0$ for any $X, Y \in \mathcal{D}$. This completes the proof. \square

Proof of Theorem 1.4. Let $\{e_i, \varphi e_i, \xi\}_{i=1}^n$ be a local orthonormal basis as considered in Theorem 1.2. If the second order parallel tensor α is symmetric, then

$$g(R(X, Y)Z, AW) + g(R(X, Y)W, AZ) = 0.$$

Putting $Y = Z = W = \xi$ in above, we get

$$(3.14) \quad g(R(X, \xi)\xi, A\xi) = 0.$$

Note that (2.6) takes the form

$$(3.15) \quad R(X, \xi)\xi = \varphi^2 X + 2\varphi hX - h^2 X.$$

Using (3.15) in (3.14), and putting $X = e_i$ and φe_i successively in the resulting equation gives

$$(1 + \lambda_i^2)g(e_i, A\xi) - 2\lambda_i g(\varphi e_i, A\xi) = 0,$$

and

$$(1 + \lambda_i^2)g(\varphi e_i, A\xi) - 2\lambda_i g(e_i, A\xi) = 0,$$

from which we obtain

$$\{(1 + \lambda_i^2)^2 - 4\lambda_i^2\}g(e_i, A\xi) = 0,$$

and

$$\{(1 + \lambda_i^2)^2 - 4\lambda_i^2\}g(\varphi e_i, A\xi) = 0.$$

Since λ_i is different from $+1$ and -1 , we get $g(X, A\xi) = 0$ for any $X \perp \xi$. Hence $A\xi = \alpha(\xi, \xi)\xi$. Now arguing in the similar manner as in Theorem 1.2, one can conclude that

$$AX = \alpha(\xi, \xi)X,$$

for any $X \in TM$, that is, α is a constant multiple of g . If α is skew-symmetric, then we have equation (3.9). Putting $Y = Z = \xi$ and $W = A\xi$ in (3.9), we find

$$g(R(X, \xi)\xi, A^2\xi) = 0.$$

Then using (3.15) in above, and putting $X = e_i$ and φe_i successively we get $g(X, A\xi) = 0$ for any $X \perp \xi$. Hence $A\xi = \alpha(\xi, \xi)\xi$, and similar to the proof of Theorem 1.2, we obtain $AX = 0$ for any $X \in TM$. This finishes the proof. \square

Proof of Theorem 1.5. Let $\{e_i, \varphi e_i, \xi\}_{i=1}^n$ be a local orthonormal basis such that $he_i = \lambda_i e_i$. The hypothesis $tr h > -2n - 2$ shows that $\sum_{i=1}^n \lambda_i^2 < 1$ which means $\lambda_i^2 < 1$, that is, $\lambda_i \neq (+1, -1)$ for each i .

Now the condition $\mathcal{L}_V R = 0$ implies

$$(\mathcal{L}_V g)(R(X, Y)Z, W) + (\mathcal{L}_V g)(R(X, Y)W, Z) = 0.$$

Let G be a $(1, 1)$ -tensor field defined by $g(GX, Y) = (\mathcal{L}_V g)(X, Y)$. Then

$$(3.16) \quad g(R(X, Y)Z, GW) + g(R(X, Y)W, GZ) = 0.$$

Taking $Y = Z = W = \xi$ in (3.16), we get

$$g(R(X, \xi)\xi, G\xi) = 0.$$

Using (3.15) in above, and putting $X = e_i$ and φe_i successively in the resulting equation gives

$$(1 + \lambda_i^2)g(e_i, G\xi) - 2\lambda_i g(\varphi e_i, G\xi) = 0,$$

and

$$(1 + \lambda_i^2)g(\varphi e_i, G\xi) - 2\lambda_i g(e_i, G\xi) = 0,$$

from which we obtain

$$\{(1 + \lambda_i^2)^2 - 4\lambda_i^2\}g(e_i, G\xi) = 0,$$

and

$$\{(1 + \lambda_i^2)^2 - 4\lambda_i^2\}g(\varphi e_i, G\xi) = 0.$$

Since λ_i is different from $+1$ and -1 , we get $g(X, G\xi) = 0$ for any $X \perp \xi$. Hence $G\xi = g(G\xi, \xi)\xi$. Now putting $Y = Z = \xi$ in (3.16), using (3.15), and taking $X = e_i$ and φe_i successively, we obtain

$$\{(1 + \lambda_i^2)^2 - 4\lambda_i^2\}(g(G\xi, \xi)g(e_i, W) - g(Ge_i, W)) = 0,$$

and

$$\{(1 + \lambda_i^2)^2 - 4\lambda_i^2\}(g(G\xi, \xi)g(\varphi e_i, W) - g(G\varphi e_i, W)) = 0.$$

So that, we have $GX = g(G\xi, \xi)X$ for any $X \perp \xi$, and hence

$$G = g(G\xi, \xi)I,$$

that is,

$$\mathcal{L}_V g = 2fg$$

for some function f . Thus V is a conformal vector field, and we have

$$(\mathcal{L}_V S)(Y, Z) = -(2n - 1)(\nabla_Y df)Z - (\text{div grad } f)g(X, Y).$$

Since $\mathcal{L}_V R = 0$ implies $\mathcal{L}_V S = 0$, the above equation gives $\nabla df = 0$, and hence

$$\nabla(df \otimes df) = 0.$$

Since $df \otimes df$ is a $(0, 2)$ -tensor, it follows from Theorem 1.2 that $df \otimes df = cg$, for some constant c . Thus

$$(3.17) \quad (Yf)\text{grad } f = cY,$$

which for $Y = \text{grad } f$ gives $\|\text{grad } f\|^2 \text{grad } f = c \text{grad } f$. Now, if $\{e_i\}$ is an orthonormal basis then putting $Y = e_i$ in (3.17), taking inner product with e_i and summing over i yields

$$\|\text{grad } f\|^2 = c(2n + 1).$$

Consequently, we obtain f is constant and hence V is homothetic. □

References

- [1] C. L. Bejan and M. Crasmareanu, *Second order parallel tensors and Ricci solitons in 3-dimensional normal paracontact geometry*, Ann Global Anal. Geom., **46**(2014), 117–127.
- [2] D. E. Blair, *Riemannian geometry of contact and symplectic manifolds*, Progress in Mathematics **203**, Birkhäuser, New York, 2010.
- [3] D. E. Blair, T. Koufogiorgos and B. J. Papantoniou, *Contact metric manifolds satisfying a nullity condition*, Israel J. Math., **91**(1995), 189–214.
- [4] C. Călin and M. Crasmareanu, *From the Eisenhart problem to Ricci solitons in f -Kenmotsu manifolds*, Bull. Malays. Math. Sci. Soc., **33**(2010), 361–368.
- [5] J. Case, Y. Shu and G. Wei, *Rigidity of quasi-Einstein metrics*, Differential Geom. Appl., **29**(2011), 93–100.
- [6] M. Crasmareanu, *Parallel tensors and Ricci solitons in $N(\kappa)$ -quasi Einstein manifolds*, Indian J. Pure Appl. Math., **43**(2012), 359–369.
- [7] U. C. De, *Second order parallel tensors on P -Sasakian manifolds*, Publ. Math. Debrecen, **49**(1996), 33–37.
- [8] U. C. De, S. K. Chaubey and Y. J. Suh, *A note on almost co-Kähler manifolds*, Int. J. Geom. Methods Mod. Phys., **17**(2020), 2050153, 14 pp.
- [9] U. C. De and K. Mandal, *Ricci solitons and gradient Ricci solitons on $N(\kappa)$ -paracontact manifolds*, Zh. Mat. Fiz. Anal. Geom., **15**(2019), 307–320.
- [10] U. C. De, Y. J. Suh and P. Majhi, *Ricci solitons on η -Einstein contact manifolds*, Filomat, **32**(2018), 4679–4687.
- [11] G. Dileo and A. M. Pastore, *Almost Kenmotsu manifolds and local symmetry*, Bull. Belg. Math. Soc. Simon Stevin, **14**(2007), 343–354.
- [12] G. Dileo and A. M. Pastore, *Almost Kenmotsu manifolds and nullity distributions*, J. Geom., **93**(2009), 46–61.
- [13] L. P. Eisenhart, *Symmetric tensors of the second order whose first covariant derivatives are zero*, Trans. Amer. Math. Soc., **25**(1923), 297–306.
- [14] A. Ghosh and R. Sharma, *Some results on contact metric manifolds*, Ann. Global Anal. Geom., **15**(1997), 497–507.
- [15] C. He, P. Petersen and W. Wylie, *On the classification of warped product Einstein metrics*, Comm. Anal. Geom., **20**(2012), 271–311.
- [16] D. Janssens and L. Vanhecke, *Almost contact structures and curvature tensors*, Kodai Math. J., **4**(1981), 1–27.
- [17] K. Kenmotsu, *A class of almost contact Riemannian manifolds*, Tohoku Math. J., **24**(1972), 93–103.
- [18] H. Levy, *Symmetric tensors of the second order whose covariant derivatives vanish*, Ann. of Math., **27**(1925), 91–98.
- [19] A. K. Mondal, U. C. De and C. Özgür, *Second order parallel tensors on (κ, μ) -contact metric manifolds*, An. St. Univ. Ovidius Const. Ser. Mat., **18**(2011), 229–238.

- [20] D. M. Naik and V. Venkatesha, η -Ricci solitons and almost η -Ricci solitons on para-Sasakian manifolds, *Int. J. Geom. Methods Mod. Phys.*, **16**(2019), 1950134, 18 pp.
- [21] D. M. Naik, V. Venkatesha and D. G. Prakasha, *Certain results on Kenmotsu pseudo-metric manifolds*, *Miskolc Math. Notes*, **20**(2019), 1083–1099.
- [22] D. M. Naik, V. Venkatesha and H. A. Kumara, *Ricci solitons and certain related metrics on almost co-Kaehler manifolds*, *J. Math. Phys. Anal. Geom.*, **16**(2020), 402–417.
- [23] D. M. Naik, V. Venkatesha and H. A. Kumara, *Some results on almost Kenmotsu manifolds*, *Note Mat.*, **40**(2020), 87–100.
- [24] R. Sharma, *Second order parallel tensor in real and complex space forms*, *Internat. J. Math. Math. Sci.*, **12**(1989), 787–790.
- [25] R. Sharma, *Second order parallel tensors on contact manifolds. I*, *Algebras Groups Geom.*, **7**(1990), 145–152.
- [26] R. Sharma, *Second order parallel tensors on contact manifolds. II*, *C. R. Math. Rep. Acad. Sci. Canada*, **13**(1991), 259–264.
- [27] R. Sharma, *On the curvature of contact metric manifolds*, *J. Geom.*, **53**(1995), 179–190.
- [28] Y. J. Suh and U. C. De, *Yamabe solitons and Ricci solitons on almost co-Kähler manifolds*, *Canad. Math. Bull.*, **62**(3)(2019), 653–661.
- [29] D. Tarafdar and U. C. De, *Second order parallel tensors on P-Sasakian manifolds*, *Northeast. Math. J.*, **11**(1995), 260–262.
- [30] M. Tarafdar and A. Mayra, *On nearly Sasakian manifold*, *An. Stiint. Univ. “Al. I. Cuza Iasi. Mat.*, **45**(1999), 291–294.
- [31] V. Venkatesha, H. A. Kumara and D. M. Naik, *Almost *-Ricci soliton on paraKenmotsu manifolds*, *Arab. J. Math.*, **9**(2020), 715–726.
- [32] V. Venkatesha and D. M. Naik, *On 3-dimensional normal almost contact pseudo-metric manifolds*, *Afr. Mat.*, (2020), (accepted).
- [33] V. Venkatesha, D. M. Naik and H. A. Kumara, **-Ricci solitons and gradient almost *-Ricci solitons on Kenmotsu manifolds*, *Math. Slovaca*, **69**(2019), 1447–1458.
- [34] Y. Wang and X. Liu, *Second order parallel tensors on an almost Kenmotsu manifolds satisfying the nullity distributions*, *Filomat*, **28**(2014), 839–847.