

ON DEFERRED CESÀRO MEAN IN PARANORMED SPACES

SINAN ERCAN

ABSTRACT. The aim of the present study is to introduce the concepts of deferred statistical convergence, deferred statistical Cauchy sequence and deferred Cesàro summability in paranormed spaces. We investigate some properties of these concepts and some inclusion relations with examples.

1. Introduction

Zygmund [26] introduced the idea of statistical convergence in 1935. Fast [8] and Steinhaus [18] introduced statistical convergence to assign a limit to sequences which are not convergent in the usual sense independently in the same year.

We begin by recalling the notion of asymptotic (or natural) density of a set $A \subset \mathbb{N}$ such that

$$\delta(A) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : k \in A\}|,$$

whenever the limit exists. $|\{\cdot\}|$ indicate the cardinality of the enclosed set. A sequence (x_k) of numbers is called statistically convergent to a number L provided that for $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : |x_k - L| \geq \varepsilon\}| = 0.$$

In this case, it is written by $S - \lim_{k \rightarrow \infty} x_k = L$. We note that throughout the paper $\mathbb{N} := \{1, 2, \dots\}$. The notion of statistical convergence is used an effective tool to resolve many problems in Ergodic theory, Fuzzy set theory, Trigonometric series and Banach spaces in the past years. Also various researchers studied applications and generalizations of this notion (see [17]- [9]).

Agnew [1] defined the deferred Cesàro mean of real (or complex) valued sequences by

$$(D_{p,q}x)_n = \frac{1}{q(n) - p(n)} \sum_{k=p(n)+1}^{q(n)} x_k,$$

Received November 12, 2020. Accepted February 16, 2021. Published online March 30, 2021.

2010 Mathematics Subject Classification: 40A05, 40A30, 41A25.

Key words and phrases: statistical convergence, deferred statistical convergence, paranormed space.

© The Kangwon-Kyungki Mathematical Society, 2021.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

where $p = \{p(n) : n \in \mathbb{N}\}$ and $q = \{q(n) : n \in \mathbb{N}\}$ are the sequences of nonnegative integers satisfying

$$(1.1) \quad p(n) < q(n) \text{ and } \lim_{n \rightarrow \infty} q(n) = \infty.$$

Recently, Küçükaslan and Yılmaztürk [13] introduced deferred statistical convergence. For some more relevant study, we refer to ([4]- [24]).

A paranorm $g : X \rightarrow \mathbb{R}$ is defined on a linear space X provided that for all $x, y, z \in X$

- (i) $g(x) = 0$ if $x = \theta$,
- (ii) $g(-x) = g(x)$,
- (iii) $g(x + y) \leq g(x) + g(y)$,
- (iv) if (α_n) is a sequence of scalars with $\alpha_n \rightarrow \alpha_0$ ($n \rightarrow \infty$) and $x_n, a \in X$ with $x_n \rightarrow a$ ($n \rightarrow \infty$) in the sense that $g(x_n - a) \rightarrow 0$ ($n \rightarrow \infty$), then $\alpha_n x_n \rightarrow \alpha_0 a$ ($n \rightarrow \infty$), in the sense that $g(\alpha_n x_n - \alpha_0 a) \rightarrow 0$ ($n \rightarrow \infty$).

A paranorm g for which $g(x) = 0$ implies that $x = \theta$ is said to be a total paranorm on X . (X, g) is said to be a total paranormed space. We recall that each seminorm (norm) g on X is a paranorm (total). However, the converse is not true.

DEFINITION 1.1 A sequence (x_k) in a paranormed space (X, g) is said to be convergent (or g -convergent) to a number L in (X, g) if for every $\varepsilon > 0$ there exists a positive integer k_0 such that $g(x_k - L) < \varepsilon$ whenever $k > k_0$. In this case we write $g - \lim_{k \rightarrow \infty} x_k = L$ and L is called the g -limit of (x_k) [2].

Recently, Alotaibi and Alroqi [2] studied strongly Cesàro summability, statistical convergence, statistical Cauchy in paranormed spaces. Then Alghamdi and Mursaleen [15] introduced λ -statistical convergence in paranormed spaces.

2. Main Results

In this section, we introduce the concepts of deferred statistical convergence, deferred statistical Cauchy and deferred Cesàro summability in paranormed space. Furthermore, we establish certain interesting results related to these concepts.

The deferred density of $K \subset \mathbb{N}$ is given by

$$\delta_{p,q}(K) = \lim_{n \rightarrow \infty} \frac{1}{q(n) - p(n)} |\{p(n) < k \leq q(n), k \in K\}|$$

whenever the limit exists. We note that the deferred density coincides with the natural density whenever $q(n) = n$ and $p(n) = 0$.

Now, we begin with the following definitions.

DEFINITION 2.1 A sequence (x_k) is said to be deferred statistically convergent (or $DS[p, q]$ -convergent) to L in (X, g) if for every $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{q(n) - p(n)} |\{p(n) < k \leq q(n) : g(x_k - L) \geq \varepsilon\}| = 0.$$

It is denoted by $g(DS[p, q]) - \lim_{k \rightarrow \infty} x_k = L$. The set of all such sequences is denoted by $g(DS[p, q])$.

(i) If we choose $q(n) = n$ and $p(n) = 0$, then Definition 2.1 is reduced to statistical convergence in paranormed spaces (cf. [2]).

(ii) If we choose $q(n) = \lambda_n$ and $p(n) = 0$, then Definition 2.1 is reduced to λ -statistical convergence in paranormed spaces (cf. [15]).

DEFINITION 2.2 A sequence (x_k) is said to be deferred statistically Cauchy sequence in (X, g) (or $g(DS[p, q])$ -Cauchy) if for every $\varepsilon > 0$ there exists a number $N = N(\varepsilon)$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{q(n) - p(n)} |\{p(n) < k \leq q(n) : g(x_k - x_N) \geq \varepsilon\}| = 0.$$

DEFINITION 2.3 A sequence (x_k) is said to be deferred strongly $D_{p,q}$ -convergent to L in (X, g) if

$$\lim_{n \rightarrow \infty} \frac{1}{q(n) - p(n)} \sum_{k=p(n)+1}^{q(n)} |g(x_k - L)| = 0$$

and we write it as $g(D[p, q]) - \lim_{k \rightarrow \infty} x_k = L$.

THEOREM 2.4 *If a sequence (x_k) is deferred statistically convergent in (X, g) then $g(DS[p, q]) - \lim$ is unique.*

Proof. Assume that $g(DS[p, q]) - \lim x = L_1$ and $g(DS[p, q]) - \lim x = L_2$. For given $\varepsilon > 0$, define the sets $K_1(\varepsilon)$ and $K_2(\varepsilon)$ by

$$\begin{aligned} K_1(\varepsilon) &= \{p(n) < k \leq q(n) : g(x_k - L_1) \geq \varepsilon/2\}, \\ K_2(\varepsilon) &= \{p(n) < k \leq q(n) : g(x_k - L_2) \geq \varepsilon/2\}. \end{aligned}$$

Since $g(D[p, q]) - \lim x = L_1$, we have $\delta(K_1(\varepsilon)) = 0$. Similarly, $g(DS[p, q]) - \lim x = L_2$ implies that $\delta(K_2(\varepsilon)) = 0$. Now, let $K(\varepsilon) = K_1(\varepsilon) \cup K_2(\varepsilon)$. Then $\delta(K(\varepsilon)) = 0$ and $\delta(K^c(\varepsilon)) = 1$. Now, if $k \in \mathbb{N} \setminus K(\varepsilon)$, then we have $g(L_1 - L_2) \leq g(x_k - L_1) + g(x_k - L_2) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$. Since $\varepsilon > 0$ is arbitrary, we get $g(L_1 - L_2) = 0$ and hence $L_1 = L_2$. □

THEOREM 2.5 *If $g - \lim x = L$. Then $g(DS[p, q]) - \lim x = L$ but the converse is not true.*

Proof. Assume that $g - \lim x = L$. Then there exists $N \in \mathbb{N}^+$ such that $g(x_n - L) < \varepsilon$ for all $n \geq N$ and $\varepsilon > 0$. Since $A(\varepsilon) = \{k \in \mathbb{N} : g(x_k - L) \geq \varepsilon\} \subset \{1, 2, 3, \dots\}$, $\delta(A(\varepsilon)) = 0$. Hence $g[DS[p, q]) - \lim x = L$. □

The following example shows that the converse need not be true.

EXAMPLE 2.6 Let $X = \ell(1/k) = \left\{ x = (x_k) : \sum_{k=1}^{\infty} |x_k|^{1/k} < \infty \right\}$ with the paranorm $g(x) = \left(\sum_{k=1}^{\infty} |x_k|^{1/k} \right)$. Define the sequence

$$x_k = \begin{cases} k, & \text{if } k = n^2, n \in \mathbb{N}; \\ 0, & \text{otherwise;} \end{cases}$$

and write

$$K(\varepsilon) = \{k \leq n : g(x_k) \geq \varepsilon\}, \quad 0 < \varepsilon < 1.$$

We see that

$$g(x_k) = \begin{cases} k^{1/k}, & \text{if } k = n^2, n \in \mathbb{N}; \\ 0, & \text{otherwise;} \end{cases}$$

and hence

$$\lim_{k \rightarrow \infty} g(x_k) = \begin{cases} 1, & \text{if } k = n^2, n \in \mathbb{N}; \\ 0, & \text{otherwise;} \end{cases}$$

Therefore $g - \lim_{k \rightarrow \infty} x_k$ does not exist. On the other hand $\delta_{p,q}(K(\varepsilon)) = 0$, that is, $g(DS[p, q]) - \lim_{k \rightarrow \infty} x_k = 0$.

THEOREM 2.7 Let $g(DS[p, q]) - \lim x = L_1$ and $g(DS[p, q]) - \lim y = L_2$. Then
 (i) $g(DS[p, q]) - \lim (x \pm y) = L_1 \pm L_2$.
 (ii) $g(DS[p, q]) - \lim \alpha x = \alpha L, \alpha \in \mathbb{R}$.

THEOREM 2.8 A sequence (x_k) in (X, g) is statistically convergent to L if and only if there exists a set $K = \{k_1 < k_2 < k_3 < \dots < k_n < \dots\} \subseteq \mathbb{N}$ with $\delta_{p,q}(K) = 1$ such that $g(x_{k_n} - L) \rightarrow 0$ ($n \rightarrow \infty$).

Proof. Assume that (x_k) is deferred statistically convergent to L , that is, $g(DS[p, q]) - \lim_{k \rightarrow \infty} x_k = L$. Now, write $K_r = \{n \in \mathbb{N} : g(x_{k_n} - L) \geq \frac{1}{r}\}$, $M_r = \{n \in \mathbb{N} : g(x_{k_n} - L) < \frac{1}{r}\}$ for $r = 1, 2, \dots$. Then $\delta_{p,q}(K_r) = 0$,

$$(2.1) \quad M_1 \supset M_2 \supset \dots \supset M_i \supset M_{i+1} \supset \dots,$$

$$(2.2) \quad \delta_{p,q}(M_r) = 1, \quad r = 1, 2, \dots$$

We have to show that, for $n \in M_r$, (x_{k_n}) is g -convergent to L . On contrary suppose that (x_{k_n}) is not g -convergent to L . Therefore, there is $\varepsilon > 0$ such that $g(x_{k_n} - L) \geq \varepsilon$ for infinitely many terms. Let $M_\varepsilon = \{n \in \mathbb{N} : g(x_{k_n} - L) < \varepsilon\}$ and $\varepsilon > 1/r, r \in \mathbb{N}$. Then $\delta_{p,q}(M_\varepsilon) = 0$ and by (2.1), $M_r \subset M_\varepsilon$. Hence, $\delta_{p,q}(M_r) = 0$. This contradicts (2.2) and hence (x_{k_n}) is g -convergent to L . \square

Now, consider the set $K = \{k_1 < k_2 < k_3 < \dots\} \subset \mathbb{N}$ with $\delta_{p,q}(K) = 1$ and $g(x_{k_n} - L) \rightarrow 0$ ($n \rightarrow \infty$). So we can find a positive integer n_0 such that $g(x_k - L) < \varepsilon$ for $n \geq n_0$. $K_\varepsilon = \{k : g(x_k - L) \geq \varepsilon\} \subseteq \mathbb{N} - \{k_{n_0+1}, k_{n_0+2}, k_{n_0+3}, \dots\}$ and therefore $\delta_{p,q}(K_\varepsilon) = 0$. This shows that (x_k) is deferred statistically convergent to L in (X, g) .

THEOREM 2.9 A sequence (x_k) in a complete paranormed space (X, g) is deferred statistically Cauchy if and only if it is deferred statistically convergent.

Proof. Assume that (x_k) is $g(DS[p, q])$ -Cauchy but not $g(DS[p, q])$ -convergent. Then, we have $m \in \mathbb{N}$ such that $\delta_{p,q}(G(\varepsilon)) = 0$, where $G(\varepsilon) = \{n \in \mathbb{N} : g(x_n - x_m) \geq \varepsilon\}$ and $\delta_{p,q}(D(\varepsilon)) = 0$, where $D(\varepsilon) = \{n \in \mathbb{N} : g(x_n - L) < \varepsilon/2\}$, i.e., $\delta_{p,q}(D^C(\varepsilon)) = 1$. If $g(x_n - L) < \frac{\varepsilon}{2}$, then $g(x_n - x_m) \leq 2g(x_n - L) < \varepsilon$. Moreover, $\delta_{p,q}(G^C(\varepsilon)) = 0$, i.e., $\delta(G(\varepsilon)) = 1$, which leads to a contradiction, since (x_k) is $g(DS[p, q])$ -Cauchy. Hence (x_k) must be $g(DS[p, q])$ -convergent.

Conversely, let us assume that $g(DS[p, q])\text{-}\lim_{k \rightarrow \infty} x_k = L$. Then, we have $\delta(K(\varepsilon)) = 0$ where

$$K(\varepsilon) = \left\{ n \in \mathbb{N} : g(x_n - L) \geq \frac{\varepsilon}{2} \right\}.$$

This implies that

$$\delta(\mathbb{N} \setminus K(\varepsilon)) = \delta\left(\left\{ n \in \mathbb{N} : g(x_n - L) < \frac{\varepsilon}{2} \right\}\right) = 1.$$

Let $m, n \notin K(\varepsilon)$, then $g(x_m - x_n) < \varepsilon$. Let

$$M(\varepsilon) = \{n \in \mathbb{N} : g(x_m - x_n) < \varepsilon\}$$

for a fix $m \notin K(\varepsilon)$. Then $\mathbb{N} \setminus K(\varepsilon) \subset M(\varepsilon)$. Hence

$$1 = \delta(\mathbb{N} \setminus K(\varepsilon)) \leq \delta(M(\varepsilon)) \leq 1.$$

This implies $\delta(\mathbb{N} \setminus M(\varepsilon)) = 0$, where $\mathbb{N} \setminus M(\varepsilon) = \{n \in \mathbb{N} : g(x_m - x_n) \geq \varepsilon\}$. This implies that (x_n) is deferred statistically Cauchy in (X, g) . \square

THEOREM 2.10 *A sequence (x_k) in (X, g) is strongly $D_{p,q}$ -convergent to L , then (x_k) is deferred statistically convergent to L .*

Proof. Assume that (x_k) is strongly $D_{p,q}$ -convergent to L . For an arbitrary $\varepsilon > 0$, following equality

$$\begin{aligned} \frac{1}{q(n) - p(n)} \sum_{k=p(n)+1}^{q(n)} g(x_k - L) &= \frac{1}{q(n) - p(n)} \left(\sum_{\substack{k=p(n)+1 \\ g(x_k - L) \geq \varepsilon}}^{q(n)} + \sum_{\substack{k=p(n)+1 \\ g(x_k - L) < \varepsilon}}^{q(n)} \right) g(x_k - L) \\ &\geq \frac{1}{q(n) - p(n)} \sum_{\substack{k=p(n)+1 \\ g(x_k - L) \geq \varepsilon}}^{q(n)} g(x_k - L) \\ &\geq \varepsilon \frac{1}{q(n) - p(n)} |p(n) < k \leq q(n), g(x_k - L) \geq \varepsilon| \end{aligned}$$

holds. As $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} \frac{1}{q(n) - p(n)} |p(n) < k \leq q(n), g(x_k - L) \geq \varepsilon| = 0.$$

Therefore, desired result is obtained. \square

THEOREM 2.11 *If $(x_k) \in (X, g)$ is deferred statistically convergent to L and $(x_k) \in \ell_\infty$, then (x_k) is strongly $D_{p,q}$ -convergent to L .*

Proof. Suppose that $(x_k) \in \ell_\infty$ and (x_k) is deferred statistically convergent to L in (X, g) . For arbitrary $\varepsilon > 0$, we have $\delta_{p,q}(K_\varepsilon) = 0$. Since $x \in \ell_\infty$, there exists $M > 0$ such that $g(x_k - L) \leq M$ ($k = 1, 2, 3, \dots$). We have

$$\begin{aligned} \frac{1}{q(n) - p(n)} \sum_{k=p(n)+1}^{q(n)} g(x_k - L) &= \frac{1}{q(n) - p(n)} \sum_{\substack{k=p(n)+1 \\ k \notin K_\varepsilon}}^{q(n)} g(x_k - L) \\ &\quad + \frac{1}{q(n) - p(n)} \sum_{\substack{k=p(n)+1 \\ k \in K_\varepsilon}}^{q(n)} g(x_k - L) \\ &= s_1(n) + s_2(n) \end{aligned}$$

where

$$s_1(n) = \frac{1}{q(n) - p(n)} \sum_{\substack{k=p(n)+1 \\ k \notin K_\varepsilon}}^{q(n)} g(x_k - L) \text{ and } s_2(n) = \frac{1}{q(n) - p(n)} \sum_{\substack{k=p(n)+1 \\ k \in K_\varepsilon}}^{q(n)} g(x_k - L).$$

Now if $k \notin K_\varepsilon$, then $s_1(n) < \varepsilon$. For $k \in K_\varepsilon$, since $\delta_{p,q}(K_\varepsilon) = 0$ we have

$$s_2(n) \leq (\sup g(x_k - L)) \left(\frac{|K_\varepsilon|}{q(n) - p(n)} \right) \leq M \frac{|K_\varepsilon|}{q(n) - p(n)} \rightarrow 0,$$

as $n \rightarrow \infty$. This inequality completes the proof. \square

THEOREM 2.12 *If (x_k) in (X, g) is statistically convergent to L and $\left(\frac{p(n)}{q(n)-p(n)}\right) \in \ell_\infty$, then it is deferred statistically convergent to L .*

Proof. Assume that (x_k) is statistically convergent to L in (X, g) . Hence we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : g(x_k - L) \geq \varepsilon\}| = 0$$

for every $\varepsilon > 0$. From (1.1), we obtain

$$(2.3) \quad \lim_{n \rightarrow \infty} \frac{|\{k \leq q(n) : g(x_k - L) \geq \varepsilon\}|}{q(n)} = 0.$$

Moreover, we have

$$\{p(n) < k \leq q(n) : g(x_k - L) \geq \varepsilon\} \subset \{k \leq q(n) : g(x_k - L) \geq \varepsilon\}$$

and

$$|\{p(n) < k \leq q(n) : g(x_k - L) \geq \varepsilon\}| \leq |\{k \leq q(n) : g(x_k - L) \geq \varepsilon\}|.$$

Consequently

$$\begin{aligned} &\frac{1}{q(n) - p(n)} |p(n) < k \leq q(n) : g(x_k - L) \geq \varepsilon| \\ &\leq \left(1 + \frac{p(n)}{q(n) - p(n)}\right) \cdot \frac{1}{q(n)} |k \leq q(n) : g(x_k - L) \geq \varepsilon|. \end{aligned}$$

Thus using (2.3), we conclude that (x_k) is deferred statistically convergent to L in (X, g) . \square

COROLLARY 2.13 *Let $q(n)$ be an arbitrary sequence with $q(n) < n$ for all $n \in \mathbb{N}$ and $\left(\frac{n}{q(n)-p(n)}\right) \in \ell_\infty$. Then, statistical convergence implies deferred statistical convergence in (X, g) .*

THEOREM 2.14 *Let $p' = (p'(n))$ and $q' = (q'(n))$ be sequences of positive natural numbers satisfying*

$$(2.4) \quad p(n) \leq p'(n) < q'(n) \leq q(n)$$

and $\{k : p(n) < k \leq p'(n)\}, \{k : q'(n) < k \leq q(n)\}$ are finite sets for all $n \in \mathbb{N}$. Then, $g(DS[p', q'])$ -convergence implies $g(DS[p, q])$ -convergence in (X, g) .

Proof. Let us consider (x_k) is $g(DS[p', q'])$ -convergent to L . For an arbitrary $\varepsilon > 0$, the inequality

$$\begin{aligned} \{k : p(n) < k \leq q(n), g(x_k - L) \geq \varepsilon\} &= \{k : p(n) < k \leq p'(n), g(x_k - L) \geq \varepsilon\} \\ &\cup \{k : p'(n) < k \leq q'(n), g(x_k - L) \geq \varepsilon\} \\ &\cup \{k : q'(n) < k \leq q(n), g(x_k - L) \geq \varepsilon\} \end{aligned}$$

and

$$\begin{aligned} &\frac{1}{q(n) - p(n)} |\{k : p(n) < k \leq q(n), g(x_k - L) \geq \varepsilon\}| \\ &\leq \frac{1}{q'(n) - p'(n)} |\{k : p(n) < k \leq p'(n), g(x_k - L) \geq \varepsilon\}| \\ &\quad + \frac{1}{q'(n) - p'(n)} |\{k : p'(n) < k \leq q'(n), g(x_k - L) \geq \varepsilon\}| \\ &\quad + \frac{1}{q'(n) - p'(n)} |\{k : q'(n) < k \leq q(n), g(x_k - L) \geq \varepsilon\}| \end{aligned}$$

hold. As $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} \frac{1}{q(n) - p(n)} |\{k : p(n) < k \leq q(n), g(x_k - L) \geq \varepsilon\}| = 0.$$

This completes the proof. □

THEOREM 2.15 *Let $p' = (p'(n))$ and $q' = (q'(n))$ be sequences of positive natural numbers satisfying (2.4) and*

$$\lim_{n \rightarrow \infty} \frac{q(n) - p(n)}{q'(n) - p'(n)} > 0.$$

Then, $g(DS[p, q])$ -statistical convergence implies $g(DS[p', q'])$ -statistical convergence in (X, g) .

Proof. The inclusion

$$\{k : p'(n) < k \leq q'(n) : g(x_k - L) \geq \varepsilon\} \subset \{k : p(n) + 1 \leq k \leq q(n) : g(x_k - L) \geq \varepsilon\}$$

holds. Then we have the inequality

$$|\{k : p'(n) < k \leq q'(n) : g(x_k - L) \geq \varepsilon\}| \leq |\{k : p(n) + 1 \leq k \leq q(n) : g(x_k - L) \geq \varepsilon\}|.$$

Hence , we have

$$\begin{aligned} & \frac{1}{q'(n) - p'(n)} |\{k : p'(n) < k \leq q'(n) : g(x_k - L) \geq \varepsilon\}| \\ & \leq \frac{q(n) - p(n)}{q'(n) - p'(n)} \cdot \frac{1}{q(n) - p(n)} |\{k : p(n) + 1 < k \leq q(n) : g(x_k - L) \geq \varepsilon\}|. \end{aligned}$$

As $n \rightarrow \infty$, the desired result is obtained. \square

References

- [1] R. P. Agnew, *On deferred Cesàro Mean*, Comm. Ann. Math. **33** (1932), 413–421.
- [2] A. Alotaibi, A. M. Alroqi, *Statistical convergence in a paranormed space*, J. Inequal. Appl. 2012, 2012:**39**, 6 pp.
- [3] M. Altınok, B. Inan, M. Küçükaslan, *On deferred statistical convergence of sequences of sets in metric space*, TJMCS, Article ID 20150050, (2015), 9 pages.
- [4] M. Candan, A. Güneş, *Paranormed sequence space of non-absolute type founded using generalized difference matrix*, Proc. Nat. Acad. Sci. India Sect. A, 85 **2** (2015), 269–276.
- [5] M. Candan, *A new perspective on paranormed Riesz sequence space of non-absolute type*, Glob. J. Math. Anal., 3 **4** (2015), 150–163.
- [6] J. S. Connor, *The statistical and strong p -Cesàro of sequences*, Analysis, **8** (1988), 47–63.
- [7] S. Ercan, Y. Altın, M. Et, V. K. Bhardwaj, *On deferred weak statistical convergence*, J. Anal., (2020), <https://doi.org/10.1007/s41478-020-00221-5>.
- [8] H. Fast, *Sur la convergence statistique*, Colloq. Math., **2** (1951), 241–244.
- [9] J. A. Fridy, *On statistical convergence*, Analysis, **5** (1985), 301–313.
- [10] M. İlkan, E. E. Kara, *A new type of statistical Cauchy sequence and its relation to Bourbaki completeness*, Cogent Math. Stat., 5:1, 1487500 (2018), 9 pages.
- [11] M. İlkan, E. E. Kara, *On statistical convergence in quasi-metric spaces*, Demonstr. Math. **52** (1) (2019), 225–236.
- [12] C. Koşar, M. Küçükaslan, M. Et, *On asymptotically deferred statistical equivalence of sequences*, Filomat, 31:**16** (2017), 5139–5150.
- [13] M. Küçükaslan, M. Yılmaztürk, *On deferred statistical convergence of sequences*, Kyungpook Math. J., no. 2, 56 (2016), 357–366.
- [14] I. J. Maddox, *Elements of Functional Analysis*, Cambridge at the University Press, (1970).
- [15] A. Mohammed, M. Mursaleen, *λ -statistical convergence in paranormed space*, Abstr. Appl. Anal. 2013, Art. ID 264520, 5 pp.
- [16] H. Roopaei, T. Yaying, *Quasi-Cesàro matrix and associated sequence space*, Turkish J. Math. **45** (1) (2021), 153–166.
- [17] T. Šalát, *On statistically convergent sequences of real numbers*, Math. Slovaca **30** (1980), 139–150.
- [18] H. Steinhaus, *Sur la convergence ordinaire et la convergence asymptotique*, Colloq. Math. **2** (1951), 73–74.
- [19] F. Temizsu, M. Et, M. Çınar, *Δ^m -deferred statistical convergence of order α* , Filomat 30 **3** (2016), 667–673.
- [20] N. Turan, E. E. Kara, M. İlkan, *Quasi statistical convergence in cone metric spaces*, Facta Univ. Ser. Math. Inform. **33** (4) (2018), 613–626.
- [21] T. Yaying, B. Hazarika, *Lacunary arithmetic statistical convergence*, Natl. Acad. Sci. Lett., **43** (2020), 547–551.
- [22] T. Yaying, B. Hazarika, M. Mursaleen, *On sequence space defined by the domain of q -Cesàro matrix in ℓ_p space and the associated operator ideal*, J. Math. Anal. Appl. **493** (1) (2021), 124453.
- [23] T. Yaying, *Paranormed Riesz difference sequence spaces of fractional order*, Kragujevac J. Math. **46** (2) (2022), 175–191.
- [24] T. Yaying, B. Hazarika, A. Esi, *Geometric properties and compact operators on fractional Riesz difference spaces*, Kragujevac J. Math. **47** (4) (2023), 545–566.

- [25] M. Yılmaztürk, M. Küçükaslan, *On strongly defereed Cesàro summability and deferred statistical convergence of the sequences*, Bitlis Eren Univ. J. Sci. Technology **3** (2011), 22–25.
- [26] A. Zygmund, *Trigonometrical Series*, Monogr. Mat. **5**. Warszawa-Lwow 1935.

Sinan Ercan

Department of Mathematics, Fırat University, Elazığ, Turkey
E-mail: snnercan@gmail.com