



A NOTE ON DEGENERATE LAH-BELL POLYNOMIALS ARISING FROM DERIVATIVES

Xiangfan Piao¹, Yunjae Kim² and Jongkyum Kwon³

¹ Department of Mathematics

Kyungpook National University, Daegu 41566, Korea

e-mail: piaoxf76@hanmail.net

² Department of Mathematics

Kyungpook National University, Daegu 41566, Korea

e-mail: kimholzi@gmail.com

³ Department of Mathematics Education

Gyeongsang National University, Jinju, 52828, Korea

e-mail: mathkjk26@gnu.ac.kr

Abstract. Recently, Kim-Kim introduced Lah-Bell polynomials and numbers, and investigated some properties and identities of these polynomials and numbers. Kim studied Lah-Bell polynomials and numbers of degenerate version. In this paper, we study degenerate Lah-Bell polynomials arising from differential equations. Moreover, we investigate the phenomenon of scattering of the zeros of these polynomials.

1. INTRODUCTION

Definition 1.1. The unsigned Lah number $L_{n,k}$ counts the number of ways a set of n elements can be partitioned into k nonempty linearly ordered subsets, and have an explicit formula (see [1, 13, 15, 16, 24, 25]).

$$L_{n,k} = \binom{n-1}{k-1} \frac{n!}{k!} = \binom{n}{k} \frac{(n-1)!}{(k-1)!}. \quad (1.1)$$

⁰Received January 28, 2021. Revised April 2, 2021. Accepted August 6, 2021.

⁰2010 Mathematics Subject Classification: 05A19, 05A40, 11B73, 11B83.

⁰Keywords: Lah number, Lah-Bell polynomials, Stirling numbers of first(second) kinds, roots distributions.

⁰Corresponding author: Jongkyum Kwon(mathkjk26@gnu.ac.kr).

The generating function of $L_{n,k}$ is defined by (see [13, 15, 24])

$$\frac{1}{k!} \left(\frac{t}{1-t} \right)^k = \sum_{n=k}^{\infty} L_{n,k} \frac{t^n}{n!} \quad (k \geq 0). \quad (1.2)$$

Recently, Kim-Kim (see [13]) introduced the Lah-Bell polynomials as follows:

$$e^{\frac{tx}{1-t}} = \sum_{n=0}^{\infty} B_n^L(x) \frac{t^n}{n!}. \quad (1.3)$$

The degenerate exponential function is defined by (see [19–22])

$$e_{\lambda}^x(t) = (1 + \lambda t)^{\frac{x}{\lambda}}, \quad e_{\lambda}(t) := e_{\lambda}^1(t) = (1 + \lambda t)^{\frac{1}{\lambda}}, \quad (1.4)$$

where λ is a nonzero parameter.

Note that

$$\lim_{\lambda \rightarrow 0} e_{\lambda}^x(t) = \lim_{\lambda \rightarrow 0} (1 + \lambda t)^{\frac{x}{\lambda}} = e^{xt}.$$

Since $e_{\lambda}^x(t)$ defined in (1.4) is infinitely differentiable at $t = 0$, Taylor expansion of $e_{\lambda}^x(t)$ at $t = 0$ gives the following series form (see [20–23]):

$$e_{\lambda}^x(t) = \sum_{n=0}^{\infty} (x)_{n,\lambda} \frac{t^n}{n!}, \quad (1.5)$$

where $(x)_{n,\lambda}$ is defined by

$$(x)_{n,\lambda} = \begin{cases} 1, & n = 0, \\ (x - (n-1)\lambda)(x)_{n-1,\lambda}, & n \geq 1. \end{cases}$$

As is well known, the Stirling numbers of the first kind are given by (see [1–5, 10, 14])

$$(x)_n = \sum_{l=0}^n S_1(n, l) x^l, \quad (1.6)$$

where $(x)_n$ are defined by

$$(x)_n = \begin{cases} 1, & n = 0, \\ (x + 1 - n)(x)_{n-1}, & n \geq 1. \end{cases}$$

From (1.6), we easily get

$$\frac{1}{k!} (\ln(1+t))^k = \sum_{n=k}^{\infty} S_1(n, k) \frac{t^n}{n!}. \quad (1.7)$$

As an inversion formula of (1.6), the Stirling numbers of the second kind are given by (see [1–5, 10])

$$x^n = \sum_{l=0}^n S_2(n, l)(x)_l. \tag{1.8}$$

From (1.8), we get

$$\frac{1}{k!} (e^t - 1)^k = \sum_{n=k}^{\infty} S_2(n, k) \frac{t^n}{n!}. \tag{1.9}$$

Moreover, as degenerate version of (1.6) and (1.8), the degenerate Stirling numbers of the first and second kinds, respectively, are given by

$$(x)_n = \sum_{l=0}^n S_{1,\lambda}(n, l)(x)_{l,\lambda}, \tag{1.10}$$

$$(x)_{n,\lambda} = \sum_{l=0}^n S_{2,\lambda}(n, l)(x)_l$$

and

$$\frac{1}{k!} (\ln_{\lambda}(1+t))^k = \sum_{n=k}^{\infty} S_{1,\lambda}(n, k) \frac{t^n}{n!}, \tag{1.11}$$

$$\frac{1}{k!} (e_{\lambda}(t) - 1)^k = \sum_{n=k}^{\infty} S_{2,\lambda}(n, k) \frac{t^n}{n!},$$

where $\ln_{\lambda}(t) = \frac{t^{\lambda}-1}{\lambda}$ (see [6–12, 17, 18]).

Recently, Kim-Kim (see [13]) introduced the degenerate Lah-Bell polynomials which are given by the generating function to be

$$e^x_{\lambda} \left(\frac{1}{1-t} - 1 \right) = \sum_{n=0}^{\infty} B_{n,\lambda}^L(x) \frac{t^n}{n!}. \tag{1.12}$$

The rest of this paper is organized as follows. In the section 2, we study the differential equations on degenerate Lah-Bell polynomials. Using these differential equations, we derive some identities and properties of the degenerate Lah-Bell polynomials. In the section 3, we investigate the phenomenon of scattering of the zeros of those polynomials. Finally, in section 4, a summary of the Lah-Bell polynomials is given.

2. SOME IDENTITIES OF THE DEGENERATE LAH-BELL POLYNOMIAL

In this section, we derive some identities of the degenerate Lah-Bell polynomials. When $x = 1$, $B_{n,\lambda}^L := B_{n,\lambda}^L(1)$ are called the degenerate Lah-Bell

numbers. The following theorem gives an explicit expression of the Lah-Bell polynomials.

Theorem 2.1. *For non-negative integer $n \geq 0$, we have*

$$B_{n,\lambda}^L(x) = \sum_{k=0}^n (x)_{k,\lambda} L_{n,k}. \quad (2.1)$$

Proof. Combining (1.12) and (1.5) and using the fact (1.2), one can obtain the following relation (see [13]):

$$\begin{aligned} \sum_{n=0}^{\infty} B_{n,\lambda}^L(x) \frac{t^n}{n!} &= e^x \left(\frac{t}{1-t} \right) = \sum_{k=0}^{\infty} (x)_{k,\lambda} \frac{1}{k!} \left(\frac{t}{1-t} \right)^k \\ &= \sum_{k=0}^{\infty} (x)_{k,\lambda} \sum_{n=k}^{\infty} L_{n,k} \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n (x)_{k,\lambda} L_{n,k} \right) \frac{t^n}{n!}. \end{aligned} \quad (2.2)$$

Since the set $\{1, t, t^2, \dots, t^n, \dots\}$ are linearly independent, the relation (2.1) is satisfied. \square

Combining Theorem 2 and the definition of Lah number (1.1), one can have the following corollary.

Corollary 2.2. *For non-negative integer $n \geq 0$, we have*

$$B_{n,\lambda}^L(x) = n! \sum_{k=0}^n \frac{(x)_{k,\lambda}}{k!} \binom{n-1}{n-k}.$$

Proof. Substituting (1.1) into (2.1) and simplifying it, one can complete the proof. \square

The following theorem gives the relation between Lah-Bell polynomials and the Stirling numbers of first kind.

Theorem 2.3. *For non-negative integer $n \geq 0$, we have*

$$B_{n,\lambda}^L(x) = \sum_{l=0}^n \sum_{k=0}^l \lambda^{l-k} x^k S_1(l, k) L_{n,l}.$$

In particular, for $x = 1$, we have

$$B_{n,\lambda}^L = \sum_{l=0}^n \sum_{k=0}^l \lambda^{l-k} S_1(l, k) L_{n,l}.$$

Proof. Using (1.4) and (1.7), we get

$$\begin{aligned}
 e^x_\lambda \left(\frac{t}{1-t} \right) &= \left(1 + \frac{\lambda t}{1-t} \right)^{\frac{x}{\lambda}} \\
 &= e^{\frac{x}{\lambda} \ln(1 + \frac{\lambda t}{1-t})} \\
 &= \sum_{k=0}^{\infty} \frac{\lambda^{-k} x^k}{k!} \left(\ln \left(1 + \frac{\lambda t}{1-t} \right) \right)^k \\
 &= \sum_{k=0}^{\infty} \lambda^{-k} x^k \sum_{l=k}^{\infty} S_1(l, k) \frac{\lambda^l}{l!} \left(\frac{t}{1-t} \right)^l \\
 &= \sum_{k=0}^{\infty} \lambda^{-k} x^k \sum_{l=k}^{\infty} S_1(l, k) \lambda^l \sum_{n=l}^{\infty} L_{n,l} \frac{t^n}{n!} \\
 &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \sum_{k=0}^l \lambda^{l-k} x^k S_1(l, k) L_{n,l} \right) \frac{t^n}{n!}.
 \end{aligned} \tag{2.3}$$

Hence, comparing Theorem 2.1 and (2.3) leads to completing of this proof. \square

Theorem 2.4. For non-negative integer $n \geq 0$, we have

$$B_{n,\lambda}^L(x) = \sum_{m=0}^n \sum_{l=0}^m \binom{l}{m} (x)_{l,\lambda} \frac{(-1)^{m-l}}{l!} \langle l \rangle_n,$$

where $\langle l \rangle_n$ is defined by

$$\langle l \rangle_n = \begin{cases} 1, & n = 0, \\ (l + n - 1) \langle l \rangle_{n-1}, & n \geq 1. \end{cases}$$

Proof. Combining (1.4) and (1.5), we can have

$$\begin{aligned}
 \sum_{n=0}^{\infty} B_{n,\lambda}^L(x) \frac{t^n}{n!} &= e^x_\lambda \left(\frac{t}{1-t} \right) \\
 &= \left(1 + \frac{\lambda t}{1-t} \right)^{\frac{x}{\lambda}}
 \end{aligned} \tag{2.4}$$

$$\begin{aligned}
&= \sum_{l=0}^{\infty} (x)_{l,\lambda} \frac{1}{l!} \left(\frac{t}{1-t} \right)^l \\
&= \sum_{l=0}^{\infty} (x)_{l,\lambda} \frac{1}{l!} \sum_{m=0}^l \binom{l}{m} \left(\frac{1}{1-t} \right)^m (-1)^{l-m} \\
&= \sum_{l=0}^{\infty} (x)_{l,\lambda} \frac{1}{l!} \sum_{m=0}^l \binom{l}{m} \sum_{n=0}^{\infty} \langle m \rangle_n \frac{t^n}{n!} (-1)^{l-m} \\
&= \sum_{n=0}^{\infty} \sum_{m=0}^n \sum_{l=0}^m \binom{l}{m} (x)_{l,\lambda} \frac{(-1)^{l-m}}{l!} \langle m \rangle_n \frac{t^n}{n!}.
\end{aligned}$$

By comparing the coefficients of the both sides of (2.4), we can finish this proof. \square

Let $F(t, x)$ be a two variable function defined by

$$F(t, x) := \sum_{n=0}^{\infty} B_{n,\lambda}^L(x) \frac{t^n}{n!}. \quad (2.5)$$

Then, the k -th differentiation gives us the following:

$$\begin{aligned}
\frac{\partial^k}{\partial t^k} F(t, x) &= \frac{\partial^k}{\partial t^k} \left(\sum_{n=0}^{\infty} B_{n,\lambda}^L(x) \frac{t^n}{n!} \right) \\
&= \sum_{n=k}^{\infty} B_{n,\lambda}^L(x) \frac{t^{n-k}}{(n-k)!} \\
&= \sum_{n=0}^{\infty} B_{n+k,\lambda}^L(x) \frac{t^n}{n!}.
\end{aligned} \quad (2.6)$$

Now, we observe that (see [16])

$$\begin{aligned}
\frac{1}{1 + \frac{\lambda t}{1-t}} &= \sum_{k=0}^{\infty} (-1)^k (\lambda)^k \left(\frac{t}{1-t} \right)^k \\
&= \sum_{k=0}^{\infty} (-1)^k \lambda^k \sum_{n=k}^{\infty} \binom{n-1}{n-k} t^n \\
&= \sum_{n=0}^{\infty} n! \sum_{k=0}^n (-1)^k \lambda^k \binom{n-1}{n-k} \frac{t^n}{n!}
\end{aligned} \quad (2.7)$$

and

$$(1 - t)^{-2} = \sum_{l=0}^{\infty} \langle 2 \rangle_l \frac{t^l}{l!}. \tag{2.8}$$

From (2.7) and (2.8), we have

$$\begin{aligned} & \frac{\partial}{\partial t} F(t, x) \\ &= \left(1 + \frac{\lambda t}{1 - t} \right)^{\frac{x}{\lambda} - 1} \frac{x}{(1 - t)^2} \\ &= \left(\sum_{l=0}^{\infty} B_{l,\lambda}^L(x) \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} m! \sum_{k=0}^m \binom{m-1}{m-k} (-1)^k \lambda^k \frac{t^m}{m!} \right) \left(x \sum_{j=0}^{\infty} \langle 2 \rangle_j \frac{t^j}{j!} \right) \\ &= \left(\sum_{l=0}^{\infty} B_{l,\lambda}^L(x) \frac{t^l}{l!} \right) \left(\sum_{i=0}^{\infty} \left(\sum_{m=0}^i \sum_{k=0}^m \binom{i}{m} \binom{m-1}{m-k} m! (-1)^k \lambda^k x \langle 2 \rangle_{i-m} \frac{t^i}{i!} \right) \right) \\ &= \sum_{n=0}^{\infty} \sum_{i=0}^n \sum_{m=0}^i \sum_{k=0}^m \binom{n}{i} \binom{i}{m} \binom{m-1}{m-k} m! (-1)^k \lambda^k x \langle 2 \rangle_{i-m} B_{n-i,\lambda}^L(x) \frac{t^n}{n!}. \end{aligned} \tag{2.9}$$

Theorem 2.5. For any real λ and non-negative integer $n \geq 0$, we have the following recurrence relation

$$B_{n+1,\lambda}^L(x) = \sum_{i=0}^n \sum_{m=0}^i \sum_{k=0}^m \binom{n}{i} \binom{i}{m} \binom{m-1}{m-k} m! (-1)^k \lambda^k x \langle 2 \rangle_{i-m} B_{n-i,\lambda}^L(x).$$

Proof. Combining (2.6) and (2.9), we can complete this proof. □

Now, we observe that

$$\frac{\partial}{\partial x} F(t, x) = \frac{\partial}{\partial x} \left(1 + \frac{\lambda t}{1 - t} \right)^{\frac{x}{\lambda}} = \frac{1}{\lambda} \left(1 + \frac{\lambda t}{1 - t} \right)^{\frac{x}{\lambda}} \ln \left(1 + \frac{\lambda t}{1 - t} \right).$$

By mathematical induction, we can obtain

$$\frac{\partial^k}{\partial x^k} F(t, x) = \frac{1}{\lambda^k} \left(1 + \frac{\lambda t}{1 - t} \right)^{\frac{x}{\lambda}} \left(\ln \left(1 + \frac{\lambda t}{1 - t} \right) \right)^k. \tag{2.10}$$

Since

$$\begin{aligned} \left(\ln \left(1 + \frac{\lambda t}{1-t} \right) \right)^k &= k! \sum_{l=k}^{\infty} S_1(l, k) \frac{1}{l!} \left(\frac{t}{1-t} \right)^l \\ &= \sum_{m=k}^{\infty} \sum_{l=0}^m \frac{m!k!}{l!} S_1(l, k) \binom{m-1}{l-1} \frac{t^m}{m!}, \end{aligned} \quad (2.11)$$

by (2.10) and (2.11), we get

$$\begin{aligned} &\frac{\partial^k}{\partial x^k} F(t, x) \\ &= \left(\frac{1}{\lambda^k} \sum_{m=k}^{\infty} \sum_{l=0}^m \frac{m!k!}{l!} S_1(l, k) \binom{m-1}{l-1} \frac{t^m}{m!} \right) \left(\sum_{j=0}^{\infty} B_{j,\lambda}^L(x) \frac{t^j}{j!} \right) \\ &= \sum_{n=k}^{\infty} \left(\sum_{m=0}^{n-k} \sum_{l=0}^m \frac{1}{\lambda^k} \binom{n}{m} \frac{m!k!}{l!} S_1(n, k) \binom{m-1}{l-1} B_{n-m,\lambda}^L(x) \right) \frac{t^n}{n!}. \end{aligned} \quad (2.12)$$

We obtain the following by k differentiations of the function $F(t, x)$ with respect to x :

$$\frac{\partial^k}{\partial x^k} F(t, x) = \frac{\partial^k}{\partial x^k} \sum_{m=0}^{\infty} B_{m,\lambda}^L(x) \frac{t^m}{m!} = \sum_{m=k}^{\infty} \frac{\partial^k}{\partial x^k} B_{m,\lambda}^L(x) \frac{t^m}{m!}. \quad (2.13)$$

Theorem 2.6. For real number λ and non-negative integers n and k with $n \geq k$, we have

$$\frac{\partial^k}{\partial x^k} B_{n,\lambda}^L(x) = \sum_{m=0}^{n-k} \sum_{l=0}^m \binom{n}{m} \frac{m!k!}{l!\lambda^k} S_1(n, k) \binom{m-1}{l-1} B_{n-m,\lambda}^L(x). \quad (2.14)$$

Proof. Comparing (2.12) and (2.13), one can proof the above recurrence relations. \square

Theorem 2.7. For $\lambda \in \mathbb{R}$ and non-negative integers $n, k \geq 0$, we have

$$\frac{d^k}{dx^k} B_{n,\lambda}^L(x) = n! \sum_{k=0}^n \sum_{l=0}^k \sum_{m=0}^l \frac{1}{k!} \binom{n-1}{n-k} S_{2,\lambda}(k, l) S_1(l, m) (m)_k x^{m-k}.$$

Proof. Using (1.10) and Corollary 2.2, we have

$$\begin{aligned}
 \frac{d^k}{dx^k} B_{n,\lambda}^L(x) &= \frac{d^k}{dx^k} n! \sum_{k=0}^n \frac{(x)_{k,\lambda}}{k!} \binom{n-1}{n-k} \\
 &= n! \sum_{k=0}^n \frac{1}{k!} \binom{n-1}{n-k} \frac{d^k}{dx^k} (x)_{k,\lambda} \\
 &= n! \sum_{k=0}^n \frac{1}{k!} \binom{n-1}{n-k} \frac{d^k}{dx^k} \sum_{l=0}^k S_{2,\lambda}(k,l)(x)_l \\
 &= n! \sum_{k=0}^n \frac{1}{k!} \binom{n-1}{n-k} \frac{d^k}{dx^k} \sum_{l=0}^k S_{2,\lambda}(k,l) \sum_{m=0}^l S_1(l,m)x^m \\
 &= n! \sum_{k=0}^n \frac{1}{k!} \binom{n-1}{n-k} \sum_{l=0}^k \sum_{m=0}^l S_{2,\lambda}(k,l) S_1(l,m) (m)_k x^{m-k}.
 \end{aligned} \tag{2.15}$$

This is the desired result of the theorem. □

3. DISTRIBUTION OF ROOTS OF THE POLYNOMIALS

Woon [26] has studied the distribution and structure of the zeros of Bernoulli polynomials. Hence, we will investigate the numerical patten of the zeros of the polynomials $B_{n,\lambda}^L(x)$. Using the mathematica tool, the polynomial $B_{n,\lambda}^L(x)$ can be determined explicitly. For example,

$$\begin{aligned}
 B_{1,\lambda}^L(x) &= x, \\
 B_{2,\lambda}^L(x) &= x^2 + (2 - \lambda)x, \\
 B_{3,\lambda}^L(x) &= x^3 + (6 - 3\lambda)x^2 + (6 - 6\lambda + 2\lambda^2)x, \\
 B_{4,\lambda}^L(x) &= x^4 + (12 - 6\lambda)x^3 + (36 - 36\lambda + 11\lambda^2)x^2 \\
 &\quad + (24 - 36\lambda + 24\lambda^2 - 6\lambda^3)x.
 \end{aligned}$$

From the definition of the Lah-Bell polynomials $B_{n,\lambda}^L(x)$, we can obtain the following properties of the roots:

- For any real number λ , the polynomials $B_{n,\lambda}^L(x)$ with $n = 1, 2$ have only real roots.
- For any real number λ and any positive integer n , all polynomials $B_{n,\lambda}^L(x)$ have a common root which is zero.




Fig. 1. The computed roots of $B_{40,\lambda}^L(x)$ with variable λ

Firstly, we observe the impact of λ on the distribution of the roots of the polynomials. For the propose, we fix the degree of polynomials as $n = 40$. Since the explicit form of the roots of $B_{n,\lambda}^L(x)$ is unknown, we calculate the roots by using the Mathematica tool with 100 working precision. The absolute numerical error is bounded as follows:

$$\sum_{i=1}^{40} |B_{40,\lambda}(x_i)| < 10^{-62},$$

where x_i denotes the root of polynomial. Hence, the results obtained from numerical computations are reliable. We compute the numerical roots of $B_{40,\lambda}^L(x)$ with four different parameters $\lambda = \frac{5}{10}, \frac{15}{10}, \frac{20}{10}$ and $\frac{25}{10}$ and the results are plotted

in Fig. 1. As observed in Fig. 1, the roots of the Lah-Bell polynomials have four patterns.




Fig. 2. The root distribution of $B_{n,\lambda}^L(x)$ with variable λ and different integer $n = 1, 2, \dots, 40$.

Secondly, we investigate the impact of the degree of polynomials on the distribution of roots of the polynomials. We compute the numerical roots of the polynomials increasing the degree of polynomials from 1 to 40 and present in Fig. 2.




Fig. 3. Real zeros of $B_{n,\lambda}^L(x)$ for $\lambda = \frac{5}{10}, \frac{15}{10}, \frac{20}{10}, \frac{25}{10}$ and $1 \leq n \leq 40$.

Thirdly, to investigate the real roots distribution structure of $B_{n,\lambda}^L(x)$, we compute the real roots and display in Fig.3.




Fig. 4. Roots distribution structures vs $\lambda \in [0, \frac{25}{10}]$.

From the results of Fig.3, we can find a remarkably regular structure of the roots of the polynomials $B_{n,\lambda}(x)$. In order to find the roots structure, we count the number of real roots for $\lambda = \frac{5}{10}, \frac{15}{10}, \frac{20}{10}$ and $\frac{25}{10}$ and $n \in [1, 99]$ within $x \in [-1000, 1000]$ and summarize as follows:

- $\lambda = \frac{5}{10}$: the number of real roots= n .
- $\lambda = \frac{20}{10}$: the number of real roots= $\begin{cases} 1, & n = \text{odd}, \\ 2, & n = \text{even}. \end{cases}$
- $\lambda = \frac{15}{10}$: the number of real roots= $\begin{cases} 1, & n = 1, 3, \dots, 13, \\ 2, & n = 2, 4, \dots, 26, \\ 3, & n = 15, 17, \dots, 37, \\ 4, & n = 28, 30, \dots, 50, \\ 5, & n = 39, 41, \dots, 63, \\ 6, & n = 52, 54, \dots, 74, \\ 7, & n = 65, 67, \dots, 87, \\ 8, & n = 76, 78, \dots, 98, \\ 9, & n = 89, 91, \dots, 99. \end{cases}$
- $\lambda = \frac{25}{10}$: the number of real roots= $\begin{cases} 1, & n = 1, 3, \dots, 27, \\ 2, & n = 2, 4, \dots, 50, \\ 3, & n = 29, 31, \dots, 75, \\ 4, & n = 52, 54, \dots, 100, \\ 5, & n = 77, 79, \dots, 99. \end{cases}$

Finally, we compute the roots of the polynomials with fixed $n = 40$ and varying parameter $\lambda = \frac{k}{10}, k = 0, 1, \dots, 25$. The numerical results are plotted in Fig. 4.

Summaring the above discussion, we can obtain the properties of the roots of $B_{n,\lambda}^L(x)$.

- When $\lambda < 2$, the real parts of the roots of the polynomials $B_{n,\lambda}^L(x)$ are non-positive.
- When $\lambda = 2$, the polynomials $B_{n,\lambda}^L(x)$ have pure imaginary roots except for zero roots.
- When $\lambda > 2$, the real parts of the roots of the polynomials $B_{n,\lambda}^L(x)$ are non-negative.

4. CONCLUSION

In this paper, we review the Lah-Bell polynomials and numbers introduced by Kim-Kim and give an explicit formula for partial derivatives. In order to more accurately understand the Lah-Bell polynomials, the distribution of roots was numerically investigated. Further, we count the number of real roots of $B_{n,\lambda}^L(x)$ with four different parameters $\lambda = \frac{5}{10}, \frac{15}{10}, \frac{20}{10}$ and $\frac{25}{10}$. Finally, we obtain the relation between the sign of the real part of the root of $B_{n,\lambda}^L(x)$ and the value λ . In the next study, we will show theoretically the above facts.

Acknowledgments: The authors would like to thank the anonymous referees for their careful reading, valuable comments, and suggestions, which helped to improve the manuscript.

REFERENCES

- [1] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series, vol. 55. U.S. Government Printing Office, Washington, (1964), xiv+1046.
- [2] E.Y. Bell, *Exponential polynomials*, Ann. of Math., **35** (1934), 258-277.
- [3] L. Carlitz, *Arithmetic properties of the Bell polynomials*, J. Math. Anal. Appl., **15** (1966), 33-52.
- [4] L. Carlitz, *Degenerate Stirling, Bernoulli and Eulerian numbers*, Util. Math., **15** (1979), 51-88.
- [5] L. Comtet, *Advanced Combinatorics. The Art of Finite and Infinite Expansions*, Reidel, Dordrecht, Revised and enlarged edn., (1974) xi+343, ISBN: 90-277-0441-4 05-02
- [6] Dmitry V. Dolgy, D.S. Kim, T. Kim and J. Kwon, *On fully degenerate Bell numbers and polynomials*, Filomat, **34**(2) (2020), 507-514.
- [7] L.-C. Jang, D.S. Kim, T. Kim and H. Lee, *Some identities involving derangement polynomials and numbers and moments of gamma random variables*, J. Funct. Spaces, **160** (2020), Article ID 6624006, 9 pages.
- [8] D.S. Kim and T. Kim, *Some identities of Bernoulli and Euler polynomials arising from umbral calculus*, Adv. Stud. Contemp. Math. (Kyungshang), **23**(1) (2013), 159-171.
- [9] D.S. Kim and T. Kim, *Some identities of Bell polynomials*, Sci. China Math., **58** (10) (2015), 2095–2104.
- [10] D.S. Kim and T. Kim, *Some identities of degenerate Daehee numbers arising from certain differential equations*. J. Nonlinear Sci. Appl., **10** (2017), 744-751.
- [11] D.S. Kim and T. Kim, *A note on polyexponential and unipoly functions*, Russ. J. Math. Phys., **26**(1) (2019), 40-49.
- [12] D.S. Kim and T. Kim, *A note on a new type of degenerate Bernoulli numbers*, Russ. J. Math. Phys., **27**(2) (2020), 227-235.

- [13] D.S. Kim and T. Kim, *Lah-Bell numbers and polynomials*, Proc. Jangjeon Math. Soc., **23**(4) (2020), 577-586.
- [14] D.S. Kim and T. Kim, *Degenerate Sheffer sequences and λ -Sheffer sequences*, J. Math. Anal. Appl., **493**(1) (2021), 124521, 21 pp.
- [15] D.S. Kim and T. Kim, *r-extended Lah-Bell numbers and polynomials associated with r-Lah numbers*, **arXiv** (2020), 2008.06155 <https://arxiv.org/abs/2008.06155>
- [16] H. Kim, *Degenerate Lah-Bell polynomials arising from degenerate Sheffer sequences*, Advances in Difference Equations, **687** (2020), 16pages.
- [17] T. Kim, *A note on degenerate Stirling polynomials of the second kind*, Proc. Jangjeon Math. Soc., **20**(3) (2017), 319-331.
- [18] D.S. Kim, *A note on nonlinear Changhee differential equations*, Russ. J. Math. Phys., **23**(1) (2016), 88–92
- [19] T. Kim and D.S. Kim, *Degenerate Laplace transform and degenerate gamma function*, Russ. J. Math. Phys., **24**(2) (2017), 241–248.
- [20] T. Kim and D.S. Kim, *Some identities on λ -analogue of r-Stirling numbers of the first kind*, Filomat, **34**(2) (2020), 451-460.
- [21] T. Kim and D.S. Kim, *Note on degenerate gamma function*, Russ. J. Math. Phys., **27**(3) (2020), 352–358.
- [22] T. Kim, D.S. Kim, H. Kim and J. Kwon, *Some identities of degenerate Bell polynomials*, Mathematics, **8**(40) (2020), 8pages.
- [23] T. Kim, D.S. Kim, H. Lee and L. Jang, *Degenerate Bell polynomials associated with umbral calculus*, J. Inequal. Appl., **226** (2020), 15pages.
- [24] Y. Ma, D.S. Kim, T. Kim, H. Kim and H. Lee, *Some identities of Lah-Bell polynomials*, Adv. Differ. Equ., **510** (2020), 10pages.
- [25] S. Tauber, *Lah numbers for Fibonacci and Lucas polynomials*, Fibonacci Quart., **6**(5) (1968), 93–99.
- [26] S.C. Woon, *Analytic continuation of Bernoulli numbers, a New formula for the Riemann Zeta function, and the phenomenon of scattering of zeros*, Physics, **arXiv** (1997), 9705021v2