

Short communication

New Records of Two *Dorylaimoides* Species (Nematoda: Dorylaimida: Mydonomidae) from Korea

Taeho Kim¹, Jiyeon Kim², Joong-Ki Park^{3,*}

¹Department of Ecology and Conservation, National Marine Biodiversity Institute of Korea,
Seocheon 33662, Korea

²Animal & Plant Research Department, Nakdonggang National Institute of Biological Resources,
Sangju 37242, Korea

³Division of EcoScience, Ewha Womans University, Seoul 03760, Korea

ABSTRACT

The genus *Dorylaimoides* Thorne and Swanger, 1936 comprises over 70 species and predominantly inhabits soil; however, some species are found in semiaquatic or aquatic habitats. In this study, *D. elegans* (de Man, 1880) Thorne and Swanger, 1936 and *D. limnophilus* (de Man, 1880) Loof, 1964, collected from sediments in the Hangang River, are reported for the first time in Korea. These two *Dorylaimoides* species are distinguished from other *Dorylaimoides* species by their morphological characteristics: a didelphic-amphidelphic reproductive system, short tail, and digitate terminus in *D. elegans* and a mono-opisthodelphic reproductive system and long tail in *D. limnophilus*. Here, we provide detailed information on the morphological characteristic (description and illustration) and morphometrics of the two *Dorylaimoides* species, using optical microscopy. Additionally, 18S rDNA sequences of the two species from the Korean isolates are provided as molecular evidence.

Keywords: Nematode, Dorylaimida, *Dorylaimoides elegans*, *Dorylaimoides limnophilus*, new record, freshwater, Korea

INTRODUCTION

The genus *Dorylaimoides* Thorne and Swanger, 1936 belongs to the family Mydonomidae Thorne, 1964. To date, the genus comprises over 70 species; however, only three species have previously been reported in Korea: *D. leptus* Husain and Khan, 1968; *D. micoletzkyi* (de Man, 1921) Thorne and Swanger, 1936; *D. punctatus* Khan and Park, 1999 (de Man, 1921; Husain and Khan, 1968; Khan and Park, 1999). In this study, we report *D. elegans* (de Man, 1880) Thorne and Swanger, 1936 and *D. limnophilus* (de Man, 1880) Loof, 1964 collected from Korea for the first time and provide detailed descriptions of their morphological characters, morphometrics, and 18S rDNA sequences.

Live specimens were collected from freshwater and sediment samples (Wangdae-ri, Neungseo-myeon, Yeoju-si, Gyeonggi-do, Korea [GPS coordinates: 37°19'43.82"N, 127°36'15.40"E]) from the Hangang River and then isolated by sieving and the Baermann funnel method (Baermann, 1917).

Each specimen was transferred to 2 mL of water, to which 4 mL of 80°C triethanolamine-formalin (TAF, 2% triethanolamine and 7% formaldehyde) solution was added for fixation. Fixed nematodes were dehydrated using glycerin (Seinhorst, 1959) and mounted on hydroglyceric slides (Shirayama et al., 1993). Under an Imager A2 optical microscope, with differential interference contrast, equipped with an AxioCam 506 color camera, morphological and morphometric characters of the specimens were observed and measured from digital photographs using the Axiovision SE64 Rel. 4.9.1 software (Zeiss, Oberkochen, Germany).

Total genomic DNA was extracted using a nematode lysis buffer (Holterman et al., 2006) consisting of 0.2 M NaCl, 0.2 M Tris-HCl (pH 8.0), 1% (v/v) β-mercaptoethanol, and 800 µg/mL proteinase-K. DNA was stored at -20°C if not used immediately. The 18S rDNA gene primer sets used in this study were 988-F (5'-CTCAAAGATTAAGCCATGC-3')/1096-R (5'-GGTAATTCTGGAGCTAATAC-3')/1912R (5'-TTTACGGTCAGAACTAGGG-3'), 1813-F (5'-CTGC

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*To whom correspondence should be addressed
Tel: 82-2-3277-5948, Fax: 82-2-3277-2385
E-mail: jkpark@ewha.ac.kr




Fig. 1. *Dorylaimoides elegans* (de Man, 1880) Thorne and Swanger, 1936. A, Entire female; B, Female neck region; C, Female reproductive system; D, Female posterior region; E, Male posterior region. Scale bars: A-E=100 µm.

GTGAGAGGGTGAAT-3')/2646-R (5'-GCTACCTTGTAC GACTTTT-3') (Holterman et al., 2006). PCR (50 µL) was performed using 2 µL template DNA, 10 pmol of each primer, 10× Ex Taq buffer, 0.2 mM dNTP mixture, and 1.25 U of TaKaRa Ex Taq polymerase (TaKaRa, Otsu-Shiga, Japan). The conditions were as follows: initial denaturation at 95°C for 1 min, 40 cycles of denaturation at 95°C for 30 s, annealing at 50°C for 30 s, extension at 72°C for 1 min, and final extension at 72°C for 10 min. PCR products were purified with a QIAquick PCR Purification Kit (Qiagen, Hilden, Germany) and then sequenced using a 3730xl DNA Analyzer (Thermo Fisher Scientific, Waltham, MA, USA). The resulting 18S rDNA sequences were deposited in GenBank. Sequence analyses were performed using Geneious v11.0.5 (Biomatters, Auckland, New Zealand) (Kearse et al., 2012) and aligned with

sequences of available congeneric nematode species using Clustal X with default options (Thompson et al., 1997).

SYSTEMATIC ACCOUNTS

Order Dorylaimida Pearse, 1942

Superfamily Tylencholaimoidea Filipjev, 1934

Family Mydonomidae Thorne, 1964

Genus *Dorylaimoides* Thorne and Swanger, 1936

^{1*}***Dorylaimoides elegans* (de Man, 1880)**

Thorne and Swanger, 1936 (Table 1, Fig. 1)

Dorylaimus elegans de Man, 1880: 86.

Dorylaimoides elegans: Thorne and Swanger, 1936: 129–130, Pl. XXIX, fig. 174.

Korean name: ^{1*}예쁜창선충(신칭)

Table 1. Morphometrics of *Dorylaimoides elegans* and *D. limnophilus*

Character	<i>D. elegans</i>		<i>D. limnophilus</i>
	♀, n=1	♂, n=1	♀, n=3
L	1,318.2	1,241.2	1,207.3±154.0 (1,037.8–1,338.8)
Body width	36.6	35.5	32.3±3.8 (28.6–36.1)
Pharynx length	192.7	209.1	209.1±16.4 (192.7–225.6)
Tail length	42.0	32.6	126.9±20.3 (105.0–145.1)
Anal body diameter	25.3	25.1	19.5±2.5 (16.8–21.6)
a	36.1	35.0	37.4±1.3 (36.4–38.8)
b	6.8	5.9	5.8±0.3 (5.4–6.0)
c	31.4	38.1	9.5±0.3 (9.2–9.9)
c'	1.7	1.3	6.5±0.2 (6.3–6.7)
V	45.6	—	31.0±0.9 (29.9–31.5)
Lip region diameter	9.2	8.2	8.2±0.2 (8.0–8.3)
Lip region height	4.1	3.9	3.4±0.4 (3.0–3.8)
Amphid aperture	6.7	6.0	6.0±0.1 (6.0–6.1)
Odontostyle (ventral side)	7.7	7.6	5.7±0.3 (5.4–6.0)
Odontophore	15.8	14.9	15.8±0.5 (15.2–16.2)
Pharyngeal bulb length	63.7	58.6	65.2±2.8 (62.5–68.0)
Cardia diameter	7.3	7.1	12.0±1.0 (11.1–13.0)
Cardia length	7.5	7.6	8.1±0.7 (7.5–8.8)
Guiding ring from anterior end	8.1	8.0	6.3±0.2 (6.1–6.5)
Nerve ring from anterior end	105.4	96.2	99.3±4.4 (94.8–103.5)
Nerve ring (% pharynx)	54.7	46.0	47.6±1.7 (45.9–49.2)
Vulva from anterior end	601.7	—	375.0±57.4 (311.1–422.2)
Vulva to anus	674.5	—	705.5±76.4 (621.7–771.5)
Vulva to anus/tail length	16.1	—	5.6±0.3 (5.3–5.9)
Vagina diameter	18.3	—	14.5±1.7 (13.0–16.4)
Vagina length	20.9	—	15.3±1.6 (13.9–17.0)
Vagina/body diameter	0.6	—	0.5
Reproductive tract length (G1)	185.1	—	—
Reproductive tract length (G2)	207.5	—	210.7±58.0 (147.9–262.1)
G1 (%)	14.0	—	—
G2 (%)	15.7	—	17.2±2.7 (14.3–19.6)
Spicules	—	37.1	—
Spicules/anal body diameter	—	1.5	—
Spicules/tail length	—	1.2	—
Ventromedian supplements	—	5.0	—
Rectum	33.2	32.2	22.6±2.2 (20.6–25.0)
Rectum/anal body diameter	1.3	1.3	1.2±0.1 (1.1–1.2)

All measurements are in μm and in the form mean±SD (range).

L, body length; a, body length/body diameter; b, body length/distance from anterior to base of esophageal glands; c, body length/tail length; c', tail length/diameter at anus region; V, % distance of vulva from anterior end/body length; G1, % length of anterior female gonad in relation to body length; G2, % length of posterior female gonad in relation to body length.

Material examined. 1♀, 1♂, Korea: Gyeonggi-do, Yeoujusi, Neungseo-myeon, Wangdae-ri, 37°19'43.82"N, 127°36'15.40"E, 25 Feb 2019. Voucher specimens were deposited in the Nakdonggang National Institute of Biological Resources (NNIBR), Korea.

Measurements. See Table 1.

Description. Female: Body slender, ventrally curved after

fixation, length 1,318.2 μm , width 36.6 μm (maximum width at level of vulva). Cuticle with fine transverse striations, 1.2 μm thick at anterior region, 2.0 μm at mid-body, and 5.1 μm at tail. Lip region rounded, slightly offset, about 2.2 times as wide as high. Amphid funnel-shaped aperture 6.7 μm , about 0.7 times as wide as lip region width. Odontostyle asymmetrical and slightly arched, ventral side length 7.7 μm . Odon-




Fig. 2. *Dorylaimoides limnophilus* (de Man, 1880) Loof, 1964. A, Entire female; B, Female neck region; C, Female reproductive system; D, Female posterior region. Scale bars: A-D=100 μ m.

tophore arcuated, narrowing posteriorly, about 2.1 times as long as odontostyle. Guiding ring distinct and single, located 8.1 μ m from anterior end. Pharynx with a slender anterior and cylindrical bulb, 63.7 μ m in length, and about 28% of total neck length. Cardia rounded and distally enclosed by intestinal tissue, about 2.2 times as long as body width. Nerve ring located 105.4 μ m from anterior end, 54.7% of pharynx length. Excretory pore obscure. Female reproductive system didelphic-amphidelphic. Vulva transverse, located at 45.6% of body length. Vagina length 0.5 times body diameter. Uterus length 2.3–3.3 times body diameter. Oviduct length 1.8–3.0 times body diameter. Ovary reflexed, usually reaching junction of oviduct and uterus, anterior reproductive tract (Q1) 185.1 μ m, posterior (Q2) 207.5 μ m long. Rectum length about 1.3 times anal body diameter. Tail dorsally convex conoid,

with digitate tip. Male: General morphology similar to female, except for posterior region. Body ventrally hooked after fixation, length 1,241.2 μ m, width 35.5 μ m (maximum width at mid-body). Cuticle annuli 1.0 μ m thick at anterior region, 1.5 μ m at mid-body, and 4.9 μ m at tail. Lip region width 8.2 μ m. Amphid aperture 6.0 μ m. Odontostyle ventral side length 7.6 μ m. Odontophore 1.9 times odontostyle length. Nerve ring located at 46% of pharynx length. Genital system opposite. Spicules dorylaimoid, 37.1 μ m long, 1.1 times anal body diameter, with 11.0 μ m long lateral accessory pieces. Supplements mammiform, five ventromedians. Tail similar to female, with digitate terminus.

Habitat. Freshwater and sediment.

Distribution. America, Canada, Estonia, India, Korea (this study), the Netherlands, Romania.

^{1*}***Dorylaimoides limnophilus* (de Man, 1880)**
Loof, 1964 (Table 1, Fig. 2)

Dorylaimus limnophilus de Man, 1880: 96.
Thornenema limnophilum Andrassy, 1959b: 195.
Dorylaimoides riparius Andrassy, 1962: 6–8, Abb. 3.
Dorylaimoides (Tarjania) limnophilus: Loof, 1964: 287.

Material examined. 3♀♀, Korea: Gyeonggi-do, Yeoju-si, Neungseo-myeon, Wangdae-ri, 37°19'43.82"N, 127°36'15.40"E, 25 Feb 2019. Voucher specimens were deposited in the Nakdonggang National Institute of Biological Resources (NNIBR), Korea.

Measurements. See Table 1.

Description. Female: Body slender, ventrally curved after fixation, length 1,037.8–1,338.8 µm, width 28.6–36.1 µm (maximum width at level of vulva). Cuticle with fine transverse striations, 1.5–1.7 µm thick at anterior region, 2.0–2.3 µm at mid-body, and 4.2–4.6 µm at tail. Lip region angular, offset by a constriction, about 2.6 times as wide as high. Amphid cup-shaped, aperture 6.0–6.1 µm, about 0.7 times as wide as lip region width. Odontostyle asymmetrical and slightly arched, ventral side length 5.4–6.0 µm. Odontophore arcuated, narrowing posteriorly, about 2.8 times as long as odontostyle. Guiding ring distinct and single, located 6.1–6.5 µm from anterior end. Pharynx with a slender anterior and cylindrical bulb, length 62.5–68.0 µm and about 30% of total neck length. Cardia rounded and distally enclosed by intestinal tissue. Nerve ring located at 94.8–103.5 µm from anterior end, 45.9–49.2% of pharynx length. Excretory pore obscure. Reproductive system mono-opisthodelphic. Vulva transverse, located at 29.9–31.5% of body length. Vagina length 0.5 times body diameter. Uterus short. Oviduct short. Ovary reflexed, with numerous oocytes. Rectum length about 1.2 times anal body diameter. Tail long, filiform, tapering to terminus. Male: Not found.

Habitat. Freshwater and sediment.

Distribution. Belgium, Germany, Holland, Hungary, Italy, Japan, Korea (this study), the Netherlands, Romania, Slovakia, Spain, Sweden, Switzerland, Uzbekistan.

Molecular sequence information. Molecular sequences (partial 18S rDNA sequences) deposited in GenBank: *D. elegans* (GenBank accession no: MN880408) and *D. limnophilus* (GenBank accession no: MN880407).

Diagnosis and molecular analysis. Morphological and morphometric characters reported herein generally match those previously reported for these two *Dorylaimoides* species (de Man, 1880; Thorne and Swanger, 1936; Jairajpuri and Ahmad, 1992 for *D. elegans*; de Man, 1880; Andrassy, 1959a, 1959b, 1962; Loof, 1964; Peralta and Peña-Santiago, 1995b; Loof,

1990 for *D. limnophilus*), except for spicule length (37.1 vs. 30–32 µm) (Goseco et al., 1976) for *D. elegans*. The two *Dorylaimoides* species reported in this study were distinguished by their morphological characteristics: a didelphic-amphidelphic reproductive system, short tail, and digitate terminus in *D. elegans* and a mono-opisthodelphic reproductive system and long tail in *D. limnophilus* (Ahmad and Jairajpuri, 1983; Peralta and Peña-Santiago, 1995a, 1995b; Khan and Park, 1999; Ahmad et al., 2003; Ahmad and Mushtaq, 2004; Pedram et al., 2011; Gagarin and Gusakov, 2015). In addition, we obtained partial 18S rDNA sequences for the two *Dorylaimoides* species and assessed sequence similarity with other *Dorylaimoides* species from GenBank. The 18S sequences of our specimens are almost identical to those of *D. elegans* (AY146486.2, AY911976.1, AY911977.1; 99.8–100%) and *D. limnophilus* (AY284829.1, AY593950.1; 99.5–100%) deposited on GenBank.

ORCID

Taeho Kim: <https://orcid.org/0000-0003-1973-3100>
Jiyeon Kim: <https://orcid.org/0000-0003-1317-4870>
Joong-Ki Park: <https://orcid.org/0000-0002-0607-3329>

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This work was supported by a grant from the Nakdonggang National Institute of Biological Resources (NNIBR) funded by the Ministry of Environment (NNIBR202001107), Korea Research Fellowship Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science and ICT (2020R1A2C2005393) of the Republic of Korea and the National Marine Biodiversity Institute of Korea (2021M00300).

REFERENCES

- Ahmad W, Jairajpuri MS, 1983. Descriptions of new species of *Dorylaimoides* and *Culolaimus* (Dorylaimida) from India. *Revue de Nématologie*, 6:65–72.

Korean name: ^{1*}담수긴꼬리창선충(신칭)

- Ahmad W, Mushtaq P, 2004. Five new species of *Dorylaimoides* Thorne & Swanger (Nematoda: Dorylaimida) from Singapore. International Journal of Nematology, 14:99-110.
- Ahmad W, Mushtaq P, Baniyamuddin M, 2003. Descriptions of three new species of *Dorylaimoides* Thorne & Swanger, 1936 (Nematoda: Dorylaimida). Journal of Nematode Morphology and Systematics, 5:153-162.
- Andrássy I, 1959a. Taxonomische Übersicht der Dorylaimen (Nematoda), I. Acta Zoologica Academiae Scientiarum Hungaricae, 5:191-240.
- Andrássy I, 1959b. Taxonomische Übersicht der Dorylaimen (Nematoda), III. Acta Zoologica Academiae Scientiarum Hungaricae, 5:143-532.
- Andrássy I, 1962. Zwei neuen Nematoden-Arten aus dem Überschwemmungsgebiet der Donau (Danubialia Hungarica, XIII.). Opuscula Zoologica, Budapest, 4:3-8.
- Baermann G, 1917. Eine einfache methode zur auffindung von ankylostomum (Nematoden) larven in erdproben. Geneeskunding Tijdschrift voor Nederlandsch-Indië, 57:131-137.
- De Man JG, 1880. Die einheimischen, frei in der reinen Erde und im süßen Wasser lebende Nematoden. Bericht und de scriptivsystematischer Theil. Tijdschrift der Nederlandsche Dierkundige Vereeniging, 5:1-104.
- De Man JG, 1921. Nouvles recherches sur les nematodes libres terricoles de la Hollande. Capita Zoologica, 1:3-62.
- Gagarin VG, Gusakov VA, 2015. Two new species of free-living nematodes of the genus *Dorylaimoides* Thorne, Swanger, 1939 from fresh waterbodies of Vietnam. Inland Water Biology, 8:121-129. <https://doi.org/10.1134/S1995082915010071>
- Goseco CG, Ferris VR, Ferris JM, 1976. Revisions in Leptonchoidea (Nematoda: Dorylaimida) *Dorylaimoides* in Dorylaimidae, Dorylaimoidinae; *Calolaimus* and *Timmus* n. gen. in Dorylaimidae, Calolaiminae and *Miranema* in Miranematidae. Research Bulletin, Purdue University, Agriculture Experiment Station, 941:1-46.
- Holterman M, van der Wurff A, van den Elsen S, van Megen H, Bongers T, Holovachov O, Bakker J, Helder J, 2006. Phylum-wide analysis of SSU rDNA reveals deep phylogenetic relationships among nematodes and accelerated evolution toward crown clades. Molecular Biology and Evolution, 23:1792-1800. <https://doi.org/10.1093/molbev/msl044>
- Husain SI, Khan AM, 1968. *Basirotyleptus modestus* n. sp. and two new species of *Dorylaimoides* Thorne & Swanger, 1936 from India. Nematologica, 14:362-368. <https://doi.org/10.1163/187529268X00039>
- Jairajpuri MS, Ahmad W, 1992. Dorylaimida: free-living, predaceous and plant-parasitic nematodes. E.J. Brill Publishing Company, Leiden, pp. 1-458.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A, 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics, 28:1647-1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Khan Z, Park SD, 1999. Description of *Dorylaimoides punctatus* n. sp. and *Paracatinolaimus acutus* n. sp. (Nematoda: Dorylaimida) from Korea. Journal of Asia-Pacific Entomology, 2:45-50. [https://doi.org/10.1016/S1226-8615\(08\)60030-8](https://doi.org/10.1016/S1226-8615(08)60030-8)
- Loof PAA, 1964. Free-living and plant-parasitic nematodes from Venezuela. Nematologica, 10:201-300. <https://doi.org/10.1163/187529264X00042>
- Loof PAA, 1990. Systematic observations on some species of Dorylaimoididae (Nematoda: Dorylaimida). Revue de Nématologie, 13:161-170.
- Pedram M, Pourjam E, Vinciguerra MT, 2011. Description of *Dorylaimoides alborzicus* sp. n. (Dorylaimida: Nematoda) from Iran, with updated compendium and key to the species of *Dorylaimoides*. Zootaxa, 3022:58-68. <https://doi.org/10.1164/zootaxa.3022.1.4>
- Peralta M, Peña-Santiago R, 1995a. Nematodes of the order Dorylaimida from Andalucía Oriental, Spain: the genus *Dorylaimoides* Thorne & Swanger, 1936, 1. Didelphic species. Fundamental and Applied Nematology, 18:35-53.
- Peralta M, Peña Santiago R, 1995b. Nematodes of the order Dorylaimida from Andalucía Oriental, Spain: the genus *Dorylaimoides* Thorne & Swanger, 1936. 2. Pseudodidelphic-opisthodelphic species. Fundamental and Applied Nematology, 18:167-180.
- Seinhorst JW, 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. Nematologica, 4:67-69. <https://doi.org/10.1163/187529259X00381>
- Shirayama Y, Kaku T, Higgins RP, 1993. Double-sided microscopic observation of meiofauna using an HS-slide. Benthos Research, 1993:41-44. https://doi.org/10.5179/benthos.1990.1993.44_41
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG, 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Research, 25:4876-4882. <https://doi.org/10.1093/nar/25.24.4876>
- Thorne G, Swanger HH, 1936. A monograph of the nematode genera *Dorylaimus* Dujardin, *Aporcelaimus* n.g., *Dorylaimoides* n.g. and *Pungentus* n.g. Capita Zoologica, 6:1-223.

Received January 14, 2021
Revised February 25, 2021
Accepted February 26, 2021