

PROJECTIONS OF ALGEBRAIC VARIETIES WITH ALMOST LINEAR PRESENTATION II

JEAMAN AHN*

ABSTRACT. Let X be a nondegenerate reduced closed subscheme in \mathbb{P}^n . Assume that $\pi_q : X \rightarrow Y = \pi_q(X) \subset \mathbb{P}^{n-1}$ is a generic projection from the center $q \in \text{Sec}(X) \setminus X$ where $\text{Sec}(X) = \mathbb{P}^n$. Let Z be the singular locus of the projection $\pi_q(X) \subset \mathbb{P}^{n-1}$. Suppose that I_X has the almost minimal presentation, which is of the form

$$R(-3)^{\beta_{2,1}} \oplus R(-4) \rightarrow R(-2)^{\beta_{1,1}} \rightarrow I_X \rightarrow 0.$$

In this paper, we prove the followings:

- (a) Z is either a linear space or a quadric hypersurface in a linear subspace;
- (b) $H^1(\mathcal{I}_X(k)) = H^1(\mathcal{I}_Y(k))$ for all $k \in \mathbb{Z}$;
- (c) $\text{reg}(Y) \leq \max\{\text{reg}(X), 4\}$;
- (d) Y is cut out by at most quartic hypersurfaces.

1. Introduction

Let V be a vector space of dimension $n+1$ over an algebraically closed field K with a basis x_0, \dots, x_n . If $X \subset \mathbb{P}^n = \mathbb{P}(V)$ is a nondegenerate reduced subscheme then we write \mathcal{I}_X for the ideal sheaf and I_X for the defining saturated ideal of X in the homogeneous coordinate ring $R = \text{Sym}(V) = K[x_0, \dots, x_n]$. Suppose that the minimal free resolution of R/I_X is of the following form

$$(1.1) \quad \cdots \rightarrow R(-3)^{\beta_{2,1}^R} \rightarrow R(-2)^{\beta_{1,1}^R} \rightarrow R \rightarrow R/I_X \rightarrow 0.$$

The authors in [2] have proved that if $\pi_q : X \rightarrow Y \subset \mathbb{P}^{n-1}$ is a non-isomorphic generic projection with the center $q \in \mathbb{P}^n$ then

Received March 16, 2021; Accepted May 08, 2021.

2010 Mathematics Subject Classification: Primary 14N05; Secondary 13D02.

Key words and phrases: syzygy, normality, Castelnuovo-Mumford regularity, partial elimination ideal.

The author was supported by the research grant of the Kongju National University for the research year in 2018-2019.

- the singular locus $Z = \{y \in Y \mid \text{the length of } \pi_q^{-1}(y) \geq 2\}$ is a linear space;
- $H^1(\mathcal{I}_X(k)) = H^1(\mathcal{I}_Y(k))$ for all $k \in \mathbb{Z}$;
- $\text{reg}(Y) \leq \max\{\text{reg}(X), 3\}$;
- Y is cut out by at most cubic hypersurfaces.

In this paper, we slightly generalize these results to the case that I_X has an almost linear presentation, i.e., the minimal free resolution of R/I_X is of the following form:

$$\dots \rightarrow R(-3)^{\beta_{2,1}^R} \oplus R(-4) \rightarrow R(-2)^{\beta_{1,1}^R} \rightarrow R \rightarrow R/I_X \rightarrow 0.$$

In [1, Theorem 3.1], it was shown that if a generic projection π_q is an isomorphism then $H^1(\mathcal{I}_X(k)) = H^1(\mathcal{I}_Y(k))$ for all $k \geq 3$. This implies that Y is k -normal if and only if X is k -normal for $k \geq 3$.

In this paper, we will show that if a generic projection π_q is non-isomorphic then $H^1(\mathcal{I}_X(k)) = H^1(\mathcal{I}_Y(k))$ for all $k \in \mathbb{Z}$. This implies that Y is k -normal if and only if X is k -normal for all $k \in \mathbb{Z}$. Moreover, we will also prove that Y is cut out by at most quartic hypersurfaces. For the singular locus $Z = \{y \in Y \mid \text{the length of } \pi_q^{-1}(y) \geq 2\}$, it turns out that Z is either a linear space or a quadratic hypersurface in a linear subspace.

We use the partial elimination ideals introduced by M. Green ([8, Definition 6.1]) and the elimination mapping cone theorem ([2, Theorem 3.2]) to prove our results. In particular, the regularity of the first partial elimination ideal $K_1(I_X)$ will play a critical role in the proof of our result.

2. Partial elimination ideals

Let X be a nondegenerate reduced closed subscheme in \mathbb{P}^n and let $\pi_q : X \rightarrow Y = \pi_q(X) \subset \mathbb{P}^{n-1}$ be a generic projection from the center $q \in \text{Sec}(X) \setminus X$ where $\text{Sec}(X) = \mathbb{P}^n$. Considering a change of coordinates, we may assume that $q = [1, 0, \dots, 0]$. Then the i -th partial elimination ideal $K_i(I_X)$ can be defined as follows:

DEFINITION 2.1 (Partial elimination ideal). With the same notations as above, the i -th partial elimination ideal $K_i(I_X)$ is defined by

$$K_i(I_X) = \left\{ \frac{\partial^i f}{\partial x_0^i} \mid f \in I_X \text{ and } \frac{\partial^{i+1} f}{\partial x_0^{i+1}} = 0 \right\}$$

Algebraically, this can be rewritten as follows: if $f \in I_X$ has a leading term $\text{in}(f) = x_0^{d_0} \cdots x_n^{d_n}$ in the lexicographic order, we set $d_{x_0}(f) = d_0$, the leading power of x_0 in f . Let

$$\tilde{K}_i(I_X) = \bigoplus_{m \geq 0} \{f \in (I_X)_m \mid d_{x_0}(f) \leq i\}.$$

If $f \in \tilde{K}_i(I_X)$, we may write uniquely $f = x_0^i \bar{f} + g$ where $d_{x_0}(g) < i$. Then clearly $K_i(I_X)$ is the image of $\tilde{K}_i(I_X)$ in $S = K[x_1, \dots, x_n]$ under the map $f \mapsto \bar{f}$. Note that there is the following short exact sequence:

$$(2.1) \quad 0 \rightarrow \frac{\tilde{K}_{i-1}(I_X)}{I_Y} \rightarrow \frac{\tilde{K}_i(I_X)}{I_Y} \rightarrow K_i(I_X)(-i) \rightarrow 0.$$

It is known that set-theoretically $K_i(I_X)$ defines the following multiple loci [8, Proposition 6.2]

$$Z_i := \{p \in \pi_q(X) \mid \text{mult}_p(\pi_q(X)) \geq i + 1\}.$$

Moreover, there is a filtration on partial elimination ideals of I :

$$K_0(I_X) \subset K_1(I_X) \subset K_2(I_X) \subset \cdots \subset K_i(I_X) \subset \cdots \subset S = K[x_1, x_2, \dots, x_n].$$

Let us recall some definitions and basic properties of partial elimination ideals (See [4, Section 2]). If X is cut out by a homogeneous polynomial of degree d then, in generic coordinates, there exists a homogeneous polynomial $f \in I_X$ such that f is of the form

$$f = x_0^d + x_0^{d-1}g_{d-1} + \cdots + x_0g_1 + g_0$$

where g_i is a homogeneous form of degree $(d - i)$ in $S = K[x_1, \dots, x_n]$. Then we have the following exact sequence

$$(2.2) \quad 0 \rightarrow \tilde{K}_{d-1}(I_X) \rightarrow \bigoplus_{i=0}^{d-1} S(-i) \xrightarrow{\phi_0} R/I_X \rightarrow 0,$$

where the map ϕ_0 is defined by $\phi_0(e_i) = [x_0^i]$ for each free basis e_i of $S(-i)$, where $[x_0^i]$ is the quotient image of the monomial x_0^i in R/I_X (see [2], [5]). For the projection map $\pi_q : X \rightarrow Y = \pi_q(X) \subset \mathbb{P}^{n-1}$, we also have a natural map:

$$\alpha_d : 0 \rightarrow (S/I_Y)_d \rightarrow (R/I_X)_d.$$

Now we have the following commutative diagram.

LEMMA 2.2. *Let $X \subset \mathbb{P}^n = \mathbb{P}(V)$ be a nondegenerate reduced subscheme and let I_X be the defining saturated ideal of X . Then we have*

the following commutative diagram of S -modules for each $d > 0$:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \rightarrow & I_Y & \rightarrow & S & \rightarrow & S/I_Y \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \alpha \\
 (2.3) & 0 \rightarrow & \tilde{K}_{d-1}(I_X) & \rightarrow & \bigoplus_{i=0}^{d-1} S(-i) & \xrightarrow{\varphi_0} & R/I_X \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \rightarrow & \tilde{K}_{d-1}(I_X)/I_Y & \rightarrow & \bigoplus_{i=1}^{d-1} S(-i) & \rightarrow & \text{coker } \alpha \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Proof. See the proof of Lemma 2.1 in [4]. □

Finally, we remark the elimination mapping cone theorem in [2, Theorem 3.2]. Since we consider an outer projection $\pi_q : X \rightarrow Y \subset \mathbb{P}^{n-1}$, a graded S -module R/I_X is finitely generated. So we have the following long exact sequence by the map $\varphi : R/I_X(-1) \xrightarrow{\times x_0} R/I_X$ on the graded Koszul complex of R/I_X over S .

THEOREM 2.3 (Theorem 3.2 in [2]). *With the same notation as above, we have the following long exact sequence:*

$$\begin{aligned}
 \longrightarrow \text{Tor}_i^S(R/I_X, k)_{i+j} &\longrightarrow \text{Tor}_i^R(R/I_X, k)_{i+j} \longrightarrow \text{Tor}_{i-1}^S(R/I_X, k)_{i+j-1} \\
 &\xrightarrow{\delta} \text{Tor}_{i-1}^S(R/I_X, k)_{i+j} \longrightarrow \text{Tor}_{i-1}^R(R/I_X, k)_{i+j} \longrightarrow \cdots
 \end{aligned}$$

whose connecting homomorphism δ is the multiplicative map $\times x_0$.

3. Main result

THEOREM 3.1. *Let X be a nondegenerate reduced closed subscheme in \mathbb{P}^n . Assume that $\pi_q : X \rightarrow Y = \pi_q(X) \subset \mathbb{P}^{n-1}$ be a generic projection from the center $q \in \text{Sec}(X) \setminus X$ where $\text{Sec}(X) = \mathbb{P}^n$. Suppose that I_X has the almost minimal presentation, which is of the form*

$$R(-3)^{\beta_{2,1}} \oplus R(-4) \rightarrow R(-2)^{\beta_{1,1}} \rightarrow I_X \rightarrow 0.$$

Let $Z = \{y \in Y \mid \text{the length of } \pi_q^{-1}(y) \geq 2\}$ be the singular locus of the projection $\pi_q(X) \subset \mathbb{P}^{n-1}$. Then, we have

- (a) Z is either a linear space or a closed subscheme of degree two in a linear subspace;
- (b) $H^1(\mathcal{I}_X(k)) = H^1(\mathcal{I}_Y(k))$ for all $k \in \mathbb{Z}$;
- (c) $\text{reg}(Y) \leq \max\{\text{reg}(X), 4\}$;
- (d) Y is cut out by at most quartic hypersurfaces.

Proof. We may assume that I_X has the almost minimal presentation, which is of the form

$$R(-3)^{\beta_{2,1}^R} \oplus R(-4) \rightarrow R(-2)^{\beta_{1,1}^R} \rightarrow R \rightarrow R/I_X \rightarrow 0.$$

Then, it follows from Theorem 2.3 that

$$(3.1) \quad \beta_{1,2}^S \leq \beta_{2,2}^R = 1,$$

and the minimal free resolution of R/I_X as a graded S -module is of the form

$$(3.2) \quad \dots \rightarrow S(-2)^{\beta_{1,1}^S} \oplus S(-3)^{\beta_{1,2}^S} \rightarrow S \oplus S(-1) \xrightarrow{\varphi} R/I_X \rightarrow 0.$$

Now applying Lemma 2.2 for $d = 2$, we have the following diagram:

$$(3.3) \quad \begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & I_Y & \rightarrow & S & \rightarrow & S/I_Y & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow \alpha & & \\ 0 & \rightarrow & \tilde{K}_1(I_X) & \rightarrow & S \oplus S(-1) & \rightarrow & R/I_X & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & K_1(I_X)(-1) & \xrightarrow{\varphi} & S(-1) & \rightarrow & S/K_1(I_X)(-1) & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & 0 & & \end{array}$$

By the diagram chasing in (3.3) with (3.2), we obtain the following surjection map

$$\dots \rightarrow S(-2)^{\oplus \beta_{1,1}^S} \oplus S(-3)^{\oplus \beta_{1,2}^S} \rightarrow K_1(I_X)(-1) \rightarrow 0.$$

Then it follows from (3.1) that $K_1(I_X)$ is generated either by linear forms if $\beta_{1,2}^S = 0$; or by at most one quadric if $\beta_{1,2}^S = 1$. Since $K_1(I_X)$ is a radical ideal, the ideal $K_1(I_X)$ can be regarded as the singular locus Z of π_q (See [8, Proposition 6.2]). This proves (a).

Consider the following exact sequence

$$(3.4) \quad 0 \rightarrow \mathcal{O}_Y \rightarrow \pi_{q*}(\mathcal{O}_X) \rightarrow \mathcal{O}_Z(-1) \rightarrow 0.$$

Then, by taking global sections from the above sequence (3.4), we have the following commutative diagram of S -modules with exact rows and columns:

$$(3.5) \quad \begin{array}{ccccccc} & & & & & & 0 \\ & & & & & & \downarrow \\ & & & & & & \ker(\alpha) \\ & & & & & & \downarrow \\ & & 0 & & 0 & & \downarrow \\ 0 & \rightarrow & S/I_Y & \rightarrow & H_*^0(\mathcal{O}_Y) & \rightarrow & H_*^1(\mathcal{I}_Y) \rightarrow 0 \\ & & \downarrow \alpha & & \downarrow \gamma & & \downarrow \beta \\ 0 & \rightarrow & R/I_X & \rightarrow & H_*^0(\mathcal{O}_X) & \rightarrow & H_*^1(\mathcal{I}_X) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & [S/K_1(I_X)](-1) & \rightarrow & H_*^0(\mathcal{O}_Z(-1)) & \rightarrow & H_*^1(\mathcal{I}_Z(-1)) \rightarrow 0 \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

Consider the following exact sequence of graded S -modules

$$0 \rightarrow H_{\mathbf{m}}^0(S/K_1(I_X))(-1) \rightarrow [S/K_1(I_X)](-1) \rightarrow H_*^0(\mathcal{O}_Z)(-1) \rightarrow H_{\mathbf{m}}^1(S/K_1(I_X))(-1) \rightarrow 0,$$

where $H_{\mathbf{m}}^i(-)$ denotes the i -th local cohomology with respect to the irrelevant ideal $\mathbf{m} = (x_1, \dots, x_n)$ ([7, Corollary A1.12]). Since $K_1(I_X)$ is a saturated ideal, we see that $I_Z = K_1(I_X)$. Hence we have

$$\ker(\alpha) \cong H_{\mathbf{m}}^0(S/K_1(I_X))(-1) \text{ and } H_*^1(\mathcal{I}_Z(-1)) \cong H_{\mathbf{m}}^1(S/K_1(I_X))(-1).$$

Then it follows from snake lemma that

$$0 \rightarrow \ker(\alpha) \rightarrow S/K_1(I_X)(-1) \rightarrow H_*^0(\mathcal{O}_Z(-1)) \rightarrow H_*^1(\mathcal{I}_Z(-1)) \rightarrow 0.$$

Remark that the singular locus of π_q is defined by $K_1(I_X)$, which is generated by linear forms and at most one quadric polynomial. Hence Z is a complete intersection of degree ≤ 2 . Now we give a proof dividing the cases in terms of dimension of Z .

Case 1: $\dim(Z) = 0$.

Since $1 \leq \deg(Z) \leq 2$ we see that Z is a zero-dimensional closed subscheme, which is 1-regular. Hence we have

$$H^1(\mathcal{I}_Z(k)) = 0 \text{ for each } k \geq 0.$$

From (3.5), we see $H^1(\mathcal{I}_Y(k)) \cong H^1(\mathcal{I}_X(k))$ for all $k \in \mathbb{Z}$. Hence Y is m -normal if X is m -normal for each $m \geq 0$.

Case 2: $\dim(Z) \geq 1$.

In this case, note that Z is an arithmetically Cohen-Macaulay subscheme of dimension ≥ 1 . This implies that

$$H_*^1(\mathcal{I}_Z) = 0 \quad \text{and} \quad S/K_1(I_X) \cong H_*^0(\mathcal{O}_Z).$$

This implies that $H_*^1(\mathcal{I}_Y) \simeq H_*^1(\mathcal{I}_X)$. Hence X is m -normal if and only if Y is m -normal, for each $m \geq 0$. This proves (b).

Consider the left most column of exact sequence of S -modules in (3.5):

$$(3.6) \quad 0 \rightarrow S/I_Y \rightarrow R/I_X \rightarrow S/K_1(I_X)(-1) \rightarrow 0.$$

Let $d = \deg(Z) \leq 2$. Note that $S/K_1(I_X)$ is the coordinate ring of Z , which is a complete intersection scheme. Applying this to the short exact sequence (3.6), we can conclude that

$$\text{reg}(S/I_Y) \leq \max\{\text{reg}(R/I_X), d + 1\} \leq \max\{\text{reg}(R/I_X), 3\}.$$

Hence, $\text{reg}(I_Y) \leq \max\{\text{reg}(I_X), 4\}$ and thus Y is cut out by at most quartic hypersurfaces. This proves (c) and (d). \square

Let $\Sigma_d(X) := \{x \in X \mid \pi_q^{-1}(\pi_q(x)) \text{ has length } d\}$ be the d -secant locus of the projection π_q . It is known that if I_X has a minimal free presentation as in (1.1) then Z is a linear space Λ , and $\Sigma_2(X)$ is a hypersurface F of degree 2 in the linear span $\langle \Lambda, q \rangle$. This was very useful to classify non-normal del Pezzo varieties in [6] by Brodmann and Park.

COROLLARY 3.2 (Locus of 2-secant lines). *Let X be a nondegenerate reduced closed subscheme in \mathbb{P}^n . Assume that $\pi_q : X \rightarrow Y = \pi_q(X) \subset \mathbb{P}^{n-1}$ be a generic projection from the center $q \in \text{Sec}(X) \setminus X$ where $\text{Sec}(X) = \mathbb{P}^n$. Suppose that I_X has the almost minimal presentation, which is of the form*

$$R(-3)^{\beta_{2,1}} \oplus R(-4) \rightarrow R(-2)^{\beta_{1,1}} \rightarrow I_X \rightarrow 0.$$

Then we have

- (a) if $Z = \pi_q(\Sigma_2(X))$ is a linear subspace then $\Sigma_2(X)$ is a quadric hypersurface F in the linear span $\langle Z, q \rangle$;
- (b) if $Z = \pi_q(\Sigma_2(X))$ is a quadric hypersurface in linear subspace Λ then $\Sigma_2(X)$ is a hypersurface F of degree 4 in the linear span $\langle \Lambda, q \rangle$.

Proof. Note that Z is either a linear space or a quadric hypersurface in a linear subspace. We denote such a linear space by Λ . Since

$$\pi_q : \Sigma_2(X) \rightarrow Z \subset \pi_q(X)$$

is a 2 : 1 morphism, $\Sigma_2(X)$ is a hypersurface of degree $2 \deg(Z)$ in the linear span $\langle \Lambda, q \rangle$. \square

References

- [1] J. Ahn *Projections of Algebraic Varieties with Almost Linear Presentation I*, J Chungcheong Math Soc., **32** (2019), no. 1 15-21
- [2] J. Ahn and S. Kwak, *Graded mapping cone theorem, multiseccants and syzygies*, J. Algebr., **331** (2011), 243262.
- [3] J. Ahn and S. Kwak, *On syzygies, degree, and geometric properties of projective schemes with property $\mathbf{N}_{3,p}$* , J. Pure Appl. Algebr., **219** (2015) 27242739.
- [4] J. Ahn and S. Kwak, *The regularity of partial elimination ideals, Castelnuovo normality and syzygies*, J. Algebr., **533** (2019) 1-16.
- [5] J. Ahn, S. Kwak and Y. Song, *The degree complexity of smooth surfaces of codimension 2*, J. Symb. Comput., **47** (2012) 568-581.
- [6] M. Brodmann and E. Park, *On varieties of almost minimal degree I: Secant loci of rational normal scrolls*, J. Pure Appl. Algebra **214** (2010), 2033-2043.
- [7] D. Eisenbud, *The Geometry of Syzygies*, Springer-Verlag New York, **229** (2005)
- [8] M. Green, *Generic Initial Ideals*, in Six lectures on Commutative Algebra, (Elias J., Giral J.M., Miró-Roig, R.M., Zarzuela S., eds.), Progress in Mathematics **166**, Birkhäuser, (1998), 119-186.

*

Department of Mathematics Education
 Kongju National University
 Kongju 314-701, Republic of Korea
E-mail: jeamanahn@kongju.ac.kr