J. Appl. Math. & Informatics Vol. **39**(2021), No. 3 - 4, pp. 277 - 294 https://doi.org/10.14317/jami.2021.277

# WIJSMAN REGULARLY IDEAL INVARIANT CONVERGENCE OF DOUBLE SEQUENCES OF SETS

ERDINÇ DÜNDAR\*, ÖZER TALO

ABSTRACT. In this paper, we introduce the notions of Wijsman regularly invariant convergence types, Wijsman regularly  $(\mathcal{I}_{\sigma}, \mathcal{I}_{2}^{\sigma})$ -convergence, Wijsman regularly  $(\mathcal{I}_{\sigma}, \mathcal{I}_{2}^{\sigma})$ -convergence, Wijsman regularly  $(\mathcal{I}_{\sigma}, \mathcal{I}_{2}^{\sigma})$ -Cauchy double sequence and Wijsman regularly  $(\mathcal{I}_{\sigma}^{*}, \mathcal{I}_{2}^{\sigma*})$ -Cauchy double sequence of sets. Also, we investigate the relationships among this new notions.

AMS Mathematics Subject Classification : 40A05, 40A35. *Key words and phrases* : Double sequence, regularly ideal convergence, invariant convergence, regularly ideal Cauchy sequence, Wijsman convergence.

#### 1. Introduction

Throughout the paper,  $\mathbb{N}$  and  $\mathbb{R}$  denote the set of all positive integers and the set of all real numbers, respectively. The concept of convergence of real sequences has been extended to statistical convergence independently by Fast [23] and Schoenberg [39]. This concept was extended to the double sequences by Mursaleen and Edely [26]. The idea of  $\mathcal{I}$ -convergence was introduced by Kostyrko et al. [24] as a generalization of statistical convergence. Das et al. [8] introduced the concept of  $\mathcal{I}$ -convergence of double sequences in a metric space and studied some properties of this convergence. Tripathy and Tripathy [41] studied on  $\mathcal{I}$ -convergent and regularly  $\mathcal{I}$ -convergent double sequences. Dündar and Altay [10] introduced  $\mathcal{I}_2$ -convergence and regularly  $\mathcal{I}$ -convergence of double sequences. Also, Dündar [17] introduced regularly  $\mathcal{I}$ -convergence and regularly  $\mathcal{I}$ -Cauchy double sequences of functions. Recently, Dündar and Akın [22] studied regularly ideal convergence of double sequences. A lot of development have been made in this area after the works of [11, 12, 13, 16, 18, 25, 40].

Received August 31, 2020. Revised February 1, 2021. Accepted February 13, 2021.  $^{*}\mathrm{Corresponding}$  author.

<sup>@</sup> 2021 KSCAM.

Several authors have studied invariant convergent sequences (see, [7, 27, 28, 29, 31, 35, 36, 37, 38, 42]). Recently, the concepts of  $\sigma$ -uniform density of the set  $A \subseteq \mathbb{N}$ ,  $\mathcal{I}_{\sigma}$ -convergence and  $\mathcal{I}_{\sigma}^*$ -convergence of sequences of real numbers were defined by Nuray et al. [31]. The concept of  $\sigma$ -convergence of double sequences was studied by Çakan et al. [7] and the concept of  $\sigma$ -uniform density of  $A \subseteq \mathbb{N} \times \mathbb{N}$  was defined by Tortop and Dündar [42]. Dündar et al. [19] studied ideal invariant convergence of double sequences and some properties.

Now, we recall the basic definitions and concepts (See [4, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 20, 21, 18, 24, 25, 34, 41, 42, 40, 43, 44]).

Let  $\sigma$  be a mapping of the positive integers into themselves. A continuous linear functional  $\phi$  on  $\ell_{\infty}$ , the space of real bounded sequences, is said to be an invariant mean or a  $\sigma$ -mean if it satisfies following conditions:

- (1)  $\phi(x) \ge 0$ , when the sequence  $x = (x_n)$  has  $x_n \ge 0$ , for all n,
- (2)  $\phi(e) = 1$ , where e = (1, 1, 1, ...) and
- (3)  $\phi(x_{\sigma(n)}) = \phi(x_n)$ , for all  $x \in \ell_{\infty}$ .

The mappings  $\sigma$  are assumed to be one-to-one and such that  $\sigma^m(n) \neq n$ , for all positive integers n and m, where  $\sigma^m(n)$  denotes the m th iterate of the mapping  $\sigma$  at n. Thus,  $\phi$  extends the limit functional on c, the space of convergent sequences, in the sense that  $\phi(x) = \lim x$ , for all  $x \in c$ .

In the case  $\sigma$  is translation mappings  $\sigma(n) = n + 1$ , the  $\sigma$ -mean is often called a Banach limit and the space  $V_{\sigma}$ , the set of bounded sequences all of whose invariant means are equal, is the set of almost convergent sequences  $\hat{c}$ .

It can be shown that

$$V_{\sigma} = \left\{ (x_n) \in \ell_{\infty} : \lim_{m \to \infty} \frac{1}{m} \sum_{k=1}^m x_{\sigma^k(n)} = L, \text{ uniformly in } n \right\}.$$

A family of sets  $\mathcal{I} \subseteq 2^{\mathbb{N}}$  is called an ideal if and only if

(i)  $\emptyset \in \mathcal{I}$ , (ii) For each  $A, B \in \mathcal{I}$  we have  $A \cup B \in \mathcal{I}$ , (iii) For each  $A \in \mathcal{I}$  and each  $B \subseteq A$  we have  $B \in \mathcal{I}$ .

An ideal is called non-trivial if  $\mathbb{N} \notin \mathcal{I}$  and non-trivial ideal is called admissible if  $\{n\} \in \mathcal{I}$ , for each  $n \in \mathbb{N}$ .

Throughout the paper we take  $\mathcal{I}$  as an admissible ideal in  $\mathbb{N}$ .

A family of sets  $\mathcal{F} \subseteq 2^{\mathbb{N}}$  is called a filter if and only if

(i)  $\emptyset \notin \mathcal{F}$ , (ii) For each  $A, B \in \mathcal{F}$  we have  $A \cap B \in \mathcal{F}$ , (iii) For each  $A \in \mathcal{F}$  and each  $B \supseteq A$  we have  $B \in \mathcal{F}$ .

For any ideal there is a filter  $\mathcal{F}(\mathcal{I})$  corresponding with  $\mathcal{I}$ , given by  $\mathcal{F}(\mathcal{I}) = \{M \subset \mathbb{N} : (\exists A \in \mathcal{I})(M = \mathbb{N} \setminus A)\}.$ 

An admissible ideal  $\mathcal{I} \subset 2^{\mathbb{N}}$  is said to satisfy the property (AP), if for every countable family of mutually disjoint sets  $\{A_1, A_2, ...\}$  belonging to  $\mathcal{I}$ , there exists a countable family of sets  $\{B_1, B_2, ...\}$  such that  $A_j \Delta B_j$  is a finite set for  $j \in \mathbb{N}$  and  $B = \bigcup_{j=1}^{\infty} B_j \in \mathcal{I}$ .

A non-trivial ideal  $\mathcal{I}_2$  of  $\mathbb{N} \times \mathbb{N}$  is called strongly admissible ideal if  $\{i\} \times \mathbb{N}$ and  $\mathbb{N} \times \{i\}$  belong to  $\mathcal{I}_2$  for each  $i \in N$ . It is evident that a strongly admissible ideal is admissible also.

Throughout the paper, we take  $\mathcal{I}_2$  as a strongly admissible ideal in  $\mathbb{N} \times \mathbb{N}$ .  $\mathcal{I}_2^0 = \{ A \subset \mathbb{N} \times \mathbb{N} : (\exists m(A) \in \mathbb{N}) (i, j \ge m(A) \Rightarrow (i, j) \notin A) \}.$  Then,  $\mathcal{I}_2^0$  is a strongly admissible ideal and clearly an ideal  $\mathcal{I}_2$  is strongly admissible if and only if  $\mathcal{I}_2^0 \subset \mathcal{I}_2$ .

An admissible ideal  $\mathcal{I}_2 \subset 2^{\mathbb{N} \times \mathbb{N}}$  satisfies the property (AP2) if for every countable family of mutually disjoint sets  $\{E_1, E_2, ...\}$  belonging to  $\mathcal{I}_2$ , there exists a countable family of sets  $\{F_1, F_2, ...\}$  such that  $E_j \Delta F_j \in \mathcal{I}_2^0$ , i.e.,  $E_j \Delta F_j$  is included in the finite union of rows and columns in  $\mathbb{N} \times \mathbb{N}$  for each  $j \in \mathbb{N}$  and  $F = \bigcup_{i=1}^{\infty} F_i \in \mathcal{I}_2$  (hence  $F_j \in \mathcal{I}_2$ , for each  $j \in \mathbb{N}$ ).

Let  $A \subseteq \mathbb{N}$  and

$$s_m = \min_n |A \cap \{\sigma(n), \sigma^2(n), \cdots, \sigma^m(n)\}|,$$
  
$$S_m = \max_n |A \cap \{\sigma(n), \sigma^2(n), \cdots, \sigma^m(n)\}|.$$

If the limits  $\underline{V}(A) = \lim_{m \to \infty} \frac{s_m}{m}$  and  $\overline{V}(A) = \lim_{m \to \infty} \frac{S_m}{m}$  exist, then they are called a lower and upper  $\sigma$ -uniform density of the set A, respectively. If  $V(A) = \overline{V}(A)$ , then  $V(A) = \underline{V}(A) = \overline{V}(A)$  is called  $\sigma$ -uniform density of A.

Denote by  $\mathcal{I}_{\sigma}$  the class of all  $A \subseteq \mathbb{N}$  with V(A) = 0.

Let  $(X, \rho)$  be a separable metric space. For any point  $x \in X$  and any nonempty subset A of X, we define the distance from x to A by  $d(x, A) = \inf_{x \in A} \rho(x, a)$ . Throughout the paper, we let  $\mathcal{I}_{\sigma} \subset 2^{\mathbb{N}}$  be an admissible ideal,  $(X, \rho)$  be a

separable metric space and  $A, A_k$  be any non-empty closed subsets of X.

A sequence  $\{A_k\}$  is said to be Wijsman  $\mathcal{I}$ -invariant convergent or  $\mathcal{I}_W^{\sigma}$ -convergent to A if for every  $\varepsilon > 0$ , the set  $A(\varepsilon, x) = \{k : |d(x, A_k) - d(x, A)| \ge \varepsilon\} \in \mathcal{I}_{\sigma}$ , that is,  $V(A(\varepsilon, x)) = 0$ . In this case, we write  $A_k \to A(\mathcal{I}_W^{\sigma})$ .

A sequence  $\{A_k\}$  is Wijsman  $\mathcal{I}^*$ -invariant convergent or  $\mathcal{I}_W^{*\sigma}$ -convergent to A if there exists a set  $M = \{m_1 < \cdots < m_k < \cdots\} \in \mathcal{F}(\mathcal{I}_{\sigma})$  such that for each  $x \in X$ ,  $\lim_{k \to \infty} d(x, A_{m_k}) = d(x, A)$ .

A sequence  $\{A_k\}$  is said to be Wijsman  $\mathcal{I}$ -invariant Cauchy sequence or  $\mathcal{I}_W^{\sigma}$ -Cauchy sequence if for every  $\varepsilon > 0$  and for each  $x \in X$ , there exists a number  $N = N(\varepsilon, x) \in \mathbb{N}$  such that  $A(\varepsilon, x) = \{k : |d(x, A_k) - d(x, A_N)| \ge \varepsilon\} \in \mathcal{I}_{\sigma}$ , that is,  $V(A(\varepsilon, x)) = 0$ .

A sequence  $\{A_k\}$  is said to be Wijsman  $\mathcal{I}^*$ -invariant Cauchy sequence or  $\mathcal{I}_W^{*\sigma}$ -Cauchy sequence if there exists a set  $M = \{m_1 < \cdots < m_k < \ldots\} \in \mathcal{F}(\mathcal{I}_{\sigma})$  such that  $\lim_{k,p\to\infty} |d(x,A_{m_k}) - d(x,A_{m_p})| = 0$ , for each  $x \in X$ . Let  $A \subseteq \mathbb{N} \times \mathbb{N}$  and

$$s_{mn} = \min_{k,j} \left| A \cap \left\{ \left( \sigma(k), \sigma(j) \right), \left( \sigma^2(k), \sigma^2(j) \right), ..., \left( \sigma^m(k), \sigma^n(j) \right) \right\} \right|,$$
  
$$S_{mn} = \max_{k,j} \left| A \cap \left\{ \left( \sigma(k), \sigma(j) \right), \left( \sigma^2(k), \sigma^2(j) \right), ..., \left( \sigma^m(k), \sigma^n(j) \right) \right\} \right|.$$

If the limits  $\underline{V_2}(A) = \lim_{m,n\to\infty} \frac{s_{mn}}{mn}$  and  $\overline{V_2}(A) = \lim_{m,n\to\infty} \frac{S_{mn}}{mn}$  exist, then they are called a lower and an upper  $\sigma$ -uniform density of the set A, respectively. If  $\underline{V_2}(A) = \overline{V_2}(A)$ , then  $V_2(A) = \underline{V_2}(A) = \overline{V_2}(A)$  is called the  $\sigma$ -uniform density of A.

Denote by  $\mathcal{I}_2^{\sigma}$  the class of all  $A \subseteq \mathbb{N} \times \mathbb{N}$  with  $V_2(A) = 0$ .

Throughout the paper, we let  $\mathcal{I}_2^{\sigma} \subset 2^{\mathbb{N} \times \mathbb{N}}$  be a strongly admissible ideal and  $A, B_k, C_k, A_{kj}$  be any non-empty closed subsets of X.

A double sequence  $\{A_{kj}\}$  is said to be bounded if  $\sup_{k,j} d(x, A_{kj}) < \infty$ , for each

 $x \in X$ . The set of all bounded double sequences of sets will be denoted by  $L^2_{\infty}$ .

A double sequence  $\{A_{kj}\}$  is said to be Wijsman invariant convergent to A if for each  $x \in X$ ,

$$\lim_{n,n\to\infty}\frac{1}{mn}\sum_{k,j=1,1}^{m,n}d(x,A_{\sigma^k(s),\sigma^j(t)})=d(x,A), \text{ uniformly in } s,t$$

A double sequence  $\{A_{kj}\}$  is said to be Wijsman  $\mathcal{I}_2$ -invariant convergent or  $\mathcal{I}_{W_2}^{\sigma}$ -convergent to A, if for every  $\varepsilon > 0$ ,  $A(\varepsilon, x) = \{(k, j) : |d(x, A_{kj}) - d(x, A)| \ge \varepsilon\} \in \mathcal{I}_2^{\sigma}$  that is,  $V_2(A(\varepsilon, x)) = 0$ . In this case, we write  $A_{kj} \to A(\mathcal{I}_{W_2}^{\sigma})$ .

A double sequence  $\{A_{kj}\}$  is Wijsman  $\mathcal{I}_2^*$ -invariant convergent or  $\mathcal{I}_{W_2}^{*\sigma}$ -convergent to A if and only if there exists a set  $M_2 \in \mathcal{F}(\mathcal{I}_2^{\sigma}) \ (\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2^{\sigma})$  such that for each  $x \in X$  and  $(k, j) \in M_2$ ,  $\lim_{k,j \to \infty} d(x, A_{kj}) = d(x, A)$ . In this case, we write  $\mathcal{I}_{W_2}^{*\sigma} - \lim_{k,j \to \infty} d(x, A_{kj}) = d(x, A)$ .

A double sequence  $\{A_{kj}\}$  is said to be Wijsman  $\mathcal{I}_2$ -invariant Cauchy sequence or  $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy sequence, if for every  $\varepsilon > 0$  and each  $x \in X$ , there exist  $r = r(\varepsilon, x), \ s = s(\varepsilon, x) \in \mathbb{N}$  such that  $A(\varepsilon, x) = \{(k, j) : |d(x, A_{kj}) - d(x, A_{rs})| \ge \varepsilon\} \in \mathcal{I}_2^{\sigma}$ , that is,  $V_2(A(\varepsilon, x)) = 0$ .

A double sequence  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{*\sigma}$ -Cauchy if there exists a set  $M_2 \in \mathcal{F}(\mathcal{I}_2^{\sigma})$ (i.e.,  $\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2^{\sigma}$ ) such that for every  $x \in X$  and  $(k, j), (p, q) \in M_2$  $\lim_{k,j,p,q \to \infty} |d(x, A_{kj}) - d(x, A_{pq})| = 0.$ 

A double sequence  $\{A_{kj}\}$  is said to be Wijsman regularly convergent  $(R(W_2, W)$ convergent) if it is convergent in Pringsheim's sense and for each  $x \in X$  the limits  $\lim_{k \to \infty} d(x, A_{kj}), (j \in \mathbb{N}) \text{ and } \lim_{j \to \infty} d(x, A_{kj}), (k \in \mathbb{N}) \text{ exist. Note that if } \{A_{kj}\} \text{ is}$ Wijsman regularly convergent to A, then the limits

$$\lim_{k \to \infty} \lim_{j \to \infty} d(x, A_{kj}) = d(x, A) \text{ and } \lim_{j \to \infty} \lim_{k \to \infty} d(x, A_{kj}) = d(x, A)$$

exist and we write  $R(W_2, W) - \lim_{k, j \to \infty} d(x, A_{kj}) = d(x, A)$  or  $A_{kj} \xrightarrow{R(W_2, W)} A$ .

A double sequence  $\{A_{kj}\}$  is said to be regularly  $(\mathcal{I}_{W_2}, \mathcal{I}_W)$ -convergent  $(R(\mathcal{I}_{W_2}, \mathcal{I}_W)$ convergent) if it is  $\mathcal{I}_{W_2}$ -convergent in Pringsheim's sense and for every  $\varepsilon > 0$  and each  $x \in X$ ,  $\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, K_j)| \ge \varepsilon\} \in \mathcal{I}$ , for some  $K_j \in X$  and each

 $j \in \mathbb{N}$ , and  $\{j \in \mathbb{N} : |d(x, A_{kj}) - d(x, L_k)| \ge \varepsilon\} \in \mathcal{I}$ , for some  $L_k \in X$  and each  $k \in \mathbb{N}$ .

Note that if  $\{A_{kj}\}$  is  $R(\mathcal{I}_{W_2}, \mathcal{I}_W)$ -convergent to A, then we write  $R(\mathcal{I}_{W_2}, \mathcal{I}_W) - \lim_{k,j \to \infty} d(x, A_{kj}) = d(x, A)$  or  $A_{kj} \xrightarrow{R(\mathcal{I}_{W_2}, \mathcal{I}_W)} A$ .

A double sequence  $\{A_{kj}\}$  is said to be regularly  $(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*)$ -convergent  $(R(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*)$ -convergent) if there exist the sets  $M \in \mathcal{F}(\mathcal{I}_2), M_1 \in \mathcal{F}(\mathcal{I})$  and  $M_2 \in \mathcal{F}(\mathcal{I})$  (i.e.,  $\mathbb{N} \times \mathbb{N} \setminus M \in \mathcal{I}_2, \mathbb{N} \setminus M_1 \in \mathcal{I}$  and  $\mathbb{N} \setminus M_2 \in \mathcal{I}$ ) such that the limits

$$\lim_{\substack{k,j\to\infty\\(k,j)\in M}} d(x,A_{kj}), \quad \lim_{\substack{k\to\infty\\k\in M_1}} d(x,A_{kj}) \text{ and } \lim_{\substack{j\to\infty\\j\in M_2}} d(x,A_{kj})$$

exist for each fixed  $k \in \mathbb{N}$  and each fixed  $j \in \mathbb{N}$ , respectively.

Note that if  $\{A_{kj}\}$  is  $R(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*)$ -convergent to A, then for each  $x \in X$  the limits  $\lim_{k \to \infty} \lim_{j \to \infty} d(x, A_{kj})$  and  $\lim_{j \to \infty} \lim_{k \to \infty} d(x, A_{kj})$  exist and are equal to A and

we write 
$$R(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*) - \lim_{k, j \to \infty} d(x, A_{kj}) = d(x, A)$$
 or  $A_{kj} \xrightarrow{R(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*)} A$ .

A double sequence  $\{A_{kj}\}$  is said to be regularly  $(\mathcal{I}_{W_2}, \mathcal{I}_W)$ -Cauchy  $(R(\mathcal{I}_{W_2}, \mathcal{I}_W)$ -Cauchy) if it is  $\mathcal{I}_2$ -Cauchy in Pringsheim's sense and for every  $\varepsilon > 0$  and each  $x \in X$ , there exist  $m_j = m_j(\varepsilon, x), n_k = n_k(\varepsilon, x) \in \mathbb{N}$  such that

 $\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{mj})| \ge \varepsilon\} \in \mathcal{I}, (j \in \mathbb{N}), \\ \{j \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{kn_k})| \ge \varepsilon\} \in \mathcal{I}, (k \in \mathbb{N}).$ 

A double sequence  $\{A_{kj}\}$  is said to be regularly  $(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*)$ -Cauchy  $(R(\mathcal{I}_{W_2}^*, \mathcal{I}_W^*)$ -Cauchy) if there exist the sets  $M \in \mathcal{F}(\mathcal{I}_2)$ ,  $M_1 \in \mathcal{F}(\mathcal{I})$  and  $M_2 \in \mathcal{F}(\mathcal{I})$  (i.e.,  $\mathbb{N} \times \mathbb{N} \setminus M \in \mathcal{I}_2$ ,  $\mathbb{N} \setminus M_1 \in \mathcal{I}$  and  $\mathbb{N} \setminus M_2 \in \mathcal{I}$ ), and for every  $\varepsilon > 0$  and each  $x \in X$  there exist  $N = N(\varepsilon, x), s = s(\varepsilon, x), t = t(\varepsilon, x), m_j = m_j(\varepsilon, x), n_k = n_k(\varepsilon, x) \in \mathbb{N}$  such that whenever  $k, j, s, t, m_j, n_k \geq N$ , we have

$$\begin{aligned} |d(x, A_{kj}) - d(x, A_{st})| &< \varepsilon, \text{ for } (m, n), (s, t) \in M, \\ |d(x, A_{kj}) - d(x, A_{mjj})| &< \varepsilon, \text{ for each } k \in M_1 \text{ and each } j \in \mathbb{N} \\ |d(x, A_{kj}) - d(x, A_{knk})| &< \varepsilon, \text{ for each } j \in M_2 \text{ and each } k \in \mathbb{N}. \end{aligned}$$

**Lemma 1.1.** [42] Let  $\{A_{kj}\}$  be bounded sequence. If  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent to A, then  $\{A_{kj}\}$  is Wijsman invariant convergent to A.

**Lemma 1.2.** [42] Let 0 .

- (i) If  $A_{kj} \to A([W_2V_\sigma]_p)$ , then  $A_{kj} \to A(\mathcal{I}_{W_2}^{\sigma})$ .
- (ii) If  $\{A_{kj}\} \in L^2_{\infty}$  and  $A_{kj} \to A(\mathcal{I}^{\sigma}_{W_2})$ , then  $A_{kj} \to A([W_2V_{\sigma}]_p)$ .
- (iii) If  $\{A_{kj}\} \in L^2_{\infty}$ , then  $\{A_{kj}\}$  is  $\mathcal{I}^{\sigma}_{W_2}$ -convergent to A if and only if  $A_{kj} \to A([W_2V_{\sigma}]_p)$ .

**Lemma 1.3.** [42] If a sequence  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{*\sigma}$ -convergent to A, then this sequence is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent to A.

**Lemma 1.4.** [42] Let  $\mathcal{I}_2^{\sigma}$  has property (AP2). If  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent to A, then  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{*\sigma}$ -convergent to A.

**Lemma 1.5.** [42] If a double sequence  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent, then  $\{A_{kj}\}$  is an  $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy double sequence of sets.

**Lemma 1.6.** [42] If a double sequence  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{*\sigma}$ -Cauchy double sequence, then  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy double sequence of sets.

#### 2. Main Results

Now, we introduce the notions of Wijsman regularly invariant convergence, Wijsman regularly strongly invariant convergence, Wijsman regularly *p*-strongly invariant convergence, Wijsman regularly  $(\mathcal{I}_{\sigma}, \mathcal{I}_{2}^{\sigma})$ -convergence, Wijsman regularly  $(\mathcal{I}_{\sigma}^{*}, \mathcal{I}_{2}^{\sigma*})$ -convergence, Wijsman regularly  $(\mathcal{I}_{\sigma}, \mathcal{I}_{2}^{\sigma})$ -Cauchy double sequence, Wijsman regularly  $(\mathcal{I}_{\sigma}^{*}, \mathcal{I}_{2}^{\sigma*})$ -Cauchy double sequence of sets and investigate the relationship among them.

**Definition 2.1.** A double sequence  $\{A_{kj}\}$  is said to be Wijsman regularly invariant convergent  $(R(W^{\sigma}, W_2^{\sigma})$ -convergent) if it is Wijsman invariant convergent in Pringsheim's sense and the following limits hold:

$$\lim_{m \to \infty} \frac{1}{m} \sum_{k=0}^{m} d(x, A_{\sigma^k(s), \sigma^j(t)}) = d(x, B_j), \text{ uniformly in } s,$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ , and

$$\lim_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n} d(x, A_{\sigma^k(s), \sigma^j(t)}) = d(x, C_k), \text{ uniformly in } t,$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ .

Note that if  $\{A_{kj}\}$  is  $R(W^{\sigma}, W_2^{\sigma})$ -convergent to A, the following limits hold:

$$\lim_{m \to \infty} \lim_{n \to \infty} \frac{1}{mn} \sum_{k=0}^{m} \sum_{j=0}^{n} d(x, A_{\sigma^k(s), \sigma^j(t)}) = d(x, A), \text{ uniformly in } s, t$$

and

$$\lim_{n \to \infty} \lim_{m \to \infty} \frac{1}{mn} \sum_{j=0}^{n} \sum_{k=0}^{m} d(x, A_{\sigma^k(s), \sigma^j(t)}) = d(x, A), \text{ uniformly in } s, t,$$

for each  $x \in X$ . In this case, for each  $x \in X$ , we write

$$R(W^{\sigma}, W_2^{\sigma}) - \lim_{m, n \to \infty} \sum_{k, j=0, 0}^{m, n} d(x, A_{\sigma^k(s), \sigma^j(t)}) = d(x, A) \text{ or } A_{kj} \xrightarrow{R(W^{\sigma}, W_2^{\sigma})} A,$$

uniformly in s, t.

**Definition 2.2.** A double sequence  $\{A_{kj}\}$  is said to be Wijsman regularly strongly invariant convergent  $(R[W^{\sigma}, W_2^{\sigma}]$ -convergent) if it is Wijsman strongly invariant convergent in Pringsheim's sense and the following limits hold:

$$\lim_{m \to \infty} \frac{1}{m} \sum_{k=0}^{m} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, B_j)| = 0, \text{ uniformly in } s,$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ , and

$$\lim_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, C_k)| = 0, \text{ uniformly in } t,$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ .

Note that if  $\{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]$ -convergent to A, the following limits hold:

$$\lim_{m \to \infty} \lim_{n \to \infty} \frac{1}{mn} \sum_{k=0}^{m} \sum_{j=0}^{n} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, A)| = 0, \text{ uniformly in } s, t$$

and

$$\lim_{n \to \infty} \lim_{m \to \infty} \frac{1}{mn} \sum_{j=0}^{n} \sum_{k=0}^{m} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, A)| = 0, \text{ uniformly in } s, t,$$

for each  $x \in X$ . In this case, for each  $x \in X$ , we write

$$R[W^{\sigma}, W_2^{\sigma}] - \lim_{m, n \to \infty} \sum_{k, j=0, 0}^{m, n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| = 0 \text{ or } A_{kj} \xrightarrow{R[W^{\sigma}, W_2^{\sigma}]} A$$

uniformly in s, t.

**Definition 2.3.** Let  $0 . A double sequence <math>\{A_{kj}\}$  is said to be Wijsman regularly *p*-strongly invariant convergent  $(R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent), if it is Wijsman *p*-strongly invariant convergent in Pringsheim's sense and the following limits hold:

$$\lim_{m \to \infty} \frac{1}{m} \sum_{k=0}^{m} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, B_j)|^p = 0, \text{ uniformly in } s,$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$  and

$$\lim_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, C_{k})|^{p} = 0, \text{ uniformly in } t,$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ .

Note that if  $\{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent to A, the following limits hold:

$$\lim_{m \to \infty} \lim_{n \to \infty} \frac{1}{mn} \sum_{k=0}^{m} \sum_{j=0}^{n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p = 0, \text{ uniformly in } s, t$$

and

$$\lim_{n \to \infty} \lim_{m \to \infty} \frac{1}{mn} \sum_{j=0}^{n} \sum_{k=0}^{m} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, A)|^{p} = 0, \text{ uniformly in } s, t,$$

for each  $x \in X$ . In this case, for each  $x \in X$ , we write

$$R[W^{\sigma}, W_{2}^{\sigma}]_{p} - \lim_{m, n \to \infty} \sum_{k, j=0, 0}^{m, n} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, A)| = 0 \text{ or } A_{kj} \xrightarrow{R[W^{\sigma}, W_{2}^{\sigma}]_{p}} A,$$

uniformly in s, t.

**Definition 2.4.** A double sequence  $\{A_{kj}\}$  is said to be Wijsman regularly ideal invariant convergent  $(R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent), if it is Wijsman ideal invariant convergent in Pringsheim's sense and for every  $\varepsilon > 0$  and each  $x \in X$ , the following limits hold:

$$\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \varepsilon\} \in \mathcal{I}_{\sigma},$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ , and

$$\{j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| \ge \varepsilon\} \in \mathcal{I}_{\sigma},$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ .

Note that if  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent to A, then for each  $x \in X$ , we write

$$R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma}) - \lim d(x, A_{kj}) = d(x, A) \text{ or } A_{kj} \xrightarrow{R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})} A.$$

**Theorem 2.5.** Assume that  $\{A_{kj}\}$  is a bounded double sequence. If  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent, then  $\{A_{kj}\}$  is  $R(W^{\sigma}, W_2^{\sigma})$ -convergent.

*Proof.* Let  $\{A_{kj}\}$  is a bounded double sequence and  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent to A. Then,  $\{A_{kj}\}$  is Wijsman ideal invariant convergent in Pringsheim's sense and for every  $\varepsilon > 0$  and each  $x \in X$ , the followings hold:

$$\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \varepsilon\} \in \mathcal{I}_{\sigma}$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ , and

$$\{j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| \ge \varepsilon\} \in \mathcal{I}_{\sigma}$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ . Since  $\{A_{kj}\}$  is Wijsman ideal invariant convergent in Pringsheim's sense, then by Lemma 1.1  $\{A_{kj}\}$  is Wijsman invariant convergent to A.

Now, let  $\varepsilon > 0$ . We estimate

$$u(m,s) = \left|\frac{1}{m}\sum_{k=0}^{m} d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, B_j)\right|, \text{ uniformly in } s,$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ . Then, we have

$$u(m,s) \le u^1(m,s) + u^2(m,s)$$

where

$$u^{1}(m,s) = \frac{1}{m} \sum_{\substack{k=0 \ |d(x,A_{\sigma^{k}(s),\sigma^{j}(t)}) - d(x,B_{j})| \ge \varepsilon}}^{m} |d(x,A_{\sigma^{k}(s),\sigma^{j}(t)}) - d(x,B_{j})|$$

and

$$u^{2}(m,s) = \frac{1}{m} \sum_{\substack{k=0\\|d(x,A_{\sigma^{k}(s),\sigma^{j}(t)}) - d(x,B_{j})| < \varepsilon}}^{m} |d(x,A_{\sigma^{k}(s),\sigma^{j}(t)}) - d(x,B_{j})|,$$

uniformly in s, for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ . Therefore, we have  $u^2(m,s) < \varepsilon$ , for every  $s = 1, 2, \ldots$  The boundedness of  $\{A_{kj}\}$  implies that there exists K > 0 such that for each  $x \in X$ ,

$$|d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, B_j)| \le K, \ (k, s = 1, 2, \dots),$$

then this implies that

$$u^{1}(m,s) \leq \frac{K}{m} \Big| \Big\{ 1 \leq k \leq m : |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| \geq \varepsilon \Big\} \Big|$$
  
$$\leq K \frac{\max_{s} \Big| \Big\{ 1 \leq k \leq m : |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| \geq \varepsilon \Big\} \Big|}{m}$$
  
$$= K \frac{S_{m}}{m}$$

and so,  $\{A_{kj}\}$  is Wijsman  $\sigma$ -convergent to  $B_j$ .

Similarly, we can show that  $\{A_{kj}\}$  is Wijsman  $\sigma$ -convergent to  $C_k$ . Hence,  $\{A_{kj}\}$  is  $R(W^{\sigma}, W_2^{\sigma})$ -convergent. 

**Theorem 2.6.** Let 0 . Then,

- (i) If  $\{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent, then  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent.
- (i) If  $\{A_{kj}\} \in L^2_{\infty}$  and  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent, then  $\{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent. (ii) If  $\{A_{kj}\} \in L^2_{\infty}$ , then  $\{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent if and only if  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$  is convergent.

*Proof.* (i) Let  $x = \{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent. Then, it is Wijsman pstrongly invariant convergent in Pringsheim's sense and the following limits holds:

$$\lim_{m \to \infty} \frac{1}{m} \sum_{k=0}^{m} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})|^{p} = 0, \text{ uniformly in } s,$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ , and

$$\lim_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, C_{k})|^{p} = 0, \text{ uniformly in } t,$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ . Since  $\{A_{kj}\}$  is Wijsman *p*-strongly invariant convergent in Pringsheim's sense, then by Lemma 1.2  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent.

Also, for every  $\varepsilon > 0$ , some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$  we can write

$$\sum_{k=1}^{m} \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right|^{p}$$

$$\geq \sum_{\substack{k=1 \\ |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| \geq \varepsilon}^{m} |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})|^{p}$$

$$\geq \varepsilon^{p} \left| \left\{ k \leq m : |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| \geq \varepsilon \right\} \right|$$

$$\geq \varepsilon^{p} \max_{s} \left| \left\{ k \leq m : |d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| \geq \varepsilon \right\} \right|$$

and

$$\frac{1}{m} \sum_{k=1}^{m} \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right|^{p}$$

$$\geq \varepsilon^{p} \frac{\max_{s} \left| \left\{ k \leq m : \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right| \geq \varepsilon \right\} \right|}{m}$$

$$= \varepsilon^{p} \frac{S_{m}}{m},$$

for every  $s = 1, 2, \ldots$ . This implies

$$\lim_{m \to \infty} \frac{S_m}{m} = 0$$

and so  $\{A_{kj}\}$  is  $\mathcal{I}_W^{\sigma}$ -convergent to  $B_j$ .

Similarly, we can show that  $\{A_{kj}\}$  is  $\mathcal{I}_W^{\sigma}$ -convergent to  $C_k$ . Hence,  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent.

(*ii*) Let  $\{A_{kj}\} \in L^2_{\infty}$  and  $\{A_{kj}\}$  is  $R(\mathcal{I}^{\sigma}_W, \mathcal{I}^{\sigma}_{W_2})$ -convergent. Then,  $\{A_{kj}\}$  is Wijsman ideal invariant convergent in Pringsheim's sense and for every  $\varepsilon > 0$  and  $x \in X$  the followings hold:

$$\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \varepsilon\} \in \mathcal{I}_{\sigma},$$

for some  $B_j \in X$ , each  $j \in \mathbb{N}$  and each  $x \in X$ , and

$$\{j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| \ge \varepsilon\} \in \mathcal{I}_{\sigma},$$

for some  $C_k \in X$ , each  $k \in \mathbb{N}$  and each  $x \in X$ . Since  $\{A_{kj}\}$  is Wijsman ideal invariant convergent in Pringsheim's sense, then by Lemma 1.2,  $\{A_{kj}\}$  is Wijsman *p*-strongly invariant convergent. Let  $0 and <math>\varepsilon > 0$ . Since  $\{A_{kj}\}$  is bounded,  $\{A_{kj}\}$  implies that there exists K > 0 such that

$$|d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, B_j)| \le K, \ (j \in \mathbb{N}).$$

for all  $k, s \in \mathbb{N}$ , some  $B_j \in X$  and each  $x \in X$ . Then, for every  $s = 1, 2, \ldots$  we have

$$\frac{1}{m}\sum_{k=1}^{m} \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right|^{p} \\
= \frac{1}{m}\sum_{\substack{k=1\\|d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| \ge \varepsilon}^{m} \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right|^{p} \\
+ \frac{1}{m}\sum_{\substack{k=1\\|d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j})| < \varepsilon}^{m} \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right|^{p} \\
\leq K \frac{\max_{s} \left| \left\{ k \le m : \left| d(x, A_{\sigma^{k}(s), \sigma^{j}(t)}) - d(x, B_{j}) \right| \ge \varepsilon \right\} \right|}{m} + \varepsilon^{p} \\
\leq K \frac{S_{m}}{m} + \varepsilon^{p}.$$

Hence, we obtain

$$\lim_{m \to \infty} \frac{1}{m} \sum_{k=1}^{m} \left| d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, B_j) \right|^p = 0, \text{ uniformly in } s$$

and so,  $\{A_{kj}\}$  is Wijsman *p*-strongly invariant convergent to  $B_j$ .

Similarly, we show that  $\{A_{kj}\}$  is Wijsman *p*-strongly invariant convergent to  $C_k$ . Hence,  $\{A_{kj}\}$  is  $R[W^{\sigma}, W_2^{\sigma}]_p$ -convergent.

(*iii*) This is immedate consequence of (i) and (ii).  $\Box$ 

**Definition 2.7.** A double sequence  $\{A_{kj}\}$  is said to be regularly  $(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ convergent  $(R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -convergent) if and only if there exist the sets  $M \in \mathcal{F}(\mathcal{I}_2^{\sigma})$ ,  $M_1 \in \mathcal{F}(\mathcal{I}_{\sigma})$  and  $M_2 \in \mathcal{F}(\mathcal{I}_{\sigma})$  (i.e.,  $\mathbb{N} \times \mathbb{N} \setminus M \in \mathcal{I}_2^{\sigma}$ ,  $\mathbb{N} \setminus M_1 \in \mathcal{I}_{\sigma}$  and  $\mathbb{N} \setminus M_2 \in \mathcal{I}_{\sigma}$ ) such that for each  $x \in X$ , the following limits hold:

$$\lim_{\substack{k,j\to\infty\\(k,j)\in M}} d(x,A_{kj}), \lim_{\substack{k\to\infty\\k\in M_1}} d(x,A_{kj}) \ (j\in\mathbb{N}) \text{ and } \lim_{\substack{j\to\infty\\j\in M_2}} d(x,A_{kj}) \ (k\in\mathbb{N}).$$

Note that if  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -convergent to A, then for each  $x \in X$ , we write

$$R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*}) - \lim_{k,j \to \infty} d(x, A_{kj}) = d(x, A) \text{ or } d(x, A_{kj}) \xrightarrow{R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})} A$$

**Theorem 2.8.** If a double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -convergent, then  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent.

*Proof.* Let  $\{A_{kj}\}$  be  $R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -convergent. Then,  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma*}$ -convergent and so, by Lemma 1.3  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent. Also, there exist the sets

 $M_1, M_2 \in \mathcal{F}(\mathcal{I}_{\sigma})$  such that for each  $x \in X$ ,

 $(\forall \varepsilon > 0) \ (\exists k_0 \in \mathbb{N}) \ (\forall k \ge k_0) \ (k \in M_1) \ |d(x, A_{kj}) - d(x, B_j)| < \varepsilon,$ 

for some  $B_j$  and each  $j \in \mathbb{N}$ , and

$$(\forall \varepsilon > 0) \ (\exists j_0 \in \mathbb{N}) \ (\forall j \ge j_0) \ (j \in M_2) \ |d(x, A_{kj}) - d(x, C_k)| < \varepsilon,$$

for some  $C_k$  and each  $k \in \mathbb{N}$ . Hence, we have

$$A(\varepsilon) = \{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \varepsilon\} \subset H_1 \cup \{1, \dots, (k_0 - 1)\}, (j \in \mathbb{N}), B(\varepsilon) = \{j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| \ge \varepsilon\} \subset H_2 \cup \{1, \dots, (j_0 - 1)\}, (k \in \mathbb{N}), C_k \in \mathbb{N}\}$$

for  $H_1, H_2 \in \mathcal{I}_{\sigma}$ . Since  $\mathcal{I}_{\sigma}$  is an admissible ideal we get

$$H_1 \cup \{1, 2, \dots, (k_0 - 1)\} \in \mathcal{I}_\sigma \text{ and } H_2 \cup \{1, 2, \dots, (j_0 - 1)\} \in \mathcal{I}_\sigma$$

and therefore  $A(\varepsilon) \in \mathcal{I}_{\sigma}$  and  $B(\varepsilon) \in \mathcal{I}_{\sigma}$ . This shows that the double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent.

**Theorem 2.9.** Let  $\mathcal{I}_{\sigma}$  has property (AP) and  $\mathcal{I}_{2}^{\sigma}$  has property (AP2). If a double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_{W}^{\sigma}, \mathcal{I}_{W_{2}}^{\sigma})$ -convergent, then  $\{A_{kj}\}$  is  $R(\mathcal{I}_{W}^{\sigma*}, \mathcal{I}_{W_{2}}^{\sigma*})$ -convergent.

*Proof.* Let a double sequence  $\{A_{kj}\}$  be  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent. Then,  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent and so, by Lemma 1.4  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma*}$ -convergent. Also, for each  $\varepsilon > 0$  and each  $x \in X$  we have

$$A(\varepsilon) = \{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \varepsilon\} \in \mathcal{I}_{\sigma},$$

for some  $B_j$  and each  $j \in \mathbb{N}$ , and

$$B(\varepsilon) = \{ j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| \ge \varepsilon \} \in \mathcal{I}_{\sigma},$$

for some  $C_k$  and each  $k \in \mathbb{N}$ .

Now, for each  $x \in X$  we put

$$A_1 = \{k \in \mathbb{N} : |d(x, A_{kj} - B_j)| \ge 1\},\$$
  
$$A_t = \left\{k \in \mathbb{N} : \frac{1}{t} \le |d(x, A_{kj}) - d(x, B_j)| < \frac{1}{t - 1}\right\},\$$

for  $t \geq 2$ , some  $B_j$  and each  $j \in \mathbb{N}$ . It is clear that  $A_m \cap A_n = \emptyset$ , for  $m \neq n$  and  $A_m \in \mathcal{I}_{\sigma}$ , for each  $m \in \mathbb{N}$ . By the property (AP) there is a countable family of sets  $\{B_1, B_2, \ldots\}$  in  $\mathcal{I}_{\sigma}$  such that  $A_n \bigtriangleup B_n$  is a finite set for each  $n \in \mathbb{N}$  and  $\infty$ 

$$B = \bigcup_{n=1}^{n=1} B_n \in \mathcal{I}_{\sigma}.$$

We prove that

$$\lim_{\substack{k \to \infty \\ k \in M}} d(x, A_{kj}) = d(x, B_j), \text{ some } B_j \text{ and each } j \in \mathbb{N},$$

for  $M = \mathbb{N} \setminus B \in \mathcal{F}(\mathcal{I}_{\sigma})$ . Let  $\delta > 0$  be given. Choose  $t \in \mathbb{N}$  such that  $1/t < \delta$ . Then, we have for each  $x \in X$ ,

$$\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \delta\} \subset \bigcup_{n=1}^{l} x_n$$
, for some  $B_j$  and each  $j \in \mathbb{N}$ .

Since  $A_n \triangle B_n$  is a finite set for  $n \in \{1, 2, ..., t\}$ , there exists  $k_0 \in \mathbb{N}$  such that

$$\left(\bigcup_{n=1}^{t} B_n\right) \cap \{k : k \ge k_0\} = \left(\bigcup_{n=1}^{t} A_n\right) \cap \{k : k \ge k_0\}.$$

If  $k \ge k_0$  and  $k \notin B$ , then

$$k \notin \bigcup_{n=1}^{t} B_n$$
 and so  $k \notin \bigcup_{n=1}^{t} A_n$ .

Thus, we have

$$|d(x, A_{kj}) - d(x, B_j)| < \frac{1}{t} < \delta,$$

for some  $B_j$ , each  $j \in \mathbb{N}$  and each  $x \in X$ . This implies that

$$\lim_{k \to \infty} d(x, A_{kj}) = d(x, B_j).$$

Hence, we have

$$\mathcal{I}_{\sigma}^* - \lim_{k \to \infty} d(x, A_{kj}) = d(x, B_j),$$

for some  $B_j$ , each  $j \in \mathbb{N}$  and each  $x \in X$ .

Similarly, for the set

$$B(\varepsilon) = \{ j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| \ge \varepsilon \} \in \mathcal{I}_{\sigma},$$

we have

$$\mathcal{I}_{\sigma}^* - \lim_{j \to \infty} d(x, A_{kj}) = d(x, C_k),$$

for some  $C_k$ , each  $m \in \mathbb{N}$  and each  $x \in X$ . Hence, a double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma_*}, \mathcal{I}_{W_2}^{\sigma_*})$ -convergent.

**Definition 2.10.** A double sequence  $\{A_{kj}\}$  is said to be regularly  $(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -Cauchy sequence  $(R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -Cauchy sequence), if it is  $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy in Pringsheim's sense and also for every  $\varepsilon > 0$  and each  $x \in X$  there exist numbers  $m_j = m_j(\varepsilon), n_k = n_k(\varepsilon) \in \mathbb{N}$  such that

$$A_1(\varepsilon) = \left\{ k \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{m_j j})| \ge \varepsilon \right\} \in \mathcal{I}_{\sigma}, (j \in \mathbb{N})$$

and

$$A_2(\varepsilon) = \left\{ j \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{kn_k})| \ge \varepsilon \right\} \in \mathcal{I}_{\sigma}, (k \in \mathbb{N})$$

holds.

**Theorem 2.11.** If a double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent, then  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -Cauchy sequence.

*Proof.* Let  $\{A_{kj}\}$  be  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -convergent. Then,  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma}$ -convergent and by Lemma 1.5, it is  $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy sequence. Also, for every  $\varepsilon > 0$  and each  $x \in X$  we have

$$A_1\left(\frac{\varepsilon}{2}\right) = \left\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| \ge \frac{\varepsilon}{2}\right\} \in \mathcal{I}_{\sigma},$$

for some  $B_j$  and each  $j \in \mathbb{N}$ , and

$$A_2\left(\frac{\varepsilon}{2}\right) = \left\{ j \in \mathbb{N} : \left| d(x, A_{kj}) - d(x, C_k) \right| \ge \frac{\varepsilon}{2} \right\} \in \mathcal{I}_{\sigma},$$

for some  $C_k$  and each  $k \in \mathbb{N}$ . Since  $\mathcal{I}_{\sigma}$  is an admissible ideal, for each  $x \in X$  the sets

$$A_1^c\left(\frac{\varepsilon}{2}\right) = \left\{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, B_j)| < \frac{\varepsilon}{2}\right\},\$$

for some  $B_j$  and each  $j \in \mathbb{N}$ , and

$$A_2^c\left(\frac{\varepsilon}{2}\right) = \left\{ j \in \mathbb{N} : |d(x, A_{kj}) - d(x, C_k)| < \frac{\varepsilon}{2} \right\},\$$

for some  $C_k$  and each  $k \in \mathbb{N}$ , are nonempty and belong to  $\mathcal{F}(\mathcal{I}_{\sigma})$ . For  $m_j \in A_1^c(\frac{\varepsilon}{2}), (j \in \mathbb{N} \text{ and } m_j > 0)$  we have

$$|d(x, A_{m_j j}) - d(x, B_j)| < \frac{\varepsilon}{2}$$

for some  $B_j$ , each  $j \in \mathbb{N}$  and each  $x \in X$ . Now, for each  $\varepsilon > 0$  we define the set

$$B_1(\varepsilon) = \{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{m_j j})| \ge \varepsilon\}, \ (j \in \mathbb{N})$$

where  $m_j = m_j(\varepsilon) \in \mathbb{N}$ . We must prove  $B_1(\varepsilon) \subset A_1(\frac{\varepsilon}{2})$ . Let  $k \in B_1(\varepsilon)$ . Then, for  $m_j \in A_1^c(\frac{\varepsilon}{2}), (j \in \mathbb{N} \text{ and } m_j > 0)$  we have

$$\varepsilon \le |d(x, A_{kj}) - d(x, A_{m_j j})| \le |d(x, A_{kj}) - d(x, B_j)| + |d(x, A_{m_j j}) - d(x, B_j)| < |d(x, A_{kj}) - d(x, B_j)| + \frac{\varepsilon}{2},$$

for some  $B_j$  and each  $j \in \mathbb{N}$ . This shows that

$$\frac{\varepsilon}{2} < |d(x, A_{kj}) - d(x, B_j)|$$

and so  $k \in A_1(\frac{\varepsilon}{2})$ . Hence, we have  $B_1(\varepsilon) \subset A_1(\frac{\varepsilon}{2})$ . Similarly for each  $\varepsilon > 0$  and  $n_1 \in A^c(\frac{\varepsilon}{2})$   $(k \in \mathbb{N} \text{ and } n_1 > 0)$ 

Similarly, for each 
$$\varepsilon > 0$$
 and  $n_k \in A_2^c(\frac{\varepsilon}{2})$   $(k \in \mathbb{N} \text{ and } n_k > 0)$  we have

$$|d(x, A_{kn_k}) - d(x, C_k)| < \frac{\varepsilon}{2},$$

for some  $C_k$ , each  $k \in \mathbb{N}$  and each  $x \in X$ . Therefore, it can be seen that

$$B_2(\varepsilon) \subset A_2(\frac{\varepsilon}{2}),$$

where

$$B_2(\varepsilon) = \{ j \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{kn_k})| \ge \varepsilon \}, \ (k \in \mathbb{N}),$$

where  $n_k = n_k(\varepsilon) \in \mathbb{N}$ .

Therefore, we have  $B_1(\varepsilon) \in \mathcal{I}_{\sigma}$  and  $B_2(\varepsilon) \in \mathcal{I}_{\sigma}$ . This shows that  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -Cauchy sequence.

**Definition 2.12.** A double sequence  $\{A_{kj}\}$  is regularly  $(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -Cauchy sequence  $(R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -Cauchy sequence) if there exist the sets  $M \in \mathcal{F}(\mathcal{I}_2^{\sigma}), M_1 \in \mathcal{F}(\mathcal{I}_{\sigma})$  and  $M_2 \in \mathcal{F}(\mathcal{I}_{\sigma})$  (that is  $\mathbb{N} \times \mathbb{N} \setminus M = H \in \mathcal{I}_2^{\sigma}, \mathbb{N} \setminus M_1 \in \mathcal{I}_{\sigma}$  and  $\mathbb{N} \setminus M_2 \in \mathcal{I}_{\sigma}$ ) and for every  $\varepsilon > 0$  and each  $x \in X$ , there exist  $N = N(\varepsilon), s = s(\varepsilon), t = t(\varepsilon), m_j = m_j(\varepsilon), n_k = n_k(\varepsilon) \in \mathbb{N}$  such that whenever  $k, j, s, t, m_j, n_k \geq N$ , we have

$$\begin{aligned} |d(x, A_{kj}) - d(x, A_{st})| &< \varepsilon, \ (for \ (k, j), (s, t) \in M), \\ |d(x, A_{kj}) - d(x, A_{m_j j})| &< \varepsilon, \ (for \ each \ k \in M_1 \ and \ each \ j \in \mathbb{N}), \\ |d(x, A_{kj}) - d(x, A_{kn_k})| &< \varepsilon, \ (for \ each \ j \in M_2 \ and \ each \ k \in \mathbb{N}). \end{aligned}$$

**Theorem 2.13.** If a double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -Cauchy sequence, then  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -Cauchy sequence.

Proof. Since a double sequence  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -Cauchy sequence, then  $\{A_{kj}\}$  is  $\mathcal{I}_{W_2}^{\sigma*}$ -Cauchy sequence and so it is  $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy sequence by Lemma 1.6. Also, since  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma*}, \mathcal{I}_{W_2}^{\sigma*})$ -Cauchy sequence, there exist the sets  $M_1 \in \mathcal{F}(\mathcal{I}_{\sigma})$  and  $M_2 \in \mathcal{F}(\mathcal{I}_{\sigma})$  (that is,  $\mathbb{N} \setminus M_1 \in \mathcal{I}_{\sigma}$  and  $\mathbb{N} \setminus M_2 \in \mathcal{I}_{\sigma}$ ) and for every  $\varepsilon > 0$  and each  $x \in X$ , there exist  $N = N(\varepsilon), m_j = m_j(\varepsilon), n_k = n_k(\varepsilon) \in \mathbb{N}$  such that we have

$$\begin{aligned} |d(x, A_{kj}) - d(x, A_{m_j j})| &< \varepsilon, \ (for \ each \ k \in M_1 \ and \ each \ j \in \mathbb{N}), \\ |d(x, A_{kj}) - d(x, A_{kn_k})| &< \varepsilon, \ (for \ each \ j \in M_1 \ and \ each \ k \in \mathbb{N}), \end{aligned}$$

whenever  $k, j, m_j, n_k \geq N$ . Therefore,  $H_1 = \mathbb{N} \setminus M_1 \in \mathcal{I}_{\sigma}$  and  $H_2 = \mathbb{N} \setminus M_2 \in \mathcal{I}_{\sigma}$ we have

$$A_1(\varepsilon) = \{k \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{m_j j})| \ge \varepsilon\} \subset H_1 \cup \{1, \dots, (N-1)\}, (j \in \mathbb{N})\}$$

for each  $k \in M_1$  and each  $x \in X$ , and

$$A_{2}(\varepsilon) = \{ j \in \mathbb{N} : |d(x, A_{kj}) - d(x, A_{kn_{k}})| \ge \varepsilon \} \subset H_{2} \cup \{1, \dots, (N-1)\}, (k \in \mathbb{N})$$

for each  $j \in M_2$  and each  $x \in X$ . Since  $\mathcal{I}_{\sigma}$  is an admissible ideal,

$$H_1 \cup \{1, 2, \dots, (N-1)\} \in \mathcal{I}_{\sigma} \text{ and } H_2 \cup \{1, 2, \dots, (N-1)\} \in \mathcal{I}_{\sigma},$$

and so,  $A_1(\varepsilon) \in \mathcal{I}_{\sigma}$  and  $A_2(\varepsilon) \in \mathcal{I}_{\sigma}$ . Hence,  $\{A_{kj}\}$  is  $R(\mathcal{I}_W^{\sigma}, \mathcal{I}_{W_2}^{\sigma})$ -Cauchy sequence.

# 3. Conclusions and Future Work

We investigated the concepts of Wijsman regularly invariant convergence typies and Wijsman regularly ideal invariant convergence and Cauchy sequence typies. These concepts can also be studied for the lacunary sequence in the future.

# 4. Competing Interests

The authors declare that there is not any conflict of interests regarding the publication of this manuscript.

# References

- 1. M. Baronti, and P. Papini, *Convergence of sequences of sets*, In: Methods of functional analysis in approximation theory, ISNM 76, Birkhauser-Verlag, Basel, 1986.
- G. Beer, On convergence of closed sets in a metric space and distance functions, Bull. Aust. Math. Soc. 31 (1985), 421-432.
- 3. G. Beer, Wijsman convergence: A survey, Set-Valued Var. Anal. 2 (1994), 77-94.
- N. Pancaroğlu Akın, E. Dündar and F. Nuray, Wijsman *I*-invariant convergence of sequences of sets, Bulletin of Mathematical Analysis and Applications 11 (2019), 1-9.
- N. Pancaroğlu Akın, Regularly ideal invariant convergence of double sequences, in review.
   B. Altay, F. Başar, Some new spaces of double sequences, J. Math. Anal. Appl. 309 (2005),
- 70-90. 7. C. Çakan, B. Altay, M. Mursaleen, The  $\sigma$ -convergence and  $\sigma$ -core of double sequences,
- 7. C. Çakan, B. Altay, M. Mursaleen, The  $\sigma$ -convergence and  $\sigma$ -core of double sequences, Applied Mathematics Letters **19** (2006), 1122-1128.
- P. Das, P. Kostyrko, W. Wilczyński, P. Malik, *I* and *I*<sup>\*</sup>-convergence of double sequences, Math. Slovaca 58 (2008), 605-620.
- 9. K. Dems, On *I*-Cauchy sequences, Real Anal. Exchange **30** (2005), 123-128.
- E. Dündar, B. Altay, *I*<sub>2</sub>-convergence and *I*<sub>2</sub>-Cauchy of double sequences, Acta Mathematica Scientia **34** (2014), 343-353.
- E. Dündar, B. Altay, On some properties of I<sub>2</sub>-convergence and I<sub>2</sub>-Cauchy of double sequences, Gen. Math. Notes 7 (2011), 1-12.
- E. Dündar, B. Altay, *I*<sub>2</sub>-convergence of double sequences of functions, Electronic Journal of Mathematical Analysis and Applications 3 (2015), 111-121.
- E. Dündar, B. Altay *I*<sub>2</sub>-uniform convergence of double sequences of functions, Filomat **30** (2016), 1273-1281.
- E. Dündar, B. Altay, Multipliers for bounded I<sub>2</sub>-convergent of double sequences, Math. Comput. Modelling 55 (2012), 1193-1198.
- E. Dündar, Ö. Talo, *I*<sub>2</sub>-convergence of double sequences of fuzzy numbers, Iranian Journal of Fuzzy Systems **10** (2013), 37-50.
- 16. E. Dündar, M.R. Türkmen and N. Pancaroğlu Akın, Regularly ideal convergence of double sequences in fuzzy normed spaces, in review.
- E. Dündar, Regularly (I<sub>2</sub>, I)-Convergence and (I<sub>2</sub>, I)-Cauchy Double Sequences of Functions, Pioneer Journal of Algebra, Number Theory and its Applications 1 (2011), 85-98.
- E. Dündar, Ö. Talo and F. Başar, Regularly (I<sub>2</sub>, I)-convergence and regularly (I<sub>2</sub>, I)-Cauchy double sequences of fuzzy numbers, International Journal of Analysis 2013 (2013), 1-7.
- E. Dündar, U. Ulusu, F. Nuray, On ideal invariant convergence of double sequences and some properties, Creative Mathematics and Informatics 27 (2018), 161-169.
- E. Dündar, U. Ulusu, N. Pancaroğlu, Strongly I<sub>2</sub>-lacunary convergence and I<sub>2</sub>-lacunary Cauchy double sequences of sets, The Aligarh Bulletin of Mathematics 35 (2016), 1-15.
- E. Dündar, N. Pancaroğlu, U. Ulusu, Wijsman lacunary I-invariant convergence of sequences of sets, Proceedings of The National Academy Of Sciences India Section A-Physical Sciences 90 (2020), DOI: https://doi.org/10.1007/s40010-020-00694-w.
- E. Dündar and N. Pancaroğlu Akın, Wijsman regularly ideal convergence of double sequences of sets, Journal of Intelligent and Fuzzy Systems 37 (2019), 8159-8166.
- 23. H. Fast, Sur la convergence statistique, Colloq. Math. 2 (1951), 241-244.
- P. Kostyrko, T. Šalát, W. Wilczyński, *I-convergence*, Real Anal. Exchange 26 (2000), 669-686.

- V. Kumar, On I and I\*-convergence of double sequences, Math. Commun. 12 (2007), 171-181.
- M. Mursaleen, O.H.H. Edely, Statistical convergence of double sequences, J. Math. Anal. Appl. 288 (2003), 223-231.
- M. Mursaleen, Matrix transformation between some new sequence spaces, Houston J. Math. 9 (1983), 505-509.
- M. Mursaleen, On finite matrices and invariant means, Indian J. Pure Appl. Math. 10 (1979), 457-460.
- M. Mursaleen, O.H.H. Edely, On the invariant mean and statistical convergence, Appl. Math. Lett. 22 (2009), 1700-1704.
- A. Nabiev, S. Pehlivan, M. Gürdal, On *I*-Cauchy sequences, Taiwanese J. Math. 11 (2007), 569-576.
- 31. F. Nuray, H. Gök, U. Ulusu,  $\mathcal{I}_{\sigma}$ -convergence, Math. Commun. 16 (2011), 531-538.
- N. Pancaroğlu, F. Nuray, Statistical lacunary invariant summability, Theoret. Math. Appl. 3 (2013), 71-78.
- N. Pancaroğlu, F. Nuray, On invariant statistically convergence and lacunary invariant statistically convergence of sequences of sets, Prog. Appl. Math. 5 (2013), 23-29.
- A. Pringsheim, Zur theorie der zweifach unendlichen Zahlenfolgen, Math. Ann. 53 (1900), 289-321.
- R.A. Raimi, Invariant means and invariant matrix methods of summability, Duke Math. J. 30 (1963), 81-94.
- E. Savaş, Some sequence spaces involving invariant means, Indian J. Math. 31 (1989), 1-8.
- 37. E. Savaş, Strongly  $\sigma$ -convergent sequences, Bull. Calcutta Math. 81 (1989), 295-300.
- P. Schaefer, Infinite matrices and invariant means, Proc. Amer. Math. Soc. 36 (1972), 104-110.
- I.J. Schoenberg, The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.
- Y. Sever, E. Dündar, Regularly ideal convergence and regularly ideal Cauchy double sequences in 2-normed spaces, Filomat 28 (2015), 907-915.
- B. Tripathy, B.C. Tripathy, On *I*-convergent double sequences, Soochow J. Math. 31 (2005), 549-560.
- 42. Ş. Tortop, E. Dündar, Wijsman  $\mathcal{I}_2$ -invariant convergence of double sequences of sets, Journal of Inequalities and Special Functions **9** (2018), 90-100.
- U. Ulusu, E. Dündar, Asymptotically lacunary I<sub>2</sub>-invariant equivalence, Journal of Intelligent and Fuzzy Systems 36 (2019), 467-472.
- 44. U. Ulusu, E. Dündar, F. Nuray, Lacunary *I*<sub>2</sub>-invariant convergence and some properties, International Journal of Analysis and Applications 16 (2018), 317-327.
- R.A. Wijsman, Convergence of sequences of convex sets, cones and functions, Bull. Amer. Math. Soc. 70 (1964), 186-188.
- 46. R.A. Wijsman, Convergence of sequences of convex sets, cones and functions II, Trans. Amer. Math. Soc. 123 (1966), 32-45.

**Erdinç Dündar** is a Associate Professor in Department of Mathematics, Afyon Kocatepe University. He received the Ph.D. degree in Mathematics for İnönü University, Malatya, Turkey. His area of expertise includes: Summability theory, sequences spaces and fuzzy sequence spaces through functional analysis. He is the author of many international published papers.

Department of Mathematics, Afyon Kocatepe University, Afyonkarahisar, Turkey. e-mail: edundar@aku.edu.tr

Özer Talo is a Associate Professor. He received the Ph.D. degree in Mathematics for İnönü University, Malatya, Turkey. His area of expertise includes: Summability theory, sequences spaces and fuzzy sequence spaces through functional analysis. He has many international published papers.

Manisa Celal Bayar Universitesi Küme evleri, 45140, Yunusemre, Manisa, Turkey. e-mail: ozertalo@hotmail.com