

DOUBLE WIJSMAN LACUNARY STATISTICAL CONVERGENCE OF ORDER α

ESRA GÜLLE*, UĞUR ULUSU

ABSTRACT. In this paper, we introduce the concepts of Wijsman strongly p -lacunary summability of order α , Wijsman lacunary statistical convergence of order α and Hausdorff lacunary statistical convergence of order α for double set sequences. Also, we investigate some properties of these new concepts and examine the existence of some relationships between them. Furthermore, we study the relationships between these new concepts and some concepts in the literature.

AMS Mathematics Subject Classification : 40A05, 40A35.

Key words and phrases : Double set sequences, Wijsman convergence, Hausdorff convergence, Lacunary summability, Statistical convergence, Order α .

1. Introduction

The concept of statistical convergence was introduced by Fast [13] and Steinhaus [34], and later reintroduced by Schoenberg [32], independently. Then, this concept has been developed by many researchers until recently (see, [5, 7, 8, 15, 17, 27, 35, 40]).

In [14], Freedman et al. established the connection between the strongly Cesàro summable sequences space $|\sigma_1|$ and the strongly lacunary summable sequences space N_θ defined by a lacunary sequence θ . Then, using lacunary sequence concept, Fridy and Orhan [16] defined the concept of lacunary statistical convergence. Recently, the concepts of lacunary statistical convergence of order α and strongly p -lacunary summability of order α were studied by Şengül and Et [36]. For more detail, see [12].

In [26], Pringsheim introduced the concept of convergence for double sequences. Then, Mursaleen and Edely [19] extended this concept to statistical convergence. Also, using double lacunary sequence concept, the concept of

Received August 25, 2020. Revised February 25, 2021. Accepted March 30, 2021.

*Corresponding author.

© 2021 KSCAM.

lacunary statistical convergence was studied by Patterson and Savař [25]. Recently, olak and Altın [9] defined the concept of statistical convergence of order α for double sequences. Also, the concepts of almost statistical and almost lacunary statistical convergence of order α for double sequences were studied by Savař [29, 30]. More developments on double sequences can be found in [4, 6, 10, 11, 18, 20, 28, 39].

The concepts of convergence for number sequences were transferred to the concepts of convergence for set sequences by many authors. The concepts of Wijsman convergence and Hausdorff convergence are two of these transfers (see, [1, 2, 3, 43]). Nuray and Rhoades [21] extended the concepts of Wijsman convergence and Hausdorff convergence to statistical convergence for set sequences and gave some basic theorems. Then, using lacunary sequence concept, the concept of lacunary statistical convergence for set sequences was introduced by Ulusu and Nuray [41]. Recently, using ideal concept, the concept of Wijsman \mathcal{I} -lacunary statistical convergence of order α was studied by both Savař [31] and řengl and Et [37], independently.

In [22, 23, 24], Nuray et al. introduced the concepts of Wijsman strongly Cesàro summability, Wijsman statistical convergence, Wijsman strongly lacunary summability and Wijsman lacunary statistical convergence for double set sequences. Then, the concept of Hausdorff statistical convergence for double set sequences was studied by Talo et al. [38].

Lately, the concepts of Wijsman strongly p -Cesàro summability of order α , Wijsman statistical convergence of order α and Hausdorff statistical convergence of order α for double set sequences were studied by Ulusu and Glle [42].

2. Definitions and Notations

Firstly, we recall the basic concepts that need for a good understanding of our study (see, [1, 2, 22, 23, 24, 26, 28, 33, 38, 42]).

A double sequence (x_{ij}) is said to be convergent to L in Pringsheim's sense if for every $\varepsilon > 0$, there exists $N_\varepsilon \in \mathbb{N}$ such that $|x_{ij} - L| < \varepsilon$, whenever $i, j > N_\varepsilon$.

Let X be any non-empty set. The function $f : \mathbb{N} \rightarrow P(X)$ is defined by $f(i) = U_i \in P(X)$ for each $i \in \mathbb{N}$, where $P(X)$ is power set of X . The sequence $\{U_i\} = \{U_1, U_2, \dots\}$, which is the range's elements of f , is said to be set sequences.

Let (X, d) be a metric space. For any point $x \in X$ and any non-empty subset U of X , the distance from x to U is defined by

$$\rho(x, U) = \inf_{u \in U} d(x, u).$$

Throughout the study, (X, d) will be taken as a metric space and U, U_{ij} be any non-empty closed subsets of X .

A double sequence $\{U_{ij}\}$ is said to be Wijsman convergent to U if for each $x \in X$,

$$\lim_{i, j \rightarrow \infty} \rho(x, U_{ij}) = \rho(x, U).$$

A double sequence $\{U_{ij}\}$ is said to be Hausdorff convergent to U if

$$\lim_{i,j \rightarrow \infty} \sup_{x \in X} |\rho(x, U_{ij}) - \rho(x, U)| = 0.$$

A double sequence $\{U_{ij}\}$ is said to be Wijsman statistical convergent to U if for every $\varepsilon > 0$ and each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \left| \{(i, j) : i \leq n, j \leq m, |\rho(x, U_{ij}) - \rho(x, U)| \geq \varepsilon\} \right| = 0.$$

The class of all Wijsman statistical convergent sequences denotes by simply $W(S_2)$.

A double sequence $\{U_{ij}\}$ is said to be Hausdorff statistical convergent to U if for every $\varepsilon > 0$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \left| \{(i, j) : i \leq n, j \leq m, \sup_{x \in X} |\rho(x, U_{ij}) - \rho(x, U)| \geq \varepsilon\} \right| = 0.$$

A double sequence $\theta_2 = \{(k_r, j_u)\}$ is called double lacunary sequence if there exist two increasing sequence of integers such that

$$k_0 = 0, h_r = k_r - k_{r-1} \rightarrow \infty \text{ and } j_0 = 0, \bar{h}_u = j_u - j_{u-1} \rightarrow \infty \text{ as } r, u \rightarrow \infty.$$

We use the following notations in the sequel:

$$k_{ru} = k_r j_u, h_{ru} = h_r \bar{h}_u, I_{ru} = \{(i, j) : k_{r-1} < i \leq k_r \text{ and } j_{u-1} < j \leq j_u\},$$

$$q_r = \frac{k_r}{k_{r-1}} \text{ and } q_u = \frac{j_u}{j_{u-1}}.$$

Throughout the study, $\theta_2 = \{(k_r, j_u)\}$ will be taken as a double lacunary sequence.

A double sequence $\{U_{ij}\}$ is said to be Wijsman lacunary convergent to U if for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}} \sum_{(i,j) \in I_{ru}} \rho(x, U_{ij}) = \rho(x, U).$$

Let $0 < p < \infty$. A double sequence $\{U_{ij}\}$ is said to be Wijsman strongly p -lacunary convergent to U if for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}} \sum_{(i,j) \in I_{ru}} |\rho(x, U_{ij}) - \rho(x, U)|^p = 0.$$

The class of all Wijsman strongly p -lacunary convergent sequences denotes by simply $[W_2 N_\theta]^p$.

A double sequence $\{U_{ij}\}$ is said to be Wijsman lacunary statistically convergent to U if for every $\varepsilon > 0$ and each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}} \left| \{(i, j) \in I_{ru} : |\rho(x, U_{ij}) - \rho(x, U)| \geq \varepsilon\} \right| = 0.$$

The class of all Wijsman lacunary statistically convergent sequences denotes by simply $W_2 S_\theta$.

Let $0 < \alpha \leq 1$. A double sequence $\{U_{ij}\}$ is said to be Wijsman strongly Cesàro summable of order α to U or $W[C_2^\alpha]$ -summable to U if for each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{(mn)^\alpha} \sum_{i,j=1,1}^{m,n} |\rho(x, U_{ij}) - \rho(x, U)| = 0.$$

The class of all $W[C_2^\alpha]$ -summable sequences denotes by simply $W[C_2^\alpha]$.

Let $0 < \alpha \leq 1$. A double sequence $\{U_{ij}\}$ is said to be Wijsman statistically convergent of order α to U or $W(S_2^\alpha)$ -convergent to U if for every $\varepsilon > 0$ and each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{(mn)^\alpha} \left| \{(i, j) : i \leq m, j \leq n, |\rho(x, U_{ij}) - \rho(x, U)| \geq \varepsilon\} \right| = 0.$$

The class of all $W(S_2^\alpha)$ -convergent sequences denotes by simply $W(S_2^\alpha)$.

From now on, for short, we use $\rho_x(U)$ and $\rho_x(U_{ij})$ instead of $\rho(x, U)$ and $\rho(x, U_{ij})$, respectively.

3. New Concepts

In this section, we introduce the concepts of Wijsman strongly p -lacunary summability of order α , Wijsman lacunary statistical convergence of order α and Hausdorff lacunary statistical convergence of order α for double set sequences.

Definition 3.1. Let $0 < \alpha \leq 1$. A double sequence $\{U_{ij}\}$ is Wijsman lacunary summable of order α to U or $W_2N_\theta^\alpha$ -summable to U if for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} \rho_x(U_{ij}) = \rho_x(U).$$

In this case, we write $U_{ij} \xrightarrow{W_2N_\theta^\alpha} U$ or $U_{ij} \rightarrow U(W_2N_\theta^\alpha)$.

Definition 3.2. Let $0 < \alpha \leq 1$. A double sequence $\{U_{ij}\}$ is Wijsman strongly p -lacunary summable of order α to U or $[W_2N_\theta^\alpha]^p$ -summable to U if for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)|^p = 0.$$

In this case, we write $U_{ij} \xrightarrow{[W_2N_\theta^\alpha]^p} U$ or $U_{ij} \rightarrow U([W_2N_\theta^\alpha]^p)$. If $p = 1$, then the double sequence $\{U_{ij}\}$ is simply said to be Wijsman strongly lacunary summable of order α to U and we write $U_{ij} \xrightarrow{[W_2N_\theta^\alpha]} U$ or $U_{ij} \rightarrow U([W_2N_\theta^\alpha])$.

The class of all $[W_2N_\theta^\alpha]^p$ -summable sequences will be denoted by simply $[W_2N_\theta^\alpha]^p$.

Example 3.3. Let $X = \mathbb{R}^2$ and a double sequence $\{U_{ij}\}$ be defined as following:

$$U_{ij} := \begin{cases} \left\{ (x, y) : (x - 1)^2 + y^2 = \frac{1}{ij} \right\} & ; \text{ if } (i, j) \in I_{ru}, i \text{ and } j \text{ are} \\ & \text{square integers} \\ \{(0, 1)\} & ; \text{ otherwise.} \end{cases}$$

Then, the double sequence $\{U_{ij}\}$ is Wijsman strongly lacunary summable of order α to the set $U = \{(0, 1)\}$.

Remark 3.1. For $\alpha = 1$, the concepts of $W_2N_\theta^\alpha$ -summability and $[W_2N_\theta^\alpha]^p$ -summability coincide with the concepts of Wijsman lacunary convergence and Wijsman strongly p -lacunary convergence for double set sequences in [22], respectively.

Definition 3.4. Let $0 < \alpha \leq 1$. A double sequence $\{U_{ij}\}$ is Wijsman lacunary statistically convergent of order α to U or $W_2S_\theta^\alpha$ -convergent to U if for every $\varepsilon > 0$ and each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}^\alpha} \left| \left\{ (i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon \right\} \right| = 0.$$

In this case, we write $U_{ij} \xrightarrow{W_2S_\theta^\alpha} U$ or $U_{ij} \rightarrow U(W_2S_\theta^\alpha)$.

The class of all $W_2S_\theta^\alpha$ -convergent sequences will be denoted by simply $W_2S_\theta^\alpha$.

Example 3.5. Let $X = \mathbb{R}^2$ and a double sequence $\{U_{ij}\}$ be defined as following:

$$U_{ij} := \begin{cases} \left\{ (x, y) : (x + i)^2 + (y - j)^2 = 1 \right\} & ; \text{ if } (i, j) \in I_{ru}, i \text{ and } j \text{ are} \\ & \text{square integers} \\ \{(1, -1)\} & ; \text{ otherwise.} \end{cases}$$

Then, the double sequence $\{U_{ij}\}$ is Wijsman lacunary statistically convergent of order α to the set $U = \{(1, -1)\}$.

Remark 3.2. For $\alpha = 1$, the concept of $W_2S_\theta^\alpha$ -convergence coincides with the concept of Wijsman lacunary statistical convergence for double set sequences in [23].

Definition 3.6. Let $0 < \alpha \leq 1$. A double sequence $\{U_{ij}\}$ is Hausdorff lacunary statistically convergent of order α to U or $H_2(S_\theta^\alpha)$ -convergent to U if for every $\varepsilon > 0$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_{ru}^\alpha} \left| \left\{ (i, j) \in I_{ru} : \sup_{x \in X} |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon \right\} \right| = 0.$$

In this case, we write $U_{ij} \xrightarrow{H_2(S_\theta^\alpha)} U$ or $U_{ij} \rightarrow U(H_2(S_\theta^\alpha))$.

The class of all $H_2(S_\theta^\alpha)$ -convergent sequences will be denoted by simply $H_2(S_\theta^\alpha)$.

Remark 3.3. For $\alpha = 1$, the concept of $H_2(S_\theta^\alpha)$ -convergence coincides with the concept of Hausdorff lacunary statistical convergence for double set sequences, which has not been studied till now.

4. Inclusion Theorems

In this section, firstly, we investigate some properties of the new concepts that we introduced in Section 3 and examine the existence of some relationships between them.

Theorem 4.1. *If $0 < \alpha \leq \beta \leq 1$, then $[W_2N_\theta^\alpha]^p \subseteq [W_2N_\theta^\beta]^p$ for every double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$.*

Proof. Let $0 < \alpha \leq \beta \leq 1$ and suppose that $U_{ij} \xrightarrow{[W_2N_\theta^\alpha]^p} U$. For each $x \in X$, we have

$$\frac{1}{h_{ru}^\beta} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)|^p \leq \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)|^p.$$

Hence, by our assumption, we get $U_{ij} \xrightarrow{[W_2N_\theta^\beta]^p} U$. Consequently, $[W_2N_\theta^\alpha]^p \subseteq [W_2N_\theta^\beta]^p$. \square

If we take $\beta = 1$ in Theorem 4.1, then we obtain the following corollary.

Corollary 4.2. *If a double sequence $\{U_{ij}\}$ is Wijsman strongly p -lacunary summable of order α to U for some $0 < \alpha \leq 1$, then the double sequence is Wijsman strongly p -lacunary summable to U , i.e., $[W_2N_\theta^\alpha]^p \subseteq [W_2N_\theta]^p$.*

Now, without proof, we shall state a theorem that gives a relation between $[W_2N_\theta^\alpha]^p$ and $[W_2N_\theta^\alpha]^q$, where $0 < \alpha \leq 1$ and $0 < p < q < \infty$.

Theorem 4.3. *Let $0 < \alpha \leq 1$ and $0 < p < q < \infty$. Then, $[W_2N_\theta^\alpha]^q \subseteq [W_2N_\theta^\alpha]^p$ for every double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$.*

Theorem 4.4. *If $0 < \alpha \leq \beta \leq 1$, then $W_2S_\theta^\alpha \subseteq W_2S_\theta^\beta$ for every double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$.*

Proof. Let $0 < \alpha \leq \beta \leq 1$ and suppose that $U_{ij} \xrightarrow{W_2S_\theta^\alpha} U$. For every $\varepsilon > 0$ and each $x \in X$, we have

$$\begin{aligned} \frac{1}{h_{ru}^\beta} \left| \left\{ (i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon \right\} \right| \\ \leq \frac{1}{h_{ru}^\alpha} \left| \left\{ (i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon \right\} \right|. \end{aligned}$$

Hence, by our assumption, we get $U_{ij} \xrightarrow{W_2S_\theta^\beta} U$. Consequently, $W_2S_\theta^\alpha \subseteq W_2S_\theta^\beta$. \square

If we take $\beta = 1$ in Theorem 4.4, then we obtain the following corollary.

Corollary 4.5. *If a double sequence $\{U_{ij}\}$ is Wijsman lacunary statistically convergent of order α to U for some $0 < \alpha \leq 1$, then the double sequence is Wijsman lacunary statistically convergent to U , i.e., $W_2S_\theta^\alpha \subseteq W_2S_\theta$.*

Theorem 4.6. *Let $0 < \alpha \leq \beta \leq 1$ and $0 < p < \infty$. If a double sequence $\{U_{ij}\}$ is Wijsman strongly p -lacunary summable of order α to U , then the double sequence is Wijsman lacunary statistically convergent of order β to U .*

Proof. Let $0 < \alpha \leq \beta \leq 1$ and we suppose that the double sequence $\{U_{ij}\}$ is Wijsman strongly p -lacunary summable of order α to U . For every $\varepsilon > 0$ and each $x \in X$, we have

$$\begin{aligned} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)|^p &= \sum_{\substack{(i,j) \in I_{ru} \\ |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon}} |\rho_x(U_{ij}) - \rho_x(U)|^p \\ &+ \sum_{\substack{(i,j) \in I_{ru} \\ |\rho_x(U_{ij}) - \rho_x(U)| < \varepsilon}} |\rho_x(U_{ij}) - \rho_x(U)|^p \\ &\geq \sum_{\substack{(i,j) \in I_{ru} \\ |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon}} |\rho_x(U_{ij}) - \rho_x(U)|^p \\ &\geq \varepsilon^p \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \end{aligned}$$

and so

$$\begin{aligned} \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)|^p &\geq \frac{\varepsilon^p}{h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ &\geq \frac{\varepsilon^p}{h_{ru}^\beta} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right|. \end{aligned}$$

Hence, by our assumption, we get that the double sequence $\{U_{ij}\}$ is Wijsman lacunary statistically convergent of order β to U . \square

If we take $\beta = \alpha$ in Theorem 4.6, then we obtain the following corollary.

Corollary 4.7. *Let $0 < \alpha \leq 1$ and $0 < p < \infty$. If a double sequence $\{U_{ij}\}$ is Wijsman strongly p -lacunary summable of order α to U , then the double sequence is Wijsman lacunary statistically convergent of order α to U .*

Theorem 4.8. *If $0 < \alpha \leq \beta \leq 1$, then $H_2(S_\theta^\alpha) \subseteq H_2(S_\theta^\beta)$ for every double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$.*

Proof. Let $0 < \alpha \leq \beta \leq 1$ and suppose that $U_{ij} \xrightarrow{H_2(S_\theta^\alpha)} U$. For every $\varepsilon > 0$, we have

$$\begin{aligned} \frac{1}{h_{ru}^\beta} \left| \{(i, j) \in I_{ru} : \sup_{x \in X} |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ \leq \frac{1}{h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : \sup_{x \in X} |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right|. \end{aligned}$$

Hence, by our assumption, we get $U_{ij} \xrightarrow{H_2(S_\theta^\beta)} U$. Consequently, $H_2(S_\theta^\alpha) \subseteq H_2(S_\theta^\beta)$. \square

If we take $\beta = 1$ in Theorem 4.8, then we obtain the following corollary.

Corollary 4.9. *If a double sequence $\{U_{ij}\}$ is Hausdorff lacunary statistically convergent of order α to U for some $0 < \alpha \leq 1$, then the double sequence is Hausdorff lacunary statistically convergent to U .*

Theorem 4.10. *Let $0 < \alpha \leq \beta \leq 1$. If a double sequence $\{U_{ij}\}$ is Hausdorff lacunary statistically convergent of order α to U , then the double sequence is Wijsman lacunary statistically convergent of order β to U .*

Proof. Let $0 < \alpha \leq \beta \leq 1$ and suppose that the double sequence $\{U_{ij}\}$ is Hausdorff lacunary statistically convergent of order α to U . For every $\varepsilon > 0$ and each $x \in X$, we have

$$\begin{aligned} \frac{1}{h_{ru}^\beta} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ \leq \frac{1}{h_{ru}^\beta} \left| \{(i, j) \in I_{ru} : \sup_{x \in X} |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ \leq \frac{1}{h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : \sup_{x \in X} |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right|. \end{aligned}$$

Hence, by our assumption, we get that the double sequence $\{U_{ij}\}$ is Wijsman lacunary statistically convergent of order β to U . \square

If we take $\beta = \alpha$ in Theorem 4.10, then we obtain the following corollary.

Corollary 4.11. *Let $0 < \alpha \leq 1$. If a double sequence $\{U_{ij}\}$ is Hausdorff lacunary statistically convergent of order α to U , then the double sequence is Wijsman lacunary statistically convergent of order α to U .*

Now, secondly, we study the relationships between the new concepts that we introduced in Section 3 and some concepts in the literature.

Theorem 4.12. *Let $0 < \alpha \leq 1$. If $\liminf_r q_r^\alpha > 1$ and $\liminf_u q_u^\alpha > 1$ for any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, then $W[C_2^\alpha] \subseteq [W_2N_\theta^\alpha]$.*

Proof. Let $0 < \alpha \leq 1$ and suppose that $\liminf_r q_r^\alpha > 1$ and $\liminf_u q_u^\alpha > 1$. Then, there exist $\eta > 0$ and $\mu > 0$ such that $q_r^\alpha \geq 1 + \eta$ and $q_u^\alpha \geq 1 + \mu$ for all r and u , which implies that

$$\frac{k_{ru}^\alpha}{h_{ru}^\alpha} \leq \frac{(1 + \eta)(1 + \mu)}{\eta\mu} \quad \text{and} \quad \frac{k_{(r-1)(u-1)}^\alpha}{h_{ru}^\alpha} \leq \frac{1}{\eta\mu}.$$

For each $x \in X$, we can write

$$\begin{aligned} & \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)| \\ &= \frac{1}{h_{ru}^\alpha} \sum_{m,n=1,1}^{k_r, j_u} |\rho_x(U_{mn}) - \rho_x(U)| \\ & \quad - \frac{1}{h_{ru}^\alpha} \sum_{m,n=1,1}^{k_{r-1}, j_{u-1}} |\rho_x(U_{mn}) - \rho_x(U)| \\ &= \frac{k_{ru}^\alpha}{h_{ru}^\alpha} \left(\frac{1}{k_{ru}^\alpha} \sum_{m,n=1,1}^{k_r, j_u} |\rho_x(U_{mn}) - \rho_x(U)| \right) \\ & \quad - \frac{k_{(r-1)(u-1)}^\alpha}{h_{ru}^\alpha} \left(\frac{1}{k_{(r-1)(u-1)}^\alpha} \sum_{m,n=1,1}^{k_{r-1}, j_{u-1}} |\rho_x(U_{mn}) - \rho_x(U)| \right). \end{aligned}$$

If $U_{ij} \xrightarrow{W[C_2^\alpha]} U$, then for each $x \in X$

$$\frac{1}{k_{ru}^\alpha} \sum_{m,n=1,1}^{k_r, j_u} |\rho_x(U_{mn}) - \rho_x(U)| \rightarrow 0 \quad \text{and} \quad \frac{1}{k_{(r-1)(u-1)}^\alpha} \sum_{m,n=1,1}^{k_{r-1}, j_{u-1}} |\rho_x(U_{mn}) - \rho_x(U)| \rightarrow 0.$$

Thus, when the above equality is considered, for each $x \in X$ we get

$$\frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)| \rightarrow 0,$$

that is, $U_{ij} \xrightarrow{[W_2 N_\theta^\alpha]} U$. Consequently, $W[C_2^\alpha] \subseteq [W_2 N_\theta^\alpha]$. □

Theorem 4.13. *Let $0 < \alpha \leq 1$. If $\limsup_r q_r < \infty$ and $\limsup_u q_u < \infty$ for any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, then $[W_2 N_\theta^\alpha] \subseteq W[C_2^\alpha]$.*

Proof. Let $0 < \alpha \leq 1$ and suppose that $\limsup_r q_r < \infty$ and $\limsup_u q_u < \infty$. Then, there exist $M, N > 0$ such that $q_r < M$ and $q_u < N$ for all r and u . Let $U_{ij} \xrightarrow{[W_2 N_\theta^\alpha]} U$ and $\varepsilon > 0$ be given. Then, for each $x \in X$ we can find $R, U > 0$ and $H > 0$ such that

$$\sup_{i \geq R, j \geq U} \tau_{ij} < \varepsilon \quad \text{and} \quad \tau_{ij} < H \quad \text{for all } i, j = 1, 2, \dots$$

where

$$\tau_{ru} = \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)|.$$

If m and n are any integers satisfying $k_{r-1} < m \leq k_r$ and $j_{u-1} < n \leq j_u$ where $r > R$ and $u > U$, then for each $x \in X$ we can write

$$\begin{aligned}
& \frac{1}{(mn)^\alpha} \sum_{i,j=1,1}^{m,n} |\rho_x(U_{ij}) - \rho_x(U)| \\
& \leq \frac{1}{k_{(r-1)(u-1)}^\alpha} \sum_{i,j=1,1}^{k_r, j_u} |\rho_x(U_{ij}) - \rho_x(U)| \\
& = \frac{1}{k_{(r-1)(u-1)}^\alpha} \left(\sum_{I_{11}} |\rho_x(U_{ij}) - \rho_x(U)| \right. \\
& \quad + \sum_{I_{12}} |\rho_x(U_{ij}) - \rho_x(U)| + \sum_{I_{21}} |\rho_x(U_{ij}) - \rho_x(U)| \\
& \quad + \sum_{I_{22}} |\rho_x(U_{ij}) - \rho_x(U)| \\
& \quad \left. + \cdots + \sum_{I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)| \right) \\
& = \frac{h_{11}^\alpha}{k_{(r-1)(u-1)}^\alpha} \tau_{11} + \frac{h_{12}^\alpha}{k_{(r-1)(u-1)}^\alpha} \tau_{12} + \frac{h_{21}^\alpha}{k_{(r-1)(u-1)}^\alpha} \tau_{21} \\
& \quad + \frac{h_{22}^\alpha}{k_{(r-1)(u-1)}^\alpha} \tau_{22} + \cdots + \frac{h_{ru}^\alpha}{k_{(r-1)(u-1)}^\alpha} \tau_{ru} \\
& \leq \sum_{i,j=1,1}^{R,U} \frac{h_{ij}}{k_{(r-1)(u-1)}^\alpha} \tau_{ij} + \sum_{i,j=R+1,U+1}^{r,u} \frac{h_{ij}}{k_{(r-1)(u-1)}^\alpha} \tau_{ij} \\
& \leq \left(\sup_{i \geq 1, j \geq 1} \tau_{ij} \right) \frac{k_{RU}}{k_{(r-1)(u-1)}^\alpha} \\
& \quad + \left(\sup_{i \geq R, j \geq U} \tau_{ij} \right) \frac{(k_r - k_R)(j_u - j_U)}{k_{(r-1)(u-1)}^\alpha} \\
& \leq H \frac{k_{RU}}{k_{(r-1)(u-1)}^\alpha} + \varepsilon M N.
\end{aligned}$$

Since $k_{r-1}, j_{u-1} \rightarrow \infty$ as $m, n \rightarrow \infty$, it follows that for each $x \in X$

$$\frac{1}{(mn)^\alpha} \sum_{i,j=1,1}^{m,n} |\rho_x(U_{ij}) - \rho_x(U)| \rightarrow 0.$$

Thus, $U_{ij} \xrightarrow{W[C_2^\alpha]} U$. Consequently, $[W_2 N_\theta^\alpha] \subseteq W[C_2^\alpha]$. \square

Theorem 4.14. *Let $0 < \alpha \leq 1$. For any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, if*

$$1 < \liminf_r q_r^\alpha \leq \limsup_r q_r < \infty \quad \text{and} \quad 1 < \liminf_u q_u^\alpha \leq \limsup_u q_u < \infty,$$

then $[W_2N_\theta^\alpha] = W[C_2^\alpha]$.

Proof. This can be obtained from Theorem 4.12 and Theorem 4.13, immediately. □

Theorem 4.15. *Let $0 < \alpha \leq 1$. If $U_{ij} \xrightarrow{[W_2N_\theta^\alpha]} U$ and $U_{ij} \xrightarrow{W[C_2^\alpha]} V$, where $\{U_{ij}\} \in [W_2N_\theta^\alpha] \cap W[C_2^\alpha]$ for any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, then $U = V$.*

Proof. Let $0 < \alpha \leq 1$, $\{U_{ij}\} \in [W_2N_\theta^\alpha] \cap W[C_2^\alpha]$, $U_{ij} \xrightarrow{[W_2N_\theta^\alpha]} U$ and $U_{ij} \xrightarrow{W[C_2^\alpha]} V$. Also, we suppose that $U \neq V$. For each $x \in X$, we have

$$\begin{aligned} v_{ru} + \tau_{ru} &= \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)| + \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(V)| \\ &\geq \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U) - \rho_x(V)| \\ &= \frac{h_{ru}}{h_{ru}^\alpha} |\rho_x(U) - \rho_x(V)| \\ &\geq |\rho_x(U) - \rho_x(V)|, \end{aligned}$$

where

$$v_{ru} = \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(U)| \quad \text{and} \quad \tau_{ru} = \frac{1}{h_{ru}^\alpha} \sum_{(i,j) \in I_{ru}} |\rho_x(U_{ij}) - \rho_x(V)|.$$

Since $U_{ij} \xrightarrow{[W_2N_\theta^\alpha]} U$, then $v_{ru} \rightarrow 0$ for each $x \in X$. Thus, for each $x \in X$ and sufficiently large r and u , we must have

$$\tau_{ru} > \frac{1}{2} |\rho_x(U) - \rho_x(V)|.$$

Observe that for each $x \in X$ and sufficiently large r and u

$$\begin{aligned} \frac{1}{k_{ru}^\alpha} \sum_{m,n=1,1}^{k_r, j_u} |\rho_x(U_{mn}) - \rho_x(V)| &\geq \frac{1}{k_{ru}^\alpha} \sum_{(m,n) \in I_{ru}} |\rho_x(U_{mn}) - \rho_x(V)| \\ &= \frac{h_{ru}^\alpha}{k_{ru}^\alpha} \tau_{ru} \\ &= \left(1 - \frac{1}{q_r}\right)^\alpha \left(1 - \frac{1}{q_u}\right)^\alpha \tau_{ru} \\ &> \frac{1}{2} \left(1 - \frac{1}{q_r}\right)^\alpha \left(1 - \frac{1}{q_u}\right)^\alpha |\rho_x(U) - \rho_x(V)|. \end{aligned}$$

Since $U_{ij} \xrightarrow{W[C_2^\alpha]} V$, for each $x \in X$ the left side of the above inequality convergent to 0. Hence, we have $q_r \rightarrow 1$ and $q_u \rightarrow 1$, and by Theorem 4.13 this implies that

$$[W_2N_\theta^\alpha] \subseteq W[C_2^\alpha],$$

i.e.,

$$U_{ij} \xrightarrow{[W_2N_\theta^\alpha]} U \Rightarrow U_{ij} \xrightarrow{W[C_2^\alpha]} U$$

and therefore for each $x \in X$, we can write

$$\frac{1}{(mn)^\alpha} \sum_{i,j=1,1}^{m,n} |\rho_x(U_{ij}) - \rho_x(U)| \rightarrow 0.$$

For each $x \in X$, we have

$$\begin{aligned} \frac{1}{(mn)^\alpha} \sum_{i,j=1,1}^{m,n} |\rho_x(U_{ij}) - \rho_x(U)| + \frac{1}{(mn)^\alpha} \sum_{i,j=1,1}^{m,n} |\rho_x(U_{ij}) - \rho_x(V)| \\ \geq \frac{mn}{(mn)^\alpha} |\rho_x(U) - \rho_x(V)| > 0. \end{aligned}$$

Since both terms on the left side of the above inequality convergent to 0, then for each $x \in X$ we get

$$|\rho_x(U) - \rho_x(V)| = 0.$$

This situation causes a contradiction to our assumption. Consequently, we get $U = V$. \square

Theorem 4.16. *Let $0 < \alpha \leq 1$. If $\liminf_r q_r^\alpha > 1$ and $\liminf_u q_u^\alpha > 1$ for any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, then $U_{ij} \xrightarrow{W[S_2^\alpha]} U$ implies that $U_{ij} \xrightarrow{W_2S_\theta^\alpha} U$.*

Proof. Let $0 < \alpha \leq 1$ and suppose that $\liminf_r q_r^\alpha > 1$ and $\liminf_u q_u^\alpha > 1$. Then, there exists $\eta, \mu > 0$ such that $q_r^\alpha \geq 1 + \eta$ and $q_u^\alpha \geq 1 + \mu$ for all r and u , which implies that

$$\frac{h_{ru}^\alpha}{k_{ru}^\alpha} \geq \frac{\eta\mu}{(1+\eta)(1+\mu)}.$$

For every $\varepsilon > 0$ and each $x \in X$, we can write

$$\begin{aligned} \frac{1}{k_{ru}^\alpha} \left| \{(i, j) : i \leq k_r, j \leq j_u, |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ \geq \frac{1}{k_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ = \frac{h_{ru}^\alpha}{k_{ru}^\alpha h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ \geq \frac{\eta\mu}{(1+\eta)(1+\mu)} \frac{1}{h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right|. \end{aligned}$$

If $U_{ij} \xrightarrow{W(S_2^\alpha)} U$, then for each $x \in X$ the term on the left side of the above inequality convergent to 0 and this implies that

$$\frac{1}{h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \rightarrow 0.$$

Thus, we get $U_{ij} \xrightarrow{W_2 S_\theta^\alpha} U$. □

Theorem 4.17. *Let $0 < \alpha \leq 1$. If $\limsup_r q_r < \infty$ and $\limsup_u q_u < \infty$ for any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, then $U_{ij} \xrightarrow{W_2 S_\theta^\alpha} U$ implies that $U_{ij} \xrightarrow{W(S_2^\alpha)} U$.*

Proof. Let $0 < \alpha \leq 1$ and suppose that $\limsup_r q_r < \infty$ and $\limsup_u q_u < \infty$. Then, there exist $M, N > 0$ such that $q_r < M$ and $q_u < N$ for all r and u . Let $U_{ij} \xrightarrow{W_2 S_\theta^\alpha} U$ and $\varepsilon > 0$ be given, and let

$$T_{ru} := \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right|.$$

Then, there exist $r_0, u_0 \in \mathbb{N}$ such that for every $\varepsilon > 0$, each $x \in X$ and all $r \geq r_0, u \geq u_0$

$$\frac{T_{ru}}{h_{ru}^\alpha} < \varepsilon.$$

Now, let

$$H := \max\{T_{ru} : 1 \leq r \leq r_0, 1 \leq u \leq u_0\},$$

and let m and n be any integers satisfying $k_{r-1} < m \leq k_r$ and $j_{u-1} < n \leq j_u$. Then, for each $x \in X$ we can write

$$\begin{aligned} & \frac{1}{(mn)^\alpha} \left| \{(i, j) : i \leq m, j \leq n, |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ & \leq \frac{1}{k_{(r-1)(u-1)}^\alpha} \left| \{(i, j) : i \leq k_r, j \leq j_u : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\ & = \frac{1}{k_{(r-1)(u-1)}^\alpha} \{T_{11} + T_{12} + T_{21} + T_{22} + \dots + T_{r_0 u_0} + \dots + T_{ru}\} \\ & \leq \frac{r_0 u_0}{k_{(r-1)(u-1)}^\alpha} \left(\max_{\substack{1 \leq j \leq u_0 \\ 1 \leq i \leq r_0}} \{T_{ij}\} \right) \\ & \quad + \frac{1}{k_{(r-1)(u-1)}^\alpha} \left\{ h_{r_0(u_0+1)}^\alpha \frac{T_{r_0(u_0+1)}}{h_{r_0(u_0+1)}^\alpha} + h_{(r_0+1)u_0}^\alpha \frac{T_{(r_0+1)u_0}}{h_{(r_0+1)u_0}^\alpha} \right. \\ & \quad \left. + h_{(r_0+1)(u_0+1)}^\alpha \frac{T_{(r_0+1)(u_0+1)}}{h_{(r_0+1)(u_0+1)}^\alpha} + \dots + h_{ru}^\alpha \frac{T_{ru}}{h_{ru}^\alpha} \right\} \end{aligned}$$

$$\begin{aligned}
&\leq \frac{r_0 u_0 H}{k_{(r-1)(u-1)}^\alpha} + \frac{1}{k_{(r-1)(u-1)}^\alpha} \left(\sup_{\substack{r > r_0 \\ u > u_0}} \frac{T_{ru}}{h_{ru}^\alpha} \right) \left(\sum_{i,j \geq r_0, u_0}^{r,u} h_{ij}^\alpha \right) \\
&\leq \frac{r_0 u_0 H}{k_{(r-1)(u-1)}^\alpha} + \frac{1}{k_{(r-1)(u-1)}^\alpha} \left(\sup_{\substack{r > r_0 \\ u > u_0}} \frac{T_{ru}}{h_{ru}^\alpha} \right) \left(\sum_{i,j \geq r_0, u_0}^{r,u} h_{ij} \right) \\
&\leq \frac{r_0 u_0 H}{k_{(r-1)(u-1)}^\alpha} + \varepsilon \frac{(k_r - k_{r_0})(j_u - j_{u_0})}{k_{(r-1)(u-1)}} \\
&\leq \frac{r_0 u_0 H}{k_{(r-1)(u-1)}^\alpha} + \varepsilon q_r q_u \\
&\leq \frac{r_0 u_0 H}{k_{(r-1)(u-1)}^\alpha} + \varepsilon M N.
\end{aligned}$$

Since $k_{r-1}, j_{u-1} \rightarrow \infty$ as $m, n \rightarrow \infty$, it follows that for each $x \in X$

$$\frac{1}{(mn)^\alpha} \left| \{(i, j) : i \leq m, j \leq n : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \rightarrow 0.$$

Thus, we get $U_{ij} \xrightarrow{W(S_2^\alpha)} U$. \square

Theorem 4.18. Let $0 < \alpha \leq 1$. For any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, if

$$1 < \liminf_r q_r^\alpha \leq \limsup_r q_r < \infty \quad \text{and} \quad 1 < \liminf_u q_u^\alpha \leq \limsup_u q_u < \infty,$$

then $U_{ij} \xrightarrow{W_2 S_\theta^\alpha} U \Leftrightarrow U_{ij} \xrightarrow{W(S_2^\alpha)} U$.

Proof. This can be obtained from Theorem 4.16 and Theorem 4.17, immediately. \square

Theorem 4.19. Let $0 < \alpha \leq 1$. If $\liminf_{r,u \rightarrow \infty} \frac{h_{ru}^\alpha}{k_{ru}} > 0$ for any double lacunary sequence $\theta_2 = \{(k_r, j_u)\}$, then $W(S_2) \subseteq W_2 S_\theta^\alpha$.

Proof. For every $\varepsilon > 0$ and each $x \in X$, it is obvious that

$$\{i \leq k_r, j \leq j_u, |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \supseteq \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\}.$$

Thus, we get

$$\begin{aligned}
&\frac{1}{k_{ru}} \left| \{(i, j) : i \leq k_r, j \leq j_u, |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\
&\geq \frac{1}{k_{ru}} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \\
&= \frac{h_{ru}^\alpha}{k_{ru} h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right|.
\end{aligned}$$

If $U_{ij} \xrightarrow{W(S_2)} U$, then for each $x \in X$ the term on the left side of the above inequality convergent to 0 and this implies that for each $x \in X$,

$$\frac{1}{h_{ru}^\alpha} \left| \{(i, j) \in I_{ru} : |\rho_x(U_{ij}) - \rho_x(U)| \geq \varepsilon\} \right| \rightarrow 0.$$

Thus, we get $U_{ij} \xrightarrow{W_2 S_\theta^\alpha} U$. Consequently, $W(S_2) \subseteq W_2 S_\theta^\alpha$. \square

5. Conclusions and Future Work

We introduced new convergence concepts for double set sequences, also we studied the relationships between them. These concepts can also be studied for the ideal convergence and invariant convergence in the future.

6. Competing Interests

The authors declare that there is not any conflict of interests regarding the publication of this manuscript.

REFERENCES

1. M. Baronti and P. Papini, *Convergence of sequences of sets*, Methods of Functional Analysis in Approximation Theory **76** (1986), 133-155.
2. G. Beer, *On convergence of closed sets in a metric space and distance functions*, Bull. Aust. Math. Soc. **31** (1985), 421-432.
3. G. Beer, *Wijsman convergence: A survey*, Set-Valued Var. Anal. **2** (1994), 77-94.
4. S. Bhunia, P. Das and S.K. Pal, *Restricting statistical convergence*, Acta Math. Hungar. **134** (2012), 153-161.
5. J.S. Connor, *The statistical and strong p -Cesàro convergence of sequences*, Analysis **8** (1988), 47-63.
6. H. Çakalli and E. Savaş, *Statistical convergence of double sequences in topological groups*, J. Comput. Anal. Appl. **12** (2010), 421-426.
7. R. Çolak, *Statistical convergence of order α* , Modern Methods in Analysis and Its Applications **1** (2010), 121-129.
8. R. Çolak, Ç.A. Bektaş, *λ -statistical convergence of order α* , Acta Math. Sci. Ser. B **31** (2011), 953-959.
9. R. Çolak and Y. Altın, *Statistical convergence of double sequences of order α* , J. Funct. Spaces Appl. **2013** (2013), 1-5.
10. E. Dündar and B. Altay, *Multipliers for bounded \mathcal{L}_2 -convergent of double sequences*, Math. Comput. Modelling **55** (2012), 1193-1198.
11. E. Dündar and N. Pancaroğlu Akın, *Wijsman regularly ideal convergence of double sequences of sets*, J. Intell. Fuzzy Systems **37** (2019), 8159-8166.
12. M. Et and H. Şengül, *Some Cesàro-type summability spaces of order α and lacunary statistical convergence of order α* , Filomat **28** (2014), 1593-1602.
13. H. Fast, *Sur la convergence statistique*, Colloq. Math. **2** (1951), 241-244.
14. A.R. Freedman, J.J. Sember and M. Raphael, *Some Cesàro-type summability spaces*, Proc. Lond. Math. Soc. **37** (1978), 508-520.
15. J.A. Fridy, *On statistical convergence*, Analysis **5** (1985), 301-313.
16. J.A. Fridy and C. Orhan, *Lacunary statistical convergence*, Pacific J. Math. **160** (1993), 43-52.
17. A.D. Gadjiev and C. Orhan, *Some approximation theorems via statistical convergence*, Rocky Mountain J. Math. **32** (2001), 129-138.

18. F. Móricz, *Statistical convergence of multiple sequences*, Arch. Math. **81** (2003), 82-89.
19. M. Mursaleen and O.H.H. Edely, *Statistical convergence of double sequences*, J. Math. Anal. Appl. **288** (2003), 223-231.
20. M. Mursaleen, C. Çakan, S.A. Mohiuddine and E. Savaş, *Generalized statistical convergence and statistical core of double sequences*, Acta Math. Sin. **26** (2010), 2131-2144.
21. F. Nuray and B.E. Rhoades, *Statistical convergence of sequences of sets*, Fasc. Math. **49** (2012), 87-99.
22. F. Nuray, U. Ulusu and E. Dündar, *Cesàro summability of double sequences of sets*, Gen. Math. Notes **25** (2014), 8-18.
23. F. Nuray, U. Ulusu and E. Dündar, *Lacunary statistical convergence of double sequences of sets*, Soft Comput. **20** (2016), 2883-2888.
24. F. Nuray, E. Dündar and U. Ulusu, *Wijsman statistical convergence of double sequences of sets*, Iran. J. Math. Sci. Inform. **16** (2021), 55-64.
25. R.F. Patterson and E. Savaş, *Lacunary statistical convergence of double sequences*, Math. Commun. **10** (2005), 55-61.
26. A. Pringsheim, *Zur theorie der zweifach unendlichen Zahlenfolgen*, Math. Ann. **53** (1900), 289-321.
27. E. Savaş, *Some sequence spaces and statistical convergence*, Int. J. Math. Math. Sci. **29** (2002), 303-306.
28. E. Savaş and R.F. Patterson, *Lacunary statistical convergence of multiple sequences*, Appl. Math. Lett. **19** (2006), 527-534.
29. E. Savaş, *Double almost statistical convergence of order α* , Adv. Difference Equ. **2013** (2013), 1-9.
30. E. Savaş, *Double almost lacunary statistical convergence of order α* , Adv. Difference Equ. **2013** (2013), 1-10.
31. E. Savaş, *On \mathcal{I} -lacunary statistical convergence of order α for sequences of sets*, Filomat **29** (2015), 1223-1229.
32. I.J. Schoenberg, *The integrability of certain functions and related summability methods*, Amer. Math. Monthly **66** (1959), 361-375.
33. Y. Sever, Ö. Talo and B. Altay, *On convergence of double sequences of closed sets*, Contemporary Analysis and Applied Mathematics **3** (2015), 30-49.
34. H. Steinhaus, *Sur la convergence ordinaire et la convergence asymptotique*, Colloq. Math. **2** (1951), 73-74.
35. T. Šalát, *On statistically convergent sequences of real numbers*, Math. Slovaca **30** (1980), 139-150.
36. H. Şengül and M. Et, *On lacunary statistical convergence of order α* , Acta Math. Sci. Ser. B **34** (2014), 473-482.
37. H. Şengül and M. Et, *On \mathcal{I} -lacunary statistical convergence of order α of sequences of sets*, Filomat **31** (2017), 2403-2412.
38. Ö. Talo, Y. Sever and F. Başar, *On statistically convergent sequences of closed sets*, Filomat **30** (2016), 1497-1509.
39. Ş. Tortop and E. Dündar, *Wijsman \mathcal{I}_2 -invariant convergence of double sequences of sets*, Journal of Inequalities and Special Functions **9** (2018), 90-100.
40. B.C. Tripathy, *On statistical convergence*, Proc. Est. Acad. Sci. **47** (1998), 299-303.
41. U. Ulusu and F. Nuray, *Lacunary statistical convergence of sequences of sets*, Progress in Applied Mathematics **4** (2012), 99-109.
42. U. Ulusu and E. Gülle, *Some statistical convergence types for double set sequences of order α* , Facta Universitatis Ser. Math. Inform. **35** (2020), 595-603.
43. R.A. Wijsman, *Convergence of sequences of convex sets, cones and functions*, Bull. Amer. Math. Soc. **70** (1964), 186-188.

Esra Gülle is working as research assistant in Department of Mathematics, Afyon Kocatepe University. She received Ph.D. in Mathematics from Afyon Kocatepe University, Afyonkarahisar, Turkey. Her area of expertise includes: Summability theory and sequences spaces through functional analysis. She has many international published papers.

Department of Mathematics, Afyon Kocatepe University, Afyonkarahisar, Turkey.
e-mail: egulle@aku.edu.tr

Uğur Ulusu is working as Associate Professor in Sivas Cumhuriyet University. He received Ph.D. in Mathematics from Afyon Kocatepe University, Afyonkarahisar, Turkey. His area of expertise includes: Summability theory and sequences spaces through functional analysis. He is the author of many international published papers.

Sivas Cumhuriyet University, 58140, Sivas, Turkey.
e-mail: ugurulusu@cumhuriyet.edu.tr