

Daytime Napping and Nighttime Sleep During Pregnancy and Preterm Birth in Iran

Farnaz Shaliha¹, Maryam Mozaffari¹, Faeze Ramezani¹, Hamideh Hajnasiri², Farnoosh Moafi²

¹Student Research Committee, Qazvin University of Medical Sciences, Qazvin, Iran; ²Social Determinants of Health Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

Objectives: This study investigated the relationship between sleep quality during pregnancy and preterm birth.

Methods: This longitudinal study was conducted between August 2018 and May 2019. The participants were 150 pregnant women who had been referred to 7 healthcare centers in the city of Qazvin, Iran and met the inclusion criteria. The Petersburg Sleep Quality Index, the Epworth Sleepiness Scale, and 2 questions about daytime sleep status and a demographic questionnaire were administered at 14-18 weeks and 28-32 weeks of gestation. Data were analyzed using the Mann-Whitney test, the Fisher exact test, and univariate and multivariable logistic regression.

Results: In the present study, poor sleep quality affected 84.7% of the participants at 14-18 weeks and 93.3% at 28-32 weeks of gestation. The final model for preterm birth prediction incorporated age and the Petersburg Sleep Quality Index score in the second and third trimesters. Preterm birth increased by 14% with each unit increase in age. With each unit increase in the Petersburg Sleep Quality Index score in the second and third trimesters, preterm birth increased by 42% and 28%, respectively, but the *p*-values of these factors were not significant.

Conclusions: Although a significant percentage of pregnant women had poor sleep quality, no significant relationship was found between sleep quality during pregnancy and preterm birth.

Key words: Pregnancy, Premature birth, Sleep, Iran

INTRODUCTION

Preterm birth, which is defined as prenatal birth before 37 weeks of gestation, is the leading cause of infant mortality and is associated with many complications [1]. Each year, approxi-

mately 15 million babies are born prematurely worldwide [2]. The preterm birth rate in the United States has been on the rise since 2014, reaching 9.93% of births in 2017 [3]. In developing countries, especially in South Asia and sub-Saharan Africa, the preterm birth rate is estimated to be more than 15% [4]. Approximately 90% of preterm deliveries occur in developing countries, of which 85% are reported in Asia and Africa [5]. The preterm birth rate in Iran is estimated to be between 5.40% and 19.85%, indicating that preterm birth is a relatively common problem [6].

The etiological, epidemiological, and clinical nature of preterm birth in many cases is unidentified and idiopathic. Researchers have identified preterm birth as a multifactorial complication and have attributed it to individual, social, environ-

Received: August 1, 2020 **Accepted:** March 11, 2021

Corresponding author: Farnoosh Moafi
Social Determinants of Health Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, P.O. Box 34199-15315, Qazvin, Iran

E-mail: f.moafi.sbm@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

mental, psychological, medical, genetic, biological, and infertility-related factors [7,8]. Numerous studies have been conducted in this field, and factors reported to affect preterm birth include infection, premature rupture of membranes, history of preterm birth, multiple births, pre-eclampsia, use of assisted reproductive techniques, pregnancy at an early age, low education, inadequate care during pregnancy, obesity, stress, gestational diabetes, placenta previa, and reproductive system disorders [8-10]. Given that the cause of two-thirds of preterm births remains unknown, it seems that the best way to prevent this complication is to identify pregnant women with risk factors and to seek out risk factors among biological and psychosocial causes [11].

Pregnancy is a special period in life that is associated with significant changes in sleep patterns and sleep quality. It can result in sleep disorders such as insomnia, snoring, nocturnal awakening, restless legs syndrome, and obstructive sleep apnea during the day [12-14]. The prevalence of these disorders has been estimated to be 58% in the second trimester of pregnancy and 66% in the third trimester [15]. Sleep quality during pregnancy and postpartum is affected by hormonal and physiological changes. Sex hormones and gonadotropins may also contribute to sleep disorders [16]. Frequent urination, inability to find a comfortable position, pelvic pain, leg cramps, back pain, hunger, and itchy skin are some of the factors that affect the sleep quality of pregnant women [17]. Sleep disorders significantly affect the quality of life and reduce daily physical, psychological, and social functioning [18-20], and can also cause neuroendocrine, metabolic, and inflammatory changes [21]. Sleep deprivation and sleep disorders during pregnancy may increase rates of pre-eclampsia, gestational diabetes, anxiety, postpartum depression, reduced pain tolerance, prolonged labor, and cesarean delivery [22-24].

One of the undesirable consequences of poor sleep quality during pregnancy may be preterm birth [25]. In this regard, studies have reported contradictory results. Some studies have found a positive association between poor sleep quality and preterm birth [25-28], but this finding was not reported in other studies [29,30]. These studies have also been criticized; for example, the quantity and quality of sleep during pregnancy were examined by asking only 2 questions [27]. In another studies [25,29,30], sleep disorders were examined only in the third trimester. Most of these studies also investigated maternal sleep during the night and did not examine the relationship between maternal daytime sleep and preterm birth. There-

fore, due to the high prevalence of sleep disorders during pregnancy, and considering the contradictory information about the relationship between poor sleep quality and preterm birth and the importance of the effects of preterm birth on infant health, this study was conducted to determine the relationship between sleep quality during pregnancy and preterm birth.

METHODS

Study Design and Population

In this longitudinal study, multi-stage sampling was used, and the study population was pregnant women referred to healthcare centers in Qazvin, Iran, between August 2018 and May 2019. The cluster random sampling method was used in this study. Quotas were allocated to each of the 7 healthcare centers in Qazvin according to the number of pregnant women referred to each center within 1 month before sampling. The samples were selected using convenience sampling and they entered the study based on the inclusion criteria. The inclusion criteria were being Iranian; being able to read and write; not consuming alcohol, drugs, or tobacco; being 18-35 years old; being at 14-18 weeks of gestation; being a primigravida, having a low-risk and singleton pregnancy; and not having any chronic diseases or history of a mental illness (according to the patient and based on the patient's records). In this study, according to the findings of other studies [31], the ratio of p -value=0.5 was considered to find the sample size, and by using $\alpha=0.05$, $Z_{1-\alpha/2}=1.96$, and $d=0.05$, the minimum sample size was estimated at 150.

Instruments

Data collection tools included a demographic questionnaire, the Petersburg Sleep Quality Index (PSQI), and the Epworth Sleepiness Scale (ESS). In addition, daytime sleep status was examined with 2 questions: "Do you take a nap during the day?" and "How long do you nap during the day?"

The PSQI is a self-rated questionnaire that evaluates sleep quality. The reported sensitivity and specificity of the PSQI are 89.6% and 86.5%, respectively, for identifying sleep disorders in adults. The index includes 7 items: subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication, and daytime dysfunction. Each item is scored on a scale from 0 to 3 using a 4-point Likert scale. The total score is 0-21, and a total score of

5 or higher is indicative of poor sleep quality. The reliability of the Iranian version of the PSQI has been examined and the Cronbach's alpha has been reported to be 0.79 [32].

The ESS is a self-administered questionnaire measuring sleepiness during 8 common situations in daytime life. The range of the score for each item is 0-4 and the total score is 32-42. A score of 11 or higher is indicative of poor daily sleep quality. Cronbach's alpha of this questionnaire was reported to be 0.77 for the population of Iranian pregnant women [29].

Procedure

Data collection was performed in 3 stages after obtaining the necessary permissions. In the first stage, at 14-18 weeks of gestation, the demographic questionnaires, the PSQI, the ESS, and the daytime sleep status questions were completed. In the second stage, at the 28-32 weeks of gestation, except for the demographic questionnaire, the rest of the questionnaires were completed again. In the third stage, after delivery, information about the delivery was collected over the phone. Moreover, the patients' records were also checked to confirm the information that they provided. Preterm birth was defined as less than 37 weeks of gestation based on the last menstrual period or based on the ultrasound measurements taken in the first trimester.

Statistical Analysis

Data analysis was performed using SPSS version 23 (IBM Corp., Armonk, NY, USA). Descriptive statistics (e.g., frequency and percentage to describe qualitative data and mean and standard deviation to describe quantitative data) were used. The Kolmogorov-Smirnov test was used to assess the normality of the data, and it showed that the quantitative variables of the study were not normally distributed ($p < 0.05$). Therefore, to compare the sleep quality in the second and third trimesters of pregnancy, the Mann-Whitney non-parametric test was used to describe qualitative variables, and the Fisher exact test was used to compare them. The significance level was considered to be p -value < 0.05 .

Univariate and multivariable logistic regression methods were used to investigate the relationship between predictive variables and preterm birth. In order to determine the variables affecting preterm birth, each of the demographic and sleep variables was individually investigated in the univariate logistic regression analysis, and variables with a significance level less than 0.2 were entered into the multivariable logistic re-

gression model, in which the dependent variable was preterm birth. Independent variables included factors affecting preterm birth that were significant in the univariate logistic regression analysis. These variables were entered into the model using the forward Wald (i.e., stepwise) method.

Ethics Statement

The present study was approved with the code of IR.QUMS.REC.1396.310 by the Ethics Committee of Qazvin University of Medical Sciences, Iran. Informed consent was obtained from all women participating at the time of sampling. They were also provided with adequate explanations about the aim and the type of the study. They were assured that they could leave the study at any time.

RESULTS

In this study, 158 pregnant women were invited to participate, of whom 5 were excluded from the study due to pregnancy-related complications and 3 were excluded due to incomplete questionnaires. Data from 150 pregnant women were ultimately analyzed. The response rate in the present study was 94.9%.

Of the 150 pregnant women surveyed, 6 (4.0%) had preterm births and 144 (96.0%) had term births. The mean gestational age in the second trimester was 16.04 ± 1.47 weeks, while in the third trimester, it was 29.91 ± 1.38 weeks. The mean age of the pregnant women was 25.81 ± 4.15 years. The majority of participants had academic education (72.0%) and were housewives (82.0%). A sufficient income was reported by 80.0% of participants. The results of the univariate logistic regression analysis are presented in detail in Table 1.

The mean PSQI score in the second trimester of pregnancy was 6.53 ± 2.17 , and in the third trimester of pregnancy, it was 6.89 ± 2.12 . The mean ESS score was 12.59 ± 3.76 in the second trimester and 12.00 ± 3.11 in the third trimester of pregnancy. A total of 80.0% of pregnant women in the second trimester and 68.7% in the third trimester of pregnancy reported daytime napping. The mean nap duration in the second trimester was 65.40 ± 54.30 minutes, and in the third trimester, it was 43.90 ± 39.89 minutes. The results of the univariate logistic regression analysis are presented in detail in Table 1.

Table 2 presents the final model for preterm birth predictors, which included age and the PSQI score in the second and third trimesters. The results showed that as age and the PSQI

Table 1. The result of univariate logistic regression analysis

Variables	Term	Preterm	Total	Univariate logistic regression analysis		
				p-value	Standard error	β
Demographic						
Education						
Academic degree	102 (70.8)	6 (100)	108 (72.0)	-	-	-
None academic degree	42 (29.2)	0 (0.0)	42 (28.0)	0.998	6201.91	-18.37
Occupation						
Housewife	117 (81.2)	6 (100)	123 (82.0)	-	-	-
Employed	27 (18.7)	0 (0.0)	27 (18.0)	0.998	7735.14	18.23
Family income status						
Under sufficient	16 (11.0)	0 (0.0)	16 (11.0)	-	-	-
Sufficient	115 (80.0)	5 (83.0)	120 (80.0)	0.999	1004.24	-18.63
In term of savings	13 (9.0)	1 (17.0)	14 (9.0)	0.253	1.13	-0.57
Age ¹	25.72 ± 4.08	28.00 ± 5.62	25.81 ± 4.15	0.192	0.10	0.13
Sleep status						
14-18 wk						
Petersburg Sleep Quality Index ¹	6.44 ± 2.17	8.67 ± 0.51	6.53 ± 2.17	0.023	0.18	0.41
Epworth Sleepiness Scale	12.53 ± 3.75	14.00 ± 4.09	12.59 ± 3.76	0.348	0.09	0.09
Daily nap duration (min)	66.46 ± 54.87	40.00 ± 30.98	65.40 ± 54.30	0.252	0.01	-0.01
Daily nap						
Yes	116 (80.6)	4 (66.7)	120 (80.0)	-	-	-
No	28 (19.4)	2 (33.3)	30 (20.0)	0.414	0.89	0.72
28-32 wk						
Petersburg Sleep Quality Index ¹	6.81 ± 2.01	9.00 ± 3.57	6.89 ± 2.12	0.019	0.17	0.39
Epworth Sleepiness Scale	11.97 ± 3.10	12.67 ± 3.61	12.00 ± 3.11	0.593	0.11	0.06
Daily nap duration (min)	43.65 ± 39.34	50.00 ± 55.85	43.90 ± 39.89	0.702	0.01	0.00
Daily nap						
Yes	99 (68.8)	4 (66.7)	103 (68.7)	-	-	-
No	45 (31.3)	2 (33.3)	47 (31.3)	0.914	0.88	0.09

Values are presented as number (%) or mean ± standard deviation.

¹Variables selected to enter into the multivariate regression model.

Table 2. Multivariate regression analysis of preterm birth predictors

Variables	β	SE	Wald	p-value	OR (95% CI)
Age	0.13	0.109	1.59	0.206	1.14 (0.92, 1.42)
Petersburg Sleep Quality Index (wk)					
14-18	0.35	0.227	2.42	0.119	1.42 (0.91, 2.22)
28-32	0.25	0.209	1.48	0.224	1.28 (0.85, 1.94)

Hosmer and Lemeshow test: $\chi^2=5.95$ df=8 Sig=0.65

Model summary: -2 log likelihood=40.827; Cox-Snell $R^2=0.062$; Nagelkerke $R^2=0.216$

Omnibus test of model coefficients: $\chi^2=9.55$ df=3 Sig=0.02

SE, standard error; OR, odds ratio; CI, confidence interval; df, degrees of freedom; Sig, significance.

score in the second and third trimesters increased (with a higher score indicating poor sleep quality), the rate of preterm birth increased. The PSQI in the second trimester, with a ratio

of 1.42, was the strongest predictor in the model; however, none of the variables included in the final model were significant.

In addition, considering the features of the model, based on the Hosmer and Lemeshow test ($p > 0.05$) and the significance of omnibus tests of model coefficients ($p < 0.05$), the final model had good explanatory power and efficiency. This model explained between 0.6% and 22.0% of preterm birth variance.

Table 3 depicts the frequency distribution of sleep variables based on the trimester of pregnancy. The results showed that as pregnancy progressed, the PSQI score increased, meaning that sleep quality became worse over the course of pregnancy. Thus, 84.7% of participants had poor sleep quality in the second trimester and 93.3% had poor sleep quality in the third trimester, reflecting a significant difference ($p < 0.05$). However, the ESS score decreased in the third trimester of pregnancy compared to that in the second trimester, meaning as the pregnancy progressed, the quality of daytime sleep improved. Thus, 77.3% of pregnant women in the second trimester and 66.0%

of those in the third trimester had poor daytime sleep quality ($p = 0.01$). In the second and third trimesters, 80.0% and 68.7% of pregnant women reported daytime napping, respectively ($p < 0.05$). The nap duration was 54.30 ± 65.40 minutes in the second trimester; and in the third trimester, it was 39.89 ± 43.90 minutes ($p = 0.001$).

As secondary outcomes, data on mode of delivery and birth weight were collected and analyzed in the same way as for preterm birth. In these analyses, preterm birth was examined as a confounding factor in the relationship between sleep variables and mode of delivery and birth weight. In the univariate logistic regression analysis, daytime napping and the ESS score in the second trimester, and nap duration and the PSQI score in the third trimester had a significance level less than 0.2, so they were entered into the multivariable regression analysis. In the final analysis, daytime napping in the second trimester and the duration of naps in the third trimester significantly predicted the mode of delivery ($p < 0.05$, respectively). Thus, if pregnant women nap more during the day in the second trimester and the nap duration is increased in the third trimester, the incidence of cesarean section would be reduced (odds ratio, 0.32 and 0.98, respectively). The final model for the mode of delivery is presented in Table 4. For birth weight, in the univariate regression analysis, only the PSQI score in the second trimester predicted low birth weight (under 2500 g), but the relationship was not significant ($p = 0.196$).

Table 3. Sleep variables across pregnancy

Variables	Pregnancy (wk)		p-value
	14-18	28-32	
Petersburg Sleep Quality Index			
<5	23 (15.3)	10 (6.7)	0.031
≥ 5	127 (84.7)	140 (93.3)	
Epworth Sleepiness Scale			
<11	34 (22.7)	51 (34.0)	0.010
≥ 11	116 (77.3)	99 (66.0)	
Daily nap			
Yes	120 (80.0)	103 (68.7)	0.021
No	30 (20.0)	47 (31.3)	
Daily nap duration	54.30 ± 65.40	39.89 ± 43.90	0.001

Values are presented as number (%) or mean \pm standard deviation.

DISCUSSION

The present study investigated the relationship between sleep quality during pregnancy and preterm birth. Based on the results of the initial data, the risk of preterm birth increased

Table 4. Multivariate regression analysis of cesarean section predictors

Variables	β	SE	Wald	p-value	OR (95% CI)
Sleep status					
14-18 wk					
Daily nap	-1.12	0.48	5.33	0.021	0.32 (0.12, 0.84)
Epworth Sleepiness Scale	-0.48	0.05	0.94	0.332	0.95 (0.86, 1.05)
28-32 wk					
Daily nap duration (min)	-0.12	0.00	5.88	0.015	0.98 (0.97, 0.99)
Petersburg Sleep Quality Index	-0.18	0.09	0.03	0.844	0.98 (0.82, 1.17)
Hosmer and Lemeshow test: $\chi^2 = 14.142$	df = 4	Sig = 0.007			
Model summary: -2 log likelihood = 192.774; Cox-Snell $R^2 = 0.006$; Nagelkerke $R^2 = 0.008$					
Omnibus test of model coefficients: $\chi^2 = 0.833$	df = 1	Sig = 0.361			

SE, standard error; OR, odds ratio; CI, confidence interval; df, degrees of freedom; Sig, significance.

as the PSQI and the ESS score decreased, and decreased as daytime napping and nap duration increased; however, these associations were not significant. Most previous studies have examined the relationship of respiratory problems of pregnant women while sleeping with pregnancy outcomes, but few studies have been conducted on sleep quality during pregnancy. Some of those studies have assessed sleep quality in all trimesters of pregnancy [20,28,33]. Although those studies identified relationships between sleep quality and preterm birth, the results varied according to how each study defined the trimesters of pregnancy. In contrast, other studies examined sleep quality in just 1 trimester of pregnancy and found different results. In 2 studies, the relationship between sleep disorder and preterm birth was significant [25,26]. However, Dolatian et al. [29] did not report a significant relationship between sleep quality and preterm birth. Using different methods can play a role as a confounder. In most studies, sleep quality was assessed with the PSQI [25,28,33], but the Insomnia Severity Index [29], the ESS [25,29], and the International Classification of Diseases, 9th revision, Clinical Modification codes [26] have also been used. Moreover, different inclusion criteria and sample sizes can contribute to differences in results.

The results of this study showed that according to the PSQI, the quality of sleep decreased as pregnancy progressed and a significant percentage of pregnant women had poor sleep quality. This finding has been confirmed in other studies [16,28,31]. Although Mindell et al. [17] did not report a significant difference in PSQI scores throughout pregnancy, they reported that sleep disorders increased as gestational age increased. The fact that most studies on sleep quality during pregnancy used the PSQI [34] improves the generalizability of the findings.

According to the results of the present study, a significant percentage of pregnant women had poor daytime sleep quality, but they showed an improvement in daytime sleep quality as gestational age increased and daytime napping and nap duration decreased. In contrast, Hutchison et al. [16] reported an increase in daytime napping as pregnancy progressed, but this result was not supported by other studies [17,35]. According to Sedov et al. [34], the effect of gestational age on sleep disorders during pregnancy has yet to be fully understood. The reason for differences among study results can be attributed to differences in gestational age at the time of data collection, the use of different methods to assess daytime sleep quality, and different modes of reporting. In addition, lifestyle

differences throughout the world may be another contributing factor [16].

Furthermore, the present study found that an increase in daytime napping in the second trimester and in nap duration in the third trimester was associated with a reduced risk of cesarean section. Women suffering from poor sleep quality during pregnancy feel more tired and therefore have less energy to have a normal delivery [36]. Zhang et al. [20] reported that poor sleep quality in the first, second, and third trimesters of pregnancy were associated with an increased risk of cesarean section; however, this finding has not been reported in all studies [29,37]. Discrepancies among studies can be attributed to cultural and social differences, as well as differences in sleep definitions, sleep assessment tools, and gestational age at the time of sampling. None of the sleep variables was significantly related to birth weight, and likewise, other studies found no significant relationship between sleep quality and birth weight [25,29]. Given the relative lack of studies on this issue, more research is needed to reach a definitive conclusion; however, the relationships between sleep quality in pregnancy and the mode of delivery and birth weight were not the main focus of our study, so we did not investigate these variables in detail.

One of the strengths of this study is the sampling method. Qazvin, the setting of the research, is a city with a high proportion of immigrants due to its geographical and strategic status. Therefore, people from different social, economic, educational, and ethnic groups live together. Randomized cluster sampling from different areas of the city provided the opportunity to include participants from different socioeconomic classes in this study. Therefore, the authors believe that the results are generalizable to pregnant women in Iran. Using 2 questionnaires and questions about daily sleep disorders made this study different from previous studies. However, due to the limitations of the assessment, in this study, data were collected at 2 time periods, and no information was gathered on maternal sleep quality in the first trimester and before pregnancy. In addition, given the accuracy of polysomnography in assessing sleep status, it is recommended that this method be used alone or in combination with other methods in future studies. In this study, the relationship between sleep quality in pregnancy and preterm birth was not significant. Therefore, it is suggested that future studies should be conducted with larger samples of pregnant women.

A high percentage of pregnant women suffered from poor

nighttime and daytime quality of sleep. Therefore, it is important to find these women and improve maternal sleep quality by simple advice and interventions (e.g., behavioral interventions). Poor sleep quality in the second and third trimesters of pregnancy was not associated with preterm birth or birth weight. However, increased daytime napping in the second trimester and nap duration in the third trimester showed associations with a lower risk of cesarean section, implying that improving sleep quality in pregnancy might reduce the rate of cesarean sections.

CONFLICT OF INTEREST

The authors have no conflicts of interest associated with the material presented in this paper.

FUNDING

None.

ACKNOWLEDGEMENTS

The authors would like to express their appreciation for the pregnant women who took part in this study.

AUTHOR CONTRIBUTIONS

Conceptualization: MM, FM. Data curation: FS, FR. Formal analysis: FM. Funding acquisition: None. Methodology: FM, HH, FS. Writing – original draft: FM, FR, FS. Writing – review & editing: FM, MM, HH.

ORCID

Farnaz Shaliha <https://orcid.org/0000-0001-5579-8505>
Maryam Mozaffari <https://orcid.org/0000-0002-1194-9989>
Faeze Ramezani <https://orcid.org/0000-0001-6082-5889>
Hamideh Hajnasiri <https://orcid.org/0000-0003-2527-1504>
Farnoosh Moafi <https://orcid.org/0000-0002-3650-9293>

REFERENCES

1. Manuck TA, Rice MM, Bailit JL, Grobman WA, Reddy UM, Wapner RJ, et al. Preterm neonatal morbidity and mortality by gestational age: a contemporary cohort. *Am J Obstet Gynecol* 2016;215(1):103.e1-103.e14.

2. Purisch SE, Gyamfi-Bannerman C. Epidemiology of preterm birth. *Semin Perinatol* 2017;41(7):387-391.
3. Martin JA, Hamilton BE, Osterman MJ, Driscoll AK, Drake P. Births: final data for 2017. *Natl Vital Stat Rep* 2018;67(8):1-50.
4. World Health Organization. Born too soon: the global action report on preterm birth; 2012 [cited 2020 Jul 1]. Available from: https://www.who.int/pmnch/media/news/2012/201204_borntoosoon-report.pdf.
5. Beck S, Wojdyla D, Say L, Betran AP, Merialdi M, Requejo JH, et al. The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity. *Bull World Health Organ* 2010;88(1):31-38.
6. Sharifi N, Khazaeian S, Pakzad R, Fathnezhad Kazemi A, Chehreh H. Investigating the prevalence of preterm birth in Iranian population: a systematic review and meta-analysis. *J Caring Sci* 2017;6(4):371-380.
7. de Freitas AS, Dobbler PC, Mai V, Procianoy RS, Silveira RC, Corso AL, et al. Defining microbial biomarkers for risk of preterm labor. *Braz J Microbiol* 2020;51:151-159.
8. Tellapragada C, Eshwara VK, Bhat P, Acharya S, Kamath A, Bhat S, et al. Risk factors for preterm birth and low birth weight among pregnant Indian women: a hospital-based prospective study. *J Prev Med Public Health* 2016;49(3):165-175.
9. Leal MD, Esteves-Pereira AP, Nakamura-Pereira M, Torres JA, Theme-Filha M, Domingues RM, et al. Prevalence and risk factors related to preterm birth in Brazil. *Reprod Health* 2016; 13(Suppl 3):127.
10. Shoja M, Shoja E, Gharaei M. Prevalence and affecting factors on preterm birth in pregnant women referred to Bentolhoda hospital-Bojnurd. *J North Khorasan Univ Med Sci* 2015;7(4): 855-863 (Persian).
11. Mehraban Z, Alizadeh L, Sadeghniyat HK. Surveying the relationship between sleep apnea and preterm birth in women referring to health centers of Ardabil. *Urmia Med J* 2016;26(11): 931-940 (Persian).
12. Gupta R, Dhyani M, Kendzerska T, Pandi-Perumal SR, BaHamam AS, Srivanitchapoom P, et al. Restless legs syndrome and pregnancy: prevalence, possible pathophysiological mechanisms and treatment. *Acta Neurol Scand* 2016;133(5):320-329.
13. Kocsis I, Szilágyi T, Turos J, Bakó A, Frigy A. Effect of a gymnastics program on sleep characteristics in pregnant women. *Taiwan J Obstet Gynecol* 2017;56(2):204-209.
14. Mehraban Z, Yahya S, Sadeghniyat K. Restless legs syndrome during pregnancy and preterm birth in women referred to

- health centers of Ardabil. *Iran Red Crescent Med J* 2015;17(12): e24438.
15. Hung HM, Ko SH, Chen CH. The association between prenatal sleep quality and obstetric outcome. *J Nurs Res* 2014;22(3): 147-154.
 16. Hutchison BL, Stone PR, McCowan LM, Stewart AW, Thompson JM, Mitchell EA. A postal survey of maternal sleep in late pregnancy. *BMC Pregnancy Childbirth* 2012;12:144.
 17. Mindell JA, Cook RA, Nikolovski J. Sleep patterns and sleep disturbances across pregnancy. *Sleep Med* 2015;16(4):483-488.
 18. Abdollahzadeh M, Naji SA. The effect of matricaria chamomilla on sleep quality of elderly people admitted to nursing homes. *Iran J Nurs* 2014;27(89):69-79 (Persian).
 19. Ram S, Seirawan H, Kumar SK, Clark GT. Prevalence and impact of sleep disorders and sleep habits in the United States. *Sleep Breath* 2010;14(1):63-70.
 20. Zhang J, Li F, Lin Y, Sheng Q, Yu X, Zhang X. Subjective sleep quality in perimenopausal women and its related factors. *J Nanjing Med Univ* 2007;21(2):116-119.
 21. Nodine PM, Matthews EE. Common sleep disorders: management strategies and pregnancy outcomes. *J Midwifery Womens Health* 2013;58(4):368-377.
 22. Chang JJ, Pien GW, Duntley SP, Macones GA. Sleep deprivation during pregnancy and maternal and fetal outcomes: is there a relationship? *Sleep Med Rev* 2010;14(2):107-114.
 23. Gunduz S, Kosger H, Aldemir S, Akcal B, Tevriciz H, Hizli D, et al. Sleep deprivation in the last trimester of pregnancy and inadequate vitamin D: is there a relationship? *J Chin Med Assoc* 2016;79(1):34-38.
 24. Malekzadegan A, Moradkhani M, Ashayeri H, Haghani H. Effect of relaxation on insomnia during third trimester among pregnant women. *Iran J Nurs* 2010;23(64):52-58 (Persian).
 25. Najar S, Sharafi F, Afshari P, Haghhighizadeh MH. The relationship between sleep disorders during pregnancy and premature labor and low birth weight. *Iran J Obstet Gynecol Infertil* 2017;20(7):44-49 (Persian).
 26. Felder JN, Baer RJ, Rand L, Jelliffe-Pawlowski LL, Prather AA. Sleep disorder diagnosis during pregnancy and risk of preterm birth. *Obstet Gynecol* 2017;130(3):573-581.
 27. Li R, Zhang J, Zhou R, Liu J, Dai Z, Liu D, et al. Sleep disturbances during pregnancy are associated with cesarean delivery and preterm birth. *J Matern Fetal Neonatal Med* 2017;30(6): 733-738.
 28. Ota H, Hasegawa J, Sekizawa A. Effect of sleep disorders on threatened premature delivery. *J Perinat Med* 2017;45(1):57-61.
 29. Dolatian M, Mehraban Z, Sadeghniat K. The effect of impaired sleep on preterm labour. *West Indian Med J* 2014;63(1):62-67.
 30. Tomfohr-Madsen L, Cameron EE, Dunkel Schetter C, Campbell T, O'Beirne M, Letourneau N, et al. Pregnancy anxiety and preterm birth: the moderating role of sleep. *Health Psychol* 2019; 38(11):1025-1035.
 31. Seyedahmadinejad FS, Golmakani N, Asgharipour N, Shakeri MT. Investigation of sleep quality during the third trimester of pregnancy and some related factors in primigravida women referred to health care centers in Mashhad-2014. *Avicenna J Nurs Midwifery Care* 2014;22(4):53-60 (Persian).
 32. Afkham Ebrahimi A, Bandi G, Salehi M, Tafti K, Vakili Y, Farsi A. Sleep parameters and the factors affecting the quality of sleep in patients attending selected clinics of Rasoul-e-Akram hospital. *Razi J Med Sci* 2008;15:31-38 (Persian).
 33. Okun ML, Luther JF, Wisniewski SR, Wisner KL. Disturbed sleep and inflammatory cytokines in depressed and nondepressed pregnant women: an exploratory analysis of pregnancy outcomes. *Psychosom Med* 2013;75(7):670-681.
 34. Sedov ID, Cameron EE, Madigan S, Tomfohr-Madsen LM. Sleep quality during pregnancy: a meta-analysis. *Sleep Med Rev* 2018;38:168-176.
 35. Loubé DI, Poceta JS, Morales MC, Peacock MD, Mitler MM. Self-reported snoring in pregnancy. Association with fetal outcome. *Chest* 1996;109(4):885-889.
 36. Beebe KR, Lee KA. Sleep disturbance in late pregnancy and early labor. *J Perinat Neonatal Nurs* 2007;21(2):103-108.
 37. Tsai SY, Kuo LT, Lee CN, Lee YL, Landis CA. Reduced sleep duration and daytime naps in pregnant women in Taiwan. *Nurs Res* 2013;62(2):99-105.