



ARTICLE

Effect of Deer Antler Extract on Muscle Differentiation and 5-Aminoimidazole-4-Carboxamide Ribonucleoside (AICAR)-Induced Muscle Atrophy in C2C12 Cells

Kyungae Jo¹, Woo Young Jang¹, Beom Sik Yun², Jin Soo Kim², Hyun-Sun Lee³, Yeok Boo Chang¹, and Hyung Joo Suh^{1,*}

OPEN ACCESS

Received October 8, 2020
Revised January 13, 2021
Accepted April 8, 2021

¹Department of Integrated Biomedical and Life Sciences, Graduate School, Korea University, Seoul 02841, Korea

²R&D Center, Kwangdong Pharm Co., Ltd, Seoul 08381, Korea

³Agency for Korea National Food Cluster, Iksan 54576, Korea

*Corresponding author : Hyung Joo Suh
Department of Integrated Biomedical and Life Sciences, Graduate School, Korea University, Seoul 02841, Korea
Tel: +82-2-940-2764
Fax: +82-2-921-7207
E-mail: suh1960@korea.ac.kr

*ORCID
Kyungae Jo
<https://orcid.org/0000-0003-4056-6706>
Woo Young Jang
<https://orcid.org/0000-0002-5104-4949>
Beom Sik Yun
<https://orcid.org/0000-0002-0065-5555>
Jin Soo Kim
<https://orcid.org/0000-0001-8218-439X>
Hyun-Sun Lee
<https://orcid.org/0000-0003-0435-0107>
Yeok Boo Chang
<https://orcid.org/0000-0002-2445-5750>
Hyung Joo Suh
<https://orcid.org/0000-0001-8869-3929>

Abstract The effect of deer antler extract on muscle differentiation and muscle atrophy were evaluated to minimize muscle loss following aging. Various deer antler extracts (HWE, hot water extract of deer antler; FE, HWE of fermented deer antler; ET, enzyme-assisted extract of deer antler; UE, extract prepared by ultrasonication of deer antler) were evaluated for their effect on muscle differentiation and inhibition of 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR)-induced muscle atrophy in C2C12 cells. Morphological changes according to the effect of antler extracts on muscle differentiation were confirmed by Jenner-Giemsa staining. In addition, the expression levels of genes related to muscle differentiation and atrophy were confirmed through qRT-PCR. In the presence of antler extracts, the length and thickness of myotubes and myogenin differentiation 1 (MyoD1) and myogenic factor 5 (Myf5) gene expression were increased compared to those in the control group (CON). Gene expression of AMP-activated protein kinase (AMPK), MyoD1, and myogenin, along with the muscle atrophy factors muscle RING finger-1 (MuRF-1) and forkhead box O3a (FoxO3a) upon addition of deer antler extracts to muscle-atrophied C2C12 cells was determined by qRT-PCR after treatment with AICAR. The expression of MuRF-1 and FoxO3a decreased in the groups treated with antler extracts compared to that in the group treated with AICAR alone. In addition, gene expression of MyoD1 and myogenin in the muscle atrophy cell model was significantly increased compared that into the CON. Therefore, our findings indicate that antler extract can increase the expression of MyoD1, Myf5 and myogenin, inhibit muscle atrophy, and promote muscle differentiation.

Keywords deer antler, muscle differentiation, muscle atrophy, 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR), C2C12 cells

Introduction

The aging process involves the natural degeneration of all the organs of the body over time. In this process, muscular atrophy is an important influencing factor. Muscular atrophy involves the impairment of the major nerves of muscles or a decrease in the absolute volume and strength of muscles under prolonged under certain conditions (Bonewald, 2019; Szentesi et al., 2019).

Muscle cells are involved in the regeneration of muscle fibers containing mechanical, chemical, or degenerative lesions through activation, differentiation, and proliferation (Ceafalan et al., 2014; Karalaki et al., 2009). Differentiation of myoblasts is essential for the formation of muscle fibers involved in the development and regeneration of skeletal muscle (Grefte et al., 2007). These monocyte myoblasts proliferate and differentiate, following which, they fuse with existing muscle fibers to form multinucleated myotubes and myofibers. Proliferation and differentiation of muscle cells occur similarly during development and postnatal birth (Schiaffino et al., 2013). Therefore, promotion of myoblast proliferation and differentiation and induction of myotube hypertrophy should be beneficial for muscle regeneration and control of muscle mass.

Muscle atrophy with aging is accompanied by muscle damage caused by oxidative damage and malnutrition caused by lack of muscle metabolism. Thus, muscle atrophy is triggered by increasing muscle protein degradation and decreased protein synthesis (Zhang et al., 2018). In recent studies, activation of AMP-activated protein kinase (AMPK) increased muscle atrophy F-box (MAFbx)/atrogen-1 and muscle RING finger-1 (MuRF-1) via the transcription factor forkhead box O3a (FoxO3a) (Jaitovich et al., 2015). This process has been reported to be directly involved in muscle atrophy by muscle protein degradation through the activation of the ubiquitin-proteasome pathway (Jaitovich et al., 2015; Nakashima and Yakabe, 2007; Tong et al., 2009). Since muscle mass represents a balance between muscle cell replication, protein synthesis, muscle cell death and protein degradation (Scicchitano et al., 2018), increased muscle mass can prevent muscle atrophy by causing an increase in muscle differentiation and inhibiting muscle loss.

Deer antler (*Cervi parvum* Cornu) is a non-osteolytic horn of *Cervus nippon* Temminck or Marcus (*Cervus elaphus* L) and has long been used in oriental countries, such as China and Korea. Deer antler has been reported to contain hexose, pentose, uronic acid, sialic acid, free amino acids, minerals, prostaglandins, and gangliosides in glycolipids (Jhon et al., 1999; Sunwoo et al., 1995). Water-soluble proteins, polypeptides, and free amino acids have been shown to be the main biologically active components of deer antler (Moreau et al., 2004). In addition, there is evidence that deer antler extract has strong potential for promoting bone growth and development (Chen et al., 2015; Shi et al., 2010). Additionally, Chen et al. (2014) estimated through microarray analysis that the anti-fatigue effect of collagen and protein from deer antler extracts is mediated by the increased expression of genes involved in muscle contraction and development.

By evaluating the effect of strengthening muscle differentiation and inhibiting muscle atrophy by antler extract having various activities, it was attempted to investigate the possibility of application as a functional material to suppress muscle atrophy due to aging. Therefore, the effects of deer antler extract on the promotion of myoblast differentiation and 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR)-induced muscle atrophy were examined using C2C12 fibroblast.

Materials and Methods

Materials

Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), horse serum (HS), and penicillin/streptomycin (PS) were purchased from WELGENE (Daegu, Korea) and were used for cell culture and differentiation. AICAR and 3-(4,5-

dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) were obtained from Sigma-Aldrich Chemical (St. Louis, MO, USA). Dried antler (*Cervus canadensis*) from adult male Korean elk was obtained from Kwang Dong Pharmaceutical (Seoul, Korea). According to the previous report, antler water extract (HWE) and fermented antler (FE) were prepared. The antler powder was extracted by refluxing for 3 h at 95°C after adding 8 times the amount of water. Fermented antler was extracted by adding 8 times the amount of phosphate buffer (PBS, 50 mM, pH 6.0) the antler powder, fermenting it with *Bacillus subtilis* for 48 h, and refluxing for 3 h at 95°C. The ultrasonic extract (UE) was extracted by adding 8 times the amount of PBS to the antler powder for 30 minutes at 120 kHz ultrasonic waves (AUG-R3-900, Asia Ultrasonic, Bucheon, Korea). After ultrasonic extraction, reflux extraction was performed at 95°C to obtain an UE. After adding 8 times the amount of PBS (50 mM, pH 7.8) to the antler powder and suspending it, Alcalase equivalent to 0.5% of the antler powder was added, followed by hydrolysis at 50°C for 8 h. After hydrolysis, reflux extraction was performed at 95°C to obtain an antler enzyme hydrolysate (ET). The antler extracts were concentrated and lyophilized to be used in the further experiment.

Analysis of collagen

According to the Sunwoo et al. (1998) method, the collagen content was calculated by multiplying the content of hydroxyproline by 7. According to the method of Stegemann and Stalder (1967), the content of hydroxyproline before and after acid hydrolysis with 6 N HCl was measured using HPLC. A Phenomenex Luna C18 Column (250×4.6 mm, 5 μm) was employed, using following mobile phase: sodium acetate buffer (pH 4.3) containing 3% glacial acetic acid and acetonitrile were mixed with 650 and 350 mL, respectively. The flow rate was 1 mL/min, hydroxyproline was identified at a wavelength of excitation 260/emission 330 nm, and the injection quantity was 20 μL. The collagen content was calculated by multiplying the difference by 7 to the hydroxyproline content before and after acid hydrolysis.

Cell culture and differentiation

The myoblast cell line C2C12 used in this study was purchased from the American Type Culture Collection (Manassas, VA, USA). C2C12 myoblasts were maintained at 37°C and 5% CO₂ using a growth medium containing 90% DMEM, 10% FBS, and 100 unit/mL PS. To maintain cells at an appropriate density, they were passaged every 48 h. Differentiation medium containing 90% DMEM, 10% HS, and 100 unit/mL PS was used to differentiate C2C12 myoblasts into myotubes. To confirm the morphological changes during the differentiation process, an inverted microscope (Inverted microscope, Carl Zeiss, Gottingen, Germany) was used to observe cells at a magnification of 200×.

Cell viability

The cell viability of C2C12 cells was measured using an MTT assay. C2C12 cells were seeded into a 96-well plate at a concentration of 1×10^6 cells/mL the day before the experiment. Deer antler extracts (0–1,000 μg/mL) were incubated with cells for 24 h. Subsequently, the culture supernatant was discarded and 5 mg/mL MTT dissolved in DMEM was added to each well; the plate was then covered with aluminum foil to create dark conditions and incubated for 2 h. After removing the MTT solution, 100 μL of dimethyl sulfoxide was added to the cells and left for 2 h. The absorbance was measured at 540 nm using a microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Jenner-Giemsa staining and measurement of myotube length and diameter

Myotube length and diameter were measured using images obtained through Jenner-Giemsa staining of C2C12 differentiated

cells (Veliça and Bunce, 2011). The cultured cells were washed with PBS, fixed with 100% methanol for 5 min, and air dried for 10 min. The Jenner staining solution (Sigma-Aldrich, St. Louis, MO) was diluted 3-fold with 1 mM sodium PBS (pH 5.6); 1 mL of this solution was incubated in the wells for 5 min and then the cells were washed with distilled water. The cells were then incubated for 10 min at 25°C with 1 mL Giemsa stain diluted 1:20 with 1 mM sodium PBS (pH 5.6) and then washed again with water. Cells in each well were observed using a MM-400 microscope (Nikon, Tokyo, Japan) and photographed in nine randomly selected areas using a digital microscope camera JENOPTIK ProgRes speed Xt^{core} 3 and the ProgRes Image Capture Software (Jenoptik Optical Systems GmbH, Jena, Germany). Myotube length and diameter were measured and quantified using the Image-J software (Scion, Frederick, MD, USA).

RNA isolation and quantitative real-time reverse-transcription-polymerase chain reaction (qRT-PCR)

Total RNA from C2C12 cells was extracted using the TRIzol® reagent (Invitrogen, CA, USA), according to the manufacturer's protocol. Quality controlled RNA samples with high optical density ratios were treated with RQ1 RNase-free DNase I (Promega, WI, USA). One microgram of total RNA was reverse transcribed using SuperScript® III Reverse Transcriptase (Invitrogen) with oligo d (T) primers. qRT-PCR was performed using the Taqman Gene Expression Master Mix (Applied Biosystems, Foster City, CA, USA), and quantitative analyses were conducted using the StepOne plus Software V. 2.0 (Applied Biosystems). All results were normalized against the expression of the control gene, GAPDH (NM_008084.2), using the $\Delta\Delta C_t$ method (Livak and Schmittgen, 2001). The information of primers for the qRT-PCR of tested genes related to muscle differentiation and atrophy were as follows: myogenin differentiation 1 (MyoD1) (NM_010866.2), myogenic factor 5 (Myf5) (NM_008656.5), AMPK (NM_001013367.3), forkhead box O3 (FoxO3a) (NM_019740.2), muscle RING-finger protein 1 (MuRF-1) (NM_001039048.2), and myogenin (NM_031189.2).

Statistical analysis

All experimental results were obtained using the SPSS ver. 18.0 (SPSS, Chicago, IL, USA) statistical program. Data represent the mean±SD. Differences among the groups were evaluated by one-way analysis of variance (ANOVA) and Tukey's multiple test. Student's *t*-test was used to analyze differences between groups with normally distributed data.

Results and Discussion

Collagen composition and C2C12 cells viability

The collagen content of HWE, ET, UE, and FE, it was 20.09±0.19, 29.07±0.12, 27.32±0.09, 23.58±0.11 µg/mg of extract, respectively, and it was confirmed that the collagen content of the extract increased due to fermentation, enzyme, and ultrasonication treatment (Table 1). In particular, the content of hydroxyproline (4.15±0.02 µg/mg of extract) and protein (295.05±11.22 mg/g of extract) as well as collagen in ET was higher than that of other antler extracts.

Deer antler extracts, including HWE, FE, ET, and UE were used for the evaluation of myoblast C2C12 cells differentiation activity. These extracts had protein contents of 225.1–295.1 mg/g extract (data not shown). Upon evaluation of the cytotoxicity of the antler extracts against C2C12 cells (Fig. 1), all extracts except the UE did not show a cytotoxicity of up to 1,000 µg/mL. UE-treated cells showed a cell viability of 90.7%–88.9% upon treatment with 200–1,000 mg/mL extract. Deer antler extracts were found to have no or low cytotoxicity at 1,000 µg/mL. Therefore, further experiments were conducted using up to 200 µg/mL of extracts, with little cytotoxicity.

Table 1. Protein, hydroxyproline, and collagen content in antler extracts

Sample	Protein (mg/g)	Hydroxyproline ($\mu\text{g}/\text{mg}$)	Collagen ($\mu\text{g}/\text{mg}$)
HWE	240.66 \pm 10.31	2.87 \pm 0.03	20.09 \pm 0.19
ET	295.05 \pm 11.22	4.15 \pm 0.02	29.07 \pm 0.12
UE	225.08 \pm 7.22	3.90 \pm 0.01	27.32 \pm 0.09
FE	284.68 \pm 6.15	3.37 \pm 0.02	23.58 \pm 0.11

Content value represents the mean \pm SD.

HWE, hot water extract; ET, enzyme hydrolysate; UE, ultrasonic extract; FE, how water extract of fermented deer antler.




Fig. 1. Cell viability of deer antler extracts on C2C12 cells. Values are presented as the mean \pm SD for each group. CON, control group; HWE, hot water extract; FE, how water extract of fermented deer antler; ET, enzyme-derived extract; UE, extract prepared by ultrasonication of deer antler.

Deer antler extracts have been widely used in traditional Chinese medicine for centuries and are generally believed to provide nourishment, strengthen the kidneys, strengthen the spleen, strengthen bones and muscles, and promote blood flow (Wu et al., 2013). Previous studies have reported that the collagen and protein components of deer antler extracts are major bioactive substances with effects, such as anti-fatigue, anti-stress (Fengyan et al., 2001), and bone growth effects (Niu et al., 2012) and stimulation of hematopoietic function (Lee et al., 2012). Kitakaze et al. (2016) reported that collagen hydrolysates promoted differentiation and increased size of myoblast C2C12 cells. Hydroxypropyl-glycine, a collagen hydrolysate, has been reported to induces myogenic differentiation and root canal hypertrophy by increasing the size of the root canal and the expression of root canal specific myosin heavy chain (MyHC) and tropomyosin structural proteins. Hyp-Gly presumably promotes myogenic differentiation by activating the PI3K/Akt/mTOR signaling pathway according to peptide/histidine transporter 1 for entry into cells (Kitakaze et al., 2016; Li et al., 2016; Niu et al., 2012).

Changes in myotube length and diameter caused by C2C12 cells differentiation

To determine the degree of myotube differentiation during C2C12 cells differentiation, Jenner-Giemsa staining was

performed on the second and fourth days of C2C12 cells differentiation to measure changes in myotube morphology, such as myotube length and diameter (Fig. 2 and 3).

On the second day of cell differentiation, myotube length was increased in cells treated with deer antler extracts compared with the control group (CON). On the fourth day, myotube length was increased in cells treated with all deer antler extracts except for FE. The cells treated with FE showed increased myotube length at extract concentrations of 50 and 100 µg/mL and shortened myotube length when treated with extract concentration of 200 µg/mL (Fig. 3).




Fig. 2. Changes in myotube structure during C2C12 cell differentiation, as observed through Jenner-Giemsa staining. Values are presented as the mean±SD for each group. CON, control group; HWE, hot water extract; ET, enzyme-derived extract; UE, extract prepared by ultrasonication of deer antler; FE, how water extract of fermented deer antler.




Fig. 3. Changes in myotube length and diameter during C2C12 cell differentiation. Values are presented as the mean±SD for each group. The different letters indicate statistically significant ($p < 0.05$) differences among groups, as determined through Tukey's test. CON, control group; HWE, hot water extract; ET, enzyme-derived extract; UE, extract prepared by ultrasonication of deer antler; FE, how water extract of fermented deer antler.

On day two of cell differentiation, cells treated with deer antler extract showed a similar diameter to that of CON, but on day four of cell differentiation, cells showed a tendency to have increased myotube diameter compared to CON (Fig. 3). Upon treatment of cells with deer antler extract, myotube length was increased by promoting the myotube differentiation capacity, and the myotube diameter also tended to increase with increasing myotube length.

In the differentiation phase, the length of myoblasts become increased, fused with localized cells and converted into tubular multinuclear cells. When differentiation is induced from myoblasts to myotubes, the number of muscle fibers increases due to the fusion between cells and the thickening of muscle fibers also occurs (Bentzinger et al., 2012; Burattini et al., 2004). Net protein balance affects the size of individual myotubes and myofibers, where an increase in net protein levels leads to muscle hypertrophy (White et al., 2010) and contributes to an increase in skeletal muscle mass (Ontell et al., 1984).

Effect of deer antler extract on the expression of myogenic differentiation-related genes in C2C12 myoblasts

Muscle differentiation is regulated by myogenic regulatory factors (MRFs), such as MyoD1 and Myf5 (Braun et al., 1992; Hyatt et al., 2003). The effect of the antler extract on the expression of muscle differentiation factors MyoD1 and Myf5 was measured on days two and four of C2C12 cell differentiation, respectively (Fig. 4). The expression levels of MyoD1 were not significantly affected by myoblast differentiation on days two and four, whereas Myf5 expression levels were high on days




Fig. 4. Effects of deer extracts on the relative expression of MyoD1 and Myf5 mRNA in C2C12 cells at two and four days of cell differentiation. Values are presented as the mean±SD for each group. The different letters indicate statistically significant ($p < 0.05$) differences among groups, as determined by Tukey's test. CON, normal control; HWE, hot water extract; ET, enzyme-derived extract; UE, extract prepared by ultrasonication of deer antler; FE, how water extract of fermented deer antler; MyoD1, myogenin differentiation 1; Myf5, myogenic factor 5.

two and four compared with the CON. Deer antler extracts significantly increased the expression level of Myf5 ($p < 0.05$). In particular, it was found that the level of Myf5 increased in a concentration-dependent manner when the HWE was added on day four of myogenic differentiation. However, increasing concentrations of UE and FE tended to decrease the expression level of Myf5. These results indicated that antler extract was not only involved in increasing the size (length and diameter) of myotubes, but also in increasing the expression level of myogenic differentiation factor that promotes myoblast C2C12 cell differentiation.

Muscle differentiation is regulated mainly by MRFs, such as MyoD1 and Myf5, which are involved in establishing myogenic lineages (Braun et al., 1992). Later in muscle differentiation, there is an increase in the expression of MyHC, a major structural protein of myotubes (Soundharrajan et al., 2019). Deer antler extracts were also found to be involved in the increase in the expression levels of muscle differentiation factors (Fig. 4), which may increase the size of the myotubes (Fig. 3).

Inhibitory activity of deer antler extracts on muscle atrophy factors in AICAR-induced muscle atrophy in C2C12 cells

As a result of a 0.25–2 mM treatment of AICAR to create a muscle atrophy model in myoblast cells, AMPK expression increased in an AICAR concentration-dependent manner (Fig. 5). C2C12 cells showed no cytotoxicity when treated with 2 mM of AICAR (data not shown). Therefore, C2C12 cells were treated with 2 mM AICAR to induce muscle atrophy, and the effect of the inhibition of muscle atrophy through the addition of antler extracts was confirmed.

The effects of deer antler extracts (50 and 200 $\mu\text{g/mL}$) on the expression levels of muscle atrophy factors (FoxO3a and MuRF-1) were measured in AICAR-induced muscle-atrophy cells (Fig. 6). Upon AICAR treatment, the expression of AMPK increased in C2C12 cells and deer antler extract-treated muscle-atrophy cells compared to the CON. The expression levels of AMPK in deer antler extract-treated cells were higher than those in cells treated with AICAR alone (CON group). The muscle




Fig. 5. Effects of AICAR on relative AMPK mRNA expression in C2C12 cells. Values are presented as the mean \pm SD for each group. The different letters indicate statistically significant ($p < 0.05$) differences among groups, as determined by Tukey's test. CON, control group; AICAR, 5-aminoimidazole-4-carboxamide ribonucleoside; AMPK, AMP-activated protein kinase.




Fig. 6. Effects of deer antler extract on relative mRNA expression of AMPK, FoxO3a, and MuRF-1 in AICAR-induced muscle-atrophied C2C12 cells. Values are presented as the mean±SD for each group. Asterisks indicate significant differences: *** $p < 0.001$ indicates significance of difference compared with the control, as determined by Student's t-test. The different letters indicate statistically significant ($p < 0.05$) differences among groups, as determined by Tukey's test. AMPK, AMP-activated protein kinase; NOR, normal group; CON, control group; HWE, hot water extract; FE, how water extract of fermented deer antler; ET, enzyme-derived extract; UE, extract prepared by ultrasonication of deer antler; FoxO3a, forkhead box O3a; MuRF-1, muscle RING finger-1; AICAR, 5-aminoimidazole-4-carboxamide ribonucleoside.

atrophy factors FoxO3a and MuRF-1 showed significantly higher expression levels in the AICAR treated group (CON) than in the normal group (NOR). In deer antler extract-treated muscle-atrophied cells, the increase in the expression level of the muscle atrophy factor MuRF-1 tended to be lower than that in the CON group cells. In particular, the expression level of FoxO3a in deer antler extract-treated atrophied cells was significantly lower than that in cells treated with AICAR alone ($p < 0.05$).

Recent studies have shown that muscle atrophy occurs due to increased proteolysis due to the activation of the ubiquitin-proteasome pathway (Bodine and Baehr, 2014; Jagoe and Goldberg, 2001; Liu et al., 2016). In particular, muscle atrophy F-box (MAFbx)/atrogen-1 and MuRF-1, which are muscle-specific ubiquitin ligases, are expressed early in the muscle atrophy process and are directly involved in muscle protein degradation (Bodine and Baehr, 2014; Liu et al., 2016). In addition, when muscle atrophy is induced, expression of genes, such as myogenin and MyoD, muscle-specific transcription factors involved in myogenic differentiation, is reduced by activating muscle-specific gene expression (Hyatt et al., 2003; Tintignac et al., 2005).

Inhibitory activity of deer antler extracts on muscle differentiation factors in AICAR-induced muscle atrophy in C2C12 cells

The expression levels of MyoD1 and myogenin, which are muscle differentiation factors, were lower than those of normal cells upon treatment with AICAR alone (Fig. 7). In deer antler extract-treated muscle-atrophied cells, the expression levels of muscle differentiation factors MyoD1 and myogenin were higher than those in cells treated with AICAR alone. The expression levels of MyoD1 in the enzyme-treated deer antler extract (ET, 200 µg/mL)-treated muscle-atrophied cells were significantly higher than those of normal cells ($p < 0.05$). In addition, the expression levels of myogenin in the enzyme-treated deer antler extract (ET, 50 and 200 µg/mL) and fermented deer antler extract (FE, 200 µg/mL)-treated muscle-atrophied cells were significantly higher than that in normal cells. The treatment of AICAR-induced atrophied cells with deer antler extract inhibited the expression of the muscle atrophy factors FoxO3a and MuRF1. In addition, deer antler extract treatment restored the expression levels of myoD1 and myogenin to the same or higher levels as those of normal cells.




Fig. 7. Effects of deer antler extract on relative mRNA expression of MyoD1 and Myogenin in AICAR-induced muscle atrophy in C2C12 cells. Values are presented as the mean±SD for each group. Asterisks indicate significant differences: *** $p < 0.001$ indicates significance of difference compared with the control, as determined by Student's t-test. The different letters indicate statistically significant ($p < 0.05$) differences among groups, as determined by Tukey's test. NOR, normal group; CON, control group; HWE, hot water extract; FE, how water extract of fermented deer antler; ET, enzyme-derived extract; UE, extract prepared by ultrasonication of deer antler; AICAR, 5-aminoimidazole-4-carboxamide ribonucleoside.

In Fig. 6, muscle atrophy factors FoxO3a and MuRF-1 can be seen to be increased upon AICAR treatment, and their increase results in muscle atrophy. In addition, in Fig. 7, the expression levels of MyoD and myogenin, which are myogenic factors, can be seen to decrease upon AICAR treatment. However, MuRF-1 expression decreased upon treatment with antler extract, and the expression factor of myogenesis markers increased. Deer antler extracts seemed to contribute more to the inhibition of muscle atrophy by increasing the expression of muscle differentiation factors than by decreasing the expression of muscle atrophy factors.

The increase in skeletal muscle mass occurs due to an increase in muscle protein synthesis (MPS) compared to muscle protein breakdown (Rennie, 2007). To increase skeletal muscle or inhibit muscle loss, it is important to have both an increase in myogenic capacity and an inhibition of muscle loss. Therefore, the supply of high-quality dietary protein, which is an essential nutrient for promoting muscle and overall metabolic health, is important (Arentson-Lantz et al., 2015). Deer horn extract is an excellent source of protein that is especially rich in collagen type I (Li et al., 2016). Therefore, the active substances of antler extract to myogenesis and inhibit the muscle atrophy are presumed to be protein, collagen and hydroxyproline. It has also been reported that collagen breakdown products, which form a major component of antlers, are involved in the activation of mammalian target of rapamycin (mTORC1) and MPS. The mechanistic/mTORC1 pathway is the central molecular pathway of MPS and is activated by a variety of stimuli, including, but not limited to, resistance exercise (Bolster et al., 2003), insulin (Conejo et al., 2001), and dietary amino acids (Gordon et al., 2013). We found four distinct effects of treating C2C12 cells with antler extract: 1) increase in myotube size, 2) increase in muscle differentiation factor (MyoD1 and Myf5) expression, 3) suppression of muscular atrophy factor (FoxO3a and MuRF-1) expression, and 4) increase in the expression of the muscle differentiation factors (MyoD1 and myogenin), in a muscle atrophy model.

Conclusion

Several studies have reported that collagen, a major component of deer antler, helps myogenesis. In this study, experimental

results on C2C12 cells suggested that antler extract has the ability to inhibit muscle atrophy and promote muscle differentiation by increasing the expression of myoblast differentiation factors MyoD, Myf5 and myogenin. Further experiments will be conducted to identify muscle differentiation promoting substances.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by the Kwang Dong Pharmaceutical Co., Ltd (Q1823491).

Author Contributions

Conceptualization: Jo K, Suh HJ. Data curation: Jang WY, Chang YB. Formal analysis: Yun BS, Kim JS, Lee HS. Methodology: Jo K, Suh HJ. Software: Jang WY, Chang YB. Validation: Yun BS, Lee HS. Investigation: Kim JS, Lee HS. Writing - original draft: Jo K, Suh HJ. Writing - review & editing: Jo K, Jang WY, Yun BS, Kim JS, Lee HS, Chang YB, Suh HJ.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Arentson-Lantz E, Clairmont S, Paddon-Jones D, Tremblay A, Elango R. 2015. Protein: A nutrient in focus. *Appl Physiol Nutr Metab* 40:755-761.
- Bentzinger CF, Wang YX, Rudnicki MA. 2012. Building muscle: Molecular regulation of myogenesis. *Cold Spring Harb Perspect Biol* 4:a008342.
- Bodine SC, Baehr LM, 2014. Skeletal muscle atrophy and the E3 ubiquitin ligases MuRF1 and MAFbx/atrogen-1. *Am J Physiol Endocrinol Metab* 307:E469-E484.
- Bolster DR, Kubica N, Crozier SJ, Williamson DL, Farrell PA, Kimball SR, Jefferson LS. 2003. Immediate response of mammalian target of rapamycin (mTOR)-mediated signalling following acute resistance exercise in rat skeletal muscle. *J Physiol Soc* 553:213-220.
- Bonewald L. 2019. Use it or lose it to age: A review of bone and muscle communication. *Bone* 120:212-218.
- Braun T, Rudnicki MA, Arnold HH, Jaenisch R. 1992. Targeted inactivation of the muscle regulatory gene *Myf-5* results in abnormal rib development and perinatal death. *Cell* 71:369-382.
- Burattini S, Ferri P, Battistelli M, Curci R, Luchetti F, Falcieri E. 2004. C2C12 murine myoblasts as a model of skeletal muscle development: Morpho-functional characterization. *Eur J Histochem* 48:223-234.
- Ceafalan LC, Popescu BO, Hinescu ME. 2014. Cellular players in skeletal muscle regeneration. *BioMed Res Int* 2014:957014.

- Chen J, Yang Y, Abbasi S, Hajinezhad D, Kontulainen S, Honaramooz A. 2015. The effects of elk velvet antler dietary supplementation on physical growth and bone development in growing rats. *Evid Based Complement Alternat Med* 2015:819520.
- Chen JC, Hsiang CY, Lin YC, Ho TY. 2014. Deer antler extract improves fatigue effect through altering the expression of genes related to muscle strength in skeletal muscle of mice. *Evid Based Complement Alternat Med* 2014:540580.
- Conejo R, Valverde AM, Benito M, Lorenzo M. 2001. Insulin produces myogenesis in C2C12 myoblasts by induction of NF- κ B and downregulation of AP-1 activities. *J Cell Physiol* 186:82-94.
- Fengyan S, Huiping L, Yanmei W, Yuxia H, Souming X, Yuqi X. 2001. Protein component extraction and its bioactivity determination of sika deer antler base. *Anim Sci Vet Med* 18:18-20.
- Gordon BS, Kelleher AR, Kimball SR. 2013. Regulation of muscle protein synthesis and the effects of catabolic states. *Int J Biochem Cell Biol* 45:2147-2157.
- Grefte S, Kuijpers-Jagtman AM, Torensma R, Von den Hoff JW. 2007. Skeletal muscle development and regeneration. *Stem Cells Dev* 16:857-868.
- Hyatt JPK, Roy RR, Baldwin KM, Edgerton VR. 2003. Nerve activity-independent regulation of skeletal muscle atrophy: Role of MyoD and myogenin in satellite cells and myonuclei. *Am J Physiol Cell Physiol* 285:C1161-C1173.
- Jagoe RT, Goldberg AL. 2001. What do we really know about the ubiquitin-proteasome pathway in muscle atrophy? *Curr Opin Clin Nutr Metab Care* 4:183-190.
- Jaitovich A, Angulo M, Lecuona E, Dada LA, Welch LC, Cheng Y, Gusarova G, Ceco E, Liu C, Shigemura M, Barreiro E, Patterson C, Nader GA, Sznajder JI. 2015. High CO₂ levels cause skeletal muscle atrophy via AMP-activated kinase (AMPK), FoxO3a protein, and muscle-specific ring finger protein 1 (MuRF1). *J Biol Chem* 290:9183-9194.
- Jhon GJ, Park SY, Han SY, Lee S, Kim Y, Chang YS. 1999. Studies of the chemical structure of gangliosides in deer antler, *Cervus nippon*. *Chem Pharm Bull* 47:123-127.
- Karalaki M, Fili S, Philippou A, Koutsilieris M. 2009. Muscle regeneration: Cellular and molecular events. *In Vivo* 23:779-796.
- Kitakaze T, Sakamoto T, Kitano T, Inoue N, Sugihara F, Harada N, Yamaji R. 2016. The collagen derived dipeptide hydroxyprolyl-glycine promotes C2C12 myoblast differentiation and myotube hypertrophy. *Biochem Biophys Res Commun* 478:1292-1297.
- Lee SH, Suh HJ, Lee HS, Park Y, Park JW, Jung EY. 2012. Hematopoietic effect of *Bacillus subtilis*-fermented antler extract on phenylhydrazine-induced hemolytic anemia in Sprague-Dawley rats. *J Med Food* 15:774-780.
- Li N, Zhang M, Drummen GPC, Zhao Y, Tan YF, Luo S, Qu XB. 2016. Sika deer antler collagen type I-accelerated osteogenesis in bone marrow mesenchymal stem cells via the Smad pathway. *Evid Based Complement Alternat Med* 2016:2109204.
- Liu Y, Du M, Li X, Chen L, Shen Q, Tian J, Zhang D. 2016. Role of the ubiquitin-proteasome pathway on proteolytic activity in postmortem proteolysis and tenderisation of sheep skeletal muscle. *Int J Food Sci Technol* 51:2353-2359.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{- $\Delta\Delta$ C_T} method. *Methods* 25:402-408.
- Moreau M, Dupuis J, Bonneau NH, Lécuyer M. 2004. Clinical evaluation of a powder of quality elk velvet antler for the treatment of osteoarthritis in dogs. *Can Vet J* 45:133-139.
- Nakashima K, Yakabe Y. 2007. AMPK activation stimulates myofibrillar protein degradation and expression of atrophy-

- related ubiquitin ligases by increasing FOXO transcription factors in C2C12 myotubes. *Biosci Biotechnol Biochem* 71:1650-1656.
- Niu F, Zhao Y, Xu Y, Zhang H, Lin Y. 2012. Therapeutic effects of collagen of antler plate on osteoporosis in ovariectomized rats. *Chin J Mod Appl Pharm* 29:93-97.
- Ontell M, Feng KC, Klueber K, Dunn RF, Taylor F. 1984. Myosatellite cells, growth, and regeneration in murine dystrophic muscle: A quantitative study. *Anat Rec* 208:159-174.
- Rennie MJ. 2007. Exercise- and nutrient-controlled mechanisms involved in maintenance of the musculoskeletal mass. *Biochem Soc Trans* 35:1302-1305.
- Schiaffino S, Dyar KA, Ciciliot S, Blaauw B, Sandri M. 2013. Mechanisms regulating skeletal muscle growth and atrophy. *FEBS J* 280:4294-4314.
- Scicchitano BM, Dobrowolny G, Sica G, Musaro A. 2018. Molecular insights into muscle homeostasis, atrophy and wasting. *Curr Genomics* 19:356-369.
- Shi B, Li G, Wang P, Yin W, Sun G, Wu Q, Yu G. 2010. Effect of antler extract on corticosteroid-induced avascular necrosis of the femoral head in rats. *J Ethnopharmacol* 127:124-129.
- Soundharrajan I, Kim DH, Kuppusamy P, Choi KC. 2019. Modulation of osteogenic and myogenic differentiation by a phytoestrogen formononetin via p38MAPK-dependent JAK-STAT and Smad-1/5/8 signaling pathways in mouse myogenic progenitor cells. *Sci Rep* 9:9307.
- Stegemann H, Stalder K. 1967. Determination of hydroxyproline. *Clin Chim Acta* 18:267-273.
- Sunwoo HH, Nakano T, Hudson RJ, Sim JS. 1995. Chemical composition of antlers from wapiti (*Cervus elaphus*). *J Agric Food Chem* 43:2846-2849.
- Sunwoo HH, Nakano T, Sim JS. 1998. Isolation and characterization of proteoglycans from growing antlers of wapiti (*Cervus elaphus*). *Comp Biochem Physiol B Biochem Mol Biol* 121:437-442.
- Szentesi P, Csernoch L, Dux L, Keller-Pintér A. 2019. Changes in redox signaling in the skeletal muscle with aging. *Oxid Med Cell Longev* 2019:4617801.
- Tintignac LA, Lagirand J, Batonnet S, Sirri V, Leibovitch MP, Leibovitch SA. 2005. Degradation of MyoD mediated by the SCF (MAFbx) ubiquitin ligase. *J Biol Chem* 280:2847-2856.
- Tong JF, Yan X, Zhu MJ, Du M. 2009. AMP-activated protein kinase enhances the expression of muscle-specific ubiquitin ligases despite its activation of IGF-1/Akt signaling in C2C12 myotubes. *J Cell Biochem* 108:458-468.
- Veliça P, Bunce CM. 2011. A quick, simple and unbiased method to quantify C2C12 myogenic differentiation. *Muscle Nerve* 44:366-370.
- White RB, Biérinx AS, Gnocchi VF, Zammit PS. 2010. Dynamics of muscle fibre growth during postnatal mouse development. *BMC Dev Biol* 10:21.
- Wu F, Li H, Jin L, Li X, Ma Y, You J, Li S, Xu Y. 2013. Deer antler base as a traditional Chinese medicine: A review of its traditional uses, chemistry and pharmacology. *J Ethnopharmacol* 145:403-415.
- Zhang Y, Li C, Zhang H, Li W, Li J, Wang X. 2018. Color-coded digital subtraction angiography for assessing acute skeletal muscle ischemia-reperfusion injury in a rabbit model. *Acad Radiol* 25:1609-1616.