

Staphylococcus saprophyticus and *Escherichia coli*: Tracking from sperm fertility potential to assisted reproductive outcomes

Fatemeh Ghasemian, Shahin Esmaeilnezhad, Mohammad Javad Mehdipour Moghaddam

Department of Biology, Faculty of Sciences, University of Guilan, Rasht, Iran

Objective: Bacteriospermia and urogenital infections are common problems in male infertility. This study aimed to evaluate the effects of bacteriospermia on sperm parameters and clinical outcomes in semen samples infected with two common bacteria (*Staphylococcus saprophyticus* and *Escherichia coli*) in northern Iran.

Methods: Microbiological tests were performed to isolate and identify organisms from 435 semen samples from infertile couples. Semen samples were assessed according to the World Health Organization criteria. The protamine status, chromatin structure, chromatin condensation, and acrosome reaction of sperm and assisted reproductive outcomes were determined in couples with different male infertility factors.

Results: Among the total cases, the two most prevalent pathogens were considered: *S. saprophyticus* (38.2%) and *E. coli* (52.9%). In the semen samples infected with *E. coli*, the spontaneous acrosome reaction and abnormal chromatin condensation were more common ($p < 0.05$). Significant increases in abnormal chromatin condensation and deprotamination were seen in the presence of *S. saprophyticus*. In washed semen, tight adhesion between the sperm midpiece and *S. saprophyticus* was observed. There was also a significant decrease in the fertilization rate using semen samples infected with *S. saprophyticus* and *E. coli* during *in vitro* fertilization cycles ($p < 0.001$). In addition, the presence of *S. saprophyticus* and *E. coli* in semen samples was associated with a lower likelihood of clinical pregnancy in couples with various factors of male infertility.

Conclusion: Poor results of assisted reproductive techniques may be correlated with semen samples infected with two common bacteria in northern Iran.

Keywords: Assisted reproductive technique; Bacterial Infections; Fertilization; Pregnancy; Semen analysis

Introduction

Approximately 15% of cases of male infertility are due to urogenital tract inflammation and infection [1]. Clinical data show that local inflammation or infection can be seen in up to 60% of patients re-

ceiving treatment with assisted reproductive technology (ART) [2]. Infections are the main origin of inflammatory disorders in the male genital tract. However, controversy exists regarding the role of different types of bacteria in decreasing sperm parameters. Therefore, male fertility may be inhibited by the negative effects of bacteriospermia and leukocytospermia on sperm parameters [2]. Existing evidence suggests a connection between bacteriospermia and alterations of semen quality. In recent years, the detrimental effects of different types of bacteria strains and/or leukocytes on some sperm parameters were studied [3]. However, the premise that the direct effects of bacteria are the reason for decreasing the chance of success in ART has not been confirmed.

Several studies have investigated the effects of different bacteria

Received: October 20, 2020 · Revised: January 18, 2021 · Accepted: February 14, 2021

Corresponding author: **Fatemeh Ghasemian**
Department of Biology, Faculty of Sciences, University of Guilan, Rasht 41335-1914, Iran
Tel: +98-13-33343630 Fax: +98-13-33367066 E-mail: ghasemian@guilan.ac.ir

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

on semen parameters and male fertility. However, to the best of our knowledge, no study in the literature has yet investigated the impact of *Staphylococcus saprophyticus* on sperm quality and male fertility potential during *in vitro* fertilization (IVF) techniques. However, the effects of *Escherichia coli* have been explored in some research. Therefore, a comparative study of these two bacteria on semen parameters and fertility results of sperm in laboratory conditions would help to better understand the potential effects of bacteria on sperm fertility potential. Since the assisted reproductive outcomes of ejaculated semen with bacteriospermia are an interesting subject, we conducted this study to determine the prevalence of bacteriospermia in infertile men in northern Iran. In addition, the effects of bacteriospermia on sperm quality and assisted reproductive outcomes were studied among couples with different factors of male infertility.

Methods

1. Semen sample collection and preparation

After excluding infertile couples with female-factor infertility, semen samples were collected from 607 infertile couples with male factor infertility who were undergoing IVF-intracytoplasmic sperm

injection (IVF-ICSI) cycles at the Alzahra Educational and Remedial Center (IVF center) from May 2017 to June 2019 (Figure 1). The exclusion criteria were men having at least one of the following conditions: (1) taking hormone-containing medications; (2) diabetes; (3) thyroid disease; (4) a history of diseases affecting the reproductive tract (e.g., cryptorchidism, varicocele, testicular torsion, chitic, or other causes of infertility that had been medically proven); (5) a history of surgery of the reproductive system, male factor infertility with previous failed IVF-ICSI cycles, and/or hormonal abnormalities; (6) heavy smoking or alcohol use; and (7) exposure to physical or chemical agents with known negative reproductive effects that may influence the semen quality. In addition, semen samples infected with leukocytes were excluded from this study. Samples infected with other bacterial species than *S. saprophyticus* and *E. coli* were excluded because the number of semen samples infected with other bacterial species was very small. Hence, statistically significant associations with clinical outcomes would not have been found (Figure 1).

The semen samples were analyzed according to the World Health Organization (WHO) criteria [4]. The semen volume, pH, sperm motility, concentration, and viability were evaluated. According to the WHO criteria [4], the semen samples were classified as showing nor-




Figure 1. Flowchart of the study design. IVF, *in vitro* fertilization; ICSI, intracytoplasmic sperm injection. ^{a)}All normal and infected semen samples were categorized according to World Health Organization criteria into four groups as following: normozoospermia, asthenozoospermia, teratozoospermia, and oligoasthenoteratozoospermia. ICSI and/or IVF procedures were performed for each group based on sperm parameters, and assisted reproductive outcomes were followed until the live birth stage for each group; ^{b)}Due to the small number of semen samples infected with these bacteria, statistically significant associations with clinical outcomes would not have been found.

mozoospermia, teratozoospermia, asthenozoospermia, or oligoasthenoteratozoospermia. The severity of abnormalities in the semen samples in each studied group (e.g., asthenozoospermia, teratozoospermia, and oligoasthenoteratozoospermia) was similar. This study was approved by the Guilan University of Medical Sciences committee, and informed consent was obtained from all volunteers to participate in this study.

2. Seminal leukocyte quantification

To distinguish white blood cells (WBCs) in semen samples, samples were studied in high-power fields. Only samples with ≥ 5 round cells were assessed with peroxidase staining based on the Endtz test [5]. Peroxidase-positive leukocytes were indicated as stained brown, counted, and reported as $\times 10^6$ WBC/mL semen.

3. Microbiological analysis

Initially, patients received comprehensive information about the method of semen sample collection, the period of sexual abstinence, and the need to avoid taking antibiotics for at least 1 week before semen collection. Semen samples were collected by masturbation after 3–4 days of sexual abstinence. Patients were asked to urinate and wash their hands, penis, and scrotum before ejaculation to avoid possible contamination from the urine or external genitalia. Therefore, semen collection was done in a sterile container, following all given guidance. The semen samples were transported within 1 hour from collection to the Microbiology Laboratory of the University of Guilan to screen for a comprehensive range of microbiological organisms such as anaerobic and aerobic organisms as well as fungi and mycoplasma. First, the semen samples diluted with sterile saline (1:10) were centrifuged at $300 \times g$ for 15 minutes. The sediment was sown using 10- μ L calibrated loops on selective specific media. The elimination of seminal plasma and condensation of bacteria increased culture sensitivity [6].

All cultures were incubated at 37°C and the culture media, incubation times, and conditions were as follows: (1) blood agar to culture aerobic bacteria under aerobic conditions for 24 hours; (2) Sabouraud agar to culture fungi under aerobic conditions for 48 hours; and (3) chocolate agar to culture other microorganisms at 5% CO₂ for 48 hours. A number of isolated colonies of pathogenic bacteria higher than 1×10^3 colony-forming units per milliliter was considered as a positive result. Microorganisms were determined using antimicrobial sensitivity testing according to standard conventional techniques such as Gram staining, oxidase, catalase, and other biochemical tests.

4. Sperm assays

1) Toluidine blue stain

Toluidine blue (TB) and aniline blue (AB) staining have been introduced as sensitive tests to distinguish abnormalities in sperm chromatin structure and abnormal condensation, respectively. Thin smears were prepared on silane-coated slides. In TB staining, the sperm heads with normal chromatin structure were observed as light blue, while those with diminished integrity and abnormal chromatin structure were shown as deep violet/purple. In addition, in AB staining, at least 200 spermatozoa were counted and recorded as sperm with dark-blue (abnormal) and colorless (normal) heads. The positive group (test group; $n=6$) was also evaluated following the incubation of semen samples with *S. saprophyticus* and *E. coli* *in vitro*.

2) Acrosome reaction assessment

Triple staining was used to evaluate the acrosome reaction (acrosome integrity or reacted acrosome) according to the Talbot and Chacon method [7]. In brief, 2% trypan blue was added to spermatozoa (1:1), and incubated at 37°C for 15 minutes. Fixation was done using glutaraldehyde (3% glutaraldehyde in 0.1 M cacodylate buffer at pH 7.4) for 30 to 60 minutes. After centrifugation (at $600 \times g$ for 5 minutes), the smear was prepared from fixed sperm and stained with Bismarck Brown Y at 40°C for 5 minutes. The slides were then stained in Rose Bengal at 24°C for 20–45 minutes. Washing and dehydration were performed in water and an alcohol series, respectively. A total of 300 spermatozoa in each slide cleared with xylene and protected with a coverslip were examined using a light microscope. The positive group (test group; $n=6$) was also evaluated following the incubation of semen samples with *S. saprophyticus* and *E. coli* *in vitro*.

3) Determination of sperm chromatin packaging by chromomycin A3

Chromomycin A3 (CMA3) stains guanosine-cytosine-rich sequences as a polymerase inhibitor fluorochrome, which distinguishes the degree of sperm protamination [8]. In this way, air-dried smears of spermatozoa were fixed in methanol/glacial acetic acid (3:1) for 20 minutes at 4°C. The slides were then air-dried at room temperature and stained with 100 μ L of CMA3 solution (0.25 mg/mL CMA3 in McIlvaine's buffer, containing 10 μ M MgCl₂) (Figure 2). The positive group (test group; $n=5$) was also evaluated following the incubation of semen samples with *S. saprophyticus* and *E. coli* *in vitro*.

4) Scanning electron microscopy

The semen samples prepared by density gradient centrifugation were fixed with Karnovsky solution for 30 minutes at 4°C. Centrifuga-




Figure 2. Sperm deprotamination. Spermatozoa with negative (dull yellow/normal) and positive (bright yellow/abnormal) chromomycin A3 staining. Scale bar=20 μ m.

tion was done at $4,000 \times g$ for 15 minutes. After washing the samples, post-fixation was performed with 1% osmium tetroxide for 30 minutes. Dehydration was then done in ascending grades of ethanol, with drying at a critical point using CO_2 (Balzers CPD-010). Gold-coated specimens (Balzers MED-010) were examined using a Philips FEM 515 scanning electron microscope (SEM). The positive group (test group; $n=4$) was also evaluated following the incubation of semen samples with *S. saprophyticus* and *E. coli* *in vitro*. The adhesion of bacteria to sperm structures was compared to the sham group (human spermatozoa incubated with *S. saprophyticus* and *E. coli* under *in vitro* conditions) [2].

5. ICSI and IVF laboratory procedures

On the day of oocyte retrieval, semen samples were collected via masturbation, liquefied, and prepared by density gradient centrifugation (90 and 45% SpermGrade; Vitrolife, Sweden, Göteborg). Semen parameters were assessed according to the WHO [4] criteria. The ICSI or IVF procedure was performed up to 3–4 hours after sperm preparation. The selection of sperm for IVF or ICSI was based on sperm morphology, motility, concentration, patient's history, infertility etiology, and female age. The evaluation of fertilization was performed 16–18 hours after ICSI/IVF with the observation of the 2-pronuclear stage. The embryo development rate was also evaluated at 2–3 days. Intrauterine embryo transfer (ET) was performed at 2–3 days after microinjection. Increasing serum beta-human chorionic gonadotropin concentrations and detection of the fetal heart-beat were defined as biochemical and clinical pregnancy at 2 and 4 weeks after ET, respectively.

6. Statistical analysis

The prevalence of bacteriospermia was calculated from the proportion of positive cases relative to the total number of the study population and expressed as a percentage. Using linear regression, the relationship between bacteriospermia and other semen parameters was analyzed. The Fisher exact test was performed to assess the fertilization and cleavage rate, clinical pregnancy rate, and live birth rate. The statistical analysis was done using IBM SPSS ver. 20 (IBM Corp., Armonk, NY, USA). The P -values < 0.05 were considered to indicate statistical significance.

Results

1. Semen samples

In this study, the women and men had a mean age of 26.0 ± 3.2 years (range, 22–32 years) and 32.0 ± 5.9 years (range, 26–38 years), respectively. Statistically significant differences were not found for the age ($p > 0.05$) or body mass index (BMI; $p > 0.05$) of the female partners in the normozoospermia. In addition, these similar results were seen about groups of asthenozoospermia, teratozoospermia, and oligoasthenoteratozoospermia ($p > 0.05$). Men's age and BMI also did not show significant differences in different groups ($p > 0.05$). Therefore, positive or negative effects of these parameters on ART outcomes were excluded.

Microbiological analyses were done for semen samples from 435 infertile men, of which 197 semen cultures were negative (45.2%) and 238 (54.7%) were positive. *E. coli* and *S. saprophyticus* were the most commonly isolated pathogens (52.9% and 38.2%, respectively). Semen samples infected with *Enterococcus faecalis* ($n=7$, 2.9%), *Staphylococcus aureus* ($n=6$, 2.5%), *Ureaplasma urealyticum* ($n=4$, 1.6%), *Streptococcus agalactiae* ($n=3$, 1.2%), and *Gardnerella vaginalis* ($n=1$, 0.4%) were also detected. Those bacterial species other than *S. saprophyticus* and *E. coli* were excluded to trace reproductive outcomes due to the small number of semen samples, as a result of which statistical significance would not be expected for associations with clinical outcomes. The infection severity of semen samples with *S. saprophyticus* and *E. coli* was almost identical. Mixed infections with both bacteria (*S. saprophyticus* and *E. coli*) were not observed in any of the cultured semen samples. Therefore, the quality of semen samples infected with each type of bacteria (*S. saprophyticus* or *E. coli*) and their clinical outcomes were analyzed.

Among the 414 specimens analyzed based on the WHO criteria [4], 141 (34.05%) had normozoospermia, 100 (24.15%) had teratozoospermia, 103 (24.87%) had asthenozoospermia, and 70 (19.9%) had oligoasthenoteratozoospermia.

2. Sperm parameters

As seen in Table 1, the percentage of sperm deprotection (seen by CMA3 staining) in normozoospermia samples infected with *S. saprophyticus* (258/950, 27.15%) was significantly higher ($p < 0.05$) than that of the non-bacteriospermia samples (116/825, 14.06%). This rate was higher, at 37.8% ($p < 0.01$) and 36.8% ($p < 0.05$), in the asthenozoospermia and teratozoospermia samples infected with *S. saprophyticus*, respectively. The evaluation of acrosome integrity showed that the rate of premature acrosome reaction was higher in the oligoasthenoteratozoospermia samples infected with *E. coli* (279/856, 33.5%; $p < 0.05$). Sperm viability significantly declined in all samples infected with *E. coli* ($p < 0.05$). A decrease of viability was also seen in the asthenozoospermia ($p < 0.001$) and teratozoospermia ($p < 0.05$) samples infected with *S. saprophyticus* (54.9% and 59.02%, respectively). In addition, the evaluation of abnormalities in sperm chromatin condensation using the AB stain showed that the rate of abnormal sperm chromatin condensation (dark sperm) was higher in both bacteriospermia groups in the samples with different infertility factors (Table 1).

3. SEM images of bacteriospermia

SEM evaluation was performed for each sample prepared by density gradient centrifugation. The tight adhesion between bacteria and spermatozoa was found. Adhesion of *S. saprophyticus* to the tail

of the sperm, especially to the sperm midpiece, was observed. However, the frequency of these adhesions was significantly higher in the asthenozoospermia and teratozoospermia samples from patients with male factor infertility. The adhesion of both bacteria types to spermatozoa was found in the unwashed semen samples; however, after washing, adhesion of *S. saprophyticus* was still observed, but less frequently (Figure 3).




Figure 3. Interaction between bacteria and human spermatozoa. Scanning electron micrographs of prepared human spermatozoa after density gradient centrifugation. Adhesion of *Staphylococcus saprophyticus* to the sperm midpiece (arrows) was observed.

Table 1. The effect of bacteriospermia on semen quality in men with various infertility factors

Variable	No. of patients	CMA3 (%)	Spontaneous AR (%)	AB (%)	TB (%)	Viability (%)
Normozoospermia						
<i>Staphylococcus saprophyticus</i>	41	27.2 ± 3.2 ^{a)}	20.1 ± 2.2	22.6 ± 2.8	25.7 ± 2.6	68.6 ± 10.8
<i>Escherichia coli</i>	36	20.1 ± 2.2	26.2 ± 2.9	29.7 ± 2.4 ^{a)}	27.8 ± 2.4	64.1 ± 9.4 ^{b)}
Nonbacteriospermia	64	14.1 ± 2.6	18.1 ± 1.9	16.8 ± 1.8	20.1 ± 2.9	82.2 ± 10.9
Asthenozoospermia						
<i>Staphylococcus saprophyticus</i>	19	37.8 ± 2.5 ^{b)}	26.4 ± 2.2	31.6 ± 2.7	25.8 ± 2.6	54.9 ± 9.2 ^{d)}
<i>Escherichia coli</i>	30	23.8 ± 2.2	29.9 ± 2.3	29.1 ± 2.1	26.9 ± 2.4	53.4 ± 9.1 ^{d)}
Nonbacteriospermia	54	19.1 ± 1.9	19.2 ± 2.4	21.8 ± 2.2	22.4 ± 2.6	79.5 ± 10.0
Teratozoospermia						
<i>Staphylococcus saprophyticus</i>	15	36.8 ± 3.6 ^{a)}	28.6 ± 2.9	36.4 ± 3.3 ^{a)}	30.4 ± 3.8	59.0 ± 9.5 ^{a)}
<i>Escherichia coli</i>	35	26.6 ± 2.4	31.8 ± 3.3	34.7 ± 3.6	33.6 ± 3.4	53.1 ± 9.1 ^{b)}
Nonbacteriospermia	50	22.3 ± 3.2	24.3 ± 2.4	23.6 ± 2.8	26.4 ± 2.6	70.8 ± 8.3
Oligoasthenoteratozoospermia						
<i>Staphylococcus saprophyticus</i>	16	39.3 ± 3.4	27.4 ± 2.2	38.7 ± 3.5 ^{a)}	35.7 ± 3.6	53.4 ± 9.4
<i>Escherichia coli</i>	25	32.4 ± 3.1	33.5 ± 2.7 ^{a)}	37.4 ± 2.7 ^{a)}	37.4 ± 2.5	51.9 ± 8.3 ^{a)}
Nonbacteriospermia	29	28.6 ± 2.6	20.9 ± 2.2	21.6 ± 2.3	30.1 ± 2.6	61.6 ± 9.1

Values are presented as mean ± standard deviation. The semen samples infected with *S. saprophyticus* showed significant differences in terms of sperm quality such as sperm deprotection (normozoospermia, asthenozoospermia, and teratozoospermia), abnormalities in sperm chromatin condensation (teratozoospermia and oligoasthenoteratozoospermia), and viability (asthenozoospermia and teratozoospermia) in comparison to the control group (non-bacteriospermia). In addition, the semen samples infected with *E. coli* showed significant differences in sperm quality such as abnormal chromatin condensation (normozoospermia and oligoasthenoteratozoospermia), spontaneous acrosome reaction (oligoasthenoteratozoospermia), and viability (normozoospermia, asthenozoospermia, teratozoospermia, and oligoasthenoteratozoospermia) in comparison to the control group (non-bacteriospermia). CMA3, chromomycin A3; AR, acrosome reaction; AB, aniline blue; TB, toluidine blue.

^{a)} $p < 0.05$; ^{b)} $p < 0.01$; ^{c)} $p < 0.001$.

4. ICSI and IVF outcomes

Table 2 shows the clinical outcomes of IVF and ICSI cycles. The clinical results revealed that there was no significant difference in the fertilization rate of ICSI cycles from normozoospermia samples. However, the fertilization rate of normozoospermia samples infected with *S. saprophyticus* ($p < 0.001$) and *E. coli* ($p < 0.05$) was lower in the IVF cycles. No significant difference was found in the embryo development rate of groups (normozoospermia, asthenozoospermia, and oligoasthenoteratozoospermia, $p > 0.05$), except for the teratozoospermia samples infected with *S. saprophyticus* ($p < 0.05$). In contrast, the clinical pregnancy rate was lower for the semen samples infected with *E. coli* from ICSI/IVF cycles in each group. Therefore, in the normozoospermia group, the clinical pregnancy rate decreased in samples infected with *E. coli* during ICSI cycles ($p < 0.001$) and in both bacteriospermia groups during IVF cycles ($p < 0.001$). Bacteriospermia significantly influenced the clinical pregnancy rate in the asthenozoospermia (*S. saprophyticus* and *E. coli*, $p < 0.001$) and teratozoospermia (*E. coli*, $p < 0.05$) groups (Table 2).

Discussion

The present study investigated the effects of the most common bacteria (*S. saprophyticus* and *E. coli*) on sperm parameters and assisted reproductive outcomes in infertile couples with male factor infertility in northern Iran. Teratozoospermia was the most common abnormality in semen samples infected with *S. saprophyticus*, as manifested by sperm deprotonation, sperm chromatin condensation, and impaired viability. Other semen samples were also influenced by infection with *S. saprophyticus* and *E. coli*. Therefore, semen samples infected with *S. saprophyticus* showed significant differences in terms of sperm quality such as sperm deprotonation (normozoospermia and asthenozoospermia), abnormalities in sperm chromatin condensation (oligoasthenoteratozoospermia), and impaired viability (asthenozoospermia). In addition, semen samples infected with *E. coli* showed significant differences in terms of sperm quality such as abnormal chromatin condensation (normozoospermia and oligoasthenoteratozoospermia), spontaneous acrosome reaction (oligoasthenoteratozoospermia), and impaired viability (normozoospermia, asthenozoospermia, teratozoospermia, and oligoasthenoteratozo-

Table 2. The effect of bacteriospermia on assisted reproductive outcomes in men with different infertility factors

Variable		Fertilization rate (%)	Cleavage rate (%)	Clinical pregnancy rate P/ET (%)	Live birth LB/IE (%)
Normozoospermia					
<i>Staphylococcus saprophyticus</i>	ICSI	72.2	83.4	8/14 (57.1)	4/8 (50.0)
	IVF	56.2 ^{b)}	78.1	8/20 (40.0) ^{b)}	4/8 (50.0)
<i>Escherichia coli</i>	ICSI	67.2	85.9	8/17 (47.1) ^{b)}	3/8 (37.5) ^{a)}
	IVF	67.3 ^{a)}	81.6	6/13 (46.2) ^{b)}	3/6 (50.0)
Nonbacteriospermia	ICSI	74.0	83.5	17/26 (65.4)	10/17 (58.8)
	IVF	79.6	85.5	22/30 (73.3)	13/22 (59.1)
Asthenozoospermia					
<i>Staphylococcus saprophyticus</i>	ICSI	68.4	82.1	7/15 (46.7) ^{b)}	3/7 (42.8)
<i>Escherichia coli</i>	ICSI	69.1	83.4	12/25 (48.0) ^{b)}	4/12 (33.3) ^{a)}
Nonbacteriospermia	ICSI	72.7	89.3	33/49 (67.3)	18/33 (54.5)
Teratozoospermia					
<i>Staphylococcus saprophyticus</i>	ICSI	55.9	72.8 ^{a)}	6/13 (46.2)	2/6 (33.3) ^{a)}
<i>Escherichia coli</i>	ICSI	57.1	78.2	9/28 (32.1) ^{a)}	3/9 (33.3) ^{a)}
Nonbacteriospermia	ICSI	59.1	85.0	19/41 (46.3)	10/19 (52.6)
Oligoasthenoteratozoospermia					
<i>Staphylococcus saprophyticus</i>	ICSI	55.7	74.1	5/14 (35.7)	1/5 (20.0) ^{a)}
<i>Escherichia coli</i>	ICSI	60.3	79.8	8/21 (38.1)	2/8 (25.0)
Nonbacteriospermia	ICSI	58.1	76.2	10/26 (38.5)	3/10 (30.0)

Values are presented as number (%) unless otherwise indicated. The semen samples infected with *S. saprophyticus* showed significant differences in terms of assisted reproductive outcomes such as the fertilization rate (normozoospermia), embryo cleavage rate (teratozoospermia), clinical pregnancy rate (normozoospermia and asthenozoospermia), and live birth rate (oligozoospermia and oligoasthenoteratozoospermia) in comparison to the control group (non-bacteriospermia). In addition, the semen samples infected with *E. coli* showed significant differences in the fertilization rate (normozoospermia), pregnancy rate (normozoospermia, asthenozoospermia, and teratozoospermia), and live birth rate (asthenozoospermia and teratozoospermia) in comparison to the control group (non-bacteriospermia).

P, positive cycle; ET, embryo transfer; LB, live birth; IE, implanted embryo; ICSI, intracytoplasmic sperm injection; IVF, *in vitro* fertilization.

^{a)} $p < 0.05$; ^{b)} $p < 0.001$.

ospermia). The poor sperm quality in the semen samples infected with bacteria may have affected the assisted reproductive outcomes. Furthermore, tight adhesion of *S. saprophyticus* to sperm after density gradient centrifugation was shown, which is reflected in the IVF outcomes (fertilization and clinical pregnancy rates). Although many pathogenic bacteria have been recognized as potential causes of infertility, the resultant clinical outcomes in IVF/ICSI procedures have not yet been elucidated [9]. This study showed that *S. saprophyticus* can interact with spermatozoa, causing reduced acrosomal function, abnormal chromatin condensation (AB staining), sperm deprotonation (CMA3 staining), and tight adhesion to different parts of the tail. Reduced motility was seen in this group. These changes were reflected in IVF outcomes, as the fertilization rate of IVF cycles was lower than that of ICSI cycles for the normozoospermia samples.

The relationship between bacteriospermia and fertilization potential remains unclear [10]. The impact of genital tract infections and bacterial contamination of semen on male fertility potential has been examined in many studies; nevertheless, the impact of bacteriospermia on fertility potential has yet to be elucidated, and there is not complete agreement on the detrimental effect of bacterial infections on sperm quality [10,11]. The most frequently isolated microorganisms in semen have been reported to be *U. urealyticum*, *E. coli*, *E. faecalis*, and *S. faecalis* [10], but few studies have investigated the role of *S. saprophyticus* in semen quality and male fertility potential. Therefore, the clinical significance of the presence of bacteria in the seminal fluid is still a matter of debate [12]. The present study demonstrates that *E. coli* and *S. saprophyticus* reduce sperm fertility potential and impair assisted reproductive outcomes.

Some researchers have reported that no semen parameters were significantly influenced in specimens with bacteriospermia [13-17]. In addition, Sanocka-Maciejewska et al. [18] observed no effect of bacteria isolated from the genitourinary tract on semen parameters in normozoospermia samples; nevertheless, a reduced antioxidant capacity of sperm was found in infertile patients. In contrast, other studies have reported that bacteriospermia was significantly associated with semen parameters such as sperm count, motility, and morphology [19]. Attachment between *E. coli* and sperm, morphological abnormalities, and sperm agglutination have been reported in some studies [20]. A direct negative effect of certain pathogens on the function of sperm and oocyte quality has been proven [21]. In some works, it has also been reported that sperm parameters such as sperm count, motility (both *in vitro* and *in vivo*), morphology, and viability decreased in the presence of *E. coli* [3,18,19], which is in line with our results. As the results of this study show, the quality and function of sperm decreased in the presence of bacteria, such as a declined protamination rate (positive CMA3 staining), a higher rate of premature acrosome reaction, reduced viability, and abnormal

chromatin condensation (AB staining). This negative correlation between sperm quality and bacterial infection may be due to increased reactive oxygen species (ROS) production associated with inflammatory processes in semen infections, as some studies demonstrated that enhanced ROS production led to impairment of sperm DNA and male fertility [9].

In the present study, a diminished quality of sperm infected with *S. saprophyticus* and *E. coli* was followed by a decrease in the clinical pregnancy rate during ICSI/IVF cycles. It can be inferred that bacterial infections of the semen affect the fertility potential of men, even in ART under *in vitro* conditions. Therefore, the existence of bacteria and interactions between bacteria and sperm may explain why the fertilization rate and clinical pregnancy rate were lower in the semen samples infected with some bacteria.

In conclusion, bacterial infections of the male urogenital tract have been correlated with subfertility and infertility. However, these effects on fertility potential are multifunctional and complex. One relevant factor is microbial infection of the semen. This study showed negative correlations between two prevalent bacteria in the north of Iran (*E. coli* and *S. saprophyticus*) and sperm quality and assisted reproductive outcomes such as reduced viability, premature acrosome reaction, a lower protamination rate using CMA3 staining, abnormal chromatin condensation using AB staining, adhesion to sperm, change of sperm function, and lower fertilization and clinical pregnancy rates. To the best of our knowledge, no studies have previously evaluated the presence and effect of these bacteria on sperm quality and fertility potential during ICSI/IVF cycles. Therefore, the results of this study can be useful for promoting assisted reproductive outcomes.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

The authors of this study acknowledge the cooperation of the co-workers at the IVF center of Alzahra Hospital (Rasht, Iran) and at the Guilan Microbiology Laboratory (Rasht, Iran).

ORCID

Fatemeh Ghasemian <https://orcid.org/0000-0001-9284-6545>
Shahin Esmaeilnezhad <https://orcid.org/0000-0001-6279-6209>
Mohammad Javad Mehdipour Moghaddam
<https://orcid.org/0000-0002-0392-065X>

Author contributions

Conceptualization: FG. Data curation: SE, MJMM. Formal analysis: FG. Methodology: SE. Project administration: FG, MJMM. Visualization & Writing—original draft: FG. Writing—review & editing: FG, MJMM.

References

- Pellati D, Mylonakis I, Bertoloni G, Fiore C, Andrisani A, Ambrosini G, et al. Genital tract infections and infertility. *Eur J Obstet Gynecol Reprod Biol* 2008;140:3–11.
- Fraczek M, Kurpisz M. Mechanisms of the harmful effects of bacterial semen infection on ejaculated human spermatozoa: potential inflammatory markers in semen. *Folia Histochem Cytobiol* 2015;53:201–17.
- Fraczek M, Wiland E, Piasecka M, Boksa M, Gaczarzewicz D, Szumala-Kakol A, et al. Fertilizing potential of ejaculated human spermatozoa during in vitro semen bacterial infection. *Fertil Steril* 2014;102:711–9.e1.
- World Health Organization. WHO laboratory manual for the examination and processing of human semen. 5th ed. Geneva: World Health Organization; 2010.
- Shekarriz M, Sharma RK, Thomas AJ Jr, Agarwal A. Positive myeloperoxidase staining (Endtz test) as an indicator of excessive reactive oxygen species formation in semen. *J Assist Reprod Genet* 1995;12:70–4.
- Vicari E, Mongioi A, Speciale A, Caccamo F, Calogero A, Gulizia S, et al. Enhancing detection of gonococcus in ejaculates of adult males using sperm dilution. *Arch Androl* 1986;16:19–23.
- Talbot P, Chacon RS. A triple-stain technique for evaluating normal acrosome reactions of human sperm. *J Exp Zool* 1981;215: 201–8.
- Franco JG Jr, Mauri AL, Petersen CG, Massaro FC, Silva LF, Felipe V, et al. Large nuclear vacuoles are indicative of abnormal chromatin packaging in human spermatozoa. *Int J Androl* 2012;35:46–51.
- Jedrzejczak P, Fraczek M, Szumala-Kakol A, Taszarek-Hauke G, Pawelczyk L, Kurpisz M. Consequences of semen inflammation and lipid peroxidation on fertilization capacity of spermatozoa in vitro conditions. *Int J Androl* 2005;28:275–83.
- Fraczek M, Hryhorowicz M, Gill K, Zarzycka M, Gaczarzewicz D, Jedrzejczak P, et al. The effect of bacteriospermia and leukocytospermia on conventional and nonconventional semen parameters in healthy young normozoospermic males. *J Reprod Immunol* 2016;118:18–27.
- Golshani M, Taheri S, Eslami G, Suleimani Rahbar AA, Fallah F, Goudarzi H. Genital tract infection in asymptomatic infertile men and its effect on semen quality. *Iran J Public Health* 2006;35:81–4.
- Ombelet W, Bosmans E, Janssen M, Cox A, Vlasselaer J, Gyselaers W, et al. Semen parameters in a fertile versus subfertile population: a need for change in the interpretation of semen testing. *Hum Reprod* 1997;12:987–93.
- McGowan MP, Burger HG, Baker HW, de Kretser DM, Kovacs G. The incidence of non-specific infection in the semen in fertile and sub-fertile males. *Int J Androl* 1981;4:657–62.
- Berger RE, Karp LE, Williamson RA, Koehler J, Moore DE, Holmes KK. The relationship of pyospermia and seminal fluid bacteriology to sperm function as reflected in the sperm penetration assay. *Fertil Steril* 1982;37:557–64.
- Huwe P, Diemer T, Ludwig M, Liu J, Schiefer HG, Weidner W. Influence of different uropathogenic microorganisms on human sperm motility parameters in an in vitro experiment. *Andrologia* 1998;30 Suppl 1:55–9.
- Cottell E, Harrison RF, McCaffrey M, Walsh T, Mallon E, Barry-Kinsel C. Are seminal fluid microorganisms of significance or merely contaminants? *Fertil Steril* 2000;74:465–70.
- Domes T, Lo KC, Grober ED, Mullen JB, Mazzulli T, Jarvi K. The incidence and effect of bacteriospermia and elevated seminal leukocytes on semen parameters. *Fertil Steril* 2012;97:1050–5.
- Sanocka-Maciejewska D, Ciupinska M, Kurpisz M. Bacterial infection and semen quality. *J Reprod Immunol* 2005;67:51–6.
- Isaiah IN, Nche BT, Nwagu IG, Nnanna II. Current studies on bacterospermia the leading cause of male infertility: a protege and potential threat towards mans extinction. *N Am J Med Sci* 2011; 3:562–4.
- Kaur K, Prabha V. Spermagglutinating *Escherichia coli* and its role in infertility: in vivo study. *Microb Pathog* 2014;69-70:33–8.
- Diemer T, Ludwig M, Huwe P, Hales DB, Weidner W. Influence of urogenital infection on sperm function. *Curr Opin Urol* 2000;10: 39–44.