\*Corresponding author: Sun Joo Park

Introduction

Prostate cancer is the most common cancer among men worldwide and a leading cause of cancer-related deaths in men in

Received: Oct 30, 2020 Revised: Nov 23, 2020 Accepted: Nov 27, 2020

Copyright © 2021 The Korean Society of Fisheries and Aquatic Science

Department of Chemistry, Pukyong National University, Busan 48513, Korea

Western countries (Quinn & Babb, 2002). Since the incidence rates of prostate cancer have recently increased dramatically in less-developed countries such as in Asia and Africa, which have traditionally had the lowest incidence rates (Baade et al., 2009),

Aspergillus fumigatus-derived demethoxyfumitremorgin C inhibits proliferation of PC3 human prostate cancer cells through p53/p21-dependent G1 arrest and apoptosis induction

Young-Sang Kim, Sun Joo Park\*

Department of Chemistry, Pukyong National University, Busan 48513, Korea

#### Abstract

Human prostate cancer is the second most frequently diagnosed cancer worldwide, and its incidence rate continues to increase. Advanced prostate cancer is more difficult to treat than early forms due to its chemotherapy resistance. There is need for more effective agents that can inhibit the progression of advanced prostate cancer. Demethoxyfumitremorgin C (DMFTC) was isolated from the fermentation extract of the marine fungus *Aspergillus fumigatus*. Antiproliferative activity of DMFTC against human prostate cancer PC3 cells was examined through cell cycle analysis by flow cytometry, the fluorescent nuclear imaging analysis with propidium iodide (PI), and proteins expression related to cell cycle arrest and apoptosis induction. It activated the tumor suppressor p53 and the Cdk inhibitor p21, which regulate the cell progression into the G1 phase. Additionally, PI-positive late apoptotic non-viable cells were increased and the expression levels of the G1-positive downstream regulators cyclin D, cyclin E, Cdk2, and Cdk4 were decreased by DMFTC treatment. These results suggest that DMFTC induces G1 arrest and apoptosis induction through regulation of p53/p21-dependent cyclin-Cdk complexes, and it may be a useful therapeutic agent for the treatment of human advanced prostate cancer.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-

nc/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Keywords: Demethoxyfumitremorgin C, Aspergillus fumigates, Prostate cancer, G1 cell cycle arrest, Cell death

Tel: +82-51-629-5598, Fax: +82-51-629-5583, E-mail: parksj@pknu.ac.kr, ORCID: https://orcid.org/0000-0002-3458-8418



RESEARCH ARTICLE Fish Aquat Sci. 2021;24(1):1-9 https://doi.org/10.47853/FAS.2021.e1



the effective treatment of prostate cancer has gained increased attention. Since prostate cancer usually develops slower than other cancers and has a long latency period of more than 10 vears before developing into a symptomatic and clinical disease, the incidence of prostate cancer may be reduced by blocking the multi-step process involving the progression from small and low-grade lesions to large and high-grade carcinomas (Lee et al., 2008). However, advanced prostate cancer is more difficult to treat than early prostate cancer because the cancer cells do not respond to most cytotoxic chemotherapeutic agents. Therefore, there is a need for a more effective therapy for advanced prostate cancer. Prostate cancer has been known to be associated with the abnormal growth of cells in the prostate gland (Kyprianou et al., 1996). Many studies have focused on various signaling pathways that regulate the growth and proliferation of prostate cancer cells and on the development of the related chemoprevention agents (Gupta et al., 2003).

Recently, interest in bioactive ingredients from marine fungi has grown because some marine fungal-derived compounds have various biological activities, including control of cell growth (Swathi et al., 2013). Specifically, marine fungi Aspergillus fumigatus has been reported to generate many structurally and biologically diverse secondary metabolites (Finefield et al., 2012), and their derivatives have been mainly studied for their potential use in the treatment of microsporidiosis (Molina et al., 2002) and amebiasis (Killough et al., 1952) and their antiangiogenic properties (Ben-Ami et al., 2009). However, there were few reports on the anticancer activity of compounds from A. fumigatus against cancer. Recently, we reported the isolation of demethoxyfumitremorgin C (DMFTC) from the ethyl acetate (EtOAc) extract of A. fumigatus strain MFS150 and the compound inhibits growth of human prostate cancer PC3 cells (Kim et al., 2017); however, the detailed data regarding to cell cycle arrest related to apoptosis has never been examined. In the present study, we revaluated the antiproliferative activity of DMFTC in human prostate cancer PC3 cells and investigated its functional mechanism on cell cycle progression. The results indicated that this compound induces G1 phase arrest during cell cycle progression and apoptotic cell death.

# **Materials and Methods**

### **Materials and chemicals**

The PC3 human prostate cancer cell line was obtained from the Korean Cell Line Bank (KCLB, Seoul, Korea). RPMI 1640 medium was obtained from Lonza (Walkersville, MD, USA), and fetal bovine serum (FBS) and penicillin/streptomycin were bought from Gibco BRL, Life Technology (Carlsbad, CA, USA). All antibodies used for the western blot analysis were from Santa Cruz Biotechnology (Santa Cruz, CA, USA). 3-(4,5-Dimethyl-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reagent was obtained from Sigma Chemical (St. Louis, MO, USA). Other chemicals and reagents were of analytical grade and commercially available.

#### **Extraction and isolation**

DMFTC was isolated by the previous method described by Li et al. (2014) with slight modifications. We isolated A. fumigatus MFS-150 from surface of marine green algae, collected at 2009 at Seosaeng-myeon, Ulsan, in the Republic of Korea. The fungal strain was cultured at 25°C and stored in 20% glycerol with YPG medium (1% yeast extract, 1% peptone, 10% glucose, 4% agar, 50% seawater, and 50% distilled water). The fungus was cultured (30 L) for 30 days at 25 °C and pH 7.6 in YPG medium. The culture broth was extracted (2.3 g) with EtOAc (1:1.5 v/v, 1:1 v/v, broth-EtOAc) twice and was fractionated by silica gel flash chromatography [n-hexane-EtOAc 100%-0% and CH<sub>2</sub>Cl<sub>2</sub>-MeOH 1:1] to yield several fractions. The major fraction (1,251.4 mg) was further purified using octadecyl functionalized silica gel (ODS; H<sub>2</sub>O-MeOH, 100%-100%) with 1 mL/min flow rate and Sephadex LH-20 chromatography (H<sub>2</sub>O-MeOH, 100%-100%). Finally, we performed high-performance liquid chromatography (HPLC) (YMC ODS-A column; 250  $\times \times$  10 mm l.D, S-5 µm, 12 nm, methyl alcohol [MeOH]) to isolate DMFTC (final 7.2 mg). The structure and molecular formula of DMFTC, compared to previous data (Kim et al., 2017), were ascertained from <sup>1</sup>H, <sup>13</sup>C NMR (nuclear magnetic resonance), and low resolution electron impact mass spectrometry (LREI-MS) data. LREIMS m/z: 349 [M]<sup>+</sup> (C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub>).

#### Cell culture and anti-proliferation assay

PC3 human prostate cancer cells were grown in 10% FBS-containing RPMI 1640 media and incubated at 37 °C in 5% CO<sub>2</sub>. In the early less than passages 5, the cell growth inhibition and cell death effects of the compounds from *A. fumigatus* on PC3 cells were analyzed using MTT [3-(4, 5-dimethyl-2-yl)-2, 5-diphenyltetrazolium bromide] assays. To examine the anti-proliferative effect of DMFTC, the cells seeded were incubated for 24 h in 96-well plates at a concentration of  $5 \times 10^4$  cells/mL. The cells were treated with different concentrations of the compounds (0, 25, 50, and 100  $\mu$ M). At the end of the incubation time, 100  $\mu$ L MTT solution (1 mg/mL in DMSO) was added to well and incubated for another 3 h. The formed formazan salt was dissolved in 100  $\mu$ L DMSO, and the absorbance was measured at 550 nm (GENiosr microplate reader, Tecan, Grödig, Austria). The data were described as means ± SD of at least three independent assays.

#### Cell cycle analysis by flow cytometry

We performed cell cycle analysis of propidium iodide (PI)stained nuclei using CellQuest software on a FACS Calibur flow cytometer (Becton-Dickinson Biosciences, Heidelberg, Germany). PC3 cells were cultured in 6-well plates at a density of  $5 \times 10^4$  cells/mL and then were treated with DMFTC after 24 h for 72 h at various concentrations (0, 25, 50, and 100 µM). The cells were harvested at each incubation time, pelleted by centrifugation for 5 min at 500 × g, and resuspended in the chilled PBS. The cells were fixed with 70% ice-cold EtOH and stored at 4°C. Prior to analysis, cells were washed again with PBS, resuspended in 1 mL of a cold PI solution with 50 µg/mL RNase A, and further incubated on ice for 20 min in the dark. We analyzed the data using single histogram statistics (Yin et al., 2014).

#### **Cell death analysis**

Cell death was assessed by fluorescent microscopy after staining cells with Hoechst 33258 and PI as expressed by Naito (1998). Cells were grown in 24-well plates at a density of  $5 \times 10^4$  cells/ mL and treated with different concentrations of DMFTC after 24 h, and further incubated for 72 h. The cells were washed with PBS twice and fixed with a 4% formaldehyde solution in PBS for 1 h at room temperature (RT). The fixed cells were washed with PBS three times and were stained with 1 µg/mL fluorescent DNA-binding dye Hoechst 33258 and incubated for 1 h at RT to reveal nuclear condensation/aggregation. Cells were treated with different concentrations of DMFTC for 72 h for PI staining. The DNA-specific fluorescent dye PI 5 µg/mL was added to the culture medium. The plates were additionally incubated for 30 min at 37 °C. The Hoechst 33258- and PI-stained cells were visualized and photographed under a fluorescence microscope (Motic AE31, MHG-100B, Jed Pella, Redding, CA, USA; DM3000, Leica, Wetzlar, Germany).

#### Western blot analysis

Western blotting was performed according to standard pro-

cedures. Cells were cultured at a density of  $5 \times 10^4$  cells/mL in 6-well plates. The cells were treated with different concentrations of DMFTC for 72 h after incubation for 24 h. Cells were lysed in RIPA buffer at 4°C for 30 min. Total proteins were obtained, and 100 µg/mL of protein was separated using 10% SDS-PAGE and transferred onto a nitrocellulose membrane (Amersham Pharmacia Biotech, Little Chalfont, UK). The membrane was blocked using TBS-T buffer containing 5% skim milk and 0.1% Tween-20 for 1.5 h at RT. The blots were incubated for 1 h with suitable antibodies at 25°C. The respective proteins were confirmed with a chemiluminescent ECL assay kit (Amersham Pharmacia) according to the manufacturer's instructions. The western blot bands were visualized using a LAS-3000 system and detected using MultiGauge V 3.0 software (Fujifilm Life Science, Tokyo, Japan).

#### **Statistical analysis**

We analyzed the data using the Instat statistics program (Graph-Pad Software, San Diego, CA, USA). Statistical comparisons were performed using one-way analysis of variance (ANOVA) with the Bonferroni multiple comparisons test.

## Results

# Isolation of secondary metabolites from the marine fungus *A. fumigatus*

To identify compounds with cytotoxic effect against human advanced prostate cancer cells from marine fungus *A. fumigatus*, DMFTC was purified from a pool of secondary metabolites. The chemical structure and molecular formula of DMFTC ( $C_{21}H_{23}N_3O_2$ ) were ascertained from <sup>1</sup>H, <sup>13</sup>C NMR, and LREI-MS spectroscopic data together with comparison to the data published previously (Cui et al., 1996, 1997; Kim et al., 2017). The compound was identified as DMFTC, as illustrated in Fig. 1 and Table 1.

# Anti-proliferation effect of DMFTC in PC3 human prostate cancer cells

The anti-proliferation activity of the isolated DMFTC was confirmed in PC3 human prostate cancer cells. PC3 cells were treated with various concentrations (0, 25, 50, and 100  $\mu$ M) of DMFTC for 24, 48, and 72 h. Cell viability at each condition was measured by MTT assay. Fig. 2 shows that DMFTC delayed the growth of viable cells and suppressed the growth of PC3 cells in a dose- and time-dependent manner. The IC<sub>50</sub> value of



**Fig. 1. Chemical structure of DMFTC isolated from the marine-derived fungus** *Aspergillus fumigatus*. DMFTC, demethoxyfumitremorgin C.

Table 1. <sup>1</sup> H and <sup>13</sup> C NMR spectral data for demethoxyfumitre	morgin
Cin DMSO- $d_6^{10}$	_

Position	$\delta_{H}$	δ <sub>c</sub>
1 (NH)	11.1 (1H, br, s)	
2		132.6 (s)
3	5.93 (1H, d, <i>J</i> = 9.5)	49.9 ( <i>d</i> )
4		
5		169.1 (s)
6	4.24 (1H, dd, <i>J</i> = 8.1)	58.4 ( <i>d</i> )
7	2.04, 2.22 (2H, m)	27.8 ( <i>t</i> )
8	1.86 (2H, m)	22.7 ( <i>t</i> )
9	3.45 (2H, m)	44.8 ( <i>t</i> )
10		
11		165.4 (s)
12	4.28 (1H, m)	56.1 ( <i>d</i> )
13	2.86, 3.34 (2H, dd, <i>J</i> = 11.7)	21.3 ( <i>t</i> )
14		104.9 (s)
15		125.9 (s)
16	7.52 (1H, d, <i>J</i> = 7.7)	117.8 ( <i>d</i> )
17	6.98 (1H, d, <i>J</i> = 7.3)	118.7 ( <i>d</i> )
18	7.05 (1H, d, <i>J</i> = 7.0)	120.9 ( <i>d</i> )
19	7.32 (1H, d, <i>J</i> = 8.1)	111.4 ( <i>d</i> )
20		136.1 (s)
21	4.85 (1H, d, <i>J</i> = 9.5)	124.6 ( <i>d</i> )
22		134.3 (s)
23	1.57 (3H, br, s)	25.4 ( <i>t</i> )
24		17.9 ( <i>t</i> )

<sup>1)</sup>Recorded at 400 MHz for <sup>1</sup>H and 100 MHz for <sup>13</sup>C and <sup>13</sup>C DEPT. NMR, nuclear magnetic resonance.



**Fig. 2. Anti-proliferation effect of DMFTC in PC3 human prostate cancer cells.** DMFTC suppresses the proliferation of PC3 human prostate cancer cells. PC3 cells were treated with various concentrations (0, 25, 50, and 100  $\mu$ M) of DMFTC for different time intervals (24, 48, and 72 h). After the incubation, 100  $\mu$ L of MTT solution and 100  $\mu$ L of DMSO was subsequently added to each well and to solubilize the formazan salt. The absorbance was measured at 550 nm. Each value is represented as the mean  $\pm$  SD of triplicate independent experiments. <sup>\*\*\*</sup>p<0.001 compared to the control. DMFTC, demethoxyfumitremorgin C; MTT, 3-(4,5-Dimethyl-2-yl)-2,5diphenyltetrazolium bromide.

DMFTC was estimated to be  $73.35 \,\mu\text{M}$  at  $72 \,\text{h}$ .

#### DMFTC causes cell cycle arrest at G1 phase

To explore the mechanism responsible for the inhibitory effect of DMFTC on cell growth, we first examined whether DMFTC regulates the cell cycle progression of PC3 cells. PC3 cells were treated with various concentrations (0, 25, 50, and 100 µM) of DMFTC and stained with PI, and the cell cycle progression then was analyzed by flow cytometry. To identify the location of the major cell cycle stage where DMFTC inhibits the cell cycle progression in PC3 cells, the dead cells and debris were excluded. As shown in Fig. 3, the cell cycle progression was arrested in G1 phase when cells were treated with DMFTC. Compared with vehicle-treated control cells, DMFTC treatment increased the number of G1 phase cells and decreased the number of S phase cells in a concentration-dependent manner. In the presence of 100 µM, 70.44% of cells remained in G0/G1 phase compared with 59.77% of cells in the vehicle-treated control group. This result suggests that DMFTC decreases cell cycle progression through cell cycle arrest at G1 phase.



**Fig. 3. Cell cycle arrest at G1 phase by DMFTC.** DMFTC affects cell cycle progression in PC3 cells. PC3 cells were treated with various concentrations (0, 25, 50, and 100 μM) of D DMFTC for 72 h. Cells were fixed in ethanol and then stained with propidium iodide. Cells were analyzed using flow cytometry with single histogram statistics. DMFTC, demethoxyfumitremorgin C.

#### Apoptotic cell death induced by DMFTC

We next examined whether the inhibition of PC3 cell growth by DMFTC is related to apoptotic cell death. Hoechst 33258 and PI were used together for the fluorescence imaging analysis of the apoptotic cells. PI crosses the plasma membrane of cells in the later stages of apoptosis or that are already dead. Fig. 4 shows that DMFTC induces nuclear margination and obvious morphological changes. Late apoptotic cells with fragmented chromatin were also observed. The number and intensity of PI-positive non-viable cells also increased with exposure to DMFTC in a dose-dependent manner. This result indicates that DMFTC induces apoptotic cell death in PC3 cells.

# Effect of DMFTC on p53-dependent cell cycle regulatory protein levels

We also analyzed the effect of DMFTC on the expression of cell-cycle regulatory proteins that could be involved in G1 arrest and apoptosis. The tumor suppressor p53 functions an important role in the cell cycle checkpoint system responsible for progression from G1 into S phase (Livingstone et al., 1996). Many chemotherapeutic drugs used to treat cancer increase the level of p53 and activate the protein, leading to G1 phase cell cycle arrest and apoptosis (Bykov et al., 2002; Clarke et al., 1993; Di et al., 1994). The p53-dependent G1 arrest is associated with the activation of p21 and the subsequent inhibition of G1 cyclin-Cdk complexes, such as cyclin D-Cdk4/CdK6 and cyclin



**Fig. 4. Apoptotic cell death induced by DMFTC.** For nuclei staining, PC3 cells were cultured in 24-well plates for 24 h. After incubation, different concentrations of DMFTC (0, 25, 50, and 100  $\mu$ M) were added to the wells followed by incubation for 72 h. The cells were then fixed with 3% formaldehyde for 20 min at RT and then stained with Hoechst 33258 or PI for another 30 min. Finally, the nuclei of the cells were observed under an inverted fluorescent microscope (40 ×). DMFTC induces significant apoptotic morphological changes. The number of PI-positive non-viable cells and their intensities were increased by DMFTC in a dose-dependent manner. DMFTC, demethoxyfumitremorgin C; RT, room temperature.

E-Cdk2, leading to G1 cell cycle arrest and apoptosis induction (Sherr, 1996). Therefore, we examined whether the p53 and p21 tumor suppressor factors and the downstream cell cycle-related proteins are involved in the anti-proliferative effect of DMFTC. The treatment of PC3 cells with DMFTC induced the activation of p53 and p21. DMFTC treatment increased relative ratio of the intensity of phospho-p53 or phosphor-p21 to total p53 or p21 protein on dose-dependent manner.

The levels of cyclin D, cyclin E, Cdk2, and Cdk4 decreased following DMFTC treatment (Fig. 5). These results indicate that *A. fumigatus*-derived DMFTC reduces the proliferation of PC3 human prostate cancer cells through p53/p21-dependent G1 cell cycle arrest and the induction of apoptosis.

# Discussion

The most effective systemic therapy for prostate cancer is an-

drogen deprivation or the removal or blocking of the male hormone testosterone because androgen-dependent testosterone is essential for the growth and survival of prostate cells, whereas androgen deprivation causes the apoptosis of prostate cells (Hussain et al., 2013). However, some men do still develop prostate cancer. Aggressive genetic alterations cause cells to multiply, and these cells then eventually disrupt the internal structure of the prostate by breaking out of the gland and invading the stromal tissue and extending outside the prostate, acquiring advanced metastatic properties. However, in such cases of advanced prostate cancer, unfortunately, the cancer cells become resistant to apoptosis and do not respond to cytotoxic chemotherapeutic agents (Pilat et al., 1998). Therefore, agents that inhibit cell growth and induce apoptotic death in the hormone-refractory prostate cancer cells would be useful for the prevention and treatment of advanced prostate cancer. In the present study, we showed that DMFTC derived from the marine fungi A.



**Fig. 5. Effect of DMFTC on the p53/p21-dependent cell cycle regulatory proteins.** Western blotting analysis of phosphorylated p53 and p21 (A) and cyclin D, E, Cdk2, and Cdk4 (B) from PC3 cells treated with different concentrations (0, 25, 50, and 100  $\mu$ M) of DMFTC. Each value was represented as the mean ± SD of triplicate experiment. <sup>\*\*\*</sup>*p*<0.001 as compared to the control. DMFTC, demethoxyfumitremorgin C.

*fumigatus* delays cell cycle progression and induces apoptotic cell death in PC3 advanced human prostate cancer cells. This result supports well our previous data, in which we indicated that DMFTC causes apoptosis as determined by Annexin V/PI staining and decreases mitochondrial membrane potential (Kim et al., 2017). We have also reported that DMFTC inhibits preferentially the proliferation of PC3 cells rather than other human normal and cancer cells lines, including MRC5, Hela, MCF-7, and HepG2 (Kim et al., 2017). Therefore, all results suggest that DMFTC has the potential to be a significant therapeutic mechanism for the prevention and treatment of advanced prostate cancer. DMFTC regulates cell growth by p53/p21-dependent G1 phase cell cycle arrest and apoptotic pathways. In fact, the inhibition of cell cycle progression is an appropriate target for cancer treatment. Many drugs that target the cell cycle have

been tested in clinical trials (Cuzick et al., 2011). When a cell needs to stop the cell cycle in G1, kinases such as Chk2 and Chk1 phosphorylate the tumor suppressor p53, which results in the transcriptional activation of p21, an inhibitor of the G1-to-S promoting complex cyclin E-Cdk2 and/or cyclin D-Cdk4 complex, thus keeping the cell from transitioning to S phase (Sherr, 1996). p53 also plays a vital role in cancer cell apoptosis through the regulation of proteins such as the Bax (a cell death inducer) and Bcl-2 (a cell death inhibitor) families (Cory et al., 2003; Karnak & Xu, 2010). Our data show that the inhibition of cell growth and induction of apoptosis by DMFTC in PC3 cells is most likely associated with the role of p53/p21-dependent cyclin-Cdk complexes. Therefore, these results suggest that DM-FTC obtained from the fermentation extract of the marine fungus *A. fumigatus* MFS-150 may supply new therapeutic insights

to prevent and treat advanced prostate cancer that is resistant to conventional treatments.

Cui et al. (1996) previously reported that DMFTC isolated from the A. fumigatus strain BM939 exhibits an inhibitory activity on the M phase cell cycle progression of mouse mammary carcinoma tsFT210 cells. The tsFT210 cells are a temperature-sensitive p34<sup>cdc2</sup> mutant cell line. In their study, they first synchronized the cells in G2 phase and then released them to screen for G2/M phase inhibitors that trigger M phase-promoting factors related to cdc2 kinase and cyclin B for mitosis. Therefore, the effect of DMFTC on the M phase arrest of tsFT210 cells may be stronger than its effect on PC3 cells in our study. In the assays of randomly cultured PC3 cells in present study, we did not assess M phase cell cycle arrest by DMFTC. This difference in the effect of DMFTC on cell cycle arrest might be due to the different molecular determinants between PC3 human advanced prostate cancer cells and tsFT210 mouse mammary carcinoma cells. More studies may be needed to confirm the anti-proliferation effect of DMFTC in various cancer cells.

# Conclusion

In the present study, we found that DMFTC from *A. fumigatus* MFS-150 inhibits PC3 cells growth through G1 phase cell cycle arrest and apoptosis induction. DMFTC activated the tumor suppressor p53 and the Cdk inhibitor p21, which regulate the cell progression into the G1 phase, whereas decreased the levels of the G1-positive downstream regulators cyclin D, cyclin E, Cdk2, and Cdk4. DMFTC could therefore be considered as a useful agent for treatment of human prostate cancer.

#### **Competing interests**

No potential conflict of interest relevant to this article was reported.

#### **Funding sources**

This work was supported by a Research Grant of Pukyong National University (2019).

#### Acknowledgements

Not applicable.

#### Availability of data and materials

All datasets generated and/or analyzed during the current study

are available from the corresponding author on reasonable request.

#### Ethics approval and consent to participate

This article does not require IRB/IACUC approval because there are no human and animal participants.

### References

- Baade PD, Youlden DR, Krnjacki LJ. International epidemiology of prostate cancer: geographical distribution and secular trends. Mol Nutr Food Res. 2009;53:171-84.
- Ben-Ami R, Lewis RE, Leventakos K, Kontoyiannis DP. Aspergillus fumigatus inhibits angiogenesis through the production of gliotoxin and other secondary metabolites. Blood. 2009;114:5393-9.
- Bykov VJ, Issaeva N, Shilov A, Hultcrantz M, Pugacheva E, Chumakov P, et al. Restoration of the tumor suppressor function to mutant p53 by a low-molecular-weight compound. Nat Med. 2002;8:282-8.
- Clarke AR, Purdie CA, Harrison DJ, Morris RG, Bird CC, Hooper ML, et al. Thymocyte apoptosis induced by p53-dependent and independent pathways. Nature. 1993;362:849-52.
- Cory S, Huang DC, Adams JM. The Bcl-2 family: roles in cell survival and oncogenesis. Oncogene. 2003;22:8590-607.
- Cui CB, Kakeya H, Okada G, Onose R, Osada H. Novel mammalian cell cycle inhibitors, tryprostatins A, B and other diketopiperazines produced by Aspergillus fumigatus. I. taxonomy, fermentation, isolation and biological properties. J Antibiot. 1996;49:527-33.
- Cui CB, Kakeya H, Osada H. Novel mammalian cell cycle inhibitors, cyclotroprostatins A–D, produced by Aspergillus fumigatus, which inhibit mammalian cell cycle at G2/M phase. Tetrahedron. 1997;53:59-72.
- Cuzick J, Swanson GP, Fisher G, Brothman AR, Berney DM, Reid JE, et al. Prognostic value of an RNA expression signature derived from cell cycle proliferation genes in patients with prostate cancer: a retrospective study. Lancet Oncol. 2011;12:245-55.
- Di Leonardo A, Linke SP, Clarkin K, Wahl GM. DNA damage triggers a prolonged p53-dependent G1 arrest and longterm induction of Cip1 in normal human fibroblasts. Genes Dev. 1994;8:2540-51.

Finefield JM, Frisvad JC, Sherman DH, Williams RM. Fungal

origins of the bicyclo[2,2,2]diazaoctane ring system of prenylated indole alkaloids. J Nat Prod. 2012;75:812-33.

- Gupta S, Hussain T, Mukhtar H. Molecular pathway for (–)-epigallocatechin-3-gallate-induced cell cycle arrest and apoptosis of human prostate carcinoma cells. Arch Biochem Biophys. 2003;410:177-85.
- Hussain M, Tangen CM, Berry DL, Higano CS, Crawford ED, Liu G, et al. Intermittent versus continuous androgen deprivation in prostate cancer. N Engl J Med. 2013;368:1314-25.
- Karnak D, Xu L. Chemosensitization of prostate cancer by modulating Bcl-2 family proteins. Curr Drug Targets. 2010;11:699-707.
- Killough JH, Magill GB, Smith RC. The treatment of amebiasis with fumagillin. Science. 1952;115:71-2.
- Kim YS, Kim SK, Park SJ. Apoptotic effect of demethoxyfumitremorgin C from marine fungus Aspergillus fumigatus on PC3 human prostate cancer cells. Chem Biol Interact. 2017;269:18-24.
- Kyprianou N, Tu H, Jacobs SC. Apoptotic versus proliferative activities in human benign prostatic hyperplasia. Hum Pathol. 1996;27:668-75.
- Lee DH, Szczepanski M, Lee YJ. Role of Bax in quercetin-induced apoptosis in human prostate cancer cells. Biochem Pharmacol. 2008;75:2345-55.
- Li YX, Kang KH, Kim HJ, Kim SK. In vitro induction of apoptosis by isosclerone from marine-derived fungus Aspergillus fumigatus. Bioorg Med Chem Lett. 2014;24:3923-7.
- Livingstone LR, White A, Sprouse J, Livanos E, Jacks T, Tlsty TD. Altered cell cycle arrest and gene amplification potential accompany loss of wild-type p53. Cell. 1992;70:923-35.
- Molina JM, Tourneur M, Sarfati C, Chevret S, de Gouvello A, Gobert JG, et al. Fumagillin treatment of intestinal microsporidiosis. N Engl J Med. 2002;346:1963-9.
- Naito Y, Saito K, Shiiba K, Ohuchi A, Saigenji K, Nagura H, et al. CD8+ T cells infiltrated within cancer cell nests as a prognostic factor in human colorectal cancer. Cancer Res. 1998;58:3491-4.
- Pilat MJ, Kamradt JM, Pienta KJ. Hormone resistance in prostate cancer. Cancer Metastasis Rev. 1998;17:373-81.
- Quinn M, Babb P. Patterns and trends in prostate cancer incidence, survival, prevalence and mortality. Part I: international comparisons. 2002;90:162-73.
- Sherr CJ. Cancer cell cycles. Science. 1996;274:1672-7.
- Swathi J, Narendra K, Sowjanya K, Satya AK. Marine fungal metabolites as a rich source of bioactive compounds. Afri-

can J Biochem Res. 2013;7:184-96.

Yin XY, Grove L, Datta NS, Long MW, Prochownik EV. C-myc overexpression and p53 loss cooperate to promote genomic instability. Oncogene. 1999;18:1177-84.