

FRACTIONAL DIFFERENTIATIONS AND INTEGRATIONS OF QUADRUPLE HYPERGEOMETRIC SERIES

MAGED G. BIN-SAAD, KOTTAKKARAN S. NISAR, AND JIHAD A. YOUNIS

ABSTRACT. The hypergeometric series of four variables are introduced and studied by Bin-Saad and Younis recently. In this line, we derive several fractional derivative formulas, integral representations and operational formulas for new quadruple hypergeometric series.

1. Introduction

Hypergeometric functions have been attracted the attention of many researchers due to their importance and applications in diverse areas of mathematical, physical, engineering and statistical sciences [2, 4, 5]. Multiple hypergeometric functions occur in various fields of pure and applied mathematics such as approximation theory, partition theory, representation theory, group theory, mirror symmetry, difference equations and mathematical physics etc. They possess important properties such as recurrence and explicit relations, summation formulae, symmetric and convolution identities, algebraic properties etc. In recent years, several multivariable hypergeometric functions and their properties have been considered by many authors [1, 3, 7–11, 15, 19, 23].

For our purpose, we begin by recalling some known functions and earlier works.

The Gaussian hypergeometric function defined by [22],

$$(1.1) \quad {}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}, \quad (|x| < 1),$$

where, $(a)_m$ is the Pochhammer symbol defined by

$$(a)_m = \frac{\Gamma(a+m)}{\Gamma(a)} = a(a+1) \cdots (a+m-1)$$

for $m \geq 1$, $(a)_0 = 1$, Γ being the well-known Gamma function.

Received May 31, 2020; Accepted November 16, 2020.

2010 *Mathematics Subject Classification.* 33C20, 33C65, 44A45.

Key words and phrases. Quadruple hypergeometric functions, Euler integrals, integrals of Laplace-type, fractional derivative formulas, inverse pairs of symbolic operators, triple hypergeometric functions.

The Appell hypergeometric functions F_1 , F_3 and F_4 in double variables [22], defined by

$$(1.2) \quad F_1(a, b, c; d; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_m(c)_n}{(d)_{m+n}} \frac{x^m y^n}{m! n!},$$

$$(1.3) \quad F_3(a, b, c, d; e; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_m(b)_n(c)_m(d)_n}{(e)_{m+n}} \frac{x^m y^n}{m! n!},$$

$$(1.4) \quad F_4(a, b; c, d; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_{m+n}}{(c)_m(d)_n} \frac{x^m y^n}{m! n!}.$$

The Lauricella functions of three variables $F_D^{(3)}$, F_N , F_R , F_S and F_T [16] are defined as:

$$(1.5) \quad \begin{aligned} & F_D^{(3)}(a, b_1, b_2, b_3; c; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a)_{m+n+p}(b_1)_m(b_2)_n(b_3)_p}{(c)_{m+n+p}} \frac{x^m y^n z^p}{m! n! p}, \end{aligned}$$

$$(1.6) \quad \begin{aligned} & F_N(a_1, a_2, a_3, b_1, b_2, b_1; c_1, c_2, c_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_m(a_2)_n(a_3)_p(b_1)_{m+p}(b_2)_n}{(c_1)_m(c_2)_{n+p}} \frac{x^m y^n z^p}{m! n! p}, \end{aligned}$$

$$(1.7) \quad \begin{aligned} & F_R(a_1, a_2, a_1, b_1, b_2, b_1; c_1, c_2, c_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{m+p}(a_2)_n(b_1)_{m+p}(b_2)_n}{(c_1)_m(c_2)_{n+p}} \frac{x^m y^n z^p}{m! n! p}, \end{aligned}$$

$$(1.8) \quad \begin{aligned} & F_S(a_1, a_2, a_2, b_1, b_2, b_3; c, c, c; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_m(a_2)_{n+p}(b_1)_m(b_2)_n(b_3)_p}{(c)_{m+n+p}} \frac{x^m y^n z^p}{m! n! p}, \end{aligned}$$

$$(1.9) \quad \begin{aligned} & F_T(a_1, a_2, a_2, b_1, b_2, b_1; c, c, c; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_m(a_2)_{n+p}(b_1)_{m+p}(b_2)_n}{(c)_{m+n+p}} \frac{x^m y^n z^p}{m! n! p}. \end{aligned}$$

Exton's functions X_6 , X_{13} , X_{18} , X_{19} and X_{20} are defined by [13]

$$(1.10) \quad \begin{aligned} & X_6(a, b, c; d, e; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a)_{2m+n+p}(b)_n(c)_p}{(d)_{m+n}(e)_p} \frac{x^m y^n z^p}{m! n! p!}, \end{aligned}$$

$$(1.11) \quad \begin{aligned} & X_{13}(a, b, c; d; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a)_{2m+n}(b)_{n+p}(c)_p}{(d)_{m+n+p}} \frac{x^m y^n z^p}{m! n! p!}, \end{aligned}$$

$$(1.12) \quad \begin{aligned} & X_{18}(a_1, a_2, a_3, a_4; c; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n}(a_2)_n(a_3)_p(a_4)_p}{(c)_{m+n+p}} \frac{x^m y^n z^p}{m! n! p!}, \end{aligned}$$

$$(1.13) \quad \begin{aligned} & X_{19}(a_1, a_2, a_3, a_4; c_1, c_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n}(a_2)_n(a_3)_p(a_4)_p}{(c_1)_m(c_2)_{n+p}} \frac{x^m y^n z^p}{m! n! p!}, \end{aligned}$$

$$(1.14) \quad \begin{aligned} & X_{20}(a_1, a_2, a_3, a_4; c_1, c_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n}(a_2)_n(a_3)_p(a_4)_p}{(c_1)_{m+p}(c_2)_n} \frac{x^m y^n z^p}{m! n! p!}. \end{aligned}$$

The Sharma's and Parihar's hypergeometric functions of four variables $F_{27}^{(4)}$, $F_{30}^{(4)}$, $F_{47}^{(4)}$ and $F_{48}^{(4)}$ defined by [21]

$$(1.15) \quad \begin{aligned} & F_{27}^{(4)}(a_1, a_1, a_2, a_2, a_3, a_3, a_4, a_4; c_1, c_2, c_1, c_3; x, y, z, u) \\ &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{m+n}(a_2)_{p+q}(a_3)_{m+n}(a_4)_{p+q}}{(c_1)_{m+p}(c_2)_n(c_3)_q} \frac{x^m y^n z^p u^q}{m! n! p! q!}, \end{aligned}$$

$$(1.16) \quad \begin{aligned} & F_{30}^{(4)}(a_1, a_1, a_2, a_2, a_3, a_3, a_4, a_5; c_1, c_2, c_1, c_3; x, y, z, u) \\ &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{m+n}(a_2)_{p+q}(a_3)_{m+n}(a_4)_p(a_5)_q}{(c_1)_{m+p}(c_2)_n(c_3)_q} \frac{x^m y^n z^p u^q}{m! n! p! q!}, \end{aligned}$$

$$(1.17) \quad \begin{aligned} & F_{47}^{(4)}(a_1, a_1, a_2, a_2, a_3, a_3, a_4, a_4; c_1, c_2, c_1, c_2; x, y, z, u) \\ &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{m+n}(a_2)_{p+q}(a_3)_{m+n}(a_4)_{p+q}}{(c_1)_{m+p}(c_2)_{n+q}} \frac{x^m y^n z^p u^q}{m! n! p! q!}, \end{aligned}$$

$$(1.18) \quad \begin{aligned} & F_{48}^{(4)}(a_1, a_1, a_2, a_2, a_3, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\ &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{m+n}(a_2)_{p+q}(a_3)_{m+n}(a_4)_p(a_5)_q}{(c_1)_{m+p}(c_2)_{n+q}} \frac{x^m y^n z^p u^q}{m! n! p! q!}. \end{aligned}$$

Very recently, Bin-Saad and Younis [6] defined thirty functions of four variables denoted these by $X_1^{(4)}, X_2^{(4)}, \dots, X_{30}^{(4)}$. Here, we give four of them as follows:

$$(1.19) \quad X_{19}^{(4)}(a_1, a_1, a_1, a_1, a_1, a_2, a_3, a_2; c_1, c_1, c_1, c_2; x, y, z, u)$$

$$\begin{aligned}
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n+p+q} (a_2)_{q+n} (a_3)_p}{(c_1)_{m+n+p} (c_2)_q} \frac{x^m y^n z^p u^q}{m! n! p! q!}, \\
 (1.20) \quad &X_{21}^{(4)}(a_1, a_1, a_2, a_1, a_1, a_2, a_3, a_2; c_1, c_1, c_2, c_3; x, y, z, u)
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n+q} (a_2)_{q+n+p} (a_3)_p}{(c_1)_{m+n} (c_2)_p (c_3)_q} \frac{x^m y^n z^p u^q}{m! n! p! q!}, \\
 (1.21) \quad &X_{23}^{(4)}(a_1, a_1, a_2, a_1, a_1, a_2, a_3, a_2; c_1, c_1, c_1, c_2; x, y, z, u)
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n+q} (a_2)_{q+n+p} (a_3)_p}{(c_1)_{m+n+p} (c_2)_q} \frac{x^m y^n z^p u^q}{m! n! p! q!}, \\
 (1.22) \quad &X_{28}^{(4)}(a_1, a_1, a_1, a_1, a_1, a_2, a_3, a_4; c_1, c_1, c_2, c_3; x, y, z, u) \\
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n+p+q} (a_2)_n (a_3)_p (a_4)_q}{(c_1)_{m+n} (c_2)_p (c_3)_q} \frac{x^m y^n z^p u^q}{m! n! p! q!}.
 \end{aligned}$$

Here, we consider a further quadruple hypergeometric functions as follows:

$$\begin{aligned}
 (1.23) \quad &X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{p+q} (a_3)_n (a_4)_p (a_5)_q}{(c_1)_{m+p} (c_2)_{n+q}} \frac{x^m y^n z^p u^q}{m! n! p! q!},
 \end{aligned}$$

$$\begin{aligned}
 (1.24) \quad &X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u) \\
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{p+q} (a_3)_n (a_4)_p (a_5)_q}{(c_1)_{m+p+q} (c_2)_n} \frac{x^m y^n z^p u^q}{m! n! p! q!},
 \end{aligned}$$

$$\begin{aligned}
 (1.25) \quad &X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{p+q} (a_3)_n (a_4)_p (a_5)_q}{(c_1)_{n+p+q} (c_2)_m} \frac{x^m y^n z^p u^q}{m! n! p! q!},
 \end{aligned}$$

$$\begin{aligned}
 (1.26) \quad &X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
 &= \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{p+q} (a_3)_n (a_4)_p (a_5)_q}{(c)_{m+n+p+q}} \frac{x^m y^n z^p u^q}{m! n! p! q!}.
 \end{aligned}$$

The aim of this paper is to investigate certain properties of four new hypergeometric functions of four variables $X_{80}^{(4)}$, $X_{81}^{(4)}$, $X_{82}^{(4)}$, $X_{83}^{(4)}$. In Section 2, some fractional derivative formulas involving the functions $X_i^{(4)}$ ($i = 80, 81, 82, 83$) are obtained. In Section 3, we consider several integral representations of Euler-type for our series involving triple and quadruple hypergeometric functions. Section 4 deals with the Laplace integrals for each new quadruple functions.

Section 5 gives some operational formulas with the help of certain inverse pairs of symbolic operators.

2. Fractional derivatives

Here, in this section, we consider some fractional derivative formulas for the quadruple series $X_{80}^{(4)}, X_{81}^{(4)}, X_{82}^{(4)}, X_{83}^{(4)}$. The fractional derivative operator D_w^k is defined by [18]

$$(2.1) \quad D_w^k w^a = \frac{\Gamma(a+1)}{\Gamma(a-k+1)} w^{a-k}, \quad (Re(a) > -1).$$

Theorem 2.1. *The following fractional derivative formulas holds:*

$$(2.2) \quad \begin{aligned} & D_w^{a_2-c} \left[w^{a_2-1} X_{80}^{(4)} (a_1, a_1, c, c, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; \right. \\ & \quad \left. x, y, wz, wu) \right] \\ &= \frac{\Gamma(a_2)}{\Gamma(c)} w^{c-1} X_{80}^{(4)} (a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; \\ & \quad x, y, wz, wu), \end{aligned}$$

$$(2.3) \quad \begin{aligned} & D_{w_1}^{a_2-c} D_{w_2}^{a_3-c'} \left[w_1^{a_2-1} w_2^{a_3-1} X_{80}^{(4)} (a_1, a_1, c, c, a_1, c', a_4, a_5; \right. \\ & \quad \left. c_1, c_2, c_1, c_2; x, w_2 y, w_1 z, w_1 u) \right] \\ &= \frac{\Gamma(a_2)\Gamma(a_3)}{\Gamma(c)\Gamma(c')} w_1^{c-1} w_2^{c'-1} X_{80}^{(4)} (a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; \\ & \quad c_1, c_2, c_1, c_2; x, w_2 y, w_1 z, w_1 u), \end{aligned}$$

$$(2.4) \quad \begin{aligned} & D_w^{a_1-c} D_y^{a_3-c'} \left[w^{a_1-1} y^{a_3-1} X_{80}^{(4)} (c, c, a_2, a_2, c, c', a_4, a_5; \right. \\ & \quad \left. c_1, c_2, c_1, c_2; w^2 x, wy, z, u) \right] \\ &= \frac{\Gamma(a_1)\Gamma(a_3)}{\Gamma(c)\Gamma(c')} w^{c-1} y^{c'-1} X_{80}^{(4)} (a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; \\ & \quad c_1, c_2, c_1, c_2; w^2 x, wy, z, u), \end{aligned}$$

$$(2.5) \quad \begin{aligned} & D_y^{a_3-c} D_z^{a_4-c'} D_u^{a_5-c''} \left[y^{a_3-1} z^{a_4-1} u^{a_5-1} X_{80}^{(4)} (a_1, a_1, a_2, a_2, a_1, \right. \\ & \quad \left. c, c', c''; c_1, c_2, c_1, c_2; x, y, z, u) \right] \\ &= \frac{\Gamma(a_3)\Gamma(a_4)\Gamma(a_5)}{\Gamma(c)\Gamma(c')\Gamma(c'')} y^{c-1} z^{c'-1} u^{c''-1} X_{80}^{(4)} (a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; \\ & \quad c_1, c_2, c_1, c_2; x, y, z, u). \end{aligned}$$

Proof. To prove (2.2). Making an appeal to the formula (2.1), we obtain

$$D_w^{a_2-c} \left[w^{a_2-1} X_{80}^{(4)} (a_1, a_1, c, c, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, wz, wu) \right]$$

$$= w^{c-1} \sum_{m,n,p,q=0}^{\infty} \frac{(a_1)_{2m+n} (c)_{p+q} (a_3)_n (a_4)_p (a_5)_q \Gamma(a_2 + p + q) x^m y^n (wz)^p (wu)^q}{(c_1)_{m+p} (c_2)_{n+q} \Gamma(c + p + q) m! n! p! q!}.$$

In view of (1.1) to the above equation, we get the required result. A similar argument will establish (2.3) to (2.5). The proofs of the following theorems would run parallel to the result (2.2). Therefore, we omit the details. \square

Theorem 2.2. *The following fractional derivative formulas holds:*

$$(2.6) \quad D_{w_1}^{a_1-c} D_{w_2}^{a_2-c'} D_{w_3}^{a_3-c''} \left[w_1^{a_1-1} w_2^{a_2-1} w_3^{a_3-1} X_{81}^{(4)} \left(c, c, c', c', c, c'', a_4, a_5; \right. \right. \\ \left. \left. c_1, c_2, c_1, c_1; w_1^2 x, w_1 w_3 y, w_2 z, w_2 u \right) \right] \\ = \frac{\Gamma(a_1) \Gamma(a_2) \Gamma(a_3)}{\Gamma(c) \Gamma(c') \Gamma(c'')} w_1^{c-1} w_2^{c'-1} w_3^{c''-1} X_{81}^{(4)} \left(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; \right. \\ \left. c_1, c_2, c_1, c_1; w_1^2 x, w_1 w_3 y, w_2 z, w_2 u \right),$$

$$(2.7) \quad D_y^{a_1-c} D_z^{a_2-c'} \left[y^{a_1-1} z^{a_2-1} X_{81}^{(4)} \left(c, c, c', c', c, a_3, a_4, a_5; \right. \right. \\ \left. \left. c_1, c_2, c_1, c_1; x, y, z, u \right) \right] \\ = \frac{\Gamma(a_1) \Gamma(a_2)}{\Gamma(c) \Gamma(c')} y^{c-1} z^{c'-1} X_{81}^{(4)} \left(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; \right. \\ \left. c_1, c_2, c_1, c_1; x, y, z, u \right),$$

$$(2.8) \quad D_x^{a_1-c} \left[x^{a_1-1} X_{81}^{(4)} \left(c, c, a_2, a_2, c, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x^2, xy, z, u \right) \right] \\ = \frac{\Gamma(a_1)}{\Gamma(c)} x^{c-1} X_{81}^{(4)} \left(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x^2, xy, z, u \right),$$

$$(2.9) \quad D_u^{a_5-c} \left[u^{a_5-1} X_{81}^{(4)} \left(a_1, a_1, a_2, a_2, a_1, a_3, a_4, c; c_1, c_2, c_1, c_1; x, y, z, u \right) \right] \\ = \frac{\Gamma(a_5)}{\Gamma(c)} u^{c-1} X_{81}^{(4)} \left(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u \right).$$

Corollary 2.1. *In formula (2.7), if we set $x = 0 = u$, we get the following result.*

$$(2.10) \quad D_y^{a_1-c} y^{a_1-1} {}_2F_1 \left(c, a_3; c_2; y \right) D_z^{a_2-c'} z^{a_2-1} {}_2F_1 \left(c', a_4; c_1; z \right) \\ = \frac{\Gamma(a_1) \Gamma(a_2)}{\Gamma(c) \Gamma(c')} y^{c-1} z^{c'-1} {}_2F_1 \left(a_1, a_3; c_2; y \right) {}_2F_1 \left(a_2, a_4; c_1; z \right).$$

Theorem 2.3. *The following fractional derivative formulas holds:*

$$(2.11) \quad D_{w_1}^{a_2-c} D_{w_2}^{a_3-c'} D_{w_3}^{a_4-c''} D_{w_4}^{a_5-c'''} \left[w_1^{a_2-1} w_2^{a_3-1} w_3^{a_4-1} w_4^{a_5-1} \right. \\ \left. \times X_{82}^{(4)} \left(a_1, a_1, c, c, a_1, c', c'', c'''; c_2, c_1, c_1, c_1; x, w_2 y, w_1 w_3 z, w_1 w_4 u \right) \right] \\ = \frac{\Gamma(a_2) \Gamma(a_3) \Gamma(a_4) \Gamma(a_5)}{\Gamma(c) \Gamma(c') \Gamma(c'') \Gamma(c''')} w_1^{c-1} w_2^{c'-1} w_3^{c''-1} w_4^{c'''-1}$$

$$\begin{aligned}
 & \times X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; \\
 & w_1^2 x, w_2 y, w_1 w_3 z, w_1 w_4 u), \\
 (2.12) \quad & D_x^{a_1-c} D_{w_1}^{a_2-c'} D_{w_2}^{a_3-c''} [x^{a_1-1} w_1^{a_2-1} w_2^{a_3-1} \\
 & \times X_{82}^{(4)}(c, c, c', c', c, c'', a_4, a_5; c_2, c_1, c_1, c_1; x^2, w_2 x y, w_1 z, w_1 u)] \\
 & = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)}{\Gamma(c)\Gamma(c')\Gamma(c'')} w_1^{c-1} w_2^{c'-1} w_3^{c''-1} X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, \\
 & a_5; c_2, c_1, c_1, c_1; x^2, w_2 x y, w_1 z, w_1 u),
 \end{aligned}$$

$$\begin{aligned}
 (2.13) \quad & D_w^{a_1-c} D_u^{a_2-c'} D_z^{a_4-c''} [w^{a_1-1} u^{a_2-1} z^{a_4-1} \\
 & \times X_{82}^{(4)}(c, c, c', c', c, a_3, c'', a_5; c_2, c_1, c_1, c_1; w^2 x, w y, z u, u)] \\
 & = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(a_4)}{\Gamma(c)\Gamma(c')\Gamma(c'')} w^{c-1} u^{c'-1} z^{c''-1} X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; \\
 & c_2, c_1, c_1, c_1; w^2 x, w y, z u, u),
 \end{aligned}$$

$$\begin{aligned}
 (2.14) \quad & D_x^{a_1-c} D_y^{a_3-c'} [x^{a_1-1} y^{a_3-1} \\
 & \times X_{82}^{(4)}(c, c, a_2, a_2, c, c', a_4, a_5; c_2, c_1, c_1, c_1; x^2, x y, z, u)] \\
 & = \frac{\Gamma(a_1)\Gamma(a_3)}{\Gamma(c)\Gamma(c')} x^{c-1} y^{c'-1} X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; \\
 & x^2, x y, z, u).
 \end{aligned}$$

Theorem 2.4. *The following fractional derivative formulas holds:*

$$\begin{aligned}
 (2.15) \quad & D_{w_1}^{a_1-c'} D_{w_2}^{a_2-c''} D_{w_3}^{a_3-c'''} D_{w_4}^{a_4-c''''} [w_1^{a_1-1} w_2^{a_2-1} w_3^{a_3-1} w_4^{a_4-1} \\
 & \times X_{83}^{(4)}(c', c', c'', c'', c', c''', c''''; a_5; c, c, c, c; w_1^2 x, w_1 w_3 y, w_2 w_4 z, w_2 u)] \\
 & = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_4)}{\Gamma(c')\Gamma(c'')\Gamma(c''')\Gamma(c'''')} w_1^{c'-1} w_2^{c''-1} w_3^{c'''-1} w_4^{c''''-1} \\
 & \times X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; w_1^2 x, w_1 w_3 y, w_2 w_4 z, w_2 u),
 \end{aligned}$$

$$\begin{aligned}
 (2.16) \quad & D_w^{a_5-c'} [w^{a_5-1} X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, c'; c, c, c, c; x, y, z, w u)] \\
 & = \frac{\Gamma(a_5)}{\Gamma(c')} w^{c'-1} X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, w u),
 \end{aligned}$$

$$\begin{aligned}
 (2.17) \quad & D_{w_1}^{a_1-c'} D_z^{a_2-c''} D_{w_2}^{a_3-c'''} D_u^{a_5-c''''} [w_1^{a_1-1} z^{a_2-1} w_2^{a_3-1} u^{a_5-1} \\
 & \times X_{83}^{(4)}(c', c', c'', c'', c', c''', a_4, c''''; c, c, c, c; w_1^2 x, w_1 w_2 y, z, u z)]
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_5)}{\Gamma(c')\Gamma(c'')\Gamma(c''')\Gamma(c''''')} w_1^{c'-1} z^{c''-1} w_2^{c'''-1} u^{c''''-1} \\
&\quad \times X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; w_1^2 x, w_1 w_2 y, z, uz), \\
(2.18) \quad &D_x^{a_1-c'} D_y^{a_3-c''} D_z^{a_4-c'''} D_u^{a_5-c''''} [x^{a_1-1} y^{a_3-1} z^{a_4-1} u^{a_5-1} \\
&\quad \times X_{83}^{(4)}(c', c', a_2, a_2, c', c'', c''', c''''; c, c, c, c; x^2, xy, z, u)] \\
&= \frac{\Gamma(a_1)\Gamma(a_3)\Gamma(a_4)\Gamma(a_5)}{\Gamma(c')\Gamma(c'')\Gamma(c''')\Gamma(c''''')} x^{c'-1} y^{c''-1} z^{c'''-1} u^{c''''-1} \\
&\quad \times X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x^2, xy, z, u).
\end{aligned}$$

3. Integrals of Euler-type

In this section, we establish several Euler-type integrals involving the quadruple functions $X_{80}^{(4)}, X_{81}^{(4)}, X_{82}^{(4)}, X_{83}^{(4)}$ asserted in the following theorems.

Theorem 3.1. *Each of the following integral representations holds:*

$$\begin{aligned}
(3.1) \quad &X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\
&= \frac{8M_1^{a_1} M_2^{a_4} M_3^{a_5} \Gamma(a_1 + a_3) \Gamma(a_4 + a_5) \Gamma(c_2)}{\Gamma(a_1) \Gamma(a_3) \Gamma(a_4) \Gamma(a_5) \Gamma(a) \Gamma(c_2 - a)} \\
&\quad \int_0^\infty \int_0^\infty \int_0^\infty \frac{\cosh \alpha (\sinh^2 \alpha)^{a_1 - \frac{1}{2}}}{(1 + M_1 \sinh^2 \alpha)^{a_1 + a_3}} \frac{\cosh \beta (\sinh^2 \beta)^{a_4 - \frac{1}{2}}}{(1 + M_2 \sinh^2 \beta)^{a_4 + a_5}} \\
&\quad \times \frac{\cosh \gamma (\sinh^2 \gamma)^{a - \frac{1}{2}}}{(1 + M_3 \sinh^2 \gamma)^{c_2}} F_{27}^{(4)}\left(\frac{a_1 + a_3}{2}, \frac{a_1 + a_3}{2}, a_2, a_2, \right. \\
&\quad \left. \frac{a_1 + a_3 + 1}{2}, \frac{a_1 + a_3 + 1}{2}, a_4 + a_5, a_4 + a_5; c_1, a, c_1, c_2 - a; \right. \\
&\quad \left. \frac{4M_1^2 x \sinh^4 \alpha}{(1 + M_1 \sinh^2 \alpha)^2} \frac{4M_1 M_3 y \sinh^2 \alpha \sinh^2 \gamma}{(1 + M_1 \sinh^2 \alpha)^2 (1 + M_3 \sinh^2 \gamma)}, \right. \\
&\quad \left. \frac{M_2 z \sinh^2 \beta}{(1 + M_2 \sinh^2 \beta)}, \frac{u}{(1 + M_2 \sinh^2 \beta) (1 + M_3 \sinh^2 \gamma)}\right) d\alpha d\beta d\gamma, \\
&\quad (Re(a_i) > 0, (i = 1, 3, 4, 5), Re(a) > 0, Re(c_2 - a) > 0, M_j > 0, j = (1, 2, 3)),
\end{aligned}$$

$$\begin{aligned}
(3.2) \quad &X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\
&= \frac{\Gamma(a_1 + a_3) \Gamma(c_2)}{\Gamma(a_1) \Gamma(a_3) \Gamma(a) \Gamma(c_2 - a)} \\
&\quad \int_0^\infty \int_0^\infty (e^{-\alpha})^a (1 - e^{-\alpha})^{c_2 - a - 1} (e^{-\beta})^{a_1} (1 - e^{-\beta})^{a_3 - 1} \\
&\quad \times F_{30}^{(4)}\left(\frac{a_1 + a_3}{2}, \frac{a_1 + a_3}{2}, a_2, a_2, \frac{a_1 + a_3 + 1}{2}, \frac{a_1 + a_3 + 1}{2}, a_4, a_5; c_1, a, c_1, c_2 - a; \right)
\end{aligned}$$

$$4xe^{-2\beta}, 4ye^{-(\alpha+\beta)} (1 - e^{-\beta}), z, u (1 - e^{-\alpha}) \Big) d\alpha d\beta,$$

$$(Re(a_1) > 0, Re(a_3) > 0, Re(a) > 0, Re(c_2 - a) > 0),$$

$$(3.3) \quad X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u)$$

$$= \frac{\Gamma(a_1 + a_3)\Gamma(a_4 + a_5)\Gamma(c_2)}{\Gamma(a_1)\Gamma(a_3)\Gamma(a_4)\Gamma(a_5)} \int_0^\infty \int_0^\infty \frac{\alpha^{a_3}}{(1 + \alpha)^{a_1+a_3}} \frac{\beta^{a_5}}{(1 + \beta)^{a_4+a_5}}$$

$$\times F_{47}^{(4)}\left(\frac{a_1+a_3}{2}, \frac{a_1+a_3}{2}, a_2, a_2, \frac{a_1+a_3+1}{2}, \frac{a_1+a_3+1}{2},$$

$$a_4 + a_5, a_4 + a_5; c_1, c_2, c_1, c_2; \frac{4x}{(1+\alpha)^2}, \frac{4\alpha y}{(1+\alpha)^2}, \frac{z}{(1+\beta)}, \frac{\beta u}{(1+\beta)}\right) d\alpha d\beta,$$

$$(Re(a_i) > 0, (i = 1, 3, 4, 5)),$$

$$(3.4) \quad X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u)$$

$$= \frac{\Gamma(a_1 + a_3)(1 + M)^{a_3}}{\Gamma(a_1)\Gamma(a_3)} \int_0^1 \frac{\alpha^{a_3-1}(1 - \alpha)^{a_1-1}}{(1 + M\alpha)^{a_1+a_3}}$$

$$\times F_{48}^{(4)}\left(\frac{a_1+a_3}{2}, \frac{a_1+a_3}{2}, a_2, a_2, \frac{a_1+a_3+1}{2}, \frac{a_1+a_3+1}{2}, a_4, a_5; c_1, c_2, c_1, c_2; ,$$

$$\frac{4(1-\alpha)^2 x}{(1+M\alpha)^2}, \frac{4(1+M)\alpha(1-\alpha)y}{(1+M\alpha)^2}, z, u\right) d\alpha,$$

$$(Re(a_1) > 0, Re(a_3) > 0, M > -1).$$

Corollary 3.1. *By considering equation (3.1) we obtain, when $x = 0$.*

$$(3.5) \quad F_N(a_4, a_1, a_5, a_2, a_3, a_2; c_1, c_2, c_2; z, y, u)$$

$$= \frac{8M_1^{a_1} M_2^{a_4} M_3^a \Gamma(a_1 + a_3)\Gamma(a_4 + a_5)\Gamma(c_2)}{\Gamma(a_1)\Gamma(a_3)\Gamma(a_4)\Gamma(a_5)\Gamma(a)\Gamma(c_2 - a)}$$

$$\int_0^\infty \int_0^\infty \int_0^\infty \frac{\cosh \alpha (\sinh^2 \alpha)^{a_1 - \frac{1}{2}}}{(1 + M_1 \sinh^2 \alpha)^{a_1 + a_3}}$$

$$\times \frac{\cosh \beta (\sinh^2 \beta)^{a_4 - \frac{1}{2}} \cosh \gamma (\sinh^2 \gamma)^{a - \frac{1}{2}}}{(1 + M_2 \sinh^2 \beta)^{a_4 + a_5} (1 + M_3 \sinh^2 \gamma)^{c_2}}$$

$$\times {}_2F_1\left(\frac{a_1+a_3}{2}, \frac{a_1+a_3+1}{2}; a; \frac{4M_1 M_3 y \sinh^2 \alpha \sinh^2 \gamma}{(1+M_1 \sinh^2 \alpha)^2 (1+M_3 \sinh^2 \gamma)}\right)$$

$$\times F_4\left(a_2, a_4 + a_5; c_1, c_2 - a; \frac{M_2 z \sinh^2 \beta}{(1+M_2 \sinh^2 \beta)}, \frac{u}{(1+M_2 \sinh^2 \beta)(1+M_3 \sinh^2 \gamma)}\right)$$

$$\times d\alpha d\beta d\gamma,$$

$$(Re(a_i) > 0, (i = 1, 3, 4, 5), Re(a) > 0, Re(c_2 - a) > 0, M_j > 0, j = (1, 2, 3)).$$

Theorem 3.2. *Each of the following integral representations holds true:*

$$(3.6) \quad X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u)$$

$$= \frac{\Gamma(a_1 + a_2)\Gamma(a_3 + a_5)}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_5)} \int_0^1 \int_0^1 \alpha^{a_1-1} \beta^{a_3-1} (1 - \alpha)^{a_2-1} (1 - \beta)^{a_5-1}$$

$$\begin{aligned} & \times X_{19}^{(4)}(a_1 + a_2, a_1 + a_2, a_1 + a_2, a_1 + a_2, a_1 + a_2, a_3 + a_5, a_4, a_3 + a_5; \\ & c_1, c_1, c_1, c_2; \alpha^2 x, (1 - \alpha)(1 - \beta)u, (1 - \alpha)z, \alpha\beta y) d\alpha d\beta, \\ & (Re(a_i) > 0, (i = 1, 2, 3, 5)), \end{aligned}$$

$$\begin{aligned} (3.7) \quad & X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u) \\ & = \frac{\Gamma(a_1 + a_5)\Gamma(a_2 + a_3)\Gamma(c_1)}{2^{a_1+a_2+a_3+a_5+c_1-6}\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_5)\Gamma(a)\Gamma(c_1-a)} \\ & \int_{-1}^1 \int_{-1}^1 \int_{-1}^1 \frac{[(1+\alpha)^2]^{a_1-\frac{1}{2}} [(1-\alpha)^2]^{a_5-\frac{1}{2}} [(1+\beta)^2]^{a_2-\frac{1}{2}} [(1-\beta)^2]^{a_3-\frac{1}{2}}}{(1+\alpha^2)^{a_1+a_5} (1+\beta^2)^{a_2+a_3}} \\ & \times \frac{[(1+\gamma)^2]^{a-\frac{1}{2}} [(1-\gamma)^2]^{c_1-a-\frac{1}{2}}}{(1+\gamma^2)^{c_1}} X_{21}^{(4)}(a_1 + a_5, a_1 + a_5, a_2 + a_3, \\ & a_1 + a_5, a_1 + a_5, a_2 + a_3, a_4, a_2 + a_3; a, a, c_1 - a, c_2; \frac{(1+\alpha)^4(1+\gamma)^2 x}{8(1+\alpha^2)(1+\gamma^2)}, \\ & \frac{(1-\alpha)^2(1+\beta)^2(1+\gamma)^2 u}{8(1+\alpha^2)(1+\beta^2)(1+\gamma^2)}, \frac{(1+\beta)^2(1-\gamma)^2 z}{4(1+\beta^2)(1+\gamma^2)}, \frac{(1+\alpha)^2(1-\beta)^2 y}{4(1+\alpha^2)(1+\beta^2)}) d\alpha d\beta d\gamma, \\ & (Re(a_i) > 0, (i = 1, 2, 3, 5), Re(a) > 0, Re(c_1 - a) > 0), \end{aligned}$$

$$\begin{aligned} (3.8) \quad & X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u) \\ & = \frac{\Gamma(a_1 + a_5)\Gamma(a_2 + a_3)(1 + M_1)^{a_1}(1 + M_2)^{a_2}}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_5)} \int_0^1 \int_0^1 \frac{\alpha^{a_1-1}(1-\alpha)^{a_5-1}}{(1 + M_1\alpha)^{a_1+a_5}} \\ & \times \frac{\beta^{a_2-1}(1-\beta)^{a_3-1}}{(1 + M_2\beta)^{a_2+a_3}} X_{23}^{(4)}(a_1 + a_5, a_1 + a_5, a_2 + a_3, a_1 + a_5, a_1 + a_5, \\ & a_2 + a_3, a_4, a_2 + a_3; c_1, c_1, c_1, c_2; \frac{(1+M_1)^2\alpha^2 x}{(1+M_1\alpha)^2}, \frac{(1+M_2)(1-\alpha)\beta u}{(1+M_1\alpha)(1+M_2\beta)}, \\ & \frac{(1+M_2)\beta z}{(1+M_2\beta)}, \frac{(1+M_1)\alpha(1-\beta)y}{(1+M_1\alpha)(1+M_2\beta)}) d\alpha d\beta, \\ & (Re(a_i) > 0, (i = 1, 2, 3, 5), M_1 > -1, M_2 > -1), \end{aligned}$$

$$\begin{aligned} (3.9) \quad & X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u) \\ & = \frac{\Gamma(a_1 + a_2)\Gamma(c_1)}{2^{a_1+a_2+c_1-2}\Gamma(a_1)\Gamma(a_2)\Gamma(a)\Gamma(c_1-a)} \int_{-1}^1 \int_{-1}^1 (1+\alpha)^{a_1-1}(1-\alpha)^{a_2-1} \\ & \times (1+\beta)^{a-1}(1-\beta)^{c_1-a-1} X_{28}^{(4)}(a_1 + a_2, a_1 + a_2, a_1 + a_2, a_1 + a_2, \\ & a_1 + a_2, a_4, a_3, a_5; c_1 - a, c_1 - a, c_2, a; \frac{(1+\alpha)^2(1-\beta)x}{8}, \frac{(1-\alpha)(1-\beta)z}{4}, \\ & \frac{(1+\alpha)y}{2}, \frac{(1-\alpha)(1+\beta)u}{4}) d\alpha d\beta, \\ & (Re(a_1) > 0, Re(a_2) > 0, Re(a) > 0, Re(c_1 - a) > 0), \end{aligned}$$

Theorem 3.3. *Each of the following integral representations holds:*

$$\begin{aligned}
 (3.10) \quad & X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \frac{\Gamma(a_2 + a_3)\Gamma(c_2)}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(c_2 - a_1)} \\
 &\int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \left(\frac{1}{2} + \alpha\right)^{a_1-1} \left(\frac{1}{2} + \beta\right)^{a_2-1} \left(\frac{1}{2} - \beta\right)^{a_3-1} \\
 &\times \left[\left(\frac{1}{2} - \alpha\right) + \left(\frac{1}{2} + \alpha\right)^2 x \right]^{c_2-a_1-1} F_D^{(3)}\left(a_2 + a_3, 1 + a_1 - c_2, \right. \\
 &\left. a_4, a_5; c_1; -\frac{\left(\frac{1}{2}+\alpha\right)\left(\frac{1}{2}-\beta\right)y}{\left[\left(\frac{1}{2}-\alpha\right)+\left(\frac{1}{2}+\alpha\right)^2 x\right]}, \left(\frac{1}{2} + \beta\right) z, \left(\frac{1}{2} + \beta\right) u \right) d\alpha d\beta \\
 &(Re(a_i) > 0, (i = 1, 2,), Re(c_2 - a_1) > 0),
 \end{aligned}$$

$$\begin{aligned}
 (3.11) \quad & X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \frac{4\Gamma(a_1 + a_3)\Gamma(a_4 + a_5)}{\Gamma(a_1)\Gamma(a_3)\Gamma(a_4)\Gamma(a_5)} \\
 &\int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{a_1-\frac{1}{2}} (\cos^2 \alpha)^{a_3-\frac{1}{2}} (\sin^2 \beta)^{a_4-\frac{1}{2}} \\
 &\times (\cos^2 \beta)^{a_5-\frac{1}{2}} F_R\left(\frac{a_1+a_3}{2}, a_2, \frac{a_1+a_3}{2}, \frac{a_1+a_3+1}{2}, a_4 + a_5, \frac{a_1+a_3+1}{2}; \right. \\
 &\left. c_2, c_1, c_1; 4x \sin^4 \alpha, z \sin^2 \beta + u \cos^2 \beta, y \sin^2 2\alpha \right) d\alpha d\beta, \\
 &(Re(a_i) > 0, (i = 1, 3, 4, 5)),
 \end{aligned}$$

$$\begin{aligned}
 (3.12) \quad & X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \frac{2\Gamma(c_2)(1 + M)}{\Gamma(a_1)\Gamma(c_2 - a_1)} \int_0^{\frac{\pi}{2}} \frac{(\sin^2 \alpha)^{\frac{1}{2}} (\cos^2 \alpha)^{a_1-\frac{1}{2}}}{(1 + M \sin^2 \alpha)^{2c_2-a_1-1}} \\
 &\times \left[(1 + M) \sin^2 \alpha (1 + M \sin^2 \alpha) + x \cos^4 \alpha \right]^{c_2-a_1-1} F_S(1 + a_1 - c_2, \\
 &a_2, a_2, a_3, a_4, a_5; c_1, c_1, c_1; \frac{-\cos^2 \alpha(1+M \sin^2 \alpha)y}{[(1+M) \sin^2 \alpha(1+M \sin^2 \alpha)+x \cos^4 \alpha]}, z, u) d\alpha, \\
 &(Re(a_1) > 0, Re(c_2 - a_1) > 0, M > -1),
 \end{aligned}$$

$$\begin{aligned}
 (3.13) \quad & X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \frac{4\Gamma(a_3 + a_4)\Gamma(c_2)M_1^{a_3}M_2^{a_1}}{\Gamma(a_1)\Gamma(a_3)\Gamma(a_4)\Gamma(c_2 - a_1)}
 \end{aligned}$$

$$\begin{aligned}
& \times \int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{(\sin^2 \alpha)^{a_3 - \frac{1}{2}} (\cos^2 \alpha)^{a_4 - \frac{1}{2}} (\sin^2 \beta)^{a_1 - \frac{1}{2}} (\cos^2 \beta)^{c_2 - a_1 - \frac{1}{2}}}{(\cos^2 \alpha + M_1 \sin^2 \alpha)^{a_3 + a_4} (\cos^2 \beta + M_2 \sin^2 \beta)^{2c_2 - a_1 - 1}} \\
& \times [(\cos^2 \beta + M_2 \sin^2 \beta) + M_2^2 x \sin^2 \beta \tan^2 \beta]^{c_2 - a_1 - 1} \\
& F_T(1 + a_1 - c_1, a_2, a_2, a_3 + a_4, a_5, a_3 + a_4; c_1, c_1, c_1; \\
& \quad \frac{-M_1 M_2 \sin^2 \alpha \tan^2 \beta (\cos^2 \beta + M_2 \sin^2 \beta) y}{(\cos^2 \alpha + M_1 \sin^2 \alpha) [(\cos^2 \beta + M_2 \sin^2 \beta) + M_2^2 x \sin^2 \beta \tan^2 \beta]}, \\
& \quad u, \frac{z \cos^2 \alpha}{(\cos^2 \alpha + M_1 \sin^2 \alpha)}) d\alpha d\beta, \\
& (Re(a_i) > 0, (i = 1, 3, 4), Re(c_2 - a_1) > 0, M_1 > 0, M_2 > 0).
\end{aligned}$$

Corollary 3.2. *By considering equation (3.13) we obtain, when $u = 0$.*

$$\begin{aligned}
(3.14) \quad & X_{19}(a_1, a_3, a_2, a_4; c_2, c_1; x, y, z) \\
& = \frac{4\Gamma(a_3 + a_4)\Gamma(c_2)M_1^{a_3}M_2^{a_1}}{\Gamma(a_1)\Gamma(a_3)\Gamma(a_4)\Gamma(c_2 - a_1)} \\
& \times \int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{(\sin^2 \alpha)^{a_3 - \frac{1}{2}} (\cos^2 \alpha)^{a_4 - \frac{1}{2}} (\sin^2 \beta)^{a_1 - \frac{1}{2}} (\cos^2 \beta)^{c_2 - a_1 - \frac{1}{2}}}{(\cos^2 \alpha + M_1 \sin^2 \alpha)^{a_3 + a_4} (\cos^2 \beta + M_2 \sin^2 \beta)^{2c_2 - a_1 - 1}} \\
& \times [(\cos^2 \beta + M_2 \sin^2 \beta) + M_2^2 x \sin^2 \beta \tan^2 \beta]^{c_2 - a_1 - 1} \\
& F_1(a_3 + a_4, 1 + a_1 - c_2, a_2; c_1; \\
& \quad \frac{-M_1 M_2 \sin^2 \alpha \tan^2 \beta (\cos^2 \beta + M_2 \sin^2 \beta) y}{(\cos^2 \alpha + M_1 \sin^2 \alpha) [(\cos^2 \beta + M_2 \sin^2 \beta) + M_2^2 x \sin^2 \beta \tan^2 \beta]}, \\
& \quad \frac{z \cos^2 \alpha}{(\cos^2 \alpha + M_1 \sin^2 \alpha)}) d\alpha d\beta, \\
& (Re(a_i) > 0, (i = 1, 3, 4), Re(c_2 - a_1) > 0, M_1 > 0, M_2 > 0).
\end{aligned}$$

Theorem 3.4. *Each of the following integral representations holds true:*

$$\begin{aligned}
(3.15) \quad & X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
& = \frac{\Gamma(a_1 + a_2)\Gamma(a_4 + a_5)\Gamma(c)}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_4)\Gamma(a_5)\Gamma(c - a)(S_1 - R_1)^{a_1 + a_2 - 1}(S_2 - R_2)^{a_4 + a_5 - 1}(S_3 - R_3)^{c - 1}} \\
& \times \int_{R_1}^{S_1} \int_{R_2}^{S_2} \int_{R_3}^{S_3} (\alpha - R_1)^{a_1 - 1} (S_1 - \alpha)^{a_2 - 1} (\beta - R_2)^{a_4 - 1} (S_2 - \beta)^{a_5 - 1} (\gamma - R_3)^{a - 1} \\
& \times (S_3 - \gamma)^{c - a - 1} X_6\left(a_1 + a_2, a_3, a_4 + a_5; c - a, a; \frac{(\alpha - R_1)^2(S_3 - \gamma)x}{(S_1 - R_1)^2(S_3 - R_3)}, \right. \\
& \quad \left. \frac{(\alpha - R_1)(S_3 - \gamma)y}{(S_1 - R_1)(S_3 - R_3)}, \frac{(S_1 - \alpha)(\beta - R_2)(\gamma - R_3)z + (S_1 - \alpha)(S_2 - \beta)(\gamma - R_3)u}{(S_1 - R_1)(S_2 - R_2)(S_3 - R_3)}\right) d\alpha d\beta d\gamma, \\
& (Re(a_i) > 0, (i = 1, 2, 4, 5), Re(a) > 0, Re(c - a) > 0, R_j < S_j, (j = 1, 2, 3)),
\end{aligned}$$

$$\begin{aligned}
(3.16) \quad & X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
& = \frac{\Gamma(a_3 + a_4 + a_5)(S_1 - T_1)^{a_5}(R_1 - T_1)^{a_3 + a_4}(S_2 - T_2)^{a_3}(R_2 - T_2)^{a_4}}{\Gamma(a_3)\Gamma(a_4)\Gamma(a_5)(S_1 - R_1)^{a_3 + a_4 + a_5 - 1}(S_2 - R_2)^{a_3 + a_4 - 1}}
\end{aligned}$$

$$\begin{aligned} & \times \int_{R_1}^{S_1} \int_{R_2}^{S_2} \frac{(\alpha - R_1)^{a_5-1} (S_1 - \alpha)^{a_3+a_4-1} (\beta - R_2)^{a_3-1} (S_2 - \beta)^{a_4-1}}{(\alpha - T_1)^{a_3+a_4+a_5} (\beta - T_2)^{a_3+a_4}} \\ & \times X_{13} \left(a_1, a_3 + a_4 + a_5, a_2; c; x, \frac{(R_1 - T_1)(S_2 - T_2)(S_1 - \alpha)(\beta - R_2)y}{(S_1 - R_1)(S_2 - R_2)(\alpha - T_1)(\beta - T_2)}, \right. \\ & \left. \frac{(R_1 - T_1)(R_2 - T_2)(S_1 - \alpha)(S_2 - \beta)z + (S_1 - T_1)(S_2 - R_2)(\alpha - R_1)(\beta - T_2)u}{(S_1 - R_1)(S_2 - R_2)(\alpha - T_1)(\beta - T_2)} \right) d\alpha d\beta, \\ & (Re(a_i) > 0, (i = 3, 4, 5), T_1 < R_1 < S_1, T_2 < R_2 < S_2), \end{aligned}$$

$$\begin{aligned} (3.17) \quad & X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\ & = \frac{2M^{a_4}\Gamma(a_4 + a_5)}{\Gamma(a_4)\Gamma(a_5)} \int_0^\infty \frac{\cosh \alpha (\sinh^2 \alpha)^{a_4 - \frac{1}{2}}}{(1 + M \sinh^2 \alpha)^{a_4 + a_5}} \\ & \quad \times X_{18}(a_1, a_3, a_2, a_4 + a_5; c; x, y, \frac{Mz \sinh^2 \alpha + u}{(1 + M \sinh^2 \alpha)}) d\alpha, \\ & (Re(a_4) > 0, Re(a_5) > 0, M > 0), \end{aligned}$$

$$\begin{aligned} (3.18) \quad & X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\ & = \frac{\Gamma(a_4 + a_5)\Gamma(c)}{\Gamma(a_4)\Gamma(a_5)\Gamma(a)\Gamma(c - a)} \int_0^\infty \int_0^\infty \frac{\alpha^{a-1}}{(1 + \alpha)^c} \frac{\beta^{a_5-1}}{(1 + \beta)^{a_4+a_5}} \\ & \quad \times X_{20}(a_1, a_3, a_2, a_4 + a_5; \\ & \quad a, c - a; \frac{\alpha x}{(1 + \alpha)}, \frac{y}{(1 + \alpha)}, \frac{\alpha z + \alpha \beta u}{(1 + \alpha)(1 + \beta)}) d\alpha d\beta, \\ & (\Re(a_4) > 0, \Re(a_5) > 0, \Re(a) > 0, \Re(c - a) > 0). \end{aligned}$$

Proof. It is noted that each of the integral representations of Euler-type in (3.1)-(3.18) can be proved mainly by expressing the series definition of the involved special function in each integrand and changing the order of the integral sign and the summation, and finally using the following integral representations of the Beta function (see [12, 20]):

$$\begin{aligned} B(a, b) &= \begin{cases} \int_0^1 \alpha^{a-1} (1 - \alpha)^{b-1} dt & (\Re(a) > 0, \Re(b) > 0), \\ \frac{\Gamma(a + b)}{\Gamma(a)\Gamma(b)} & (a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-), \end{cases} \\ B(a, b) &= \int_0^1 \alpha^{a-1} (1 - \alpha)^{b-1} d\alpha \\ &= (S - R)^{1-a-b} \int_R^S (\alpha - R)^{a-1} (S - \alpha)^{b-1} d\alpha \\ & \quad (Re(a) > 0, Re(b) > 0, R < S), \\ B(a, b) &= \frac{(S - T)^a (R - T)^b}{(S - R)^{a+b-1}} \int_R^S \frac{(\alpha - R)^{a-1} (S - \alpha)^{b-1}}{(\alpha - T)^{a+b}} d\alpha \end{aligned}$$

$$\begin{aligned}
&= (M+1)^a \int_0^1 \frac{\alpha^{a-1}(1-\alpha)^{b-1}}{(1+M\alpha)} d\alpha \\
&\quad (T < R < S, M > -1, \operatorname{Re}(a) > 0, \operatorname{Re}(b) > 0), \\
B(a, b) &= 2 \int_0^{\frac{\pi}{2}} (\sin \alpha)^{2a-1} (\cos \alpha)^{2b-1} d\alpha = \int_0^\infty \frac{\alpha^{a-1}}{(1+\alpha)^{a+b}} d\alpha \\
&\quad (\operatorname{Re}(a) > 0, \operatorname{Re}(b) > 0), \\
B(a, b) &= 2^{1-a-b} \int_{-1}^1 (1+\alpha)^{a-1} (1-\alpha)^{b-1} d\alpha \\
&= 2M^a \int_0^\infty \frac{\cosh \alpha (\sinh \alpha)^{2a-1}}{(1+M \sinh^2 \alpha)^{a+b}} d\alpha \\
&\quad (\operatorname{Re}(a) > 0, \operatorname{Re}(b) > 0, M > 0). \quad \square
\end{aligned}$$

4. Laplace-type integrals

The confluent hypergeometric functions ${}_0F_1$, ${}_1F_1$, Φ_2 , Φ_3 , $\Phi_2^{(3)}$ and $\Phi_3^{(3)}$ are defined, respectively, by (see [22])

$$(4.1) \quad {}_0F_1(-; c; x) = \sum_{m=0}^{\infty} \frac{1}{(c)_m} \frac{x^m}{m!},$$

$$(4.2) \quad {}_1F_1(a; c; x) = \sum_{m=0}^{\infty} \frac{(a)_m}{(c)_m} \frac{x^m}{m!},$$

$$(4.3) \quad \Phi_2(a, b; c; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_m (b)_n}{(c)_{m+n}} \frac{x^m}{m!} \frac{y^n}{n!},$$

$$(4.4) \quad \Phi_3(a; c; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_m}{(c)_{m+n}} \frac{x^m}{m!} \frac{y^n}{n!},$$

$$(4.5) \quad \Phi_2^{(3)}(a, b, c; d; x, y, z) = \sum_{m, n, p=0}^{\infty} \frac{(a)_m (b)_n (c)_p}{(d)_{m+n+p}} \frac{x^m}{m!} \frac{y^n}{n!} \frac{z^p}{p!}$$

and

$$(4.6) \quad \Phi_3^{(3)}(a, b; c; x, y, z) = \sum_{m, n, p=0}^{\infty} \frac{(a)_m (b)_n}{(c)_{m+n+p}} \frac{x^m}{m!} \frac{y^n}{n!} \frac{z^p}{p!}.$$

Theorem 4.1. For $X_{80}^{(4)}$, $X_{81}^{(4)}$, $X_{82}^{(4)}$, $X_{83}^{(4)}$, we have the following integral representations:

$$(4.7) \quad X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u)$$

$$\begin{aligned}
 &= \frac{1}{\Gamma(a_1)\Gamma(a_2)} \int_0^\infty \int_0^\infty e^{-(s+t)} s^{a_1-1} t^{a_2-1} \\
 &\quad \times \Phi_3(a_4; c_1; tz, s^2x) \Phi_2(a_3, a_5; c_2; sy, tu) dsdt, \\
 &\quad (Re(a_1) > 0, Re(a_2) > 0),
 \end{aligned}$$

$$\begin{aligned}
 (4.8) \quad &X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u) \\
 &= \frac{1}{\Gamma(a_1)\Gamma(a_4)\Gamma(a_5)} \int_0^\infty \int_0^\infty \int_0^\infty e^{-(s+t+v)} s^{a_1-1} t^{a_4-1} v^{a_5-1} \\
 &\quad \times \Phi_3(a_2; c_1; tz + uv, s^2x) {}_1F_1(a_3; c_2; sy) dsdtdv, \\
 &\quad (Re(a_1) > 0, Re(a_4) > 0, Re(a_5) > 0),
 \end{aligned}$$

$$\begin{aligned}
 (4.9) \quad &X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \frac{1}{\Gamma(a_1)\Gamma(a_2)} \int_0^\infty \int_0^\infty e^{-(s+t)} s^{a_1-1} t^{a_2-1} \\
 &\quad \times {}_0F_1(-; c_2; s^2x) \Phi_2^{(3)}(a_3, a_4, a_5; c_1; sy, tz, tu) dsdt, \\
 &\quad (Re(a_1) > 0, Re(a_2) > 0),
 \end{aligned}$$

$$\begin{aligned}
 (4.10) \quad &X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
 &= \frac{1}{\Gamma(a_1)\Gamma(a_4)\Gamma(a_5)} \int_0^\infty \int_0^\infty \int_0^\infty e^{-(s+t+v)} s^{a_1-1} t^{a_4-1} v^{a_5-1} \\
 &\quad \times \Phi_3^{(3)}(a_2, a_3; c; tz + uv, sy, s^2x) dsdtdv, \\
 &\quad (Re(a_1) > 0, Re(a_4) > 0, Re(a_5) > 0).
 \end{aligned}$$

Proof. To prove the equality (4.7), denote by Δ the right side of the equality (4.7). Then, by substituting the expression of the confluent hypergeometric functions (4.3) and (4.4) into the right hand side of (4.7), we obtain

$$\begin{aligned}
 \Delta = &\sum_{m,n,p,q=0}^\infty \frac{(a_3)_n (a_4)_p (a_5)_q}{(c_1)_{m+p} (c_2)_{n+q} \Gamma(a_1)\Gamma(a_2)} \frac{x^m y^n z^p u^q}{m! n! p! q!} \int_0^\infty \int_0^\infty e^{-s} s^{a_1+2m+n-1} \\
 &\times e^{-t} t^{a_2+p+q-1} dsdt.
 \end{aligned}$$

Using the known result (see [12])

$$\Gamma(a) = \int_0^\infty e^{-s} s^{a-1} ds, (\Re(a) > 0),$$

after a little simplification, we easily arrive at the left-hand side of (4.7). Then, we easily can obtain the equalities (4.8) to (4.10) in similar way. \square

5. Operational formulas

Hasanov and Srivastava [14] define following operators:

$$(5.1) \quad \begin{aligned} H_{t_1, \dots, t_i}(a, b) &:= \frac{\Gamma(b)\Gamma(a + \delta_1 + \dots + \delta_i)}{\Gamma(a)\Gamma(b + \delta_1 + \dots + \delta_i)} \\ &= \sum_{k_1, \dots, k_i=0}^{\infty} \frac{(b-a)_{k_1+\dots+k_i} (-\delta_1)_{k_1} \dots (-\delta_i)_{k_i}}{(b)_{k_1+\dots+k_i} k_1! \dots k_i!} \end{aligned}$$

and

$$(5.2) \quad \begin{aligned} \bar{H}_{t_1, \dots, t_i}(a, b) &:= \frac{\Gamma(a)\Gamma(a + \delta_1 + \dots + \delta_i)}{\Gamma(b)\Gamma(a + \delta_1 + \dots + \delta_i)} \\ &= \sum_{k_1, \dots, k_i=0}^{\infty} \frac{(b-a)_{k_1+\dots+k_i} (-\delta_1)_{k_1} \dots (-\delta_i)_{k_i}}{(1-a-\delta_1-\dots-\delta_i)_{k_1+\dots+k_i} k_1! \dots k_i!}, \end{aligned}$$

where $\delta_j := t_j \frac{\partial}{\partial t_j}$, $j = 1, \dots, i$; $i \in \mathbb{N} := \{1, 2, 3, \dots\}$.

Now, by using the symbolic operators (5.1) and (5.2), we establish the following operational representations.

Theorem 5.1. *The following results hold:*

$$(5.3) \quad \begin{aligned} X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\ = H_{z,u}(a_2, a) X_{80}^{(4)}(a_1, a_1, a, a, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u), \end{aligned}$$

$$(5.4) \quad \begin{aligned} X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\ = H_y(a_3, a) X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u), \end{aligned}$$

$$(5.5) \quad \begin{aligned} X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\ = H_y(a_3, a) H_z(a_4, \acute{a}) X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a, \acute{a}, a_5; c_1, c_2, c_1, c_2; x, y, z, u), \end{aligned}$$

$$(5.6) \quad \begin{aligned} X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_2; x, y, z, u) \\ = \bar{H}_y(a, a_3) \bar{H}_z(\acute{a}, a_4) X_{80}^{(4)}(a_1, a_1, a_2, a_2, a_1, a, \acute{a}, a_5; c_1, c_2, c_1, c_2; x, y, z, u). \end{aligned}$$

Theorem 5.2. *The following results hold:*

$$(5.7) \quad \begin{aligned} X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_5, a_4, a_3; c_1, c_2, c_1, c_1; x, u, z, y) \\ = H_u(a_5, c_2) (1-u)^{-a_1} F_S \left(\frac{a_1}{2}, a_2, a_2, \frac{a_1+1}{2}, a_3, a_4; c_1, c_1, c_1; \frac{4x}{(1-u)^2}, y, z \right), \end{aligned}$$

$$(5.8) \quad \begin{aligned} (1-u)^{-a_1} F_S \left(\frac{a_1}{2}, a_2, a_2, \frac{a_1+1}{2}, a_3, a_4; c_1, c_1, c_1; \frac{4x}{(1-u)^2}, y, z \right) \\ = \bar{H}_u(a_5, c_2) X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_5, a_4, a_3; c_1, c_2, c_1, c_1; x, u, z, y), \end{aligned}$$

$$(5.9) \quad X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u)$$

$$\begin{aligned}
 &= H_{x,z,u}(c, c_1) X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c_2, c, c; x, y, z, u), \\
 (5.10) \quad &X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c_2, c_1, c_1; x, y, z, u) \\
 &= H_y(c, c_2) X_{81}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_1, c, c_1, c_1; x, y, z, u),
 \end{aligned}$$

Corollary 5.1. *In formula (5.8), if we put $z = 0$, we have the following result.*

$$\begin{aligned}
 (5.11) \quad &X_{20}(a_1, a_5, a_2, a_3; c_1, c_2; x, u, y) \\
 &= H_u(a_5, c_2) (1 - u)^{-a_1} F_3\left(\frac{a_1}{2}, a_2, \frac{a_1+1}{2}, a_3; c_1; \frac{4x}{(1-u)^2}, y\right).
 \end{aligned}$$

Theorem 5.3. *The following results hold:*

$$\begin{aligned}
 (5.12) \quad &X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= H_y(a_3, a) H_z(a_4, \acute{a}) H_u(a_5, \acute{a}) \\
 &\quad \times X_{82}^{(4)}\left(a_1, a_1, a_2, a_2, a_1, a, \acute{a}, \acute{a}; c_2, c_1, c_1, c_1; x, y, z, u\right),
 \end{aligned}$$

$$\begin{aligned}
 (5.13) \quad &X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \bar{H}_y(a, a_3) \bar{H}_z(\acute{a}, a_4) \bar{H}_u(\acute{a}, a_5) \\
 &\quad \times X_{82}^{(4)}\left(a_1, a_1, a_2, a_2, a_1, a, \acute{a}, \acute{a}; c_2, c_1, c_1, c_1; x, y, z, u\right),
 \end{aligned}$$

$$\begin{aligned}
 (5.14) \quad &X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= H_x(c, c_2) X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c_1, c_1, c_1; x, y, z, u),
 \end{aligned}$$

$$\begin{aligned}
 (5.15) \quad &X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c_2, c_1, c_1, c_1; x, y, z, u) \\
 &= \bar{H}_x(c_2, c) X_{82}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c_1, c_1, c_1; x, y, z, u).
 \end{aligned}$$

Theorem 5.4. *The following results hold:*

$$\begin{aligned}
 (5.16) \quad &X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
 &= H_{z,u}(a_2, a) H_y(a_3, \acute{a}) X_{83}^{(4)}(a_1, a_1, a, a, a_1, \acute{a}, a_4, a_5; c, c, c, c; x, y, z, u),
 \end{aligned}$$

$$\begin{aligned}
 (5.17) \quad &X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
 &= H_u(a_5, a) X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a; c, c, c, c; x, y, z, u),
 \end{aligned}$$

$$\begin{aligned}
 (5.18) \quad &X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
 &= \bar{H}_u(a, a_5) X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a; c, c, c, c; x, y, z, u),
 \end{aligned}$$

$$\begin{aligned}
 (5.19) \quad &X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a_5; c, c, c, c; x, y, z, u) \\
 &= \bar{H}_{x,y,z,u}(c, \acute{c}) X_{83}^{(4)}(a_1, a_1, a_2, a_2, a_1, a_3, a_4, a; \acute{c}, \acute{c}, \acute{c}, \acute{c}; x, y, z, u).
 \end{aligned}$$

Proof. The proof of the results (5.3) to (5.19) is based upon application of Mellin and Mellin-Barnes integral representation methods for hypergeometric functions (see [17]). We omit the details involved in these derivations. \square

References

- [1] P. Agarwal, J. A. Younis, and T. Kim, *Certain generating functions for the quadruple hypergeometric series K_{10}* , Notes on Number Theory and Discrete Math. **25** (2019), 16–23.
- [2] L. C. Andrews, *Special Functions for Engineers and Applied Mathematicians*, Macmillan Publishing Company, New York, 1985.
- [3] E. Ata and O. Kıymaz, *A study on certain properties of generalized special functions defined by Fox-Wright function*, Appl. Math. Nonlinear Sci. **5** (2020), no. 1, 147–162. <https://doi.org/10.2478/amns.2020.1.00014>
- [4] W. W. Bell, *Special Functions for Scientists and Engineers*, Oxford University press, London, 1968.
- [5] L. Bers, *Mathematical Aspects of Subsonic and Transonic Gas Dynamics*, Wiley, New York, 1958.
- [6] M. G. Bin-Saad and J. A. Younis, *On connections between certain class of new quadruple and known triple hypergeometric series*, Tamap Journal of Mathematics and Statistics. Volume 2019.
- [7] ———, *Some integrals connected with a new quadruple hypergeometric series*, Universal J. Math. Appl. **3** (2020), 19–27.
- [8] ———, *Further quadruple hypergeometric series and their certain properties*, Boletim da Sociedade Paranaense de Matematica. In Press.
- [9] M. G. Bin-Saad, J. A. Younis, and R. Aktas, *New quadruple hypergeometric series and their integral representations*, Sarajevo J. Math. **14(27)** (2018), no. 1, 45–57. <https://doi.org/10.17654/ms103010021>
- [10] Yu. A. Brychkov and S. Saad, *On some formulas for the Appell function $F_2(a, b, b'; c, c', w, z)$* , Integral Transforms Spec. Funct. **25** (2014), 111–123.
- [11] J. Choi and A. K. Rathie, *General summation formulas for Kampé de Fériet function*, Montes Taurus J. Pure Appl. Math. **1** (2019) 107–128.
- [12] A. Erdélyi, W. Magnus, F. Oberhettinger, and F.G. Tricomi, *Higher Transcendental Functions. Vol. I*, McGraw-Hill Book Company, New York, Toronto and London, 1953.
- [13] H. Exton, *Hypergeometric functions of three variables*, J. Indian Acad. Math. **4** (1982), no. 2, 113–119.
- [14] A. Hasanov and H. M. Srivastava, *Decomposition formulas associated with the Lauricella multivariable hypergeometric functions*, Comput. Math. Appl. **53** (2007), no. 7, 1119–1128. <https://doi.org/10.1016/j.camwa.2006.07.007>
- [15] W. Koepf, I. Kim, and A. K. Rathie, *On a new class of Laplace-type integrals involving generalized hypergeometric functions*, Axioms **8** (2019), 87.
- [16] G. Lauricella, *Sull' funzioni ipergeometriche a pi variabili*, Rend. Cric. Mat. Palermo. **7** (1893), 111–158.
- [17] O. I. Marichev, *Handbook of integral transforms of higher transcendental functions*, translated from the Russian by L. W. Longdon, Ellis Horwood Series: Mathematics and its Applications, Ellis Horwood Ltd., Chichester, 1983.
- [18] K. S. Miller and B. Ross, *An introduction to the fractional calculus and fractional differential equations*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1993.
- [19] G. V. Milovanović, R. K. Parmar, and A. K. Rathie, *Certain Laplace transforms of convolution type integrals involving product of two special ${}_pF_p$ functions*, Demonstr. Math. **51** (2018), no. 1, 264–276. <https://doi.org/10.1515/dema-2018-0025>

- [20] E. D. Rainville, *Special Functions*, The Macmillan Company, New York, NY, USA, 1960; Reprinted by Chelsea Publishing Company, Bronx, NY, USA, 1971.
- [21] C. Sharma and C. L. Parihar, *Hypergeometric functions of four variables. I*, J. Indian Acad. Math. **11** (1989), no. 2, 121–133.
- [22] H. M. Srivastava and P. W. Karlsson, *A Treatise on Generating Functions*, Ellis Horwood Lt1., Chichester, 1984.
- [23] J. A. Younis and K. S. Nisar, *Several Euler-type integrals involving Exton's quadruple hypergeometric series*, J. Math. Computer Sci. **21** (2020), 286–295.

MAGED G. BIN-SAAD
DEPARTMENT OF MATHEMATICS
ADEN UNIVERSITY
ADEN, YEMEN
Email address: mgbinsaad@yahoo.com

KOTTAKKARAN SOOPPY NISAR
DEPARTMENT OF MATHEMATICS
COLLEGE OF ARTS AND SCIENCE-WADI ALDAWASER
11991, PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY
KINGDOM OF SAUDI ARABIA
Email address: ksnisar1@gmail.com, n.sooppy@psau.edu.sa

JIHAD A. YOUNIS
DEPARTMENT OF MATHEMATICS
ADEN UNIVERSITY
ADEN, YEMEN
Email address: jihadalsaqqaf@gmail.com