

## Healthcare Considerations for Special Populations during the COVID-19 Pandemic: A Review

Kim, Jeung-Im<sup>1,\*</sup> · Im, Yeojin<sup>2,\*</sup> · Song, Ju-Eun<sup>3</sup> · Jang, Sun Joo<sup>4</sup>

<sup>1</sup>School of Nursing, Soonchunhyang University, Cheonan

<sup>2</sup>College of Nursing Science · East-West Nursing Research Institute, Kyung Hee University, Seoul

<sup>3</sup>College of Nursing · Research Institute of Nursing Science, Ajou University, Suwon

<sup>4</sup>Red Cross College of Nursing, Chung-Ang University, Seoul, Korea

The coronavirus disease 2019 (COVID-19) has emerged as a threat to human health and public safety. People of all ages are susceptible to severe acute respiratory syndrome coronavirus 2 infection. However, the clinical manifestations of this infection differ by age. This study purposes to describe healthcare considerations for special populations, such as children, pregnant and lactating women, and older adults, who may have unique healthcare needs, in the pandemic situation. To realize the research purpose, we conducted a review of the practice guidelines of public documents and qualified studies that were published online/offline during a specific period. The review identified current knowledge on care for newborns, children in schools, pregnant women (from antenatal to postpartum care), and older adults suffering from high-risk conditions. Subsequently, we summarize vaccination guidance for special populations and, finally, discuss the issues currently affecting special populations. Therefore, this current knowledge on care for special populations helps nurses to provide accurate information on vaccinations aimed at preventing COVID-19 and protecting the masses from infection. Currently, the scarcity of information on COVID-19 variants necessitates further research on measures to reduce pandemic spread.

**Key words:** Child; Pregnant Women; Aged; COVID-19; SARS-CoV-2

### INTRODUCTION

Currently, everyone may be susceptible to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection (COVID-19). However, the specific clinical features of COVID-19 vary with age [1]. According to existing reports, most children and the majority of pregnant women with COVID-19 experience mild or no symptoms, and the symptoms are similar to those of several common illnesses, such as cold, sore throat, influenza, and allergies [2,3].

Populations at higher risk for infection, such as pregnant women, individuals with preexisting medical conditions, and older adults, need to take precautions to protect themselves from getting COVID-19. Meanwhile, various other groups such as families, children, newborns, and breastfeeding individuals, are also susceptible to infection [4]. Moreover, infants may have a higher risk of COVID-19 than older children [2]. Pregnant women with underlying health conditions, such as diabetes, are also more likely to develop severe COVID-19 [3]. Furthermore, the risk of severity is higher in

\*These authors contributed equally as first authors.

Address reprint requests to :

Song, Ju-Eun

College of Nursing · Research Institute of Nursing Science, Ajou University, 164 Worldcup-ro, Yeongtong-gu, Suwon 16499, Korea

Tel: +82-31-219-7018 Fax: +82-31-219-7020 E-mail: [songje@ajou.ac.kr](mailto:songje@ajou.ac.kr)

Jang, Sun Joo

Red Cross College of Nursing, Chung-Ang University, 221 Heukseok-dong, Dongjak-gu, Seoul 06974, Korea

Tel: +82-2-820-5934 Fax: +82-2-824-7961 E-mail: [icedcoffee@cau.ac.kr](mailto:icedcoffee@cau.ac.kr)

Received: July 29, 2021 Revised: August 23, 2021 Accepted: August 27, 2021 Published online October 31, 2021

This is an Open Access article distributed under the terms of the Creative Commons Attribution NoDerivs License. (<http://creativecommons.org/licenses/by-nd/4.0>)

If the original work is properly cited and retained without any modification or reproduction, it can be used and re-distributed in any format and medium.

people aged 65 years or older; this group has a 630 times higher risk of dying from COVID-19 than the 18- to 29-year-old group [5]. In South Korea, the lethality rates were 1.1% for individuals aged 60 years or older and 18.7% for those aged 85 years or over (Appendix 1) [6]. Therefore, the vaccination of the 65 years and older group was prioritized among all the age groups.

There is significant heterogeneity among special populations regarding the effectiveness of COVID-19 vaccination, as well as the risk and severity of COVID-19, which influences the pandemic decision-making process. Further, a risk/benefit analysis for individual patients is at the center of the collaborative clinician/patient decision-making process, since the data for vaccination in specific populations remain limited.

Nurses play an important role in providing vaccination-related information and support to prevent COVID-19 [7]. Pregnant and lactating women are encouraged to clarify any queries on health concerns, such as the negative impact of COVID-19 vaccines on oneself and fetal outcomes, with their healthcare providers [8]. Vaccination may be considered at any gestational age, including the first trimester. As indicated by published safety data, a messenger RNA (mRNA) vaccine is generally preferred over other vaccines for pregnant women [9]. Further, the American College of Obstetricians and Gynecologists recommends that COVID-19 vaccines should not be withheld from pregnant or breastfeeding individuals.

Since the identification of SARS-CoV-2 has not been completed and a vaccine developed over a short period does not guarantee high levels of protection against the virus's mutations, nurses should remain sensitive to updated information. The Ministry of Health in Canada encourages its nurses to regularly check its website for updates and recommends them to consider the website information as current information and check whether the data are up-to-date before conveying the information to patients or performing counseling [9].

Doctors report on and revise the guidelines for their areas of expertise based on the cases and evidence collected over the previous year. To ensure that each special population

group receives adequate and appropriate care, guidelines should be established for nurses on the different approaches to be adopted for outpatients and inpatients, children and pregnant women, older people, surgical patients and drug-treated patients, and oncological and non-neoplastic patients [4].

This review describes the nursing practice guidelines and specific healthcare considerations for special population groups, who may have unique healthcare needs, to prevent and manage infections during the COVID-19 pandemic period, based on reliable data and public documents. Further, we focus on such special populations, including children, pregnant women, and older adults, as high-risk groups for COVID-19.

## HEALTHCARE CONSIDERATIONS FOR CHILDREN

The main clinical symptoms of COVID-19 in children—from newborns to adolescents—are generally less severe than those in adult patients. Mild symptoms include fever, cough, and rhinorrhea; in some cases, there are no symptoms [10]. On the other hand, infants may have a higher risk of COVID-19 than older children; further, respiratory symptoms are less frequently reported among children compared to adults [10,11]. Although the severity of dyspnea symptoms in children with COVID-19 was generally lower than that of the symptoms reported in adult patients, infants with COVID-19 are more likely to develop dyspnea than others. Moreover, there have been increasing reports about cases of older school-aged children and adolescents experiencing prolonged fever, abdominal pain, shock, and cardiac dysfunction following a COVID-19 infection [10].

Thus, given that children are also vulnerable to morbidity and mortality from COVID-19, and considering that families and children are basic members of society for whom preventive and health management efforts against COVID-19 are implemented, specific health care guidelines for children, from newborns to adolescents, need to be considered.

The clinical manifestations of COVID-19 in pediatric patients include mild respiratory symptoms, such as sore

throat; cough; difficulty in breathing; gastrointestinal symptoms, including diarrhea, vomiting, or stomachache; or the onset of severe headache accompanied by fever. Often, children do not exhibit any symptoms [2,10]. Dyspnea was more common in infants than in any other age group, whereas mild gastrointestinal symptoms, such as vomiting and diarrhea, occurred more in older children [10,11].

Infants and children with underlying medical conditions have a greater probability of developing severe COVID-19 [2]. The underlying medical conditions include chronic lung disease; asthma; diabetes; and other genetic, neurologic, or metabolic conditions. Further, congenital heart disease, medical complexities, immunosuppression, and obesity were conditions that increased the risk of severe infection in children [2]. Although most pediatric patients with COVID-19 do not exhibit severe symptoms, severe COVID-19 in children often results in fatal conditions requiring Intensive Care Unit (ICU) admission [10,11].

For children with COVID-19 symptoms, the following guidelines are relevant [2]: 1) Keep the child home. 2) Consider whether the child should see a healthcare provider and be tested for COVID-19. The Centers for Disease Control and Prevention (CDC) recommend the testing of all children with COVID-19 symptoms. 3) Protect the caregivers caring for the sick child from COVID-19 by guiding them to wear masks, frequently practice hand hygiene, perform symptom monitoring on COVID-19, as well as using preventive/protective measures for symptoms of COVID-19. 4) Inform the child's school about the symptoms, notifying whether the child took a COVID-19 test, and conveying the test results. 5) Review the policies of the child's school (or childcare facility) to understand when a child who has been sick can return to the school (or childcare facility). 6) Finally, when children return to school or participate in in-person activities, they should stay safe with others [2]. In case the child shows any of the emergency warning signs, such as breathing difficulties; chest pain or pressure; confusion; cyanotic conditions like pale, gray-colored, or blue-colored lips, skin, or nail beds; and an inability to wake up or stay awake, the caregiver should call emergency services or take the child to the emergency department [2,11].

## 1. Guidelines for newborn care

COVID-19 during pregnancy is associated with only a slight increase in the risk of neonatal morbidities and no increase in neonatal mortality. Currently, it is difficult to conclude whether the rare cases of infection during pregnancy occur from horizontal transmission at birth or vertical transmission through the placenta [12].

The American Academy of Pediatrics recommends the rooming-in of infants with their mothers in accordance with hospital practices unless the mother is too unwell to care for her infant. The mother should implement infection-control precautions, including hand hygiene and mask wearing, during breastfeeding and performing rooming-in care of the newborn [13]. Further, even the most recent guidelines indicate that the breastfeeding practice is the best option to feed infants despite any risk of the mother or infant getting infected. Along with following all COVID-19 precaution measures, hospitals should provide breast milk or donor milk to sick infants admitted in the neonatal care unit [13,14].

In addition to encouraging caregivers to take personal precautions, the hospital should develop a strategy to support breastfeeding in suspected or infected cases of mothers or newborns using a systemic approach. Healthcare delivery with minimal or no contact can be ensured by promoting telemedicine or the remote monitoring of patients. The efforts of the Texas Tech University Health Science Center El Paso Children's Hospital Neonatal Intensive Care Unit to support low-cost breast milk delivery with minimal contact can be one of options that hospitals can emulate [14].

Additionally, among all mothers, including those with confirmed or suspected COVID-19, and newborns, kangaroo mother care and breastfeeding should be encouraged. If mothers are unwell, healthy family members should provide kangaroo care to newborns. In many regions, in the post-pandemic recovery period, the importance of creating opportunities to rebuild strategies to provide high-quality maternal and child health services, including kangaroo care, was emphasized [15].

## 2. Guidelines for child care centers and schools

Although strategies to combat infection may change according to variations in national strategies depending on the number of confirmed COVID-19 cases, the Ministry of Education in South Korea is trying to expand in-person education for all students at every level of school from the upcoming second fall semester. Further, the COVID-19 guidelines for kindergarten, elementary, and middle-to-high schools are presented on the Ministry of Education's webpage [16], and the updated guideline for child care centers is published on the Ministry of Health and Welfare's webpage. These guidelines pertain to the quarantining, isolation, symptom screening, and testing of children, as well as the national strategy to combat COVID-19 spread. Aspects such as attendance checking and school performance evaluation are mentioned in the guideline for schools.

The key prevention strategies for schools and child care centers include consistent and correct mask usage, hand hygiene, physical distancing and respiratory etiquette, timely sanitization of the physical environment, provision of appropriate ventilation, and facilitation of contact tracing in collaboration with community health centers [2,16]. During the COVID-19 pandemic, school nurses play an important role in ensuring the safety of schools and promoting child care programs (in case healthcare professionals are involved in health care management in child care facilities). The critical tasks of school nurses include the routine evaluation of symptoms or exposures in students and provision of support to school administrators and teachers in executing prevention strategies, such as participating in contact tracing; maintaining school-based clinics (if available); implementing school-based testing strategies; and providing medical support to students, families, and school staff in various ways [2,16].

## 3. Vaccination of children

Early information reveals that vaccines may help prevent people from spreading COVID-19 to others. The guidelines issued by the CDC in the United States recommend the Pfizer-BioNTech vaccine (Pfizer, New York, NY, USA; Bi-

oNTech, Mainz, Germany) for people aged 12 years and older [2,7]. South Korea is scheduled to begin the vaccination of senior high school students, teachers, and school staffs during July to August 2021 [16]. The CDC states that vaccination can protect children from becoming infected. Moreover, vaccination can help keep children from becoming severely sick even if they do become infected. Finally, the vaccination of all adult family members and children aged 12 years and older helps protect the entire family against COVID-19. Currently, the impact of COVID-19 vaccines on children younger than 12 years old is under study.

Recently, the CDC reported an increase in myocarditis and pericarditis cases in adolescents and young adults following COVID-19 vaccination in the United States [2]. Despite being associated with a possible risk of myocarditis or pericarditis, vaccination has several known and potential benefits that outweigh its known and potential risks. Hence, the CDC continues to recommend COVID-19 vaccination for all children aged 12 years and older [2]. Further, children should undergo timely and routine immunization as per the national immunization schedule.

## 4. Issues faced by children during the COVID-19 pandemic

Measures to diminish the spread of COVID-19, such as physical distancing, quarantining, and nationwide school closures, affect the health and well-being of children. In these circumstances, children may feel isolated, anxious, bored, and uncertain, as well as being fearful or sorrowful about the virus's impact on their families. Hence, policies that introduce helpful resources and ideas to support parents in helping their children understand the virus and protect themselves against COVID-19 should be implemented.

The suspension of in-person education in schools during the COVID-19 pandemic has raised concerns regarding the possibility of learning loss in students. In all communities, schools form an important part of the infrastructure by providing safe and supportive learning environments for students; employing teachers and other staff; and enabling parents, guardians, and caregivers to work outside their homes. Research suggests that students make no or little academic

progress through home learning and the related learning losses are more serious in countries with more fragile political and health care infrastructures or longer school closure periods compared to other countries [17]. The results of a research suggested that school closures did not contribute to epidemic control and the implementation of school closures alone would prevent only 2%~4% of deaths, which are far lower than the corresponding figures for other social distancing interventions [18]. Hence, policymakers should consider such relevant evidence and implement combinations of various social distancing measures, rather than focusing on school closures alone.

Additionally, the increase in mental health issues, including depression and anxiety, among children as a result of the social isolation practiced by people during the COVID-19 pandemic is a cause of widespread concern [19,20]. In the United Kingdom, a study on the longitudinal changes in childhood mental health that occurred during lockdowns, which promote social distancing practices and school closures, observed a significant increase in depressive symptoms and medium-to-large effect sizes of confidence intervals in a cohort of 8- to 12-year-olds compared to the period before lockdowns [19]. Therefore, policymakers should incorporate the association between school closures and social distancing and childhood mental health in the decision-making process of the education sector.

## HEALTHCARE CONSIDERATIONS FOR PREGNANT WOMEN

Pregnant women constitute a very special population because their health affects fetal health; hence, healthcare professionals should be aware of certain special considerations to prevent this group from contracting COVID-19 or manage them in case they do become infected. Generally, pregnant women are not considered a high-risk group for SARS-CoV-2 infection [21]. However, since pregnancy causes numerous physical changes, including immune and respiratory system changes [22,23], pregnant women have significantly higher risk of severe COVID-19 than non-pregnant women of similar age groups [21,24]. Therefore, preg-

nant women are considered clinically vulnerable to COVID-19 [23,25].

According to the national data in the United States, the majority of pregnant women with COVID-19 were infected during the third trimester ( $\geq 28$  weeks), followed by the second (14~27 weeks) and first trimesters ( $< 14$  weeks) [26]. Although the severity of COVID-19 in pregnant women ranges from asymptomatic to severe or critical, more than two-thirds generally remain asymptomatic and most of the symptomatic pregnant women experience only mild symptoms, for instance, a cold or the flu [21]. Further, although the common symptoms varied across studies, cough and fever were identified as the most common ones, whereas dyspnea, myalgia, loss of the sense of taste, and diarrhea were considered relatively less common [21].

Regarding the impact of COVID-19 on pregnancy, symptomatic maternal COVID-19 is related to a greater risk of preterm birth [3,27] and still birth [27] or a higher rate of caesarean birth compared to non-infected pregnant women [21,25]. Further, symptomatic pregnant women requiring hospitalization show worse overall progress than their non-symptomatic counterparts, and maternal outcomes include the risk, albeit very low, of death [22]. Further, pregnant women who are relatively older ( $\geq 35$  years); are overweight (body mass index  $\geq 25$  kg/m<sup>2</sup>); or have preexisting disease conditions, such as hypertension and diabetes, experience worse progress than the healthy pregnant women. Hence, they are more likely to receive ICU care than non-pregnant women of similar ages or non-risk pregnancy groups [22,24]. Therefore, pregnant women with high risk factors must promptly report any early symptoms of COVID-19, such as fever or cough, to their healthcare providers.

With respect to birth outcomes for individuals with COVID-19 in the US, vaginal delivery was more common than caesarean section, and the number of term births (37 weeks and over) was higher than that of preterm births (before 37 weeks) [26]. Also, today, data suggesting COVID-19's negative effects on the fetus or neonatal outcomes are limited [21-23]. Although some cases of vertical transmission (i.e., transmission from a woman to her baby during the

antenatal or intrapartum period) of SARS-CoV-2 have been reported [28], currently available data indicate that this transmission is uncommon [28–30]. Even cases of newborn infection do not appear to be affected by delivery modes, skin-to-skin contact, feeding types, or practices of rooming-in right after childbirth [21,22]. Therefore, healthcare professionals in the obstetric setting should support pregnant women to achieve successful maternal transition during pregnancy and childbirth by following the care principles or guidelines associated with the COVID-19 pandemic.

### 1. Guidelines for antenatal care of pregnant women

Healthy, uninfected pregnant women should adhere to the following general precautions to ensure protection from COVID-19: 1) frequently washing hands with an alcohol-based hand rub or soap and water; 2) maintaining a distance of at least six feet from others and avoiding crowded spaces; 3) wearing a non-medical fabric mask whenever one is unable to maintain social distancing; 4) avoiding touching the mouth, nose, and eyes; and 5) maintaining respiratory hygiene, such as covering the mouth and nose with a bent elbow or tissue when sneezing or coughing [21]. If pregnant women are not in self-isolation due to suspected or confirmed COVID-19, they should continue their routine antenatal care with modifications, if necessary [21,23]. When healthy pregnant women without any COVID-19 symptoms visit the hospital for antenatal care, healthcare professionals should use adequate personal protective equipment (PPE) before meeting them. In general, all women and anyone accompanying them should wear face coverings, such as face masks, according to national guidelines [21,22,29].

In many nations, if women report symptoms of COVID-19, such as fever or cough, or suspicions of being infected over telephone, the maternity nurse first conducts differential diagnoses regarding the symptoms. If the symptoms suggest COVID-19, pregnant women should refer themselves to the national service for SARS-CoV-2 testing. Obstetric units should have specific checklists or triage tools to assess the severity of the suspected or confirmed COVID-19 referred to over telephone [21,29,31]. For women with mild symptoms but no comorbidities, general rules of infection management,

including staying home, keeping a minimum distance of at least 6 ft from other people, and performing frequent hand-washing with soap and water or alcohol solutions, can be applied. In such cases, telephonic evaluation, rather than face-to-face prenatal visits, to monitor symptom severity is suggested [22]. However, if women experience fever exceeding 38°C or shortness of breath, which are the widely acknowledged COVID-19 alarm signs, they should visit the emergency room of the nearest hospital as soon as possible [22,31]. Further, on experiencing urgent pregnancy-related issues, such as uterine contractions, fluid leakage, or vaginal bleeding, women should directly visit the emergency department of the nearest hospital, as mentioned in the national guidelines, regardless of the severity of their COVID-19 symptoms [22].

When pregnant women show symptoms that necessitate hospitalization, the following general principles regarding the clinical management of pregnant women with COVID-19 are applicable: early isolation, strict infection control, SARS-CoV-2 and coinfection testing, oxygen therapy initiation (as needed), fluid overload avoidance, antibiotic use to prevent secondary bacterial infection, fetal condition and uterine contraction monitoring, early mechanical ventilation to avoid progressive respiratory failure, individualized delivery planning, and adoption of a team-based approach that facilitates multispecialty consultations [24]. Further, health care professionals should practice clinical management in keeping with the national guidelines when treating infected women in hospitals [21,29,32].

For women recovering from COVID-19 without hospital admission, due to the mildness or absence of symptoms, antenatal care should be maintained without any change in schedule following a period of self-isolation. However, women recovering from a critical condition that required hospital admission should plan antenatal care together with healthcare professionals before hospital discharge. Further, those experiencing severe COVID-19 symptoms should undergo ultrasound scanning 14 days after recovery to assess fetal health in case urgent clinical reasons for earlier scans are absent [21,22]. In general, all pregnant women with confirmed COVID-19 who have experienced hospitalization

should be given thromboprophylaxis for 10 days following hospital discharge, as pregnancy increases the risk of thromboembolism [21,22].

## 2. Vaccination of pregnant women

On December 30, 2020, the Joint Committee on Vaccination and Immunization (JCVI) in the United Kingdom published a guideline regarding the vaccination of pregnant women and confirmed that vaccination does not pose any safety concerns or harm to these women [33]. Nevertheless, the guideline stated that vaccination should be considered during pregnancy only when the risk of exposure to an infection is high or unavoidable [33]. However, the JCVI updated its advice on June 12, 2021 and notified that pregnant women should receive COVID-19 vaccination according to the general guidelines specified for people of the same age or risk group [22].

Only limited verified data regarding vaccination safety in pregnant women are available because of this group's exclusion from the Phase 3 COVID-19 vaccine trial [34,35]. However, according to data from the "v-safe after vaccination health checker, surveillance system" in the United States, pain at the injection site was reported as a common side effect of the COVID-19 vaccine in pregnant women compared to non-pregnant women; meanwhile, the occurrence of headache, myalgia, chills, and fever was less frequent in pregnant than non-pregnant women [34]. A comparison of adverse pregnancy and neonatal outcomes revealed similar incidence rates in pregnancy loss, live birth, and preterm birth between pregnant women receiving COVID-19 vaccination and pregnant women in the pre-pandemic period [34]. Further, in pregnant women, the receipt of a COVID-19 mRNA vaccine was immunogenic and the vaccine-elicited antibodies were transferred to infant cord blood and breastmilk [35]. However, in South Korea, no guideline has yet been established for vaccination in pregnant women; therefore, currently, vaccination is not performed in pregnant women [36]. Currently, the vaccines by Pfizer and Moderna are preferred for first-dose vaccination in pregnant women of any age in the United States. Anyone who received the first dose of a vaccine before becoming pregnant and

needs to take the second dose during pregnancy should take the second dose of the same vaccine unless they experienced any serious side effects after their first dose [22].

## 3. Issues faced during delivery and postpartum care during the COVID-19 pandemic

Regarding issues pertaining to labor and delivery care, to date, there is no evidence to suggest that the active COVID-19 virus passes through the placenta or breastmilk, since the virus has not been found in amniotic fluid or breastmilk samples [30]. Therefore, healthcare professionals should discuss delivery modes with the pregnant woman and her family, understand their preferences, and consider any obstetric or fetal condition requiring special care. In these women, caesarean section should be performed only when its obstetrical criteria are satisfied; hence, there is no need for pregnant women to give birth by caesarean section even if they are suspected of or confirmed as having COVID-19 [21,22,29].

Further, laboring women should be placed in a dedicated delivery room during labor and delivery, preferably with negative pressure. To avoid the unnecessary transfer of patients from one room to another, it should be possible to convert this delivery room to satisfy the requirements of a caesarian section, if necessary. The women in labor must wear a surgical mask throughout labor and delivery. The healthcare professionals supporting these women should wear PPE during obstetric procedures [37]. Further, electrical fetal monitoring throughout the labor process is suggested to check for any increase in the risk of fetal distress. Vaginal examination should be minimized to reduce the risk of professional exposure, as well [22].

With regards to postpartum care issues, the World Health Organization (WHO) recommends mothers with COVID-19 be roomed with their babies [3]; further, these mothers should be encouraged to practice skin-to-skin/kangaroo care if the newborn does not require additional medical care after childbirth. Hence, healthcare professionals should support women to breastfeed safely and ensure good respiratory hygiene while holding their newborns and providing skin-to-skin contact in the rooming-in environment [3,21]. More de-

tails on breastfeeding are included in the section “Guidelines for newborn care.”

Mothers showing COVID-19 symptoms should wear a medical mask during any contact with their baby. To minimize the risk of mother-to-infant transmission, everyone involved in the care of the mother and her infant should adopt a precautionary approach. Healthcare professionals should support the mother and her family in making decisions based on adequate information about breastfeeding or rooming-in even when they have COVID-19 symptoms [38].

Once women with COVID-19 are discharged from hospitals after childbirth, they should practice self-isolation at home for 14 days [25]. Those who gave birth recently and tested positive for COVID-19 after childbirth should follow all recommended guidelines, including the necessary in-person assessments conducted by healthcare professionals using appropriate PPE.

## HEALTHCARE CONSIDERATIONS FOR OLDER ADULTS

Globally, the proportion of the older population is increasing, and the pace of population aging is significantly higher today, compared to earlier figures [39]. Due to the occurrence of aging-related changes, diseases, and disabilities, the medical and nursing needs of older adults have escalated over the past several years, causing an increase in their medical expenditure [40]. In particular, the functions of the immune and respiratory systems decline with age [40,41]. Among older individuals, adaptive immune responses resulting from a cytokine storm cause uncontrollable inflammatory response and lymphocyte depletion; these effects are often aggravated by COVID-19 [41].

Older individuals are likely to experience neurocognitive disorders, hypertension, diabetes or cardiovascular disease, respiratory diseases, or disabilities caused by aging. Therefore, they form a high-risk group during the COVID-19 pandemic, and advanced age can be an independent risk factor of high mortality from COVID-19 [42]. Accordingly, the CDC, National Institute for Health, WHO, and National Health Service suggest various precautions for older people and

have established guidelines for healthcare professionals caring for older adults at high risk of COVID-19 [42–44]. Here, we integrate and summarize such guidelines to enable nurses to make optimal decisions while caring for older patients and to contribute to the prevention and prognosis of COVID-19 in this group.

### 1. Guidelines for care of older adults

Along with adhering to the guidelines specifically targeting the older population, nurses should strictly follow all general precautions while caring for this population group. The same preventive measures, including getting vaccinated, wearing a mask, practicing physical distancing (maintaining a distance of six feet from others), disinfecting surfaces, and washing hands [7], applicable to any other age group should be adhered to while caring for older individuals [44]. Normal body temperatures can be lower in older adults than in younger adults. Consequently, fever temperatures can be lower among older individuals [44]. A single temperature reading above 37.8°C, multiple readings higher than 37.2°C, or a rise in temperature greater than 1.1°C above the person’s baseline temperature can be signs of infection. Further, older people with cardiovascular disease have higher risks of developing multi-organ microvascular endothelial damage and uncontrolled systemic inflammatory response [41]. In addition, other unexplained and new symptoms include, but are not limited to, dizziness, lethargy, changes in cognition, chest pain, or loss of appetite. Older individuals may likely present with atypical signs and symptoms (delirium, increased frequency of falls, etc.), as well. Further, individuals with cognitive impairment who may not be able to describe COVID-19 symptoms may present with refusal of food and drink or an abrupt change in mental status, functional status, or behaviors [5,42–44]. In addition, the WHO’s guideline recommends strict social isolation in the geriatric population to control deaths in countries heavily affected by COVID-19 [43].

Currently, there is no evidence to suggest that taking any specific medication (e.g., blood pressure medication or ibuprofen) increases the severity of COVID-19 [44]. Therefore, nurses should encourage older adults to continue taking their



medications and following their treatment plans, as prescribed by their doctors. Any changes to medications should be made only after consulting the relevant doctor [44].

If healthcare professionals having close contact with older individuals diagnosed with COVID-19 are completely vaccinated, they will not need to stay away from others or get tested unless they show symptoms [42–44]. However, if they are not completely vaccinated, healthcare professionals should report to the manager, not attend any services at their workplace until the completion of their quarantine period, stay home, self-monitor for symptoms, and follow the national CDC's guidelines in case they develop symptoms [5,43].

## 2. Vaccination of older adults

Currently, older adults are being encouraged to get vaccinated as soon as possible. According to the CDC's morbidity and mortality report [45], older people should be prioritized over other groups for vaccination because the risk of COVID-19 severity increases with age. In the United States, COVID-19 vaccination began toward the end of 2020, and older adults were prioritized in the early phases [45]. As on May 1, 2021, 82% of older adults received at least one vaccine dose and, as on June 27, 2021, more than 90% of this group received at least one dose in the United Kingdom [46]. Consequently, the number of COVID-19 cases, emergency department visits, hospital admissions, and deaths declined in the older adult populations of the two countries [47]. In the United States, in the first 5 months after the initiation of vaccination, the rates of COVID-19 incidence in the older population decreased by 40% compared to the 18–49 years age group [47]. Additionally, the older adults' emergency department visits and deaths decreased by 50% and 66%, respectively [47]. Further, evaluation reports reveal that the risk of COVID-19-related hospitalizations significantly decreased among adults aged 65 years and older following the complete vaccination of this group [42]. However, older adults may require additional support to get vaccinated and, therefore, collaborations with other healthcare professionals or institutions that may already be working with older adults should be considered. In addition, according to a longitudinal

study [48], the COVID-19 antibody responses significantly decreased after 3 months. Therefore, after vaccination, adhering to the implementation of adequate healthcare protocols should be the most crucial strategy to be followed during the pandemic.

## 3. Issues faced by older adults during the COVID-19 pandemic

The globally implemented COVID-19 prevention strategy of social distancing can be a major cause of loneliness, which is an independent risk factor for mental health problems [49]. Social distancing is an effective preventive measure that can save lives; however, it also deprives older adults of basic human needs, such as companionship and direct communication [48].

Their bio-psychosocial vulnerabilities make older adults the group that is the most susceptible to COVID-19 pandemic-related mental health problems, therefore, social isolation can negatively affect their mental health [50]. Healthcare professionals should consider the discriminatory value of cumulative deficit frailty to identify older people [51]. While recommending older people to isolate themselves and shield themselves from infection, authorities should use Q-Frailty as a balanced measure of the risk of isolation leading to adverse health (physical, mental, and cognitive) and functional outcomes in the target population [52]. During the COVID-19 pandemic, anxiety exacerbates the severity of depression among older individuals [53]; therefore, appropriate psychological interventions to reduce anxiety and depression in this age group are recommended.

In long-term care settings, healthcare professionals should assess and improve their readiness to respond to COVID-19. Additional precautions and quarantine practices must be guaranteed for older adults and healthcare professionals in long-term care facilities. The CDC preparedness checklist for long-term acute care facilities should be utilized as a tool to develop a comprehensive COVID-19 strategy [43,54]. This checklist comprises three sections: 1) a structure for making plans and decisions; 2) the development of a written COVID-19 plan; and 3) the elements of a COVID-19 plan (a general plan to protect residents and visitors from getting

**Table 1.** Summary of the Healthcare Considerations for Special Populations

| Population group | Care considerations   | Vaccination   |
|------------------|---|---|
| Children         | <ul style="list-style-type: none"> <li>• The symptoms of children with COVID-19 are generally less severe than those of adults. Respiratory symptoms (i.e., dyspnea) was more common in infants and gastrointestinal symptoms were more common in older children.</li> <li>• School nurses play an important role in keeping the school safe during the pandemic.</li> <li>• The learning loss and mental health issues arising from prolonged social distancing and suspension of face-to-face instruction in school should be addressed.</li> </ul>   | The Pfizer-BioNTech vaccine is recommended for individuals aged 12 years and older.   |
| Pregnant women   | <ul style="list-style-type: none"> <li>• Symptomatic maternal COVID-19 is related to a higher risk of preterm birth; however, there is little evidence that it affects the fetus or causes neonatal infection.</li> <li>• Women in labor should be placed in a dedicated delivery room, preferably one with negative pressure, and they should wear a surgical mask.</li> <li>• Mothers with suspected or confirmed COVID-19 can room in with and receive support to breastfeed their infants.</li> <li>• All pregnant women who have been hospitalized and had confirmed COVID-19 should be offered thrombo-prophylaxis following hospital discharge.</li> </ul> | Pfizer and Moderna vaccines are the preferred vaccines for pregnant women of any age. |
| Older adults     | <ul style="list-style-type: none"> <li>• Older adults should be prioritized for vaccination over other age groups because the risk of severe illness from COVID-19 increases with age.</li> <li>• Healthcare professionals should consider not only balancing measures of risk of isolation but also appropriate psychological interventions to reduce anxiety and depression.</li> <li>• Additional precautions and quarantine must be guaranteed in long-term care facilities.</li> </ul>   | Older adults are encouraged to get vaccinated.  |

infected; facilitate communications; provide supplies and resources; identify and manage unwell residents, provide occupational health education and training; and focus on considerations regarding visitors and surge capacities, including a contingency staffing plan) [54]. Refer to the “Strategies to prevent the spread of COVID-19 in long-term care facilities” (available from: <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/prevent-spread-in-long-term-care-facilities.html>) and “Infection prevention and control for COVID-19: Interim guidance for long-term care homes” (available from: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/prevent-control-covid-19-long-term-care-homes.html>) for more details.

In summary, we described the guidelines and specific healthcare considerations for special populations at high risk of infection, including children, pregnant and lactating women, and older adults, during the pandemic period. Table 1 presents the core summaries of specific considerations for each population group.

## CONCLUSIONS

The COVID-19 pandemic continues to threaten human health and public safety, and people of all ages appear to be susceptible to SARS-CoV-2 infection. However, the infection’s clinical manifestations differ among groups. This review described the nursing practice guidelines and specific healthcare considerations for special populations, who might have unique healthcare needs, in preventing and managing COVID-19. Nurses should be educated with accurate information to maximize protection against COVID-19, including the Delta variants, and to prevent spreading it to others. However, this review is limited by the scarcity of information. Hence, further research is required to confirm the accuracy of the presented data before providing the information to patients or performing counseling.

## CONFLICTS OF INTEREST

The authors declared no conflict of interest.

## ACKNOWLEDGEMENTS

None.

## FUNDING

This study was supported by Soonchuhyang University.

## DATA SHARING STATEMENT

Please contact the corresponding author for data availability.

## AUTHOR CONTRIBUTIONS

Conceptualization or/and Methodology: Kim JI & Im Y & Song JE & Jang SJ.

Data curation or/and Analysis: Kim JI & Im Y & Song JE & Jang SJ.

Funding acquisition: Kim JI.

Investigation: Kim JI & Im Y & Song JE & Jang SJ.

Project administration or/and Supervision: Kim JI & Im Y & Song JE & Jang SJ.

Resources or/and Software: None.

Validation: Kim JI & Im Y & Song JE & Jang SJ.

Visualization: None.

Writing original draft or/and Review & Editing: Kim JI & Im Y & Song JE & Jang SJ.

## REFERENCES

- Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19. *Nature Reviews. Microbiology*. 2021; 19(3):141-154. <https://doi.org/10.1038/s41579-020-00459-7>
- Centers for Disease Control and Prevention (CDC). Information for Pediatric Healthcare Providers. [Internet]. Atlanta: CDC; c2021 [cited 2021 Jul 12]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/pediatric-hcp.html>.
- World Health Organization (WHO). Coronavirus disease (COVID-19): Pregnancy and childbirth [Internet]. Geneva: WHO; c2020 [cited 2021 Jul 12]. Available from: <https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-pregnancy-and-childbirth>.
- Centers for Disease Control and Prevention (CDC). Different groups of people [Internet]. Atlanta: CDC; c2021 [cited 2021 Aug 19]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/index.html>.
- National Institute on Aging (NIA). Government COVID-19 resources for older adults [Internet]. Bethesda: NIA; c2021 [cited 2021 Jun 28]. Available from: <https://www.nia.nih.gov/health/government-covid-19-resources-older-adults>.
- Ministry of Health and Welfare (MOHW). Case by age [Internet]. Sejong: MOHW; c2021 [cited 2021 Jun 26]. Available from: <http://ncov.mohw.go.kr/>.
- Kim JI, Yu M, Yu S, Park JH. Information and general guidance for healthcare professionals in the fourth wave of COVID-19. *Journal of Korean Academy of Nursing*. 2021; 51(4):395-407. <https://doi.org/10.4040/jkan.21137>
- Centers for Disease Control and Prevention (CDC). COVID-19 vaccines while pregnant or breastfeeding [Internet]. Atlanta: CDC; c2021 [cited 2021 Jul 12]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/recommendations/pregnancy.html>.
- Ministry of Health. COVID-19 vaccination recommendations for special populations [Internet]. Toronto: Ministry of Health; c2021 [cited 2021 Jul 12]. Available from: [https://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/vaccine/COVID-19\\_vaccination\\_rec\\_special\\_populations.pdf](https://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/vaccine/COVID-19_vaccination_rec_special_populations.pdf).
- Yasuhara J, Kuno T, Takagi H, Sumitomo N. Clinical characteristics of COVID-19 in children: A systematic review. *Pediatric Pulmonology*. 2020;55(10):2565-2575. <https://doi.org/10.1002/ppul.24991>
- Assaker R, Colas AE, Julien-Marsollier F, Bruneau B, Marsac L, Greff B, et al. Presenting symptoms of COVID-19 in children: A meta-analysis of published studies. *British Journal of Anaesthesia*. 2020;125(3):e330-e332. <https://doi.org/10.1016/j.bja.2020.05.026>
- Norman M, Navér L, Söderling J, Ahlberg M, Hervius Askling H, Aronsson B, et al. Association of maternal SARS-CoV-2 infection in pregnancy with neonatal outcomes. *JAMA*. 2021;325(20):2076-2086. <https://doi.org/10.1001/jama.2021.5775>
- Salvatore CM, Han JY, Acker KP, Tiwari P, Jin J, Brandler M, et al. Neonatal management and outcomes during the COVID-19 pandemic: An observation cohort study. *The Lancet. Child & Adolescent Health*. 2020;4(10):721-727. [https://doi.org/10.1016/S2352-4642\(20\)30235-2](https://doi.org/10.1016/S2352-4642(20)30235-2)
- Singh AP, Kumar VH, Panda S. Supporting breastfeeding in 2021 and beyond-lessons from the pandemic. *Pediatric Reports*. 2021;13(2):289-301. <https://doi.org/10.3390/pediatric13020037>
- Minckas N, Medvedev MM, Adejuyigbe EA, Brotherton

- H, Chellani H, Estifanos AS, et al. Preterm care during the COVID-19 pandemic: A comparative risk analysis of neonatal deaths averted by kangaroo mother care versus mortality due to SARS-CoV-2 infection. *EClinicalMedicine*. 2021;33:100733.  
<https://doi.org/10.1016/j.eclinm.2021.100733>
16. Ministry of Education (MOE). Korea Disease Control and Prevention Agency (KDCA). Manual for education in COVID-19 pandemic [Internet]. Sejong: MOE; c2021 [cited 2021 Jul 12]. Available from: <https://www.moe.go.kr/sub/info.do?m=580101&page=580101&num=05&s=moe>.
  17. Engzell P, Frey A, Verhagen MD. Learning loss due to school closures during the COVID-19 pandemic. *Proceedings of the National Academy of Sciences of the United States of America*. 2021;118(17):e2022376118.  
<https://doi.org/10.1073/pnas.2022376118>
  18. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and management practices during coronavirus outbreaks including COVID-19: A rapid systematic review. *The Lancet. Child & Adolescent Health*. 2020;4(5):397-404.  
[https://doi.org/10.1016/S2352-4642\(20\)30095-X](https://doi.org/10.1016/S2352-4642(20)30095-X)
  19. Bignardi G, Dalmaijer ES, Anwyl-Irvine AL, Smith TA, Siugzdaite R, Uh S, et al. Longitudinal increases in childhood depression symptoms during the COVID-19 lockdown. *Archives of Disease in Childhood*. 2020;106(8):791-797.  
<https://doi.org/10.1136/archdischild-2020-320372>
  20. Loades ME, Chatburn E, Higson-Sweeney N, Reynolds S, Shafran R, Brigden A, et al. Rapid systematic review: The impact of social isolation and loneliness on the mental health of children and adolescents in the context of COVID-19. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2020;59(11):1218-1239.e3.  
<https://doi.org/10.1016/j.jaac.2020.05.009>
  21. Royal College of Obstetricians and Gynaecologists (RCOG). Coronavirus (COVID-19) infection and pregnancy. Information for healthcare professionals [Internet]. London: RCOG; c2021 [cited 2021 Jul 12]. Available from: <https://www.rcog.org.uk/coronavirus-pregnancy>.
  22. Public Health England (PHE). COVID-19 vaccination: A guide for all women of childbearing age, pregnant or breastfeeding [Internet]. London: PHE; c2021 [cited 2021 Jul 12]. Available from: <https://www.gov.uk/government/publications/covid-19-vaccination-women-of-childbearing-age-currently-pregnant-planning-a-pregnancy-or-breastfeeding/covid-19-vaccination-a-guide-for-women-of-childbearing-age-pregnant-planning-a-pregnancy-or-breastfeeding>.
  23. Wastnedge EAN, Reynolds RM, van Boeckel SR, Stock SJ, Denison FC, Maybin JA, et al. Pregnancy and COVID-19. *Physiological Reviews*. 2021;101(1):303-318.  
<https://doi.org/10.1152/physrev.00024.2020>
  24. Rasmussen SA, Smulian JC, Lednicky JA, Wen TS, Jamieson DJ. Coronavirus disease 2019 (COVID-19) and pregnancy: What obstetricians need to know. *American Journal of Obstetrics and Gynecology*. 2020;222(5):415-426.  
<https://doi.org/10.1016/j.ajog.2020.02.017>
  25. National Health Service (NHS). Pregnancy and coronavirus (COVID-19) [Internet]. London: NHS; c2021 [cited 2021 Jul 12]. Available from: <https://www.nhs.uk/conditions/coronavirus-covid-19/people-at-higher-risk/pregnancy-and-coronavirus/>.
  26. Centers for Disease Control and Prevention (CDC). Data on COVID-19 during pregnancy: Birth and infant outcomes [Internet]. Atlanta: CDC; c2021 [cited 2021 Aug 20]. Available from: <https://covid.cdc.gov/covid-data-tracker/#pregnant-birth-infant>.
  27. Centers for Disease Control and Prevention (CDC). Pregnant and recently pregnant people: At increased risk for severe illness from COVID-19 [Internet]. Atlanta: CDC; c2021 [cited 2021 Jul 12]. Available from: <https://korean.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/pregnant-people.html>.
  28. The American College of Obstetricians and Gynecologists (ACOG). COVID-19 FAQs for obstetrician-gynecologists, obstetrics [Internet]. Washington, DC: ACOG; c2020 [cited 2021 Jul 12]. Available from: <https://www.acog.org/clinical-information/physician-faqs/covid-19-faqs-for-ob-gyns-obstetrics>.
  29. López M, Gonce A, Meler E, Plaza A, Hernández S, Martínez-Portilla RJ, et al. Coronavirus disease 2019 in pregnancy: A clinical management protocol and considerations for practice. *Fetal Diagnosis and Therapy*. 2020;47(7):519-528.  
<https://doi.org/10.1159/000508487>
  30. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: A retrospective review of medical records. *Lancet*. 2020;395(10226):809-815.  
[https://doi.org/10.1016/S0140-6736\(20\)30360-3](https://doi.org/10.1016/S0140-6736(20)30360-3)
  31. López M, Gonce A, Meler E, Plaza A, Hernández S, Martínez-Portilla RJ, et al. Supplementary material for: Coronavirus disease 2019 in pregnancy: A clinical management protocol and considerations for practice [Internet]. Basel: Karger Publishers; c2020 [cited 2021 Jul 12]. Available from: [https://karger.figshare.com/articles/dataset/Supplementary\\_Material\\_for\\_Coronavirus\\_Disease\\_2019\\_in\\_Pregnancy\\_A\\_Clinical\\_Management\\_Protocol\\_and\\_Considerations\\_for\\_Practice/12471776](https://karger.figshare.com/articles/dataset/Supplementary_Material_for_Coronavirus_Disease_2019_in_Pregnancy_A_Clinical_Management_Protocol_and_Considerations_for_Practice/12471776).
  32. Centers for Disease Control and Prevention (CDC). Considerations for inpatient obstetric healthcare settings [Internet].

- Atlanta: CDC; c2020 [cited 2021 Jul 12]. Available from: [www.cdc.gov/coronavirus/2019-ncov/hcp/inpatient-obstetric-healthcare-guidance.html](http://www.cdc.gov/coronavirus/2019-ncov/hcp/inpatient-obstetric-healthcare-guidance.html).
33. Department of Health & Social Care (DHSC). Joint Committee on Vaccination and Immunisation: Advice on priority groups for COVID-19 vaccination, 30 December 2020 [Internet]. London: DHSC; c2021 [cited 2021 Jul 12]. Available from: <https://www.gov.uk/government/publications/priority-groups-for-coronavirus-covid-19-vaccination-advice-from-the-jcvi-30-december-2020/joint-committee-on-vaccination-and-immunisation-advice-on-priority-groups-for-covid-19-vaccination-30-december-2020>.
  34. Shimabukuro TT, Kim SY, Myers TR, Moro PL, Oduyobo T, Panagiotakopoulos L, et al.; CDC v-safe COVID-19 Pregnancy Registry Team. Preliminary findings of mRNA Covid-19 vaccine safety in pregnant persons. *The New England Journal of Medicine*. 2021;384(24):2273-2282. <https://doi.org/10.1056/NEJMoa2104983>
  35. Collier AY, McMahan K, Yu J, Tostanoski LH, Aguayo R, Ansel J, et al. Immunogenicity of COVID-19 mRNA vaccines in pregnant and lactating women. *JAMA*. 2021;325(23):2370-2380. <https://doi.org/10.1001/jama.2021.7563>
  36. Korea Disease Control and Prevention Agency (KDCA). Vaccination for COVID-19 in Korea [Internet]. Cheongju: KDCA; c2021 [cited 2021 Jul 27]. Available from: <https://ncv.kdca.go.kr/menu.es?mid=a12206000000>.
  37. World Health Organization (WHO). Rational use of personal protective equipment for coronavirus disease (COVID-19) and considerations during severe shortages [Internet]. Geneva: WHO; c2020 [cited 2021 Jul 12]. Available from: [https://www.who.int/publications-detail/rational-use-of-personal-protective-equipment-for-coronavirus-disease-\(covid-19\)-and-considerations-during-severe-shortages](https://www.who.int/publications-detail/rational-use-of-personal-protective-equipment-for-coronavirus-disease-(covid-19)-and-considerations-during-severe-shortages).
  38. Centers for Disease Control and Prevention (CDC). Coronavirus disease (COVID-19) and breastfeeding [Internet]. Atlanta: CDC; c2021 [cited 2021 Jul 12]. Available from: [www.cdc.gov/breastfeeding/breastfeeding-special-circumstances/maternal-or-infant-illnesses/covid-19-and-breastfeeding.html](http://www.cdc.gov/breastfeeding/breastfeeding-special-circumstances/maternal-or-infant-illnesses/covid-19-and-breastfeeding.html).
  39. World Health Organization (WHO). Ageing and health [Internet]. Geneva: WHO; c2018 [cited 2021 Jun 28]. Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.
  40. Lee S, Moon Y. A study of the financial projection of health expenditures of the aged of national health insurance - focused on the healthy ageing of EU. *Journal of Critical Social Policy*. 2018;58:53-93. <https://doi.org/10.47042/ACSW.2018.02.58.53>
  41. Wang Y, Pang SC, Yang Y. A potential association between immunosenescence and high COVID-19 related mortality among elderly patients with cardiovascular diseases. *Immunity & Ageing*. 2021;18(1):25. <https://doi.org/10.1186/s12979-021-00234-z>
  42. Division of Population Health, National Center for Chronic Disease Prevention and Health Promotion. COVID-19 recommendations for older adults [Internet]. Atlanta: Centers for Disease Control and Prevention (CDC); c2021 [cited 2021 Jun 28]. Available from: <https://www.cdc.gov/aging/covid19-guidance.html>.
  43. WHO Regional Office for the Western Pacific. Guidance on COVID-19 for the care of older people and people living in long-term care facilities, other non-acute care facilities and home care [Internet]. Manila: WHO Regional Office for the Western Pacific; c2020 [cited 2021 Jun 28]. Available from: <https://apps.who.int/iris/handle/10665/331913>.
  44. National Health Service (NHS). Other conditions and coronavirus (COVID-19) [Internet]. London: NHS; c2021 [cited 2021 Jun 28]. Available from: <https://www.nhs.uk/conditions/coronavirus-covid-19/people-at-higher-risk/other-conditions-and-coronavirus>.
  45. Garg S, Kim L, Whitaker M, O'Halloran A, Cummings C, Holstein R, et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 - COVID-NET, 14 states, March 1-30, 2020. *MMWR. Morbidity and Mortality Weekly Report*. 2020; 69(15):458-464. <https://doi.org/10.15585/mmwr.mm6915e3>
  46. Public Health England (PHE). COVID-19 vaccine surveillance report - week 26 [Internet]. London: PHE; c2021 [cited 2021 Jul 6]. Available from: <https://www.gov.uk/government/publications/covid-19-vaccine-surveillance-report>.
  47. Christie A, Henley SJ, Mattocks L, Fernando R, Lansky A, Ahmad FB, et al. Decreases in COVID-19 cases, emergency department visits, hospital admissions, and deaths among older adults following the introduction of COVID-19 vaccine - United States, September 6, 2020-May 1, 2021. *MMWR. Morbidity and Mortality Weekly Report*. 2021;70(23):858-864. <https://doi.org/10.15585/mmwr.mm7023e2>
  48. Seow J, Graham C, Merrick B, Acors S, Pickering S, Steel KJA, et al. Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. *Nature Microbiology*. 2020;5(12):1598-1607. <https://doi.org/10.1038/s41564-020-00813-8>
  49. Donovan NJ, Blazer D. Social isolation and loneliness in older adults: Review and commentary of a National Academies report. *The American Journal of Geriatric Psychiatry*. 2020;28(12):1233-1244. <https://doi.org/10.1016/j.jagp.2020.08.005>

50. Santini ZI, Jose PE, York Cornwell E, Koyanagi A, Nielsen L, Hinrichsen C, et al. Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): A longitudinal mediation analysis. *The Lancet. Public Health.* 2020;5(1):e62–e70. [https://doi.org/10.1016/S2468-2667\(19\)30230-0](https://doi.org/10.1016/S2468-2667(19)30230-0)
51. Vernon MJ. Identifying older people most vulnerable to COVID-19 [Internet]. London: British Geriatrics Society; c2020 [cited 2021 Jul 6]. Available from: <https://www.bgs.org.uk/resources/identifying-older-people-most-vulnerable-to-covid-19>.
52. Hippisley-Cox J, Coupland C. Development and validation of QMortality risk prediction algorithm to estimate short term risk of death and assess frailty: Cohort study. *The BMJ.* 2017;358:j4208. <https://doi.org/10.1136/bmj.j4208>
53. Yildirim H, Işık K, Aylaz R. The effect of anxiety levels of elderly people in quarantine on depression during COVID-19 pandemic. *Social Work in Public Health.* 2021;36(2):194–204. <https://doi.org/10.1080/19371918.2020.1868372>
54. Centers for Disease Control and Prevention (CDC). Coronavirus disease 2019 (COVID-19) preparedness checklist for nursing homes and other long-term care settings [Internet]. Atlanta: CDC; c2021 [cited 2021 Jul 6]. Available from: [https://www.cdc.gov/coronavirus/2019-ncov/downloads/novel-coronavirus-2019-Nursing-Homes-Preparedness-Checklist\\_3\\_13.pdf](https://www.cdc.gov/coronavirus/2019-ncov/downloads/novel-coronavirus-2019-Nursing-Homes-Preparedness-Checklist_3_13.pdf).

#### Appendix 1. Distribution of Confirmed COVID-19 Cases and Deaths and Lethality Data across Age Groups

| Age group (yr) | Number of confirmed cases n (%) | Number of deaths n (%) | Lethality (%) |
|----------------|---------------------------------|------------------------|---------------|
| > 80           | 5,881 (3.81)                    | 1,099 (54.62)          | 18.69         |
| 70~79          | 10,274 (6.65)                   | 573 (28.48)            | 5.58          |
| 60~69          | 22,880 (14.81)                  | 241 (11.98)            | 1.05          |
| 50~59          | 28,348 (18.35)                  | 73 (3.63)              | 0.26          |
| 40~49          | 23,755 (15.38)                  | 15 (0.75)              | 0.06          |
| 30~39          | 21,608 (13.99)                  | 8 (0.40)               | 0.04          |
| 20~29          | 23,564 (15.26)                  | 3 (0.15)               | 0.01          |
| 10~19          | 11,142 (7.21)                   | 0 (0.00)               | -             |
| 0~9            | 7,005 (4.54)                    | 0 (0.00)               | -             |

- = Not available; Lethality = Number of deaths/number of confirmed cases × 100.

(Cited from the article of Ministry of Health and Welfare (MOHW). Case by age [Internet] Sejong: MOHW; c2021 [cited 2021 Jun 26]. Available from: <http://ncov.mohw.go.kr/>) [6].