



ARTICLE

Incorporation of Selected Strains of *Pediococcus* spp. on Quality Characteristics of Dry Fermented Sausage during Fermentation and Ripening

Semeneh Seleshe, Ammara Ameer, and Suk Nam Kang*

Department of Animal Resource, Daegu University, Gyeongsan 38453, Korea

OPEN ACCESS

Received July 16, 2021

Revised October 6, 2021

Accepted October 13, 2021

*Corresponding author : Suk Nam Kang
Department of Animal Resource, Daegu
University, Gyeongsan 38453, Korea
Tel: +82-53-850-6726
Fax: +82-53-850-6729
E-mail: sk-kang@daegu.ac.kr

*ORCID
Semeneh Seleshe
<https://orcid.org/0000-0002-2599-393X>
Ammara Ameer
<https://orcid.org/0000-0002-8110-2433>
Suk Nam Kang
<https://orcid.org/0000-0002-9230-3070>

Abstract This research investigated the physio-chemical and microbial quality characteristics of dry fermented sausage from selected *Pediococcus* strains: *P. acidilactici* (PE1) and *P. pentosaceus* (PE2) as compared to commercial starter culture (COS) during fermentation and ripening. Treatments showed no substantial variation ($p < 0.05$) in water activity (a_w) values across the study period. PE1 and PE2 treatments exhibited similar ($p > 0.05$) pH values and presented remarkable ($p < 0.05$) lower volatile basic nitrogen (VBN) and thiobarbituric acid reactive (TBARS) content than COS treatment throughout the ripening period. However, the pH values in COS batch were considerably lower than others. PE1 samples presented a significant highest ($p < 0.05$) counts both in lactic acid bacteria (LAB) and total plate count (TPC) than COS and PE2 treatments at 7 days fermentation, and it resulted in a similar and higher TPC count as COS after the ripening period. After the ripening process, treatments are ordered based on LAB counts as follows: COS > PE1 > PE2. All batches presented similar redness and yellowness attributes since the 7 days of fermentation and in lightness across the study period. Treatments were similar ($p > 0.05$) in springiness and chewiness traits across the study period and in hardness characteristics in the ripened products. Cohesiveness was higher in PE1 and COS batches. No variation ($p > 0.05$) in aroma and sourness sensory attributes of treatments. The color attribute was highest ($p < 0.05$) in PE1 and PE2 treatments and PE1 had the highest overall acceptability. The overall outstanding merit exhibited by PE1 can be utilized in the commercial production of high-quality dry fermented sausage.

Keywords dry fermented sausage, starter cultures, strains, ripening, water activity

Introduction

Fermented meat products have long been regarded as prestigious foods (Leroy and Vuyst, 2016), and they are still manufactured in large quantities, particularly in Europe, owing to their peculiar and unique sensory attributes, versatility, and alleged ties to culinary and traditional culture (Leroy et al., 2013). They are a component of traditional menus in certain parts of the world, where they are regarded as appealing gastronomic entities that contribute to geographical and cultural uniqueness (Leroy and Vuyst,

2016). According to Casaburi et al. (2007), complex physicochemical reactions occur during the fermentation and aging of sausages, resulting in a decline in pH and shifts in microbial communities. Among different requirements, acidification through fermentation by lactic acid bacteria (LAB) and subsequent slow drying/ripening often at room temperature enhancing long shelf-life (Barbut, 2011), more nutritious and substantial and unique flavor of dry fermented meat products.

A real breakthrough in the fermentation of meat products was achieved by the use of pure bacterial cultures in the 1950s (Incze, 1998). The use of starter cultures, along with the processing environment (temperature and relative humidity conditions), are the critical features operated by the fermented sausage industries to enhance the safety and stability of their products, both of which are amongst the major drivers of the dynamic phenomena that occur during ripening (Pasini et al., 2018). During the manufacturing of dry fermented sausage, well-fitting starter cultures must be chosen from microbes that are well suited to the ingredients used in preparation of the meat batter and processing environment, and should be more versatile due to their special metabolic capabilities (Leroy et al., 2006) in order to dominate the process.

LAB and *Staphylococcus* species are the major microbial groups of technological concern in fermented sausage production (Morot-Bizot et al., 2006). LAB play a key role by producing lactic acid and lowering pH, which affects the ultimate product's physicochemical properties as well as its microbiological stability (Drosinos et al., 2007). The pH reduction facilitates also the process of ripening which improves the moisture content and delivers the ultimate dry characteristics of fermented meat products. The production of lactic acid and other antimicrobials metabolites by LAB plays a crucial role in organoleptic attributes of aroma, taste, and texture of meat products, and are vital in hygiene of the product by inhibiting the activity of pathogenic and spoilage microorganisms (Villani et al., 1994).

Amongst the LAB species, *Lactobacillus sakei*, *Lactobacillus curvatus*, *Lactobacillus plantarum*, *Pediococcus acidilactici*, and *Pediococcus pentosaceus* are widely used starter cultures in the fermentation of meat products (Diguță et al., 2020; Hugas and Monfort, 1997). *Pediococcus* species have gained growing attention and are considered as competitive candidates in the fermented sausages industries now days due to their functional efficiency and probiotic features (Diguță et al., 2020), and their antagonistic impact on other microorganisms, including food pathogens, through the production secretion of lactic acid and bacteriocins known as pediocins (Daeschel and Klaenhammer, 1985).

There has been no comparative research on the performance traits between *P. acidilactici* (PE1) and *P. pentosaceus* (PE2) strains on quality characteristics of dry fermented sausages to date. We believe that assessing the functional characteristics of individual strains aids in the selection and production of high-quality fermented meats on an industrial level, whether they are used as a single strain or as a combination of strains. This research, therefore, investigated the effects of PE1 and PE2 strains of *Pediococcus* spp. (LAB) as compared to commercial starter culture (COS), Starterkulturen Almi Rohschinken, on physicochemical and microbial quality characteristics of dry fermented sausage during fermentation and ripening.

Materials and Methods

Starter culture preparation

The two different starter cultures from *Pediococcus* species of LAB employed in manufacturing of two different types of dry fermented sausages, used as PE1 and PE2 treatments, in the present study were obtained from the Microbial Resource Center, Korea Research Institute of Bioscience and Biotechnology, Daejeon, Korea as lyophilized stocks. The LAB strains were: PE1 and PE2. The COS, *Starterkulturen Almi Rohschinken* (Almi GmbH & Co KG, Oftring, Austria) was used as the third treatment and positive control (COS). The starter cultures were enriched with MRS broth (Difco, Becton, NJ, USA) by

incubating at 37°C for 24 h. The designed suspension incorporated into the sausage batter was one mL/kg (v/w) and each starter culture had approximately 6 Log CFU/g. The viable cell count in the starter cultures suspensions was performed using a hemocytometer (Marienfeld-Superior, Paul Marienfeld GmbH & Co.KG, Lauda-Königshofen, Germany) supported with a computer magnification system.

Dry fermented sausages manufacture and sampling

Low-temperature dry fermented sausages were produced in the pilot meat processing center, Animal Resources Department, Daegu University. Fresh loin pork was purchased from the commercial market of Geyongsan, Korea, and used for the study. After trimming the connective tissues and excess fat, the lean meat was kept in a refrigerator for further use. Chilled pork and pork fat were cut into small cubes and minced twice using a meat mincer (M-12S, Hankook Fujee Industries, Suwon, Korea). The basic sausage formulation includes lean pork meat (65%), pork fat (21.5%), ice water (10%), NPS (97:3, blend of NaCl and nitrite; 0.34%), NaCl (1.70%), sugar (0.45%), glucose (0.45%), sodium ascorbate (0.20%), and sausage seasonings (0.36%). After adequately mixing the ingredients with help of a rotary slice cutter (SF-2002, Samwoo Industry, Korea), the batter was divided into three lots (3 kg each) and randomly allocated to 3 distinct starter cultures treatments: COS, PE1, and PE2. The supreme starter cultures (LAB) concentration reached $\sim 10^6$ CFU/g when introduced to the meat batters, and one mL/kg (v/w) starter culture was added for each respective treatment. The batters and corresponding starter cultures were thoroughly blended with the help of a food mixer (SP-100A, Spar Food Machinery MFG, Taiwan) and filled into collagen casings (IKJIN, Seoul, Korea), 2.4 cm diameter and 15 cm length, with vacuum filling machine (RVF 327, Düker-REX Fleischereimaschinen GmbH, Laufach, Germany). The sausages were fermented and ripened in a digital chamber system (SMK-2000SL, Metatek, Nonsan, Korea) with a temperature and relative humidity (RH) control unit. During the fermentation process (the first seven days), the temperature was kept at 23°C and RH was varied between 90% and 95%. Subsequently, the ripening process was performed for 14 days (after the fermentation process) at 15°C (the temperature was decreased slowly from 23°C), and RH oscillated from 70% to 75%. The manufacturing process of dry fermented sausages is displayed by a schematic diagram (Fig. 1). Sampling was done at 1 and 7 days of fermentation and at 7 and 14 days of ripening for water activity (a_w), pH, salinity, microbial quality [total plate count (TPC) and MRS counts], and color attributes L^* (lightness), a^* (redness), and b^* (yellowness) of sausages samples. The volatile basic nitrogen (VBN) content, thiobarbituric acid reactive (TBARS), and texture profile analysis (TPA) were performed at 7 and 14 days of ripening. For each analysis time and batches, three replicates samples were withdrawn, and analyses were performed accordingly.

Microbial quality analysis

Microbiological quality characteristics were conducted by enumeration of TPC and LAB. About 25 g portion of a sample from each dry fermented sausage was taken aseptically with a sterile spoon, mixed with 225 mL of 0.1% peptone water, and homogenized in a Stomacher Lab Blender (model 400 Circulator, Seward Laboratory, USA) for 2 min. Serial 10-fold dilutions (10^{-1} to 10^{-7}) were prepared by diluting one mL of the sample in nine mL of 0.1% sterile peptone water. Enumerations of the grown colony of microorganisms were conducted after incubating samples with their respective selective medium: Plate Count Agar (Difco) was used for TPCs and *Lactobacillus* MRS agar (Difco) for LAB. Plates from different and appropriate dilutions were incubated in triplicate at 37°C for 48 h (Drosinos et al., 2005). The average numbers of colonies per countable plate were counted and the total numbers of colonies per gram (CFU/g) were determined, and then data were presented in Log CFU/g.




Fig. 1. Schematic diagram for the manufacturing process of dry fermented sausages.

Analyses of pH, salt content and water activity (a_w)

Analysis of a_w was performed using a_w measuring device (LabMaster- a_w , Novasina AG, Lachen, Switzerland) after slicing the core of the samples about 4 mm cubes. The pH values of samples were analyzed after homogenizing three grams of a sample with 30 mL of distilled water using a homogenizer (Model Polytron® PT 2500 E Stand Dispersion Device, Kinematica AG, Switzerland). A digital pH meter (Mettler Toledo, Columbus, OH, USA) was used for reading the values. The same homogenates of samples used in pH analysis were subjected to salt (NaCl) content analysis using a salinity measuring device (EUTECH SALT 6+, Thermo Scientific, Gul Cir, Singapore). Three readings per sample were performed and the percentage (%) of salt content was calculated.

Volatile basic nitrogen (VBN) analysis

For the determination of VBN contents, the Conway micro diffusion method (Conway, 1950) was followed after some modifications. Briefly, two cleaned Conway's apparatus were used for each sample and vaseline was applied as sealing agent to the edge of the outer ring. Three grams of the sample and 30 mL of distilled water was homogenized at 1,000×g for 1 min

by using a homogenizer. Filtration of the homogenate was performed with the help of Whatman no. 1 filter paper (GE Healthcare Life Sci., Pittsburg, PA, USA). One mL the filtrate was pipetted to the outer chamber of a Conway unit, and 1 mL of 0.01 N boric acid (H_3BO_3) and 100 μ L of Conway indicator (0.066% bromocresol green:0.066% methyl red, 1:1) were pipetted to the inner chamber. Sealing the Conway unit was done immediately after the addition of 1 mL of 50% potassium carbonate (K_2CO_3) in the outer chamber. Then, samples were incubated at 37°C for 2 h. Finally, 0.02 N sulfuric acid (H_2SO_4) was added to the inner chamber of the Conway unit, the change in color was monitored, and the VBN contents was measured. Total VBN values were expressed in mg%.

Thiobarbituric acid reactive (TBARS) analysis

Measurement of lipid oxidation, TBARS content Analysis, was conducted using the method indicated by Pikul et al. (1989) after some modification. Briefly, 5 g of sausage sample, 50 μ L of BHA (7.2% in ethanol) and 15 mL of distilled water was homogenized and then centrifuged with the help of a centrifuge (Hanil Science Industrial, Incheon, Korea) at 1,500 \times g for 15 min. About 2 mL of the supernatant and 4 mL thiobarbituric acid solution (20 mM TBA in 15% Trichloroacetic acid, TCA) was mixed, heating the mixture was performed in a water bath at 90°C for 30 min and then cooled to room temperature. Therefore, extraction of TBARS was done from cooled samples and the absorbance was measured at 532 nm using a spectrophotometer (Multiskan Go, Thermo Fisher Scientific, MA, USA). TBARS content, mg malonaldehyde per kg (mg MA/kg), of each sausage sample was calculated by multiplying the reading of the optical density with a K factor of 5.2.

Instrumental color analysis

Instrumental color analysis was performed from the inner surface of the sliced sausages using a portable chromameter (CR-400, Konica Minolta, NJ, USA). The device was calibrated with a standard calibration plate ($Y=92.80$, $x=0.3136$, and $y=0.3194$) ahead of the analysis, and five readings per sample were taken for L^* , a^* , and b^* . Average values were calculated from five readings and expressed as L^* , a^* , and b^* based on the CIE color system (CIE, 1976).

Texture profile analysis (TPA)

TPA was performed using a TPA measuring device (TA 1, Lloyd Instruments, Largo, FL, USA). During the study sessions, three lots of samples (1 cm in height and 2 cm in diameter) from each batch were compressed 80% vertically for two consecutive compression cycles by exerting a 1 N load cell at a crosshead moved at 2 mm/s. And then, the TPA characteristics were obtained from force-time deformation curves for hardness (kgf), springiness, cohesiveness, and chewiness (kgf) of sausage samples.

Sensorial analysis

Sensory quality analysis was conducted for color, aroma, sourness, and overall acceptability of dry fermented sausages using descriptive sensory analysis (scoring method). The sensory evaluation was performed by seven experienced panelists who are searchers and students in Daegu university's department of animal resources, meat science laboratory. Ahead of the actual evaluation session, the panelists were trained on sensory characteristics of dry fermented sausages using five-point scale. The intensity scale used to define the quality attributes ranged from 1 to 5 that corresponds to the sensory attributes of samples as follows "extremely pale to very dark," for color, "very weak fermented aroma to very strong fermented aroma,"

for aroma, and “light sour to strong sour” for sourness. Three different types of commercial dry fermented sausage were used during training session, and panel were given 3 slices (5 mm thickness) of samples on white plastic plates during the judgment. To avoid carryover influences, all samples were individually labeled with three digits and provided at random. Before each sample was examined, the panel were provided cold water to rinse their mouths. The sensory analysis method was certified by the life management committee of Daegu University and given an IRB number (1040621-201905-HR-004-02).

Statistical analysis

Statistical data were analyzed by using Analysis Variance (ANOVA) for the three replicates. SAS software version 9.4 (SAS Institute, Cary, NC, USA) was used for the analysis, and a significance level of $p < 0.05$ was applied for all evaluations. Differences among the means were compared according to Duncans’s Multiple Range Test.

Results and Discussion

Changes in a_w , pH and salt content

The shelf-life of a product is entirely influenced by its a_w content, and low a_w is well recognized for enhancing the storage stability of foods. Table 1 shows the changes in the a_w values of DFS incorporated with different *Pediococcus* strains (PE1 and PE2) and the control (COS) during fermentation and ripening. The two experimental treatments, PE1 and PE2, revealed no difference in a_w values throughout the study period ($p > 0.05$), and they have a similar effect to the control, COS, which is utilized as a starter culture in a commercial application. However, the fermentation and ripening time have a considerable effect ($p < 0.05$) on the a_w of all treatments. Treatments exhibited the same trend of declining in a_w as fermentation progressed from day one towards an extended period of ripening. In the early fermentation of day one, all three treatments presented the same a_w value of 0.95. Consequently, the a_w declined and ranged from 0.79 to 0.81 at the end of the ripening process. Based on the final a_w values at the end of the ripening period, according to Luecke (1997), fermented sausage can be categorized into “semi-dry” or “dry”. Sausages with a_w values in the ranges between 0.90 and 0.95 fall into the “semi-dry” category, whereas sausages having a_w less than 0.90 are referred to as “dry”. Considering this classification criterion, all three sausages of the present study can be grouped into dry sausages. Due to the lower a_w , such dry fermented sausages possibly have a long shelf life and can also be stored without cold storage.

In sausage production, a lower pH value is crucial because it helps to prevent the growth of adverse microorganisms, improves the redder color (Lorenzo et al., 2014), and enhances the development of the finished product's distinct flavor and aroma (Hugas and Monfort, 1997). The changes in pH of fermented sausages during fermentation and ripening are presented in Table 1. According to Valyasevi et al. (2001), the production of lactic acid by LAB is mainly responsible for the reduction in pH and rise in acidity during fermentation. At the early stage of day one fermentation, the type of starter culture had no variation in pH value and all three treatments had a similar acidification effect. The pH values between the test samples, PE1 and PE2, were similar ($p > 0.05$) across the fermentation and ripening period. However, inoculation of COS resulted in a considerably lower pH value than two products from the *Pediococcus* strain of starter cultures during the late stage of fermentation (seven-day fermentation) and throughout the ripening period. According to the Food Safety and Inspection Service of the USA, shelf-stable dry fermented sausages are required to reach a final pH of 5 or less (HORIBA, 2021). In the present study, all three batches achieved a pH of lower than 5 after the completion of the maturation. The lower pH values are often used to confirm the starter cultures’ ability to induce strong acidification (Yim et al., 2017). The exhibited lower pH

Table 1. Effect of different starter cultures on a_w , pH and salinity values of dry fermented sausages during fermentation and ripening

Attributes	Days	Treatments ¹⁾			SEM
		COS	PE1	PE2	
a_w	1 day of fermentation	0.95 ^A	0.95 ^A	0.95 ^A	0.01
	7 days of fermentation	0.91 ^B	0.92 ^B	0.92 ^B	0.00
	7 days of ripening	0.83 ^C	0.84 ^C	0.84 ^C	0.01
	14 days of ripening	0.81 ^D	0.79 ^D	0.80 ^D	0.00
	SEM	0.01	0.01	0.01	
pH	1 day of fermentation	5.26 ^A	5.21 ^A	5.15 ^A	0.04
	7 days of fermentation	4.62 ^{bb}	4.76 ^{ab}	4.76 ^{ab}	0.01
	7 days of ripening	4.57 ^{bc}	4.72 ^{ab}	4.71 ^{ab}	0.01
	14 days of ripening	4.49 ^{bd}	4.66 ^{ac}	4.68 ^{ab}	0.01
	SEM	0.01	0.01	0.04	
Salinity	1 day of fermentation	3.33 ^{ab}	3.25 ^{ad}	3.05 ^{bc}	0.04
	7 days of fermentation	5.71 ^{aa}	4.94 ^{abc}	4.43 ^{bb}	0.29
	7 days of ripening	5.94 ^{aa}	5.23 ^{cb}	5.40 ^{ba}	0.03
	14 days of ripening	6.07 ^A	6.15 ^A	6.29 ^A	0.34
	SEM	0.15	0.10	0.35	

¹⁾ Treatments are different strain of *Pediococcus* spp. used in the present study: COS, commercial starter culture; PE1, *P. acidilactici*; PE2, *P. pentosaceus*, *Starterkulturen Almi Rohschinken*, used as control (n=3).

^{a-c} Means in a row with different superscript are significantly different ($p < 0.05$).

^{A-D} Means in a column with different superscripts are significantly different ($p < 0.05$).

confirmed that the individual *Pediococcus* strains' effectiveness in the acidification process in the production of shelf-stable products. Reduced the pH and increased the lactic acid content were documented earlier by Zhao et al. (2011) in fermented sausages formulated from a mixture of *Pediococcus acidilactici*, *Staphylococcus xylosum*, and PE2. The current research revealed that the required pH can be achieved by independent use of *Pediococcus* strains as starter cultures. Regarding the times of fermentation and ripening, all treatment batches exhibited a substantial drop ($p < 0.05$) in their pH, and the change in the pH range was from (5.15–5.26) at day one fermentation to (4.62–4.76), (4.57–4.72) and (4.49–4.68) at seven-days fermentation, seven-days and fourteen-days of maturation, respectively. According to Baka et al. (2011) investigation, the pH reduction is attributed to the production of lactic acid by LAB, which are increasingly becoming the dominant microorganisms during fermentation and ripening. Whereas, other authors documented that low pH values benefit the starter cultures' ability to cause extreme acidification (Casaburi et al., 2007; Zhao et al., 2011). In general, the a_w and pH values of the fermented sausages in the current study were in ranges that provide stability at room temperature and enhances consumer safety.

Salt (NaCl) plays a crucial role in the production of dry fermented sausages due to its effects on the required salty taste characteristics, enhancing flavor, texture, and overall acceptability. Furthermore, salt plays an essential function in reducing water activity, controlling enzymatic and metabolic activities, and contributing to microbiological safety and stability, all of which are important in increasing product shelf life (Hoppu et al., 2017). Salt promotes the solubility of the myofibrillar protein, improves the binding properties and texture, and increases the viscosity of meat batters, all of which contribute to the technological features of meat (Terrell, 1983). Commonly, fermented sausages are characterized by their high salt content.

Based on consumer demands and technological properties, the amount of salt added during fermented sausage production is normally between 2% and 4% (Vignolo et al., 2010). The experimental samples in the current study were manufactured by adding about 2.3% salt (NaCl). During fermentation and early ripening period of 7 days, significant difference ($p < 0.05$) in salt content among the three treatments was exhibited, and the COS, yielded the highest salt content than the two *Pediococcus* starters culture samples (PE1 and PE2; Table 1) at 7 d of ripening period. However, after the completion of the ripening process, the products didn't show a difference ($p > 0.05$) in the salt content. As compared to the initial fermentation time, all three treatments had a considerably higher ($p < 0.05$) salt content by the end of the ripening period, ranging from 6.07% to 6.29%. The percentages of salt in the final products of the present study could be considered acceptable as 5.09% to 9.58% (average 7.13%) was reported by Lilic et al. (2011) as normal amount for dry fermented products.

Thiobarbituric acid reactive (TBARS) and VBN contents

Studies documented that lipid oxidation can alter nutritive values, colors, and flavors of meat products (Kim et al., 2015) and poses health risks to consumers (Grun et al., 2006). The amount of aldehydes formed by lipids' secondary oxidation is measured using TBARS values. And the malondialdehyde (MDA) is examined as an indicative product. The extent of lipid oxidation is also represented in TBARS values (Chen et al., 2017). Consequently, an investigation of TBARS is considered a standardized method for assessing secondary lipid oxidation products in meat and meat products. The addition of different starter cultures resulted in a considerable variation ($p < 0.05$) in the TBARS values throughout the processing time (Table 2). Samples added with PE1 starter culture exhibited remarkably lower ($p < 0.05$) TBARS content across the study period, and samples from PE2 presented a comparable effect with PE1 treatment during the ripening period. The TBARS content of treatments was in the following order after the completion of the ripening process: PE1 < PE2 < COS, which is similar to the trend demonstrated in the VBN investigation of treatments. Marco et al. (2006) reported that the TBARS values in the range of 0.6–2.8 mg MDA/kg are reasonable for fermented sausages. The TBARS values of all three batches in the current study did not exceed the described range. Moreover, all three batches showed a decrease in TBARS content from the completion of fermentation towards the ripening process. It may be associated with the similar effect of the starter cultures interference in the oxidation process of lipids. The decline in TBARS contents of all three batches from fermentation to the final ripening stage may be resulted from the accumulation of lactic acid and the drop in pH, the decrease in a_w , and the increase in salt concentration observed in three treatments that enhanced the hurdle effect and prevented enzymatic and microbial oxidation. An earlier study by Juncher et al. (2000) documented that low pH suppresses the oxidative activity of lipid. Bingol et al. (2014) reported that lower pH values, caused by the lactic acid activities of starter cultures, had a significant impact on TBARS values. Zhao et al. (2020) noted a decreasing trend in the TBARS values of Chinese dry sausage during the later stages of ripening, which is concordant with the current finding.

Protein degradation can occur from the spoilage bacteria or intrinsic enzymatic activities, and chemical compounds such as NH_3 , H_2S , and $\text{CH}_3\text{CH}_2\text{SH}$, which are designated as VBN constituents, could be produced (Huang et al., 2014). Furthermore, protein degradation during fermentation is another important mechanism that leads to the development of non-protein nitrogenous compounds with low molecular weights (Ruiz-Capillas and Jiménez-Colmenero, 2004). Consequently, the VBN content is usually used to assess the freshness of raw meats, as well as the shelf life and microbiological quality of processed meats (Ba et al., 2018). Results for the investigation of the VBN and TBARS contents of the experimental samples are presented in Table 2. At the end of the fermentation period, PE2 and the control treatment COS had similar lower VBN values than PE1 samples. However, both *Pediococcus* strains, PE1 and PE2, presented a substantially ($p < 0.05$) lower values

Table 2. Effect of different starter cultures on VBN and TBARS values of dry fermented sausages during fermentation and ripening

Attributes	Days	Treatments ¹⁾			SEM
		COS	PE1	PE2	
VBN (mg%)	7 days of fermentation	5.60 ^{bc}	6.72 ^{aA}	6.07 ^{bB}	0.30
	7 days of ripening	6.72 ^{aB}	5.69 ^{bB}	5.60 ^{bB}	0.19
	14 days of ripening	8.40 ^{aA}	5.60 ^{cB}	6.91 ^{bA}	0.19
	SEM	0.23	0.19	0.26	
TBARS (mg MA/kg)	7 days of fermentation	1.23 ^{bA}	0.95 ^{cA}	1.29 ^{aA}	0.01
	7 days of ripening	1.11 ^{aB}	0.56 ^{cC}	1.05 ^{bB}	0.02
	14 days of ripening	0.84 ^{aC}	0.65 ^{cB}	0.80 ^{bC}	0.01
	SEM	0.02	0.01	0.02	

¹⁾ Treatments are different strain of *Pediococcus* spp. used in the present study: COS, commercial starter culture; PE1, *P. acidilactici*; PE2, *P. pentosaceus*, *Starterkulturen Almi Rohschinken*, used as control (n=3).

^{a-c} Means in a row with different superscripts are significantly different (p<0.05).

^{A-C} Means in a column with different superscripts are significantly different (p<0.05).

VBN, volatile basic nitrogen; TBARS, thiobarbituric acid reactive substance.

compared to the COS throughout the ripening period. Based on the VBN contents of treatments at the final stage of ripening, they ordered as follows: PE1<PE2<COS. The VBN values of treatments in the contemporary study ranged from 5.60 to 8.40 mg%, which is considerably lower than the content, 20–25 mg%, identified by Rai et al. (2010) for a similar product type on the 25th day of ripening. The processing time has a profound effect on the VBN contents of samples. VBN substantially (p<0.05) increased in PE2 and COS samples and decreased in PE1 treatment from 7 d of fermentation towards the end of ripening period, and the value was substantially lower (p<0.05) in PE1 than both COS and PE2. This may be resulted from the efficiency of PE1 starter culture to neutralize the VBN content with their organic acids (e.g., lactic acid) or bacteriocin production (Yin et al., 2002) that helps in the stability of the product. In contrast, the increase in the VBN content exhibited in COS and PE2 samples due to extended processing time, according to Egan et al. (1981), could be elucidated by increased protein degradation.

Microbiological quality characteristics

The selection and application of successful starter cultures are crucial factors for the production of high-quality and shelf-stable dry fermented sausages that meet consumer demand. Effect of different starter cultures addition on TPC and LAB (MRS) counts of dry fermented sausages during the fermentation and ripening period is presented in Table 3. Statistical analysis showed that treatments from different starter cultures resulted in a substantial difference (p<0.05) in TPC and LAB counts. Likewise, Essid and Hassouna (2013) found that utilizing a selective starter significantly affects the total viable, LAB, *Staphylococci*, and Enterobacteriaceae numbers in dry fermented sausage production. At the early fermentation stage of day one, sausages added with COS and PE2 had higher TPC than PE1 samples, and the count ranged from 7.40 to 8.15 Log CFU/g. However, as the processing transformed to final stage fermentation, PE1 samples overwhelmed both COS and PE2 treatments, then resulted in a similarly higher count as the COS at the final stage of ripening. The counts were about 6.84, 6.50, and 6.05 logs CFU/g in control (COS) and treatments added with PE1 and PE2, respectively, in the final products. The TPC showed fluctuation between early fermentation and ripening and steadily decreased subsequently for all the batches after ripening. The decline in the counts for all treatments exhibited in the present study might be associated with the increase in

Table 3. Effect of different starter cultures on total plate count (TPC) and lactic acid bacteria (LAB) counts of dry fermented sausages during fermentation and ripening time

Attributes	Days	Treatments ¹⁾			SEM
		COS	PE1	PE2	
TPC	1 day of fermentation	8.15 ^{AA}	7.40 ^{BB}	7.97 ^{AA}	0.27
	7 days of fermentation	7.61 ^{BB}	8.39 ^{AA}	7.42 ^{BB}	0.40
	7 days of ripening	7.25 ^{BC}	7.33 ^B	7.76 ^A	0.28
	14 days of ripening	6.84 ^{AC}	6.50 ^{AC}	6.05 ^{BC}	0.28
	SEM	0.33	0.39	0.18	
LAB	1 day of fermentation	6.83 ^B	7.09 ^C	7.23 ^B	0.24
	7 days of fermentation	7.51 ^{BA}	8.72 ^{AA}	7.65 ^{BA}	0.59
	7 days of ripening	6.42 ^{BC}	7.81 ^{AB}	7.96 ^{AA}	0.10
	14 days of ripening	6.71 ^{ABC}	6.47 ^{BD}	5.92 ^{CC}	0.12
	SEM	0.22	0.34	0.40	

¹⁾ Treatments are different strain of *Pediococcus* spp. used in the present study: COS, commercial starter culture; PE1, *P. acidilactici*; PE2, *P. pentosaceus*, *Starterkulturen Almi Rohschinken*, used as control (n=3).

^{a-c} Means in a row with different superscripts are significantly different (p<0.05).

^{A-D} Means in a column with different superscripts are significantly different (p<0.05).

TPC, total plate count; LAB, lactic acid bacteria.

salt concentration, as the drying process advanced, and the lower a_w and pH values. Regarding LAB counts of samples, the addition of different starter cultures had a significant (p<0.05) effect during late fermentation and throughout the ripening period. At the final fermentation period, samples added with PE1 showed significantly highest LAB counts compared to both COS and PE2 treatments. The LAB counts were between 7.51 Log CFU/g for the control COS starter culture and 8.72 Log CFU/g for sausage incorporated with a PE1 starter culture. As the processing stage advanced to 7 days of ripening, both treatments added with *Pediococcus* strains, PE1 and PE2, resulted in similar and substantially higher (p<0.05) counts than a COS or the control COS. However, after the ripening period, the COS batch presented the highest count than other treatments. Treatments are ordered based on LAB counts as follows: COS>PE1>PE2 with corresponding values of 6.71, 6.47, and 5.92 Log CFU/g, respectively, at the end of the ripening process. The LAB counts at the end of the ripening were lower in all samples compared to the end of the fermentation period. The rapid growth of LAB during fermentation is normally expected as incorporated starter cultures induce the LAB to become dominant, resulting in a massive bacterial population (Lim et al., 2008). Similar to the TPC counts, all three batches showed fluctuation in LAB counts due to the effect of the processing stage, and a steady decrease in counts was exhibited between the early fermentation stage and the completion of ripening time for all three batches. TPC and LAB counts showed an almost similar trend (r=0.86, p>0.05: data not presented) that is concurring with the previous finding of Lim et al. (2008).

Instrumental color characteristics

The color attribute of meat products has a significant impact on consumer perceptions and is the most important component of product quality. Consequently, a product may be rejected solely on the basis of its color before any other attributes are considered. Different factors govern the color of dry fermented sausages, including the sausage's composition, fat-to-lean ratio, amount and type of spices and additives, and technical operations followed (Perez-Alvarezet and Fernandez-Lopez, 2011). The

results for the instrumental color characteristics of dry fermented sausages inoculated with different types of starter cultures during the fermentation and ripening period are presented in Table 4. The addition of different starter cultures showed no difference ($p>0.05$) in the lightness parameter throughout the fermentation and ripening study. Treatments displayed a similar trend in the redness and yellowness attributes and both parameters were significantly affected by the inoculation of different starter cultures during the initial fermentation period of day one. Samples from PE2 starter culture had markedly increased values than the control, COS, and PE1 treatments in both attributes. However, the difference among samples disappeared, and all batches showed similar a^* and b^* characteristics since the 7th days of the fermentation period. The fermentation and ripening time have a noticeable effect on L^* value of treatments. Substantial decrease in L^* was exhibited in all treatments due to extended fermentation and ripening time. A similar finding was reported in an earlier study by Kurt (2016) that L^* value is significantly affected due to ripening time. The possible reason for the decreased lightness value might be associated with loss of moisture, especially the surface drying of the sausage samples. The present study exhibited that both a^* and b^* characteristics were not affected ($p>0.05$) due to fermentation and ripening times. Redness is the most sensitive parameter, and its intensity indicates the meat products' stability against oxidation (Ergezer et al., 2018). An earlier study documented that the redness trait decrease during maturation (Cavalheiro, 2013). However, the current study did not exhibit the reported trend, and all batches showed no marked variation due to ripening time. The finding implies that all of the starter cultures utilized excel solely in the manufacture of dry fermented sausages in terms of their effect on color features.

Table 4. Effect of different starter cultures on color characteristics of dry fermented sausages during fermentation and ripening

Attributes	Days	Treatments ¹⁾			SEM
		COS	PE1	PE2	
L^*	1 day of fermentation	64.95 ^A	64.59 ^A	64.59 ^A	2.69
	7 day of fermentation	64.55 ^A	64.41 ^A	63.51 ^A	1.61
	7 day of ripening	59.12 ^B	54.68 ^B	57.11 ^B	1.85
	14 day of ripening	56.58 ^B	53.58 ^B	57.36 ^B	3.50
	SEM	2.31	2.47	2.77	
a^*	1 day of fermentation	6.77 ^b	7.70 ^b	8.89 ^a	0.99
	7 days of fermentation	6.81	7.45	6.98	1.31
	7 days of ripening	7.99	9.544	7.18	1.48
	14 days of ripening	7.79	8.74	7.43	1.48
	SEM	1.27	1.31	1.40	
b^*	1 day of fermentation	7.97 ^b	8.28 ^b	9.23 ^a	0.64
	7 days of fermentation	8.23	8.75	7.99	0.52
	7 days of ripening	7.65	7.84	8.14	1.31
	14 days of ripening	9.20	7.99	9.28	1.07
	SEM	1.03	0.90	0.88	

¹⁾ Treatments are different strain of *Pediococcus* spp. used in the present study: COS, commercial starter culture; PE1, *P. acidilactici*; PE2, *P. pentosaceus*, *Starterkulturen Almi Rohschinken*, used as control (n=3).

^{a,b} Means in a row with different superscripts are significantly different ($p<0.05$).

^{A,B} Means in a column with different superscripts are significantly different ($p<0.05$).

Texture profile analysis (TPA)

Effects of different starter cultures incorporation on the texture profile analysis (TPA) of dry fermented sausage during fermentation and ripening are described in Table 5. No substantial difference ($p>0.05$) was exhibited among all batches in springiness and chewiness characteristics across the fermentation and ripening period. The hardness property of all three treatments did not show marked difference ($p>0.05$) during the fermentation study. However, COS and PE2 samples showed similarly and substantially higher ($p<0.05$) hardness values than PE1 on the 7th day of ripening. Subsequently, the variation disappeared again, and all the batches showed similar hardness after the completion of the ripening period. Cohesiveness is a characteristic that shows how the sample holds together during chewing (low=fibers break easily, wad disintegrates; high=wad raises in size, compete against breakdown; Lyon and Lyon, 1990). Cohesiveness was not varied in all samples during fermentation and 7 days of ripening time. However, a noticeable difference ($p<0.05$) was observed in the cohesiveness values amongst the final ripened products, and samples in PE1 and the COS, presented similarly higher values than PE2 treated samples. Fermentation and ripening time didn't affect the springiness of COS and PE1 treatments and the chewiness of all treated batches. The springiness value in PE2 samples was increased due to the extended time of ripening. Substantially increased hardness and decreased cohesiveness ($p<0.05$) were revealed in all treatments due to the change in ripening time. As the samples get ripened, the available free water (water activity) is reduced to a significant level (Table 1), and an increase in the hardness values ($r=0.85$, $p>0.05$: data not presented) of all treatments was exhibited. The unification of ingredients may

Table 5. Effect of different starter cultures on TPA values of dry fermented sausages during fermentation and ripening

Attributes	Days	Treatments ¹⁾			SEM
		COS	PE1	PE2	
Hardness (kgf)	7 day of fermentation	1.23 ^B	1.12 ^B	1.39 ^B	0.18
	7 days of ripening	1.69 ^{ab}	1.11 ^{bb}	1.54 ^{ab}	0.17
	14 days of ripening	2.39 ^A	2.93 ^A	3.17 ^A	1.16
	SEM	0.25	0.83	0.81	
Springiness	7 days of fermentation	0.92	0.88	0.93 ^B	0.04
	7 days of ripening	0.93	0.93	0.93 ^B	0.03
	14 days of ripening	0.94	0.95	0.97 ^A	0.03
	SEM	0.02	0.05	0.02	
Cohesiveness	7 days of fermentation	0.32 ^A	0.35 ^A	0.33 ^A	0.06
	7 days of ripening	0.23 ^{AB}	0.24 ^B	0.26 ^B	0.05
	14 days of ripening	0.14 ^{ab}	0.13 ^{ab}	0.09 ^{bc}	0.34
	SEM	0.05	0.05	0.03	
Chewiness (kgf)	7 days of fermentation	0.37	0.34	0.43	0.11
	7 days of ripening	0.32	0.25	0.37	0.09
	14 days of ripening	0.32	0.36	0.26	0.10
	SEM	0.08	0.12	0.09	

¹⁾ Treatments are different strain of *Pediococcus* spp. used in the present study: COS, commercial starter culture; PE1, *P. acidilactici*; PE2, *P. pentosaceus*, *Starterkulturen Almi Rohschinken*, used as control (n=3).

^{a,b} Means in a row with different superscripts are significantly different ($p<0.05$).

^{A-C} Means in a column with different superscripts are significantly different ($p<0.05$).

TPA, texture profile analysis.

be affected due to extended ripening time and resulted in fragmentation of structural components that ended with lower cohesiveness characteristics of all batches studied.

Sensory analysis

The sensory quality traits of food products are the primary evaluation criteria of consumers before the consideration of any other quality attributes. During the ripening process, starter cultures play an important role in the acidification and generation of fermented sausages with unique and distinct quality characteristics, which contributes to sensory acceptability and physical properties (Bassi et al., 2015). Dry fermented sausages from different strains of *Pediococcus* spp. and control (COS) were subjected to sensory evaluation and judged by seven trained panelists for four attributes (color, aroma, sourness, and overall acceptability), and the results are presented in Table 6. Among the four descriptors considered in the current study, there were no significant differences ($p>0.05$) in the aroma and fermented sour taste characteristics of treatments. According to the panel judgments, the sausages added with PE1 and PE2 were perceived to have a similarity in aroma and sour taste as that of COS samples. However, the inoculation of different strains of *Pediococcus* spp. had a significant effect ($p<0.05$) on the color and overall acceptability traits of the ripened sausages. Regarding the color, sausages both in PE1 and PE2 treatments showed similar and higher ($p>0.05$) dark red color intensity than the COS batch. In the overall acceptability attribute, the sensory evaluation panelists generally preferred samples in PE1 than PE1 and COS treatments. In general, the current sensory evaluation showed the inoculation of different strains of *Pediococcus* spp. presented similar characteristics in aroma and fermented acidic taste as that of the COS and both PE1 and PE2 batches were better in color attributes. Moreover, PE1 scored highest in the overall acceptability attribute while PE2 and COS were perceived to have similarities.

Conclusion

The addition of PE1 and PE2 resulted in a similar beneficial effect in the reduction of pH value and acidification of the products. PE1 inoculation showed a substantially higher LAB count than PE2 after-ripening and pronounced high count than the COS during fermentation and early ripening period. Moreover, PE1 addition led to significantly lower TBARS and VBN contents than COS and PE2. All three starter cultures resulted in similar color characteristics (L^* , a^* , and b^*) since 7 days of fermentation. Moreover, the TPA characteristics of springiness, chewiness, and hardness were the same in all three batches of the ripened products. However, the cohesiveness property was better in PE1 and COS products than in PE2. From the grounds of the present study findings, it can be concluded that the overall performance of PE1 strain was conducive to

Table 6. Effect of different starter cultures on sensory characteristics of dry fermented sausages after the ripening period

Attributes	Treatments ¹⁾			SEM
	COS	PE1	PE2	
Color	2.47 ^b	4.31 ^a	3.46 ^a	0.85
Aroma	3.33	4.14	3.23	0.81
Sourness	4.34	3.96	3.60	0.73
Overall acceptability	3.69 ^b	4.47 ^a	3.57 ^b	0.52

¹⁾ Treatments are different strain of *Pediococcus* spp. used in the present study: COS, commercial starter culture; PE1, *P. acidilactici*; PE2, *P. pentosaceus*, *Starterkulturen Almi Rohschinken*, used as control (n=3).

^{a,b} Means in a row with different superscripts are significantly different ($p<0.05$).

improving the quality characteristics of dry fermented sausage. The inoculation of different strains of *Pediococcus* spp. and COS presented similar sensory characteristics in aroma and fermented acidic taste. PE1 and PE2 batches were better in color quality and PE1 preferred in the overall acceptability attribute.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (2017R1A2B201277).

Author Contributions

Conceptualization: Kang SN. Data curation: Seleshe S, Ameer A. Formal analysis: Seleshe S, Ameer A. Methodology: Seleshe S, Kang SN. Software: Seleshe S, Kang SN. Validation: Kang SN. Investigation: Kang SN. Writing - original draft: Seleshe S, Ameer A, Kang SN. Writing - review & editing: Seleshe S, Ameer A, Kang SN.

Ethics Approval

The sensory analysis method was certified with IRB number (1040621-201905-HR-004-02) by the life management committee of Daegu University.

References

- Ba HV, Seo HW, Seong PN, Kang SM, Kim YS, Cho SH, Park BY, Ham JS, Kim JH. 2018. *Lactobacillus plantarum* (KACC 92189) as a potential probiotic starter culture for quality improvement of fermented sausages. *Korean J Food Sci Anim Resour* 38:189-202.
- Baka AM, Papavergou EJ, Pragalaki T, Bloukas JG, Kotzekidou P. 2011. Effect of selected autochthonous starter cultures on processing and quality characteristics of Greek fermented sausages. *LWT-Food Sci Technol* 44:54-61.
- Barbut S. 2011. 11 - Saturated fat reduction in processed meat products. In *Reducing saturated fats in foods*. Talbot G (ed). Woodhead, Cambridge, UK. pp 379-392.
- Bassi D, Puglisi E, Cocconcelli PS. 2015. Comparing natural and selected starter cultures in meat and cheese fermentations. *Curr Opin Food Sci* 2:118-122.
- Bingol EB, Ciftcioglu G, Eker FY, Yardibi H, Yesil O, Bayrakal GM, Demirel G. 2014. Effect of starter cultures combinations on lipolytic activity and ripening of dry fermented sausages. *Ital J Anim Sci* 13:3422.
- Casaburi A, Aristoy MC, Cavella S, Monaco RD, Ercolini D, Toldrá F, Villani F. 2007. Biochemical and sensory characteristics of traditional fermented sausages of Vallo di Diano (Southern Italy) as affected by the use of starter cultures. *Meat Sci* 76:295-307.

- Cavalheiro CP, Piovesan N, Terra LM, Lovato M, Terra NN, Fries LLM. 2013. Colorimetric and sensory characteristics of fermented cured sausage with Brazilian ostrich meat addition. *Food Sci Technol, Campinas* 33:660-665.
- Chen Q, Kong B, Han Q, Xia X, Xu L. 2017. The role of bacterial fermentation in lipolysis and lipid oxidation in Harbin dry sausages and its flavour development. *LWT-Food Sci Technol* 77:389-396.
- Commission International de l'Eclairage [CIE]. 1976. Official recommendations on uniform color spaces. Color difference equations and metric color terms. Suppl. No. 2. CIE, No. 15 Colourimetry. Paris, France.
- Conway EJ. 1950. Microdiffusion analysis and volumetric error. 3rd ed. Crosby Lockwood and Son, London, UK.
- Daeschel MA, Klaenhammer TR. 1985. Association of a 13.6-megadalton plasmid in *Pediococcus pentosaceus* with bacteriocin activity. *Appl Environ Microbiol* 50:1538-1541.
- Diguță CF, Nițoi GD, Matei F, Luță G, Cornea CP. 2020. The biotechnological potential of *Pediococcus* spp. isolated from Kombucha microbial consortium. *Foods* 9:1780.
- Drosinos EH, Mataragas M, Xiraphi N, Moschonas G, Gaitis F, Metaxopoulos J. 2005. Characterization of the microbial flora from a traditional Greek fermented sausage. *Meat Sci* 69:307-317.
- Drosinos EH, Paramithiotis S, Kolovos G, Tsikouras I, Metaxopoulos I. 2007. Phenotypic and technological diversity of lactic acid bacteria and *Staphylococci* isolated from traditionally fermented sausages in southern Greece. *Food Microbiol* 24:260-270.
- Egan H, Kirk RS, Sawyer R. 1981. Pearson's chemical analysis of foods. 8th ed. Churchill Livingstone, London, UK. pp 534-539.
- Ergezer H, Kaya Hİ, Şimşek Ö. 2018. Antioxidant and antimicrobial potential of artichoke (*Cynara scolymus* L.) extract in beef patties. *Czech J Food Sci* 36:154-162.
- Essid I, Hassouna M. 2013. Effect of inoculation of selected *Staphylococcus xylosum* and *Lactobacillus plantarum* strains on biochemical, microbiological and textural characteristics of a Tunisian dry fermented sausage. *Food Control* 32:707-714.
- Grun IU, Ahn J, Clarke AD, Lorenzen CL. 2006. Reducing oxidation of meat. *Food Technol* 60:36-43.
- Hoppu U, Hopia A, Pohjanheimo T, Rotola-Pukkila M, Mäkinen S, Pihlanto A, Sandell M. 2017. Effect of salt reduction on consumer acceptance and sensory quality of food. *Foods* 6:103.
- HORIBA, 2021. 18 - pH measurement in the acidification process of fermented sausages. Available from: <https://www.scribd.com/document/437490575/18-PH-Measurement-in-the-Acidification-Process-of-Fermented-Sausages>. Accessed at Apr 7, 2021.
- Huang L, Zhao J, Chen Q, Zhang Y. 2014. Nondestructive measurement of total volatile basic nitrogen (TVB-N) in pork meat by integrating near infrared spectroscopy, computer vision and electronic nose techniques. *Food Chem* 145:228-236.
- Hugas M, Monfort JM. 1997. Bacterial starter cultures for meat fermentation. *Food Chem* 59:547-554.
- Incze K. 1998. Dry fermented sausages. *Meat Sci* 49:S169-S177.
- Juncher D, Vestergaard CS, Søltoft-Jensen J, Weber CJ, Bertelsen G, Skibsted LH. 2000. Effects of chemical hurdles on microbiological and oxidative stability of a cooked cured emulsion type meat product. *Meat Sci* 55:483-491.
- Kim HW, Hwang KE, Song DH, Kim YJ, Ham YK, Yeo EJ, Jeong TJ, Choi YS, Kim CJ. 2015. Effect of pre-rigor salting levels on physicochemical and textural properties of chicken breast muscles. *Korean J Food Sci Anim Resour* 35:577-584.
- Kurt S. 2016. The effects of grape seed flour on the quality of Turkish dry fermented sausage (Sucuk) during ripening and refrigerated storage. *Korean J Food Sci Anim Resour* 36:300-308.

- Leroy F, De Vuyst L. 2016. Fermented foods: Fermented meat products. In Encyclopedia of food and health. Caballero B, Finglas PM, Toldrá F (ed). Academic Press, Amsterdam, Netherland. pp 656-660.
- Leroy F, Geyzen A, Janssens M, De Vuyst L, Scholliers P. 2013. Meat fermentation at the crossroads of innovation and tradition: A historical outlook. Trends Food Sci Technol 31:130-137.
- Leroy F, Verluyten J, De Vuyst L. 2006. Functional meat starter cultures for improved sausage fermentation. Int J Food Microbiol 106:270-285.
- Lilic S, Saicic S, Vranic D, Trbovic D, Borovic B, Velebit B, Lakicevic B. 2011. Salt and sodium content in dry fermented sausages and dried meat in Serbia. Available from: https://www.researchgate.net/publication/271523538_Salt_and_sodium_content_in_dry_fermented_sausages_and_dried_meat_in_Serbia. Accessed at Apr 2, 2021.
- Lim DG, Seol KH, Jeon HJ, Jo C, Lee M. 2008. Application of electron-beam irradiation combined with antioxidants for fermented sausage and its quality characteristic. Radiat Phys Chem 77:818-824.
- Lorenzo JM, Gómez M, Fonseca S. 2014. Effect of commercial starter cultures on physicochemical characteristics, microbial counts and free fatty acid composition of dry-cured foal sausage. Food Control 46:382-389.
- Luecke FK. 1997. Fermented sausages. In Microbiology of fermented foods. Wood BJB (ed). Blackie Academic & Professional, London, UK. pp 441-483.
- Lyon BG, Lyon CE. 1990. Texture profile of broiler pectoralis major as influenced by post-mortem deboning time and heat method. Poult Sci 69:329-340.
- Marco A, Navarro JL, Flores M. 2006. The influence of nitrite and nitrate on microbial, chemical and sensory parameters of slow dry fermented sausage. Meat Sci 73:660-673.
- Morot-Bizot SC, Leroy S, Talon R. 2006. Staphylococcal community of a small unit manufacturing traditional dry fermented sausages. Int J Food Microbiol 108:210-217.
- Pasini F, Soglia F, Petracci M, Caboni MF, Marziali S, Montanari C, Gardini F, Grazia L, Tabanelli G. 2018. Effect of fermentation with different lactic acid bacteria starter cultures on biogenic amine content and ripening patterns in dry fermented sausages. Nutrients 10:1497.
- Pérez-Alvarez JA, Fernández-López J. 2011. Color measurements on muscle-based foods. In Sensory analysis of foods of animal origin. Nolet LML, Toldrá F (ed). CRC Press, Boca Raton, FL, USA. pp 3-10.
- Pikul J, Leszczynski DE, Kummerow FA. 1989. Evaluation of three modified TBA methods for measuring lipid oxidation in chicken meat. J Agric Food Chem 37:1309-1313.
- Rai KP, Zhang C, Xia WS. 2010. Effects of pure starter cultures on physico-chemical and sensory quality of dry fermented Chinese-style sausage. J Food Sci Technol 47:188-194.
- Ruiz-Capillas C, Jiménez-Colmenero F. 2004. Biogenic amines in meat and meat products. Crit Rev Food Sci Nutr 44:489-599.
- Terrell RN. 1983. Reducing the sodium content of processed meats. Food Technol 37:66-71.
- Valyasevi R, Jungsirowat P, Smitinont T, Praphilong W, Chowalitnitithum C. 2001. Improvement of starter culture for Nham fermentation. National Science and Technology Development Agency, Khlong Luang, Thailand.
- Vignolo G, Fontana C, Fadda S. 2010. Semidry and dry fermented sausages. In Handbook of meat processing. Toldr F (ed). Wiley-Blackwell, Oxford, UK. pp 379-398.
- Villani F, Pepe O, Mauriello G, Salzano G, Moschetti G, Coppola S. 1994. Antimicrobial activity of *Staphylococcus xylosus* from Italian sausages against *Listeria monocytogenes*. Lett Appl Microbiol 18:159-161.

- Yim DG, Chung YH, Nam KC. 2017. Effects of starter cultures on physicochemical properties of fermented sausages. *Korean J Food Nutr* 30:1105-1112.
- Yin LJ, Pan CL, Jiang ST. 2002. Effect of lactic acid bacterial fermentation on the characteristics of minced mackerel. *J Food Sci* 67:786-792.
- Zhao B, Zhou H, Zhang S, Pan X, Li S, Zhu N, Wu Q, Wang S, Qiao X, Chen W. 2020. Changes of protein oxidation, lipid oxidation and lipolysis in Chinese dry sausage with different sodium chloride curing salt content. *Food Sci Hum Wellness* 9:328-337.
- Zhao L, Jin Y, Ma C, Song H, Li H, Wang Z, Xiao S. 2011. Physico-chemical characteristics and free fatty acid composition of dry fermented mutton sausages as affected by the use of various combinations of starter cultures and spices. *Meat Sci* 88:761-766.