


Extermination Speed of an Imidacloprid and Flumethrin Polymer Matrix Collar against Larvae, Nymphs and Adults of *Haemaphysalis longicornis*

Kyu-Sung Ahn¹ , Ah-Jin Ahn² , Ha-Jung Kim³ , Chang-Min Lee³ , Guk-Hyun Suh³ , Eun Jeong Won⁴ ,
SungShik Shin^{1,*} 

¹Department of Parasitology, College of Veterinary Medicine, Chonnam National University, Gwangju 61186, Korea; ²Gwangju Metropolitan Government Research Institute of Public Health and Environment, Gwangju 61954, Korea; ³Department of Internal Medicine, College of Veterinary Medicine, Chonnam National University, Gwangju 61186, Korea; ⁴Department of Parasitology and Tropical Medicine, Chonnam National University Medical School, Hwasun-gun 58128, Korea

Abstract: The objective of this study was to evaluate the efficacy of an imidacloprid 10% and flumethrin 4.5% polymer matrix collar against the developmental stages of *Haemaphysalis longicornis* infesting dogs using the hair from treated dogs in a semi-in-vitro assay set. When incubated with 0.5 g of the hair collected from the dogs installed with the drug-embedded collar after 10 days, average death rate of the larval, nymphal, and adult *H. longicornis* was 21.5%, 77.9%, and 100% at 30 min, 1 hr, and 2 hr, respectively. This study showed the larval stages as well as the nymphal and adult stages of *H. longicornis* ticks are killed upon contact with the hair from dogs treated with the collar within 2 hr.

Key words: *Haemaphysalis longicornis*, Seresto, tick, imidacloprid, flumethrin, polymer, collar

INTRODUCTION

The speed and ability to kill both the juvenile and adult ticks are important factors for an acaricide to successfully prevent the transmission of tick-borne diseases in dogs. The tick-borne pathogens are transmitted from ticks to animals or to humans during a tick sucks blood and injects saliva containing pathogens into the host tissues. The transmission of diseases such as Lyme borreliosis takes 24 to 48 hr after a tick bite [1]. The Powassan virus causing encephalitis in humans can be transmitted to host animals in 15 min of tick attachment [2]. The transmission course emphasizes the importance of the speed kill of acaricides against ticks both for the prevention against tick bites and tick-borne pathogens. The ability of an acaricide to kill the larval stage of ticks as well as the nymphal and adult stages is also important, because evidence for the transovarial transmission of pathogens via the female-egg-larvae route in ticks have been well documented. Examples of

transovarial transmission include *Borrelia miyamotoi* by *Ixodes scapularis* tick, *Leishmania infantum* by *Rhipicephalus sanguineus*, *Rickettsia bellii* by *Ixodes loricatus* and *Babesia equi* by *Haemaphysalis longicornis* [3-8]. In dogs, the transmission of pathogens via the transovarial route from infected female ticks to dogs by larval ticks was documented on *Babesia gibsoni* by *Haemaphysalis hystricus* and *Rickettsia rickettsia* by *Rhipicephalus sanguineus* [5, 6].

The role of larval ticks should therefore be considered important in the transmission of tick-borne diseases. It should be ensured that an acaricide kills larvae as well as ticks in the nymph and adult stages. However, it is difficult to assess the in-vivo clinical efficacy of drugs against the larval stage of ticks on host animals due to their minute size which hinders them from being found on the host's skin, especially in the thick and/or colored hairs. For this reason, larvicidal assays of acaricides have used in-vitro systems such as Food and Agriculture Organization larval packet test (LPT) or larval immersion microassay (LIM) [9,10]. However, limitations exist with these assays to obtain information on in-vivo acaricidal efficacy on the host animal skin after acaricides are applied.

The host hair from dogs treated with certain insecticides such as pyrethroids or neonicotinoids are coated with a small amount of the chemicals, which is enough to kill ticks or fleas

•Received 18 July 2021, revised 11 October 2021, accepted 12 October 2021.

*Corresponding author (sungshik@jnu.ac.kr)

© 2021, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

upon contact with the hair [11-13]. It was reported that adult cat fleas (*Ctenocephalides felis* Bouche' 1835) died within 50-90 min on contact with hair from imidacloprid (Advantage®, Elanco Animal Health, Indianapolis, Indiana, USA)-treated dogs, whereas adult fleas, having contact with hair of untreated control dogs, were found alive even 7 days after the beginning of the experiments [11]. The hair removed from a treated dog with a contact-kill acaricide can, therefore, be used to assess the speed of kill against ticks. Seresto® (Elanco Animal Health) is a contact-kill collar product for dogs and cats that provides long term broad spectrum parasitocidal activity by combining the insecticidal properties of imidacloprid (10%) with the acaricidal properties of flumethrin (4.5%). The active ingredients spread from the site of direct contact over the entire skin surface of the treated animal and provide the long term steady and continuous release of the ingredients by the collar matrix system [14]. The product offers a breakthrough in protection against fleas and ticks for dogs. It was documented that a preventive efficacy against ticks in dogs varied between 91.2% and 100% (overall mean: 94.7%) for 8 months [15].

The purpose of this study is twofold: to assess the time taken to exterminate larvae, nymphs, and adults of the species in contact with the hair from the host dogs treated with the collar using a semi-in-vitro assay, and to evaluate the efficacy of 10% imidacloprid and 4.5% flumethrin polymer matrix collar against *Haemaphysalis longicornis* tick infestation on dogs.

MATERIALS AND METHODS

Ethics approval

A randomized, controlled efficacy trial was conducted in the animal care facility at the College of Veterinary Medicine, Chonnam National University, Gwangju, Korea. The design and experimental procedures used in this study followed the rules and regulations of the Korea Animal Protection Law (No. 13023, articles 8 and 9). This study was approved by the Institutional Animal Care and Use Committee of Chonnam National University (CNU IACUC-YB-2018-81).

Animals

Before the semi-in-vitro assay using the hair from treated dogs was performed, an in-vivo clinical efficacy of an ectoparasiticide against *H. longicornis* infestation in dogs was conducted to confirm its efficacy in field conditions. Twelve healthy

young dogs between 2 and 6 months old of both sexes with no history of tick infestation or exposure to tick habitats were enrolled in this study. Each dog received 3 doses of a vaccine against canine distemper, adenovirus Type 2, parvovirus, and parainfluenza (Vanguard® Plus 5, Zoetis, Parsippany, New Jersey, USA). All dogs were dewormed with a combination of praziquantel, pyrantel pamoate, and fenbentel (Drontal® plus; Elanco Animal Health) before being enrolled in this study. Dogs were individually housed in wire mesh cages (90×70×60 cm) and were fed commercial feed once per day, with water provided ad libitum. Six dogs were randomly assigned to the treatment group and 6 dogs in the untreated control group. On day -3, a polymer matrix collar containing 10% imidacloprid and 4.5% flumethrin (Seresto®, Elanco Animal Health) was applied to all 6 dogs in the treatment group according to the manufacturer's instructions: dogs ≤8 kg received a small collar and dogs >8 kg received a large collar.

In vivo drug efficacy evaluation

On day 0, all 12 dogs were naturally exposed to ticks by walking them for 1 hr in wooded and bushy areas with tall grass and leaf litter in Yeongkwang-gun, Jeonnam, Korea (35°14'39.5"N 126°35'08.7"E) where *H. longicornis* was found to be the dominant species of ticks in a preliminary investigation. The exposure of dogs to ticks was repeated on day 30. From day 1, each dog was clinically examined, and ticks were searched for once daily until all ticks became detached from the host. Dogs were examined daily for the presence of ticks by thumb counting, with examination of the cephalic, ventral, dorsal, and leg regions. Both nymphal and adult stages of ticks attached to the skin of the host animal were counted, and those ticks crawling around on the skin were not counted. Representative tick specimens (1 or 2 adult ticks per dog) were identified as *H. longicornis* based on morphology [16].

The efficacy of the collar against ticks was calculated as follows: $\text{Efficacy} = (\text{mean total number of ticks on control dogs} - \text{mean total number of ticks on treated dogs}) / (\text{mean total number of ticks on control dogs}) \times 100$. Collars were deemed effective against ticks if the calculated efficacy, based on arithmetic and geometric means, was at least 90% [17].

In vitro drug efficacy evaluation

For the semi-in-vitro assay, the larval, nymphal, and adult stages of ticks were co-cultured with hair from dogs collected on day 10 after the application of the collar containing 10%

imidacloprid and 4.5% flumethrin. Hair was collected from 3 dogs randomly selected from the treatment group and 3 dogs from the untreated control group at the dorsal medial line of each dog approximately 20 cm away from the collar. Collected hair was individually weighed, cut into pieces of approximately 5-10 mm long, and 0.5 g of hair from each dog was placed in a 50 ml conical centrifuge tube (Hyundai Micro Co. Ltd., Seoul, Korea). Each conical centrifuge tube contained a 1.5 ml micro-centrifuge tube sealed at the inner top wall of the 50 ml tube using a hot melt glue gun (Fig. 1). A small amount of sterilized cotton soaked with distilled water was inserted into the micro-centrifuge tube to provide humidity inside the 50 ml tube. Disposable gloves and scissors were used to prevent any contamination between hair samples.

Fifteen larvae, nymphs, or adults of *H. longicornis* were placed inside each of the 50 ml tube containing 0.5 g of the host hair collected from dogs installed with the collar or from the untreated control dogs. Larvae of *H. longicornis* were obtained from newly hatched eggs from engorged adult females in the laboratory, and additional nymphs and adults were collected from wooded fields at the study site for natural infection. The tubes were closed tightly and were incubated for 0.5, 1, 2, 4, 6, and 24 hr at room temperature (25°C). At each incubation time point, the number of live and dead ticks were counted using a dissecting microscope. Ticks were considered dead if they showed no movement and did not move or respond to prodding with a fine-tipped needle. Acaricidal efficacy was determined by comparing the mean number of live ticks remaining on the treated dog hair with the mean number of dead

mites remaining on the control dog hair.

RESULTS

Efficacy of 10% imidacloprid and 4.5% flumethrin in a polymer matrix collar against *H. longicornis* ticks in a field trial

In the untreated control group of 6 dogs on day 0 to *H. longicornis*-populated fields 3 days after the application of the polymer matrix collar containing 10% imidacloprid and 4.5% flumethrin, an average of 30.0, 38.1, and 36.1 ticks were found on the host animals on days 1, 2, and 3 after exposure to ticks (Data not shown in Table). The highest number of ticks attached to the host skin were observed on day 2 because there were some ticks still crawling around on day 1. The majority of ticks finished feeding on blood and detached from the host from day 4 after exposure, leaving an average of 6.6 ticks remaining on the host skin on day 4 and 0.3 ticks on day 7. Ticks were not found on the host skin from day 8 onwards in the control group. On the other hand, no ticks were found on any of the 6 dogs applied with the collar during the study period.

Although the average number of ticks attached to dogs was smaller than the first exposure, an average of 18.7, 21.2, and 14.9 ticks were found on the host animals on days 1, 2, and 3 after exposure to ticks (Data not shown in Table). The majority of ticks finished feeding on blood and detached from the host from day 4 after exposure, leaving an average of 4.6 ticks remaining on the host skin on day 4 and 0.4 ticks on day 7.




Fig. 1. A semi-in vitro acaricide assay set. (A) A disposable 50 ml conical centrifuge tube with a 1.5 ml micro-centrifuge tube sealed to inner top wall using a hot melt glue gun. A small amount of sterilized cotton soaked with distilled water was inserted into the micro-centrifuge tube to provide humidity in the 50 ml tube. (B) The dog hairs and 15 *Haemaphysalis longicornis* ticks (black dots) were co-incubated in the 50 ml conical tube.

Same as in the first exposure, ticks were not found from any of the 6 dogs treated with the drug-embedded collar on the second exposure. Dogs in both groups did not show any abnormal clinical signs of side-effects such as loss of appetite, vomiting, depression, or lethargy during the study period.

Efficacy of 10% imidacloprid and 4.5% flumethrin in a polymer matrix collar against the developmental stages of ticks by the semi-in-vitro assay

The efficacy of host hair collected from dogs treated with the polymer matrix collar containing 10% imidacloprid and 4.5% flumethrin against larval-, nymphal- and adult-stage ticks is summarized in Table 1. When larval stage ticks of *H. longicornis* ticks were incubated with the host hair from the drug-treated dogs, 24.4%, 95.5%, and 100.0% of ticks were killed within 30 min, 1 hr, and 2 hr after incubation respectively. Similar results were obtained when the nymphal- and adult-stages of *H. longicornis* were incubated with the host hair. Overall, averages of 21.5%, 77.9%, and 100% of ticks were killed within 30 min, 1 hr, and 2 hr after incubation with host hair from dogs in the drug-treated group respectively. On the contrary, all developmental stages of *H. longicornis* ticks incubated with the host hair from the untreated control dogs survived throughout the test period (24 hr).

DISCUSSION

Some chemicals such as pyrethrins, phrethroids, neonicotinoids, phenylpyrazoles, metaflumizone, and macrocyclic lactones of the spinosyn group are contact-kill insecticides [13]. Flumethrin, a second generation of photostable pyrethrins, and imidacloprid, one of neonicotinoid insecticides are the 2 active ingredients embedded in the polymer matrix collar that kill fleas and ticks on contact. The semi-in-vitro assay used in this study shows that the larval, nymph, and adult stages of *H. longicornis* ticks are efficiently killed upon contact with hair from dogs treated with the collar within 2 hr.

It is difficult to assess the time to kill ticks that are on the host animal down to the hourly or minute intervals in conventional clinical studies of drug efficacy using live animals because it takes considerable, and challenging, time to find live ticks that are scattered around the host body after infestation. It is especially so when measuring the time to kill nymphs or larvae due to their minute sizes that are difficult to find in the midst of the host's hair. The in-vitro evaluation of a

drug's efficacy using either a filter paper impregnated with a test drug or immersion of ticks in a drug-diluted solution before incubation in filter paper clips was therefore frequently used [18,19]. However, the filter paper or immersion tests do not represent the speed of the killing by an acaricide in situ on the host animal's skin. The semi-in-vitro assay using the hair from drug-treated dogs, on the other hand, can assess the efficacy of the speed of extermination by an acaricide down to hourly or even minute intervals.

The results of our study also indicate that tick-borne diseases by *H. longicornis* ticks can be further prevented by a polymer matrix collar containing 10% imidacloprid and 4.5% flumethrin, because ticks are expected to be killed even before they attach themselves to the host's skin and start feeding on blood. The Asian long-horned tick (or bush tick in Australia), *H. longicornis*, is prevalent in Asian countries like China, Japan, Korea, and the far-east of Russia; and it has also been found in Australia, New Zealand, several Pacific islands (Caledonia, Fiji, Western Samoa, Tonga, Vanuatu), and more recently in North America [20-23]. Besides its role as a vector for *Theileria orientalis* in livestock and severe fever with thrombocytopenia syndrome virus (SFTSV) in humans [24,25], this species is the vector of *Babesia gibsoni* for dogs in Asia [26], and is also a putative vector of *Hepatozoon canis* [27], *Rickettsia japonica* [28], *Borrelia*, *Bartonella*, *Anaplasma*, and *Ehrlichia* [29].

Both pyrethroids and neonicotinoids act through contact. Pyrethroids such as flumethrin, permethrin, and deltamethrin kill insects and mites by opening Na⁺ channels, inducing nerve cell membrane depolarization [13]. Neonicotinoids such as imidacloprid, nitenpyram, and dinotefuran act like agonists on postsynaptic nicotinic acetylcholine receptors, mainly in motoneurons, inducing nerve membrane depolarization, which causes spastic paralysis in insects. The absence of ticks on the host's skins in the treatment group in the field exposure study is therefore due to the contact killing of imidacloprid and flumethrin embedded in the collar, even before ticks inserted their hypostomes into the hosts' skins.

Although the number of dogs enrolled in this study is small to draw any statistical significance, this study clearly illustrates that the drug-treated collar (Seresto®) can successfully prevent dogs from undergoing *H. longicornis* infestation, as shown by the high efficacy against tick infestation in the 2 field exposures of dogs to areas where *H. longicornis* is populated (Data not shown in Table). Number of previous reports demonstrated the long-term efficacy of the product against ticks and fleas

Table 1. Efficacy of the hair from dogs treated with 10% imidacloprid and 4.5% flumethrin incorporated in a polymer matrix collar against the developmental stages of *Haemaphysalis longicornis*

Developmental Stages	Treated with Seresto ¹													
	Grop		0.5		1		2		4		6		24	
	Hour	Mean No. of ticks	Live	Dead	Live	Dead	L	D	L	D	L	D	L	D
Larvae			11.3	3.7	0.7	14.3	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0
	Efficacy (%)		0.0	24.4	0.0	95.5	0.0	100.0	0.0	100.0	0.0	100.0	0.0	100.0
Nymph			11.3	3.7	5.0	10.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0
	Efficacy (%)		0.0	24.4	0.0	66.7	0.0	100.0	0.0	100.0	0.0	100.0	0.0	100.0
Adult			12.7	2.3	4.3	11.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0
	Efficacy (%)		0.0	15.6	0.0	71.7	0.0	100.0	0.0	100.0	0.0	100.0	0.0	100.0
Total			35.3	9.7	10.0	35.3	0.0	45.0	0.0	45.0	0.0	45.0	0.0	45.0
	Efficacy (%)		21.5	77.9			100.0			100.0		100.0		100.0
Untreated control														
Developmental Stages	Grop		0.5		1		2		4		6		24	
	Hour	Mean No. of ticks	L	D	L	D	L	D	L	D	L	D	L	D
			L <td>D <td>L <td>D <td>L <td>D <td>L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td></td></td></td></td></td></td>	D <td>L <td>D <td>L <td>D <td>L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td></td></td></td></td></td>	L <td>D <td>L <td>D <td>L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td></td></td></td></td>	D <td>L <td>D <td>L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td></td></td></td>	L <td>D <td>L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td></td></td>	D <td>L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td></td>	L <td>D <td>L <td>D <td>L <td>D </td></td></td></td></td>	D <td>L <td>D <td>L <td>D </td></td></td></td>	L <td>D <td>L <td>D </td></td></td>	D <td>L <td>D </td></td>	L <td>D </td>	D
Larvae		15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0
	Efficacy (%)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Nymph		15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0
	Efficacy (%)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Adult		15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0	0.0	15.0
	Efficacy (%)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total		45	0.0	45	0.0	45	0.0	45	0.0	45	0.0	45	0.0	45
	Efficacy (%)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

¹Seresto® Collar, incorporated with 10% (w/v) imidacloprid and 4.5% (w/v) flumethrin.

in dogs and cats. In multicenter clinical field studies in dogs and cats in Europe, Stanneck et al. [15] reported that the imidacloprid/flumethrin collar proved to reduce tick counts by at least 90% and flea counts by at least 95% for a period of at least 7-8 months in cats and dogs under field conditions. Brianti et al. [30] also reported the efficacy of the collar in 82 dogs that the mean intensity of ticks (*Rhipicephalus sanguineus sensu lato*) decreased to 3.5 with a reduction of 92.5%. In a parallel group study on the speed of kill, repellent ability, and acaricidal efficacy of Seresto® and Broadline® (Merial) against artificially-induced infestations with *Ixodes ricinus* on cats, Seresto® proved to be 100% effective against adult *I. ricinus* at all assessment times (6, 12, 24, and 48 hr after infestation) throughout the month-long study [30]. Since the tick counts were done only on days 0 and 30 and only 12 dogs were enrolled in our study, a direct comparison of results with previous studies would not be relevant, and the study group (only 6 treated dogs and 6 as a control) is too small for a reliable statistical analysis. However, the combination of the field study using 12 dogs and the semi-in-vitro assay using the host hair from treated dogs demonstrates the excellent efficacy of the polymer matrix collar containing 10% imidacloprid and 4.5% flumethrin against all 3 developmental stages of *H. longicornis* ticks.

The semi-in-vitro assay using the hair from treated dogs may not be applicable to evaluate acaricides with systemic modes of action. While some active ingredients that are used in spot-on formulations or collars that act upon contact with the arthropods; others are systemic and/or mixed in their mode of action with a transcutaneous penetration and a plasma circulation [13]. Systemic insecticides are drugs whose active ingredient is distributed by blood circulation regardless of the route of administration. Systemic insecticides currently being used in dogs against tick infestation were identified which include imidacloprid, spinosad, fluralaner, afoxolaner, sarolaner, lufenuron, and nitenpyram [31-33].

In conclusion, this study showed that ticks are successfully prevented by applying the polymer matrix collar containing 10% imidacloprid and 4.5% flumethrin. and the semi-in-vitro assay using the hair from dogs can be a useful tool to evaluate the contact-kill efficacy of acaricides.

ACKNOWLEDGMENT

This study was supported in part by the Ministry of Agricul-

ture, Food and Rural Affairs, Republic of Korea (Grant No. 315016-3-C00).

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

REFERENCES

1. Piesman J, Mather TN, Sinsky R, Spielman A. Duration of tick attachment and *Borrelia burgdorferi* transmission. *J Clin Microbiol* 1987; 25: 557-558. <https://doi.org/10.1128/jcm.25.3.557-558.1987>
2. Ebel GD, Kramer LD. Short Report: duration of tick attachment required for transmission of powassan virus by deer ticks. *Am J Trop Med Hyg* 2004; 71: 268-271. <https://doi.org/10.4269/ajtmh.2004.71.3.0700268>
3. Rollend L, Fish D, Childs JE. Transovarial transmission of *Borrelia spirochetes* by *Ixodes scapularis*: a summary of the literature and recent observations. *Ticks Tick-borne Dis* 2013; 4: 46-51. <https://doi.org/10.1016/j.ttbdis.2012.06.008>
4. Ikadai H, Sasaki M, Ishida H, Matsuu A, Igarashi I, Fujisaki K, Oyamada T. Molecular evidence of *Babesia equi* transmission in *Haemaphysalis longicornis*. *Am J Trop Med Hyg* 2007; 76: 694-697. <https://doi.org/10.4269/ajtmh.2007.76.694>
5. Jongejan F, Su BL, Yang HJ, Berger L, Bevers J, Liu PC, Fang JC, Cheng YW, Kraakman C, Plaxton N. Molecular evidence for the transovarial passage of *Babesia gibsoni* in *Haemaphysalis hystricis* (Acari: Ixodidae) ticks from Taiwan: a novel vector for canine babesiosis. *Parasit Vectors* 2018; 11: 134. <https://doi.org/10.1186/s13071-018-2722-y>
6. Piranda EM, Faccini JL, Pinter A, Pacheco RC, Cançado PH, Labruna MB. Experimental infection of *Rhipicephalus sanguineus* ticks with the bacterium *Rickettsia rickettsii*, using experimentally infected dogs. *Vector Borne Zoonotic Dis* 2011; 11: 29-36. <https://doi.org/10.1089/vbz.2009.0250>
7. Dantas-Torres F, Martins TF, de Paiva-Cavalcanti M, Figueredo LA, Lima BS, Brandão-Filho SP. Transovarial passage of *Leishmania infantum* kDNA in artificially infected *Rhipicephalus sanguineus*. *Exp Parasitol* 2010; 125: 184-185. <https://doi.org/10.1016/j.exppara.2010.02.003>
8. Horta MC, Pinter A, Schumaker TT, Labruna MB. Natural infection, transovarial transmission, and transstadial survival of *Rickettsia bellii* in the tick *Ixodes loricatus* (Acari: Ixodidae) from Brazil. *Ann NY Acad Sci* 2006; 1078: 285-290. <https://doi.org/10.1196/annals.1374.053>
9. Miller RJ, Davey RB, George JE. Modification of the food and agriculture organization larval packet test to measure amitraz-susceptibility against Ixodidae. *J Med Entomol* 2002; 39: 645-651. <https://doi.org/10.1603/0022-2585-39.4.645>
10. White WH, Plummer PR, Kemper CJ, Miller RJ, Davey RB, Kemp DH, Hughes S, Smith CK 2nd, Gutierrez JA. An in vitro larval

- immersion microassay for identifying and characterizing candidate acaricides. *J Med Entomol* 2004; 4: 1034-1042. <https://doi.org/10.1603/0022-2585-41.6.1034>
11. Mehlhorn H, Hansen O, Mencke N. Comparative study on the effects of three insecticides (fipronil, imidacloprid, selamectin) on developmental stages of the cat flea (*Ctenocephalides felis* Bouche 1835): a light and electron microscopic analysis of in vivo and in vitro experiments. *Parasitol Res* 2001; 87: 198-207. <https://doi.org/10.1007/PL00008575>
 12. Mehlhorn H, Schmahl G, Mencke N, Bach T. In vitro and in vivo studies on the effect of a combination containing 10% imidacloprid and 50% permethrin against *Ixodes ricinus* ticks. *Parasitol Res* 2003; 89: 323-325. <https://doi.org/10.1007/s00436-002-0809-y>
 13. Beugnet F, Franc M. Insecticide and acaricide molecules and/or combinations to prevent pet infestation by ectoparasites. *Trends Parasitol* 2012; 28: 267-279. <https://doi.org/10.1016/j.pt.2012.04.004>
 14. Mehlhorn H, Mencke N, Hansen O. Effects of imidacloprid on adult and larval stages of the flea *Ctenocephalides felis* after in vivo and in vitro application: a light-and electron-microscopy study. *Parasitol Res* 1999; 85: 625-637. <https://doi.org/10.1007/s004360050607>
 15. Stanneck D, Rass J, Radeloff I, Kruedewagen E, Le Sueur C, Hellmann K, Krieger K. Evaluation of the long-term efficacy and safety of an imidacloprid 10%/flumethrin 4.5% polymer matrix collar (Seresto®) in dogs and cats naturally infested with fleas and/or ticks in multicentre clinical field studies in Europe. *Parasit Vectors* 2012; 5: 66. <https://doi.org/10.1186/1756-3305-5-66>
 16. Yamaguti N, Tipton V, Keegan H, Toshioka S. Ticks of Japan, Korea, and the Ryukyu Islands: Brigham Young Univ Sci Bull Biol Ser 1971; 45: 1-226. <https://scholarsarchive.byu.edu/byuscib/vol15/iss1/1>
 17. Marchiondo A, Holdsworth P, Fourie L, Rugg D, Hellmann K, Snyder D, et al. World Association for the Advancement of Veterinary Parasitology (WAAVP): guidelines for evaluating the efficacy of parasiticides for the treatment, prevention and control of flea and tick infestations on dogs and cats. *Parasit Vectors* 2013; 194: 84-97. <https://doi.org/10.1016/j.vetpar.2013.02.003>
 18. Hagimori I, Machida H, Goi R, Mencke N. Efficacy of imidacloprid/permethrin and fipronil/(S)-methoprene combinations against *Haemaphysalis longicornis* ticks evaluated under in vitro and in vivo conditions. *Parasitol Res* 2005; 97 (suppl): 120-126. <https://doi.org/10.1007/s00436-005-1455-y>
 19. Borges L, Ferri P, Silva W, Silva W, Silva J. In vitro efficacy of extracts of *Melia azedarach* against the tick *Boophilus microplus*. *Med Vet Entomol* 2003; 17: 228-231. <https://doi.org/10.1046/j.1365-2915.2003.00426.x>
 20. Hoogstraal H, Roberts FHS, Kohls GM, Tipton VJ. Review of *Haemaphysalis (Kaiseriana) longicornis* Neumann (resurrected) of Australia, New Zealand, New Caledonia, Fiji, Japan, Korea, and northeastern China and USSR, and its parthenogenetic and bisexual populations (Ixodoidea, Ixodidae). *J Parasitol* 1968; 54: 1197-1213. <https://doi.org/10.2307/3276992>
 21. Roberts FHS. Australian Ticks. Melbourne, Australia. Commonwealth Scientific and Industrial Research Organization. 1970. <https://www.cabdirect.org/cabdirect/abstract/19711001240>
 22. Rainey T, Occi JL, Robbins RG, Egizi A. Discovery of *Haemaphysalis longicornis* (Ixodida: Ixodidae) Parasitizing a Sheep in New Jersey, United States. *J Med Entomol* 2018; 55: 757-759. <https://doi.org/10.1093/jme/tjy006>
 23. Heath AC, Palma RL, Cane RP, Hardwick S. Checklist of New Zealand ticks (Acari: Ixodidae, Argasidae). *Zootaxa* 2011; 2995: 55-63. <https://doi.org/10.5281/zenodo.203775>
 24. Luo LM, Zhao L, Wen HL, Zhang ZI, Liu JW, Fang LZ, Xue ZF, Ma DQ, Zhang XS, Ding SJ, Lei XY, Yu XJ. *Haemaphysalis longicornis* ticks as reservoir and vector of severe fever with thrombocytopenia syndrome virus in China. *Emerg Infect Dis* 2015; 21: 1770-1776. <https://doi.org/10.3201/eid2110.150126>
 25. Heath A. Biology, ecology and distribution of the tick, *Haemaphysalis longicornis* Neumann (Acari: Ixodidae) in New Zealand. *N Z Vet J* 2016; 64: 10-20. <https://doi.org/10.1080/00480169.2015.1035769>
 26. Higuchi S, Simomura S, Yoshida H, Hoshi F, Kawamura S, Yasuda Y. Development of *Babesia gibsoni* in the gut epithelium of the tick, *Haemaphysalis longicornis*. *J Vet Med Sci* 1991; 53: 129-131. <https://doi.org/10.1292/jvms.53.129>
 27. Murata T, Inoue M, Taura Y, Nakama S, Abe H, Fujisaki K. Detection of *Hepatozoon canis* oocyst from ticks collected from the infected dogs. *J Vet Med Sci* 1995; 57: 111-112. <https://doi.org/10.1292/jvms.57.111>
 28. Lee JH, Park HS, Jung KD, Jang WJ, Koh SE, Kang SS, Lee IY, Lee WJ, Kim BJ, Kook YH, Park KH, Lee SH. Identification of the spotted fever group rickettsiae detected from *Haemaphysalis longicornis* in Korea. *Microbiol Immunol* 2003; 47: 301-304. <https://doi.org/10.1111/j.1348-0421.2003.tb03399.x>
 29. Sun J, Liu Q, Lu L, Ding G, Guo J, Fu C, Zhang J, Meng F, Wu H, Song X, Ren D, Li D, Guo Y, Wang J, Li G, Liu J, Lin H. Coinfection with four genera of bacteria (*Borrelia*, *Bartonella*, *Anaplasma*, and *Ehrlichia*) in *Haemaphysalis longicornis* and *Ixodes sinensis* ticks from China. *Vector Borne Zoonotic Dis* 2008; 8: 791-796. <https://doi.org/10.1089/vbz.2008.0005>
 30. Fourie JJ, Horak IG, De Vos C, Deuster K, Schunack B. Comparative speed of kill, repellent (anti-feeding) and acaricidal efficacy of an Imidacloprid/Flumethrin Collar (Seresto®) and a Fipronil/(S)-Methoprene/Eprinomectin/Praziquantel Spot-on (Broadline®) against *Ixodes ricinus* (Linné, 1758) on cats. *Parasitol Res* 2015; 114: 109-116. <https://doi.org/10.1007/s00436-015-4517-9>
 31. Gomez SA, Picado A. Systemic insecticides used in dogs: potential candidates for phlebotomine vector control? *Trop Med Int Health* 2017; 22: 755-764. <https://doi.org/10.1111/tmi.12870>
 32. Dobson P, Tinembart O, Fisch R, Junquera P. Efficacy of nitenpyram as a systemic flea adulticide in dogs and cats. *Vet Rec* 2000; 147: 709-713. <https://doi.org/10.1136/vr.147.25.709>
 33. Rust MK. Advances in the control of *Ctenocephalides felis* (cat flea) on cats and dogs. *Trends Parasitol* 2005; 21: 232-236. <https://doi.org/10.1016/j.pt.2005.03.010>