

Differential expression and localization of tight junction proteins in the goat epididymis

Sung Woo Kim^{1#}, Yu-Da Jeong^{2#}, Ga-Yeong Lee², Jinwook Lee¹, Jae-Yeung Lee¹, Chan-Lan Kim¹, Yeoung-Gyu Ko¹, Sung-Soo Lee¹ and Bongki Kim^{2*}

¹*Animal Genetic Resource Research Center, National Institute of Animal Science, Rural Development Administration, Hamyang 50000, Korea*

²*Department of Animal Resources Science, Kongju National University, Yesan 32439, Korea*



Received: Jan 5, 2022
 Revised: Feb 13, 2022
 Accepted: Feb 18, 2022

#These authors contributed equally to this work.

*Corresponding author

Bongki Kim
 Department of Animal Resources Science, Kongju National University, Yesan 32439, Korea.
 Tel: +82-41-330-1246
 E-mail: bkkim@kongju.ac.kr

Copyright © 2022 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Sung Woo Kim
<https://orcid.org/0000-0001-8521-3010>
 Yu-Da Jeong
<https://orcid.org/0000-0002-0569-9004>
 Ga-Yeong Lee
<https://orcid.org/0000-0001-8208-8728>
 Jinwook Lee
<https://orcid.org/0000-0001-9019-1653>
 Jae-Yeung Lee
<https://orcid.org/0000-0003-2965-7826>
 Chan-Lan Kim
<https://orcid.org/0000-0003-2826-3171>
 Yeoung-Gyu Ko
<https://orcid.org/0000-0002-6108-6325>
 Sung-Soo Lee
<https://orcid.org/0000-0003-1255-9509>

Abstract

The blood-epididymis barrier (BEB) forms a unique microenvironment that is crucial for the maturation, protection, transport, and storage of spermatozoa in the epididymis. To characterize the function of tight junctions (TJs), which are constitutive components of the BEB, we determined the expression and localization of TJ proteins such as zonula occludens (ZO)-1, 2, and 3, occludin, and claudin3 (Cldn3) during postnatal development in the goat epididymis. To assess the expression patterns of TJ proteins in immature (3 months of age) and mature (14 months of age) goat epididymides, two different experimental methods were used including immunofluorescence labeling and western blotting. We show that, ZO-1, 2, and 3, and occludin, were strictly expressed and localized to the TJs of the goat epididymis, whereas Cldn3 was present in basolateral membranes as well as TJs. All TJ proteins examined were more highly expressed in the immature epididymis compared to levels in mature tissue. In conclusion, our study indicates that at least five TJ proteins, namely ZO-1, ZO-2, ZO-3, occludin, and Cldn3, are present in TJs, and the expression strength and pattern of TJ proteins tend to be age dependent in the goat epididymis. Together, these data suggest that the distinct expression patterns of TJ proteins are essential for regulating components of the luminal contents in the epididymal epithelium and for forming adequate luminal conditions that are necessary for the maturation, protection, transport, and storage of spermatozoa in the goat epididymis.

Keywords: Blood-epididymis barrier, Tight junctions, Zonula occludens, Claudin, Occludin

INTRODUCTION

The formation of the biological microenvironment for sperm maturation and storage in the epididymis occurs via various specific events. One of the most important events is cell-cell communication among the principal, basal, and clear cells of the epididymal epithelium to create a specialized environment [1-3]. In addition, the blood-epididymis barrier (BEB) is comprised of tight junctions (TJs) between neighboring epithelial cells, which also supports the creation and maintenance of proper

Bongki Kim
<https://orcid.org/0000-0002-5229-7294>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was supported by National Research Foundation of Korea funded by the Ministry of Education (Grant No. 2019R11A3A01060532), and the Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ015583) from the Korean Rural Development Administration.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Kim SW, Jeong YD, Kim B.

Data curation: Jeong YD, Kim B.

Formal analysis: Kim SW, Jeong YD, Lee GY.

Methodology: Lee J, Lee JY, Kim CL, Ko YG, Lee SS.

Validation: Kim SW, Jeong YD.

Investigation: Kim SW, Jeong YD, Lee GY.

Writing - original draft: Kim SW, Jeong YD.

Writing - review & editing: Kim SW, Jeong

YD, Lee GY, Lee J, Lee JY, Kim CL, Ko YG, Lee SS, Kim B.

Ethics approval and consent to participate

All procedures described here were reviewed and approved by the Institutional Animal Care and Use Committee of the National Institute of Animal Science, Korea (Approval No. NIAS 2018-290).

physiological conditions for the maturation and storage of spermatozoa [4–6]. In male reproductive tracts, the distribution of various tight junctional proteins is well reported in the testis and the epididymis, especially. In the testis, the blood–testis barrier (BTB) is located close to the base of the seminiferous tubule between adjacent Sertoli cells [7,8]. In the epididymis, however, the BEB is formed at apical surfaces of the epididymal duct between epithelial cells [9,10]. The BTB and BEB show positional differences but have the same function, which is to regulate the movement of electrolytes, ions, water, nutrients, hormones, and molecules, and the permeability of the paracellular transport pathway. The BTB and BEB also maintain the immunologically privileged adluminal compartment, which is a necessary component of the functioning barrier of the testis and epididymis [11–16].

TJs are protein complexes composed of several peripheral membranes and integral transmembrane proteins. Zonula occludens-1 (ZO-1) was first discovered as a TJ component in epithelia located on the intracellular side of the plasma membrane [17]. Additionally, two more proteins, ZO-2 and ZO-3, were confirmed as members of the ZO protein family [18,19]. These ZO proteins play a critical role in forming bridges between trans-membrane associated proteins and the actin cytoskeleton [20,21]. In contrast, occludin was first confirmed as the integral transmembrane protein and has four transmembrane domains with intracellular N- and C-termini [22]. In addition, claudins (Cldns) are also known as TJ transmembrane proteins, comprising at least 27 members in mammals [23,24]. Both occludin and Cldns interact with various intracellular proteins of TJs, including ZO-1, ZO-2, and ZO-3 [25,26]. TJ complexes that form a variety of proteins can create a physical, physiological, and immunological barrier.

Recent studies have reported that hormones can regulate the expression levels and localization patterns of TJ proteins in the testis. For example, gonadotropin hormones (GH), follicle stimulating hormone (FSH), and luteinizing hormone (LH) regulate the formation of the pituitary gland and are necessary for TJ formation in rat testes because GH plays a critical role in the development and differentiation of Sertoli cells where TJs are formed [27,28]. Androgens can upregulate Cldn3 and Cldn11 in Sertoli cells [13,29], and the administration of flutamide, an androgen receptor antagonist, disrupts Cldn11 protein in rat testes [30]. Additionally, in the epididymis, we reported that the extracellular signal-regulated kinase (ERK) signaling pathway is a key player in controlling the expression and localization of TJ proteins, such as ZO-2, ZO-3, occludin, and Cldn1, 3, and 4 in TJs of the mouse epididymis [31]; however, we know less about the functional roles of TJs in the epididymis than in the testis. Moreover, there has been no study on the expression and localization of TJ proteins and changes in TJ proteins during postnatal development of the epididymis in goats. Hence, in this present study, we characterized the distribution and localization patterns of TJ proteins in the immature and mature epididymides of goats using immunofluorescence and western blotting analyses.

MATERIALS AND METHODS

Antibodies

All information of primary and secondary antibodies used for immunofluorescence staining and western blotting is described in Tables 1 and 2, respectively.

Tissue collection

Epididymides were collected using a surgical castration procedure from immature (3-month-old, $n = 4$), and mature (14-month-old, $n = 5$) black goats with an average body weight of 9.95 ± 0.66 kg and 28.04 ± 1.42 kg, respectively, at the National Institute of Animal Science. All animal

Table 1. List of primary antibodies

Antibody	Catalog	Company	Application
ZO-1 monoclonal	33-9100	Thermo Fisher Scientific	IF (1:300) and IB (1:1000)
Occludin polyclonal	71-1500	Thermo Fisher Scientific	IF (1:200) and IB (1:600)
Claudin-3 polyclonal	34-1700	Thermo Fisher Scientific	IF (1:150) and IB (1:600)
ZO-2 polyclonal	2847	Cell Signaling Technology	IF (1:150) and IB (1:1000)
ZO-3 monoclonal	3704	Cell Signaling Technology	IF (1:150) and IB (1:1000)
β-Actin monoclonal	5125	Cell Signaling Technology	IB (1:2000)

ZO, zonula occludens; IF, immunofluorescence labeling; IB, immunoblotting.

Table 2. List of secondary antibodies

Host species	Against	Conjugated to	Catalog No.	Company	Application
Donkey	Rabbit IgG	Cy3	711-166-152	Jackson ImmunoResearch	IF (1:400)
Donkey	Mouse IgG	Alexa Fluor 488	715-546-151	Jackson ImmunoResearch	IF (1:200)
Donkey	Rabbit IgG	HRP	711-036-152	Jackson ImmunoResearch	IB (1:1000)
Goat	Mouse IgG	HRP	115-036-003	Jackson ImmunoResearch	IB (1:1000)

IgG, immunoglobulin G; IF, immunofluorescence labeling; HRP, horseradish peroxidase; IB, immunoblotting.

experiments used here were performed under regulation and permission of the Institutional Animal Care and Use Committee (IACUC) of the National Institute of Animal Science (NIAS, Approval No. NIAS 2018-290).

Tissue fixation, embedding and sectioning

Following castration, the epididymides were briefly cleaned in a cold phosphate-buffered saline (PBS) buffer to remove blood. The fixation of the epididymides was achieved by immersion in 4% paraformaldehyde solution (PFA) for 48–72 hr at 4 °C, and the fixative was replaced with fresh solution every 24 hr. The samples were rinsed with PBS five times for 30 minutes each at room temperature (RT). To prevent ice crystal formation during freezing tissue samples, the epididymal tissues were immersed into 30% sucrose solution in PBS for at least 48–72 hr (until tissue sinks). The 30% sucrose solution was replaced with fresh one every 24 hr. The tissues were completely embedded in Tissue Tek compound (Sakura Finetek, Torrance, CA, USA) for 20 minutes, and freeze at –20 °C to –80 °C. 10 μm thick tissue sections were carefully cut and collected using a Leica cryostat (Leica Biosystems, Buffalo Grove, IL, USA). The cryostat temperature was between –18 °C and –25 °C. The tissue sections were mounted on gelatin coated slides (Sigma Aldrich, St. Louis, MO, USA), dried for 30 minutes at RT, and stored at –20 °C to –80 °C until further use.

Immunofluorescence staining

The cryostat sections were rehydrated in PBS for 15 min. For epitope retrieval, the sections were immersed into preheated retrieval solution (pH 9.0, 95 °C; DAKO, Carpinteria, CA, USA) for 30 minutes and then cooled to RT. The sections were then incubated in blocking buffer (1% bovine serum albumin [BSA] in PBS; Thermo Scientific, Rockford, IL, USA) for 30 minutes at RT to prevent non-specific binding issue on the target epitopes between primary antibodies and tissues. The sections were then incubated with primary antibodies diluted in the diluent buffer (DAKO) overnight at 4 °C. Thereafter, the sections were rinsed 3 times for 20 minutes in PBS and incubated with secondary antibodies diluted in the diluent buffer for 90 minutes at RT. For double immunostaining, primary and secondary antibody incubation steps were repeated without

the second block step. After this period, the sections were rinsed 3 times for 20 minutes in PBS and then placed in anti-fade mounting media containing 4',6-diamidino-2-phenylindole (DAPI; Vecta Labs, Burlingame, CA, USA) for 5 minutes at RT to counterstain the nuclei. For staining on negative controls, sections were incubated with the antibody diluent and a non-immune serum (Thermo Scientific) of the same isotype and at the same concentration as the primary antibodies. Confocal microscope (LSM800; ZEISS, Oberkochen, Germany) was used to acquire the images that analyzed by Zen Blue (ZEISS, Oberkochen, Germany).

Quantitative analysis of tight junction proteins by immunofluorescence

Epididymis tissues were double-immunostained with ZO-1 and ZO-2, ZO-3, occludin, or Cldn3. A LSM800 confocal microscope was used to acquire images. All images used for quantification were obtained in the same conditions. For comparisons of levels of TJ proteins between groups, the epididymides of three immature and mature animals was conducted and three cryostat sections from each group was labeled and analyzed. The quantification of the expression level of ZO-1 colocalization with each target protein in TJs was calculated by using the Image J software (NIH, Bethesda, MD, USA), as described earlier [31].

Protein extraction and immunoblotting

The epididymides were homogenized on ice for 30 minutes in RIPA buffer (Thermo Scientific) additive with complete protease inhibitors (Roche, Penzberg, Germany). The homogenized sample was centrifuged at 18,000×g for 20 minutes at 4°C, and then the supernatant were collected. Protein concentration was determined by the bicinchoninic acid (BCA) Protein Assay Kit (Thermo Scientific). Protein samples were diluted with 4× lithium dodecyl sulfate (LDS) Sample Buffer (Bio-Rad Laboratories, Hercules, CA, USA) in the presence of 4% β-mercaptoethanol. Samples were heated at 99°C for 10 minutes and cooled on ice before loading on gels. Proteins were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) gel electrophoresis and then wet-blotted onto polyvinylidene difluoride (PVDF) membranes for 90 minutes at 90 V. Blots were blocked for 60 minutes with 5% non-fat dry milk in Tris-buffered saline with 0.1% Tween-20 (TBST) at RT. Primary antibodies diluted in TBS were applied overnight at 4°C. After washing in TBST, secondary antibodies were applied for 60 minutes at RT. Antibody binding was visualized by Chemiluminescence reagent (Perkin Elmer Life Sciences) and were detected with iBright FL1500 (Invitrogen).

Statistical analysis

We have carried out a western blotting with the triplicated samples each group. The mean value of each target protein was normalized to the value of β-actin. Data are presented as mean ± standard error of the mean. Statistical comparison was evaluated by Student's *t*-tests.

RESULTS

Expression and localization of ZO-1, ZO-2, and ZO-3 in immature and mature goat epididymides

To confirm that ZO-1 is expressed in the goat epididymis, the epididymides at postnatal month (PNM) 3, and 14 were stained with a ZO-1 monoclonal antibody. ZO-1 was strictly expressed in the apical region (where TJs are present) of the epididymal tubules (Figs. 1A and 1B, white arrowheads). The expression level of ZO-1 was higher in the PNM3 than in the PNM14 epididymis. Quantification of the average pixel intensity of ZO-1 immunostaining in TJs showed




Fig. 1. Immunohistochemical expression and localization of zona occludens (ZO)-1 in the goat epididymis. The epididymis was immunostained for ZO-1 (green). (A) Epididymis at postnatal month (PNM) 3. (B) Epididymis at PNM14. Insets indicate ZO-1 labeling with DAPI (Bar = 20 μ m). (C) Quantification of ZO-1 labeling in TJs confirmed the decrease in ZO-1 expression in the mature epididymides compared to that in the immature ones. (D) Western blot analysis was applied to investigate ZO-1 expression in the epididymis at PNM3, and PNM14. * $p < 0.05$, ** $p < 0.001$.

a drastic decrease in the adult epididymis (Fig. 1C). The same results were observed by western blot analysis (Fig. 1D).

Next, we investigated whether the other two ZO proteins, ZO-2 and ZO-3, were present in the TJs of the goat epididymis. Double labeling with ZO-1 and ZO-2 showed that ZO-2 was co-localized with ZO-1 in TJs of the epididymal tubules (Figs. 2A and 2B, yellow arrowheads). ZO-2 expression pattern was similar to that of ZO-1 (Figs. 2C and 2D, white arrowheads). ZO-2 labeling was brighter in the PNM3 epididymides compared to that at PNM14. Quantification of the average pixel intensity of ZO-2-associated immunostaining in TJs revealed a drastic decrease in the mature epididymis (Fig. 2E).

Western blotting verified the decrease in ZO-2 in PNM14 epididymides (Fig. 2F). ZO-3 also colabeled with ZO-1 in TJs in the goat epididymis (Figs. 3A and 3B, yellow arrowheads). Further, its expression was significantly decreased at PNM14 compared to that in PNM3 epididymides (Figs. 3C and 3D, white arrowheads). The quantification of ZO-3 expression was confirmed by immunofluorescence (Fig. 3E) and western blotting (Fig. 3F). These results indicated that all three




Fig. 2. Immunohistochemical expression and localization of zona occludens (ZO)-2 in the goat epididymis. The epididymis was immunostained for ZO-1 (green) and ZO-2 (red). (A, C) Epididymis at postnatal month (PNM) 3. (B, D) Epididymis at PNM14. Yellow and white arrowheads indicate an overlap in ZO-1 and ZO-2 staining and ZO-2 single staining, respectively. Insets indicate ZO-1 single staining (Bar = 20 μ m). (E) Quantification of ZO-2 labeling in regions co-localized with ZO-1 in TJJs revealed a reduction in the ZO-2 expression level in the mature epididymides compared to that in the immature ones. (F) Western blot analysis was applied to investigate ZO-2 expression in the epididymis at PNM3 and PNM14. * $p < 0.05$, ** $p < 0.001$.




Fig. 3. Immunohistochemical expression and localization of ZO-3 in the goat epididymis. The epididymis was immunostained for zona occludens (ZO)-1 (green) and ZO-3 (red). (A, C) Epididymis at postnatal month (PNM) 3. (B, D) Epididymis at PNM14. Yellow and white arrowheads indicate in overlap of ZO-1 and ZO-3 staining and ZO-3 single staining, respectively. Insets indicate ZO-1 single staining (Bar = 20 μ m). (E) Quantification of ZO-3 labeling in regions co-localized with ZO-1 in TJ's revealed a reduction in the ZO-3 expression level in the mature epididymides compared to that in the immature ones. (F) Western blot analysis was applied to investigate ZO-3 expression in the epididymis at PNM3 and PNM14. * $p < 0.05$, ** $p < 0.001$.

ZO proteins were maintained at high expression levels during immature stages, but were drastically reduced in the mature epididymides.

Expression and localization of occludin in immature and mature goat epididymides

To determine whether occludin, a transmembrane protein, could be localized to the TJs of the goat epididymis, double-staining was performed with ZO-1 and occludin. Occludin protein colocalized with ZO-1 in TJs (Figs. 4A and 4B, yellow arrowheads). Occludin was also significantly decreased in the PNM14 epididymides compared to that of PNM3 (Figs. 4C and 4D, white arrowheads). The quantification of occludin expression was confirmed by immunofluorescence (Fig. 4E) and western blotting (Fig. 4F). This result indicated that occludin was also significantly decreased in the mature epididymis, similar to the three ZO proteins, known as peripheral membrane proteins.

Changes in claudin3 in immature and mature goat epididymides

To determine whether Cldn3, a transmembrane protein, could be expressed in the TJs of the goat epididymis, double-labeling was performed using ZO-1 and Cldn3 antibodies. Interestingly, the localization and expression of Cldn3 tended to be age-dependent. At PNM3, Cldn3 was expressed and localized to the TJs (Figs. 5A and 5C; yellow arrowheads) and basolateral membranes (Figs. 5A and 5C; white arrowheads). At PNM14, Cldn3 was significantly decreased in TJs where ZO-1 was localized (Figs. 5B and 5D; yellow arrowheads) and also showed low expression levels in the basolateral membranes (Figs. 5B and 5D; white arrowheads). Quantification of the average pixel intensity of Cldn3-associated immunostaining from TJs revealed a significantly reduced signal at PNM14 compared to that at PNM3 (Fig. 5E). The total expression level of Cldn3 was examined by western blotting (Fig. 5F). These results indicate that Cldn3 is expressed at and localized to the basolateral membrane and TJs. Moreover, the changes in Cldn3 expression patterns appear to be age-dependent.

DISCUSSION

Incomplete spermatozoa can obtain abilities such as progressive motility and fertilization capacity through the final maturation step in the epididymis. The maturation events of spermatozoa in the epididymis depend on a unique luminal environment that is provided and maintained by the formation of the BEB, which is formed by TJs composed of several proteins, including transmembrane and peripheral proteins. These specialized TJ proteins function in forming the seal between epithelial cells, which is essential for sperm maturation as well as sperm protection from the immune system. However, there is still limited information on the regulation mechanism of TJ proteins that are key factors for BEB formation. Here, we determined the expression and localization patterns of TJ proteins, including peripheral membrane proteins, three ZOs (ZO-1, ZO-2, and ZO-3), and the transmembrane proteins, occludin and Cldn3, in the immature and mature goat epididymis.

Distribution of tight junction proteins in the goat epididymis

ZO proteins comprise a family of TJ-associated proteins that have a functional role in linking transmembrane proteins to the actin-based cytoskeleton [21]. In the present study, all three ZOs (ZO-1, ZO-2, and ZO-3) were present and localized along the apical regions of adjacent epithelial cells of both the immature and mature epididymis. These results agree with those of a previous study showing that all three ZOs were present in the TJs of the adult epididymis of mice [31]. In addition, we suggest that in the goat epididymis, their expression levels are different between the




Fig. 4. Immunohistochemical expression and localization of occludin in the goat epididymis. The epididymis was immunostained for zona occludens (ZO)-1 (green) and occludin (red). (A, C) Epididymis at postnatal month (PNM) 3. (B, D) Epididymis at PNM14. Yellow and white arrowheads indicate an overlap in ZO-1 and occludin staining and occludin single staining, respectively. Insets indicate ZO-1 single staining (Bar = 20 μ m). (E) Quantification of occludin labeling in regions co-localized with ZO-1 in TJs revealed a reduction in the occludin expression level in the mature epididymides compared to that in the immature ones. (F) Western blot analysis was applied to investigate the occludin expression in the epididymis at PNM3 and PNM14. * $p < 0.05$, ** $p < 0.001$.




Fig. 5. Immunohistochemical expression and localization of claudins3 (Cldn3) in the goat epididymis. The epididymis was immunostained for zona occludens [ZO]-1 (green) and Cldn3 (red). (A, C) Epididymis at postnatal month (PNM) 3. (B, D) Epididymis at PNM14. Yellow and white arrowheads indicate an overlap in ZO-1 and Cldn3 staining and Cldn3 single staining, respectively. Insets indicate ZO-1 single staining (Bars = 20 μ m). (E) Quantification of Cldn3 labeling in regions co-localized with ZO-1 in TJs revealed a reduction in the Cldn3 expression level in the mature epididymides compared to that in the immature ones. (F) Western blot analysis was applied to investigate Cldn3 expression in the epididymis at PNM3 and PNM14. * $p < 0.05$, ** $p < 0.001$.

immature and mature stages. For example, all three ZO proteins were detected at significantly lower levels in the mature epididymides than in the immature ones (Figs. 1, 2, and 3). Previous studies have demonstrated that ZO-1/ZO-2 and ZO-1/ZO-3 complexes are present, whereas ZO-2 is not associated with ZO-3 [21]. Moreover, we previously demonstrated the protein expression compensatory relationship between ZO-2 and ZO-3 without affecting ZO-1 expression levels in the epididymis of mice [31]. However, in the present study, we did not observe any compensation between ZO-2 and ZO-3 during goat epididymis development. These unexpected findings suggest the possibilities that other TJ proteins could contribute to BEB formation in the mature epididymis instead of ZO proteins or that specific relationships among these three ZO proteins might create an optimal system in this organ. Further studies are needed to clarify the reason underlying the reduction of all three ZO proteins in the mature epididymis.

We also observed that occludin and Cldn3 proteins, transmembrane proteins associated with ZO-1, ZO-2, and ZO-3 were localized and expressed in TJs. The localization and expression of a few types of transmembrane proteins in the epididymis have been previously reported in various species, including pigs [32], canines [33], rabbits [34], mice [31], and rats [35]. Occludin expression in TJs appears from embryonic day 18.5 in the mouse epididymis and its expression is maintained until adulthood [36]. Various Cldns were also observed in the epididymis, including Cldn3, which is expressed in the TJs and basolateral membranes in mice [31], rats [37], and bats [38]. In agreement with these previous studies, we found that occludin was present in TJs and that Cldn3 was expressed in both the basolateral membranes and TJs in the goat epididymis. Moreover, our results indicated much higher expression of occludin and Cldn3 in the immature epididymis, as high levels of ZO proteins were also observed. This might be because the high expression of ZO proteins in the immature epididymis can increase the chance of an association with occludin and Cldn3. Otherwise, the reduction of ZO proteins could result in a decreased chance of binding to other transmembrane TJ proteins, such as occludin and Cldn3. All TJ proteins examined in this study were expressed at significantly higher levels in the immature epididymis than in the mature epididymis. These findings suggest that some factors regulating the expression of TJ proteins are likely to be present during epididymis development. Additional studies investigating the expression and localization patterns of other TJ proteins are needed to elucidate these unexpected issues in the goat epididymis.

Regulation of tight junction proteins in the goat epididymis

Testicular luminal factors (TLFs) might be important in regulating changes in the distribution of TJ proteins. TLFs synthesized and secreted from the testis, include hormones, growth factors, and other luminal nutrients, which are directly in contact with epididymal epithelial cells that contribute to the formation of TJs and play an important role in the development of the epididymis [1, 39–41]. According to a previous study, the secretion of testicular luminal fluid might occur only in the normal mature testes of many species [42]. Therefore, it is possible that the epithelial cells exposed to TLFs during puberty might contribute to changes in the levels of TJ proteins in the mature epididymis. We previously reported that the epithelium is not perfectly differentiated until PNM2 [43] and PNM3 (unpublished data). Thus, the undifferentiated epithelium has a stronger TJ association between epithelial cells, and those that are differentiated by contact with TLFs could reduce their TJ association at the epididymal epithelium in goats. The reduction of these TJ proteins might modify the epididymal luminal contents that make up a unique environment in the adult epididymis for sperm maturation and storage. In addition, among TLFs, an androgen hormone known to be the most important key factor is responsible for regulating and maintaining epididymal structure and function. Although not well understood, an association exists between

TJs and androgens. Several studies have shown that androgens regulate TJ molecules including Cldn3, Cldn11, occludin, and ZO-1 [44–46] and are crucial for the regulation of every aspect of spermatogenesis via formation of the BTB [47,48]. However, direct links between androgens and TJ proteins in the epididymis have not been well documented. Therefore, more research on the effect of androgens on BEB formation is needed to confirm this assumption. Moreover, estrogen, another steroid hormone, is a good candidate factor that controls the expression and localization of TJ proteins. Estrogens are known as the most important female hormone, but in males, estrogens are critical to maintain testicular and epididymal function [49,50]. Without estrogens, male animals are infertile owing to failed fluid reabsorption, which is necessary for sperm concentration maintenance in the efferent ductules of the epididymis [51]. Earlier studies also provided some evidence that estrogens negatively control BTB function. Exposure to estrogens or diethylstilbestrol (a synthetic nonsteroidal estrogen) delays BTB formation and induces spermatogenesis failure in rats [27,52]. Moreover, bisphenol A (an estrogenic environmental toxicant) treatment induces disruption of the TJ barrier by changing the expression and localization of transmembrane proteins at the BTB [53]. Therefore, it is possible that an increased exposure to estrogens in the adult epididymis might decrease the expression levels of TJ proteins in the BEB.

Based on the present results, ZO-1, ZO-2, ZO-3, occludin, and Cldns are expressed and localized at the TJs of immature and mature epididymides of goats. Moreover, all TJ proteins were expressed at much higher levels in the immature epididymis than in the mature epididymis. This finding implies that undifferentiated epithelial cells might form a stronger BEB through high expression levels of TJ proteins, and then, the expression of TJ proteins decreases with the differentiation of epithelial cells in the goat epididymis. Our findings could contribute to a better understanding of the organization of TJ proteins in the BEB during the development of the goat epididymis. Although we could not directly explain the regulatory mechanism of TJ proteins in the BEB, our findings further the understanding of the reproductive physiological characteristics of the goat epididymis.

REFERENCES

1. Breton S, Ruan YC, Park YJ, Kim B. Regulation of epithelial function, differentiation, and remodeling in the epididymis. *Asian J Androl.* 2016;18:3-9. <https://doi.org/10.4103/1008-682X.165946>
2. Shum WWC, Ruan YC, Da Silva N, Breton S. Establishment of cell-cell cross talk in the epididymis: control of luminal acidification. *J Androl.* 2011;32:576-86. <https://doi.org/10.2164/jandrol.111.012971>
3. Shum WWC, Da Silva N, Brown D, Breton S. Regulation of luminal acidification in the male reproductive tract via cell-cell crosstalk. *J Exp Biol.* 2009;212:1753-61. <https://doi.org/10.1242/jeb.027284>
4. Cyr DG, Gregory M, Dubé É, Dufresne J, Chan PTK, Hermo L. Orchestration of occludins, claudins, catenins and cadherins as players involved in maintenance of the blood-epididymal barrier in animals and humans. *Asian J Androl.* 2007;9:463-75. <https://doi.org/10.1111/j.1745-7262.2007.00308.x>
5. Mital P, Hinton BT, Dufour JM. The blood-testis and blood-epididymis barriers are more than just their tight junctions. *Biol Reprod.* 2011;84:851-8. <https://doi.org/10.1095/biolreprod.110.087452>
6. França LR, Auharek SA, Hess RA, Dufour JM, Hinton BT. Blood-tissue barriers: morphofunctional and immunological aspects of the blood-testis and blood-epididymal

- barriers. In: Cheng CY editor. *Biology and regulation of blood-tissue barriers*. New York, NY: Springer Science Business Media; 2013. pp. 237-59.
7. Dym M, Fawcett DW. The blood-testis barrier in the rat and the physiological compartmentation of the seminiferous epithelium. *Biol Reprod*. 1970;3:308-26. <https://doi.org/10.1093/biolreprod/3.3.308>
 8. Fawcett DW, Leak LV, Heidger PM Jr. Electron microscopic observations on the structural components of the blood-testis barrier. *J Reprod Fertil Suppl*. 1970;10:105-22.
 9. Hoffer AP, Hinton BT. Morphological evidence for a blood-epididymis barrier and the effects of gossypol on its integrity. *Biol Reprod*. 1984;30:991-1004. <https://doi.org/10.1095/biolreprod30.4.991>
 10. Agarwal A, Hoffer AP. Ultrastructural studies on the development of the blood-epididymis barrier in immature rats. *J Androl*. 1989;10:425-31. <https://doi.org/10.1002/j.1939-4640.1989.tb00132.x>
 11. Setchell BP. The movement of fluids and substances in the testis. *Aust J Biol Sci*. 1986;39:193-207. <https://doi.org/10.1071/BI9860193>
 12. Banks WA, McLay RN, Kastin AJ, Sarmiento U, Scully S. Passage of leptin across the blood-testis barrier. *Am J Physiol*. 1999;276:E1099-104. <https://doi.org/10.1152/ajpendo.1999.276.6.E1099>
 13. Meng J, Holdcraft RW, Shima JE, Griswold MD, Braun RE. Androgens regulate the permeability of the blood-testis barrier. *Proc Natl Acad Sci USA*. 2005;102:16696-700. <https://doi.org/10.1073/pnas.0506084102>
 14. Hinton BT, Howards SS. Rat testis and epididymis can transport [3H] 3-O-methyl-D-glucose, [3H] inositol and [3H] α -aminoisobutyric acid across its epithelia in vivo. *Biol Reprod*. 1982;27:1181-9. <https://doi.org/10.1095/biolreprod27.5.1181>
 15. Hinton BT, Hernandez H. Neutral amino acid absorption by the rat epididymis. *Biol Reprod*. 1987;37:288-92. <https://doi.org/10.1095/biolreprod37.2.288>
 16. Hedger MP. Immunophysiology and pathology of inflammation in the testis and epididymis. *J Androl*. 2011;32:625-40. <https://doi.org/10.2164/jandrol.111.012989>
 17. Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA. Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. *J Cell Biol*. 1986;103:755-66. <https://doi.org/10.1083/jcb.103.3.755>
 18. Gumbiner B, Lowenkopf T, Apatira D. Identification of a 160-kDa polypeptide that binds to the tight junction protein ZO-1. *Proc Natl Acad Sci USA*. 1991;88:3460-4. <https://doi.org/10.1073/pnas.88.8.3460>
 19. Haskins J, Gu L, Wittchen ES, Hibbard J, Stevenson BR. ZO-3, a novel member of the MAGUK protein family found at the tight junction, interacts with ZO-1 and occludin. *J Cell Biol*. 1998;141:199-208. <https://doi.org/10.1083/jcb.141.1.199>
 20. Itoh M, Nagafuchi A, Moroi S, Tsukita S. Involvement of ZO-1 in cadherin-based cell adhesion through its direct binding to α catenin and actin filaments. *J Cell Biol*. 1997;138:181-92. <https://doi.org/10.1083/jcb.138.1.181>
 21. Wittchen ES, Haskins J, Stevenson BR. Protein interactions at the tight junction: Actin has multiple binding partners, and ZO-1 forms independent complexes with ZO-2 and ZO-3. *J Biol Chem*. 1999;274:35179-85. <https://doi.org/10.1074/jbc.274.49.35179>
 22. Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S, et al. Occludin: a novel integral membrane protein localizing at tight junctions. *J Cell Biol*. 1993;123:1777-88. <https://doi.org/10.1083/jcb.123.6.1777>
 23. Furuse M, Fujita K, Hiiragi T, Fujimoto K, Tsukita S. Claudin-1 and -2: novel integral

- membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J Cell Biol.* 1998;141:1539-50. <https://doi.org/10.1083/jcb.141.7.1539>
24. Van Itallie CM, Anderson JM. Claudins and epithelial paracellular transport. *Annu Rev Physiol.* 2006;68:403-29. <https://doi.org/10.1146/annurev.physiol.68.040104.131404>
 25. Furuse M, Itoh M, Hirase T, Nagafuchi A, Yonemura S, Tsukita S, et al. Direct association of occludin with ZO-1 and its possible involvement in the localization of occludin at tight junctions. *J Cell Biol.* 1994;127:1617-26. <https://doi.org/10.1083/jcb.127.6.1617>
 26. Itoh M, Furuse M, Morita K, Kubota K, Saitou M, Tsukita S. Direct binding of three tight junction-associated MAGUKs, ZO-1, ZO-2, and ZO-3, with the COOH termini of claudins. *J Cell Biol.* 1999;147:1351-63. <https://doi.org/10.1083/jcb.147.6.1351>
 27. Vitale R, Fawcett DW, Dym M. The normal development of the blood-testis barrier and the effects of clomiphene and estrogen treatment. *Anat Rec.* 1973;176:333-44. <https://doi.org/10.1002/ar.1091760309>
 28. Russell LD, Peterson RN. Sertoli cell junctions: morphological and functional correlates. *Int Rev Cytol.* 1985;94:177-211. [https://doi.org/10.1016/S0074-7696\(08\)60397-6](https://doi.org/10.1016/S0074-7696(08)60397-6)
 29. Kaitu'u-Lino TJ, Sluka P, Foo CFH, Stanton PG. Claudin-11 expression and localisation is regulated by androgens in rat Sertoli cells in vitro. *Reproduction.* 2007;133:1169-79. <https://doi.org/10.1530/REP-06-0385>
 30. Florin A, Maire M, Bozec A, Hellani A, Chater S, Bars R, et al. Androgens and postmeiotic germ cells regulate claudin-11 expression in rat Sertoli cells. *Endocrinology.* 2005;146:1532-40. <https://doi.org/10.1210/en.2004-0834>
 31. Kim B, Breton S. The MAPK/ERK-signaling pathway regulates the expression and distribution of tight junction proteins in the mouse proximal epididymis. *Biol Reprod.* 2016;94:22. <https://doi.org/10.1095/biolreprod.115.134965>
 32. Park YJ, Kim B. The Expression and localization of ZO-1, claudin 1, and claudin 4 in the pig epididymis. *J Anim Reprod Biotechnol.* 2019;34:190-6. <https://doi.org/10.12750/JARB.34.3.190>
 33. Ahn C, Shin DH, Lee D, Kang SM, Seok JH, Kang HY, et al. Expression of claudins, occludin, junction adhesion molecule A and zona occludens 1 in canine organs. *Mol Med Rep.* 2016;14:3697-703. <https://doi.org/10.3892/mmr.2016.5725>
 34. Chen W, Hu J, Zhang Z, Chen L, Xie H, Dong N, et al. Localization and expression of zonula occludens-1 in the rabbit corneal epithelium following exposure to benzalkonium chloride. *PLOS ONE.* 2012;7:e40893. <https://doi.org/10.1371/journal.pone.0040893>
 35. Morrow CMK, Mruk D, Cheng CY, Hess RA. Claudin and occludin expression and function in the seminiferous epithelium. *Philos Trans R Soc B Biol Sci.* 2010;365:1679-96. <https://doi.org/10.1098/rstb.2010.0025>
 36. Cyr DG, Hermo L, Egenberger N, Mertineit C, Trasler JM, Laird DW. Cellular immunolocalization of occludin during embryonic and postnatal development of the mouse testis and epididymis. *Endocrinology.* 1999;140:3815-25. <https://doi.org/10.1210/endo.140.8.6903>
 37. Gregory M, Cyr DG. Identification of multiple claudins in the rat epididymis. *Mol Reprod Dev.* 2006;73:580-8. <https://doi.org/10.1002/mrd.20467>
 38. Castro MM, Kim B, Games PD, Hill E, Neves CA, Serrão JE, et al. Distribution pattern of ZO-1 and claudins in the epididymis of vampire bats. *Tissue Barriers.* 2020;8:1779526. <https://doi.org/10.1080/21688370.2020.1779526>
 39. Rodríguez CM, Hinton BT. The testicular and epididymal luminal fluid microenvironment. In: Tulsiani DRP, editor. *Introduction to mammalian reproduction.* Boston, MA: Springer; 2003.

- pp. 61-77.
40. Kim B, Roy J, Shum WWC, Da Silva N, Breton S. Role of testicular luminal factors on basal cell elongation and proliferation in the mouse epididymis. *Biol Reprod.* 2015;92:9. <https://doi.org/10.1095/biolreprod.114.123943>
 41. Gregory M, Dufresne J, Hermo L, Cyr DG. Claudin-1 is not restricted to tight junctions in the rat epididymis. *Endocrinology.* 2001;142:854-63. <https://doi.org/10.1210/endo.142.2.7975>
 42. Setchell BP. The secretion of fluid by the testes of rats, rams and goats with some observations on the effect of age, cryptorchidism and hypophysectomy. *J Reprod Fertil.* 1970;23:79-85. <https://doi.org/10.1530/jrf.0.0230079>
 43. Jeong YD, Park YJ, Ko YG, Lee SS, Lee SH, Lee J, et al. Development and differentiation of epididymal epithelial cells in Korean native black goat. *Animals.* 2020;10:1273. <https://doi.org/10.3390/ani10081273>
 44. Wang RS, Yeh S, Chen LM, Lin HY, Zhang C, Ni J, et al. Androgen receptor in Sertoli cell is essential for germ cell nursery and junctional complex formation in mouse testes. *Endocrinology.* 2006;147:5624-33. <https://doi.org/10.1210/en.2006-0138>
 45. Willems A, Batlouni SR, Esnal A, Swinnen JV, Saunders PTK, Sharpe RM, et al. Selective ablation of the androgen receptor in mouse Sertoli cells affects sertoli cell maturation, barrier formation and cytoskeletal development. *PLOS ONE.* 2010;5:e14168. <https://doi.org/10.1371/journal.pone.0014168>
 46. Willems A, De Gendt K, Allemeersch J, Smith LB, Welsh M, Swinnen JV, et al. Early effects of Sertoli cell-selective androgen receptor ablation on testicular gene expression. *Int J Androl.* 2010;33:507-17. <https://doi.org/10.1111/j.1365-2605.2009.00964.x>
 47. Verhoeven G, Willems A, Denolet E, Swinnen JV, De Gendt K. Androgens and spermatogenesis: lessons from transgenic mouse models. *Philos Trans R Soc B Biol Sci.* 2010;365:1537-56. <https://doi.org/10.1098/rstb.2009.0117>
 48. Wang RS, Yeh S, Tzeng CR, Chang C. Androgen receptor roles in spermatogenesis and fertility: lessons from testicular cell-specific androgen receptor knockout mice. *Endocr Rev.* 2009;30:119-32. <https://doi.org/10.1210/er.2008-0025>
 49. Carreau S, Hess RA. Oestrogens and spermatogenesis. *Philos Trans R Soc B Biol Sci.* 2010;365:1517-35. <https://doi.org/10.1098/rstb.2009.0235>
 50. Joseph A, Shur BD, Hess RA. Estrogen, efferent ductules, and the epididymis. *Biol Reprod.* 2011;84:207-17. <https://doi.org/10.1095/biolreprod.110.087353>
 51. Hess RA, Bunick D, Lee KH, Bahr J, Taylor JA, Korach KS, et al. A role for oestrogens in the male reproductive system. *Nature.* 1997;390:509-12. <https://doi.org/10.1038/37352>
 52. Toyama Y, Ohkawa M, Oku R, Maekawa M, Yuasa S. Neonatally administered diethylstilbestrol retards the development of the blood-testis barrier in the rat. *J Androl.* 2001;22:413-23. <https://doi.org/10.1002/j.1939-4640.2001.tb02197.x>
 53. Li MWM, Mruk DD, Lee WM, Cheng CY. Disruption of the blood-testis barrier integrity by bisphenol A in vitro: is this a suitable model for studying blood-testis barrier dynamics? *Int J Biochem Cell Biol.* 2009;41:2302-14. <https://doi.org/10.1016/j.biocel.2009.05.016>