



# Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma

Buyanbileg Sodnom-Ish, Mi Young Eo, Hoon Myoung, Jong Ho Lee, Soung Min Kim

*Department of Oral and Maxillofacial Surgery, School of Dentistry, Dental Research Institute, Seoul National University, Seoul, Korea*

**Abstract** (J Korean Assoc Oral Maxillofac Surg 2022;48:3-12)

Selection of potential disease-specific biomarkers from saliva or epithelial tissues through next generation sequencing (NGS)-based protein studies has recently become possible. The early diagnosis of oral squamous cell carcinoma (OSCC) has been difficult, if not impossible, until now due to the lack of an effective OSCC biomarker and efficient molecular validation method. The aim of this study was to summarize the advances in the application of NGS in cancer research and to propose potential proteomic and genomic saliva biomarkers for NGS-based study in OSCC screening and diagnosis programs. We have reviewed four categories including definitions and use of NGS, salivary biomarkers and OSCC, current biomarkers using the NGS-based technique, and potential salivary biomarker candidates in OSCC using NGS.

**Key words:** Next generation sequencing, Saliva, Biomarkers, Early diagnosis, Oral squamous cell carcinoma

*[paper submitted 2021. 12. 6 / accepted 2022. 1. 11]*

## I. Introduction

Oral squamous cell carcinoma (OSCC) accounts for 90%-96% incidence of whole head and neck cancers, but there are no sensitive biomarkers for detection of OSCC. Definitive diagnosis has only been possible after examination of removed specimens. This diagnosis has been based on pathological findings such as angiogenesis, proliferation, and metastasis.

Even with radical resection combined with chemotherapy and concurrent radiation therapy, many OSCC cases demonstrate metastasis or recurrence without related or predictive symptoms. Complete eradication, early diagnosis, and metastatic and prognostic factor discovery have been research aims for OSCC management; until now, however, fulfillment of these aims has been elusive.

Like many cancers, early-stage OSCC is difficult to detect.

A significant number of patients do not seek clinical care until the OSCC is in an advanced stage. The development and advancement of screening and early diagnosis approaches has been recommended as the most effective strategy for reducing the OSCC-related morbidity and mortality rate<sup>1</sup>. Biomarkers found in the saliva are an ideal non-invasive diagnostic tool for early diagnosis of cancer<sup>2-4</sup>. Compared to blood and tissue sampling, saliva sampling is a reliable, non-invasive, convenient, and economical alternative for disease diagnosis and prognosis determination<sup>5</sup>. Saliva sampling also provides an effective and easily-acquired liquid specimen for large-scale sampling, epidemiologic screening and long-term monitoring<sup>6</sup>.

Several DNA or RNA sequencing techniques using salivary specimens have been used for karyotyping or submicroscopic chromosomal copy number changes such as microdeletions. These techniques include immunoprecipitation high-performance liquid chromatography, radioimmunoassay, electrophoretic immunoassay, mass spectrometry-based proteomics, microchips or microarrays, microfluidic devices, electrochemical biosensors<sup>7,8</sup>, fluorescence *in situ* hybridization and comparative genomic hybridization microarrays. Maxam and Gilbert<sup>9</sup> DNA sequencing was first developed in 1976 and is based on nucleobase-specific partial chemical modification of DNA and subsequent cleavage of the DNA backbone at sites

---

### Soung Min Kim

*Department of Oral and Maxillofacial Surgery, School of Dentistry, Dental Research Institute, Seoul National University, 101 Daehak-ro, Jongno-gu, Seoul 03080, Korea*

*TEL: +82-2-2072-0213*

*E-mail: smin5@snu.ac.kr*

*ORCID: <https://orcid.org/0000-0002-6916-0489>*

*© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.*

*Copyright © 2022 The Korean Association of Oral and Maxillofacial Surgeons.*

adjacent to the modified nucleotides. This sequencing method was advanced by Sanger et al.<sup>10</sup> in 1977. Sanger sequencing<sup>10</sup>, also known as chain termination sequencing, was marketed commercially by Applied Biosystems in 1986<sup>11</sup> and was the most popular DNA sequencing method until now. These two methods are based on the dideoxy method, the selective incorporation of chain-terminating dideoxynucleotides by DNA polymerase during *in vitro* DNA replication<sup>10,12</sup>. These first generation sequencing methods have been improved upon in next-generation sequencing (NGS).(Fig. 1)

NGS is a massive and parallel DNA sequencing technology for large-scale, ultra-high throughput, and automated high-speed genome analyses. NGS is a less expensive method for determining the order of nucleotides in entire genomes or targeted regions of DNA or RNA and has revolutionized the biological sciences. NGS has a wide variety of applications for the study of biological systems at a new level. NGS can be used to sequence entire genomes or specific areas of interest<sup>13</sup>. Sanger sequencing continues to be useful for smaller-scale, short-read sequencing analysis and for the validation of NGS results<sup>14</sup>.


The effectiveness of NGS in whole exome sequencing by targeted sequencing of cancer-related genes and in RNA sequencing has been demonstrated. NGS-applied OSCC research has also identified various genetic alterations and detected mutations with low variant allele frequency<sup>15</sup>. NGS is also becoming an essential method in characterization of

salivary gland tumors<sup>16</sup>. However, NGS has only rarely been applied to the identification of salivary biomarkers of OSCC. We discuss four topics in this review article: definitions and use of NGS, salivary biomarkers and OSCC, current biomarkers using NGS-based technique, and potential candidates as NGS-based salivary biomarkers.

## II. Definitions and Use of NGS

In the field of functional genomics, NGS is the most useful method for DNA and RNA analysis. This highly reproducible tool can reveal single nucleotide polymorphisms, related gene variants or spliced transcripts without input of direct DNA or RNA features as in microarray procedures. NGS can also be applied to cDNA molecules for RNA sequencing by reverse transcription from candidate RNA and sequencing-library construction by massive parallel deep sequencing. The most popular NGS methods are summarized in Table 1<sup>17-31</sup>. Illumina Solexa sequencing uses a fluorescence-based Illumina platform to identify 100-150 bp of DNA by emitting a particular fluorescent signal to each chain of nucleic acid. Adaptors can fragment, ligate and anneal longer sequences randomly. Reading is carried out by polymerase chain reaction (PCR) amplification and creation of a unique spot that repeats. The product can be separated into a single strand for final sequencing.(Fig. 1)

The sequencing-by-sequence methods using pyrosequenc-



**Fig. 1.** Schematic drawing of the Maxam and Gilbert's chemical chain termination method for DNA sequencing developed in 1977 followed by Sanger's 'dideoxy method'<sup>9,10</sup>. (PCR: polymerase chain reaction) Buyanbileg Sodnom-Ish et al: Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. J Korean Assoc Oral Maxillofac Surg 2022

**Table 1.** Next generation sequencing platforms<sup>17</sup>

Platform	Template preparation	Detection method	NGS coverage (base)	Run time (day)	Gb per run	Essential descriptions
Roche 454 (Roche, Basel, Switzerland) <sup>18</sup>	Emulsion-based clonal amplification	Pyrosequencing	400*	0.42	0.40-0.60	First commercial platform for the NGS technology. The DNA amplification process is different from that of Illumina, which can sequence much longer reads <sup>19</sup> .
GS FLX Titanium (Roche) <sup>20</sup>			400*	0.42	0.035	Able to sequence 400-600 million base pairs per run with 400-500 base pair read lengths <sup>21</sup> .
Illumina MiSeq (Illumina, San Diego, CA, USA) <sup>22</sup>	Clonal bridge PCR	Reversible dye terminator	2×300	0.17-2.7	15	End-to-end sequencing solutions with reversible-terminator sequencing-by-synthesis. Smallest benchtop sequencer that can perform onboard cluster generation, amplification, genomic DNA sequencing, and data analysis in a single run. Performs both single- and paired-end runs with adjustable read lengths from 1×36 base pairs to 2×300 base pairs <sup>22</sup> .
Illumina HiSeq (Illumina) <sup>23</sup>			2×150	0.3-11	1,000	Generate up to 1,000 Gb per run with the highest yield of data greater than Phred quality score of 30 (Q30). 1 hour's cycle time can be reduced to 10 minutes <sup>23</sup> .
Illumina Genome Analyzer IIX (Illumina) <sup>24</sup>			2×150	2-14	95	Having a broad spectrum of genomic variation with short- and long-insert paired-end reads with insert sizes 200 bp to 5 kb. Used for studying the genome, epigenome, and transcriptome, and also yield greater than 85% of bases higher than Q30 at 2×50 bp <sup>24</sup> .
Life Technologies SOLiD4 (Life Technologies, Waltham, MA, USA) <sup>25</sup>	Emulsion-based clonal amplification	Oligonucleotide ligation detection	35-50	4-7	35-50	Generates 10 <sup>8</sup> -10 <sup>9</sup> small sequence reads at one time, and two-base base encoding to decode the raw data. This system utilizes four fluorescent dyes to interrogate all sixteen (4 <sup>2</sup> ) possible two-base combinations, by a number of probes. Each probe is eight nucleotides long (8-mer) <sup>26</sup> .
Life Technologies Ion Protons (Thermo Fisher Scientific, Waltham, MA, USA) <sup>27</sup>		Native deoxy-ribonucleotide triphosphates, proton detection	200	0.5	100	Does not use fluorescence or chemiluminescence. Instead, measures the H <sup>+</sup> ions released during base incorporation. The lack of any optics allows rapid expansion of the output by approximately 10-fold every six months <sup>27</sup> .
Complete Genomics (Complete Genomics, San Jose, CA, USA) <sup>28</sup>	Gridded DNA-nanoballs	Oligonucleotide ligation detection	7×10	11	3,000	A DNA nanoball sequencing, which assembles short DNA sequences into a full genome. The sequences are obtained by probe-ligation, but the clonal DNA amplification is performed by rolling circle amplification unlike the bead or emulsion amplification <sup>28</sup> .
Helicos Biosciences Heliscope (Helicos Biosciences, Cambridge, MA, USA) <sup>29</sup>	Single molecule	Reversible dye terminator	35*	8	25	A highly sensitive fluorescence detection system for direct interrogation of single DNA molecules via sequencing by synthesis <sup>30</sup> .
Pacific Biosciences SMRT (Pacific Biosciences, Menlo Park, CA, USA) <sup>31</sup>		Phospholinked fluorescent nucleotides or real-time sequencing	10,000 (N50); 30,000+ (max)	0.08	0.5	Long-read sequencing platforms with SMRT sequencing technology. Template preparation does not require any amplification steps, and the prepared library molecule is the sequencing template. The adapters have a hairpin structure (SMRT loop adapters) so that after ligation the double stranded DNA fragments will have become circular <sup>31</sup> .

(PCR: polymerase chain reaction, NGS: next generation sequencing, SMRT: single molecule real-time)

\*Average read lengths for the Roche 454 and Helicos Biosciences platforms.

Q30 is equivalent to the probability of incorrect base call 1 in 1,000 times with 99.9% base of accuracy. Run times and gigabase (Gb) output per run for single-end sequencing are noted. Run times and outputs approximately double when performing pair-end sequencing.

Buyanbileg Sodnom-Ish et al: Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. *J Korean Assoc Oral Maxillofac Surg* 2022

ing is the basis of Roche 454 platform sequencing. The release and incorporation of pyrophosphate into a new strand by polymerase is detected by fluorescence. Multiple reads over 1,000 bp long can be carried out by optical signal detection. One DNA fragment per bead can be annealed, and generic adaptors can add bases to the ends. The product is amplified by PCR as a routine adaptor-specific primers application.

The principle of measuring the direct hydrogen ion release from individual bases, not fluorescence, is the basic principle of Ion torrent and proton platform sequencing. Direct measurement of the emission of hydrogen ions by polymerase during the incorporation of deoxynucleoside triphosphate to a growing DNA strand. This is executed by uniform 200 to 400 bp fragmentation, and adapters are added to effect amplification by emulsion PCR.

### III. Salivary Biomarkers and Oral Cancer

Saliva sampling is rapidly expanding; and arrays of analytes including proteins, messenger RNA (mRNA), and DNA in saliva have been studied for their potential use as biomarkers for OSCC screening programs<sup>32,33</sup>. OSCC is a malignancy in which salivary diagnosis has the greatest potential because the OSCC environment contains saliva. One of the useful aspects of saliva is its containment of exfoliated tissues or cells from the oral cavity. This aspect suggests the existence of potential salivary biomarkers for OSCC<sup>33</sup>. A number of candidate salivary biomarkers for oral cancer, including genomic and proteomic biomarkers for OSCC, have been reported and are summarized in Table 2.

Proteomic studies using human saliva have been targeting the biological activities of various peptides and proteins in normal individuals and in those with various pathologies. Proteomic targets such as CD59, catalase, M2BP, and MRP14 were suggested from a shotgun proteome analysis for oral cancer detection<sup>34</sup>. Li et al.<sup>35</sup> reported the presence of more than 3000 RNA species, mostly mRNAs; and reports of Park et al.<sup>36</sup> and Patel et al.<sup>37</sup> support the role of microRNAs (miRNAs) in oral cancer progression and various cancers.

Several salivary biomarkers such as DAPK, TIMP3, p16, and MGMT are potential screening markers for OSCC<sup>38</sup>, and changes in DNA methylation patterns could be a useful screening tool for predicting the rate and the likelihood of malignant transformation. The methylation of DNA in saliva has been suggested as an effective biomarker for early detection of OSCC<sup>39</sup>. For example, hypermethylation of the DNA

**Table 2.** Candidate salivary biomarkers for oral cancer, based on the biomolecule markers<sup>1</sup>

Type of biomolecular markers	Biomarkers
Genomic	p53 Promoter hypermethylation of DAPK, TIMP3, p16, and MGMT genes Cyclin D1 gene amplification
mRNA	Maspin IL-8 IL-1β S100P SAT miR 31, miR 125, miR 200a
Protein	Elevated CD44 IL-6 Intermediate filament protein (Cyfra 21-1) 8-OHdG Albumin Glutathione Actin and myosin L-phenylalanine
Others	EFNB2, ANGPT1, ANGPT2, CD31, VEGF HPV and EBV Inorganic compounds: Na, Ca, F, Mg Fucose

(DAPK: death-associated protein kinase, TIMP3: tissue inhibitor of metalloproteinase-3, MGMT: O6-methylguanine-DNA methyltransferase, Maspin: mammary serine protease inhibitor, IL-8: interleukin 8, IL-1β: interleukin 1 beta, S100P: S100 calcium binding protein P, SAT: spermidine/spermineN1-acetyltransferase, miR: microRNA, CD44: cluster of differentiation 44, 8-OHdG: 8-Oxo-2'-deoxyguanosine, ANGPT: angiopoietin, VEGF: vascular endothelial growth factor, HPV: human papilloma virus, EBV: Epstein-Barr virus)

*Buyanbileg Sodnom-Ish et al: Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. J Korean Assoc Oral Maxillofac Surg 2022*

promoter of certain genes, such as p16, has been found in saliva as well as serum<sup>40,41</sup>. Promoter hypermethylation has been suggested to be an early event during OSCC genesis. Any change of transcription factor or receptor functions or other factors in tumorigenesis might be involved in loss of cell cycle control. This loss of control was reported by Liao et al.<sup>42</sup>; this group suggested the first salivary biomarker, genomic p53. Other cell cycle regulatory proteins such as mammary serine protease inhibitor and cyclin D1 were also increased in OSCC saliva<sup>43</sup>.

Due to the basic precursor characteristics of mRNA for protein expression, other potential salivary biomarkers display increased mRNA production. These include IL-8, IL-1β, S100P, and spermidine/spermineN1-acetyltransferase<sup>44</sup>. The circulating levels of CD44 are reported to be related to head and neck cancer metastasis, and CD44 is a potential salivary protein biomarker<sup>45</sup>. Actin and myosin are also regarded as promising salivary biomarkers for premalignant differentiation and malignant oral lesions<sup>33</sup> IL-6 is another premalignant

**Table 3.** NGS-based salivary genomic markers in oral cancer

Potential genomic markers <sup>1</sup>	Sample/ Collection method	NGS-based studies
p53 (TP53, P53; BCC7; LFS1; BMFS5; TRP53) <sup>46,47</sup>	Tumor tissue/ Formalin fixation and paraffin embedding	Approximately 30% of salivary gland cancers have mutations of p53 gene. A significant worse overall and disease-free survival was shown in cancers with p53 mutations.
Promoter hypermethylation of DAPK gene <sup>48,49</sup>	UWS/ Oragene DNA Self-Collection kit	DAPK methylation gene was reported with the development of cancers in women.
Promoter hypermethylation of TIMP3 gene (SFD; K222; K222TA2; HSMRK222) <sup>50-52</sup>	Salivary rinse/ 10-20 mL 0.9% NaCl, 15-60 s Salivary rinse/ rinsing and gargling with 20 mL 0.9% NaCl	Promoter hypermethylation of TIMP3 levels found in the circulating tumor DNA in saliva of disease-free survival HNSCC patients.
Promoter hypermethylation of p16 (CDKN2A, ARF; MLM; P14; P19; CMM2; INK4) <sup>47,53</sup>	Tumor tissue/ Formalin fixation and paraffin embedding Salivary rinse/ swishing with 15 mL of 0.9% normal saline, 15-30 s	The absolute frequencies of CDKN2A mutations are detected by NGS.
Promoter hypermethylation of MGMT gene <sup>54</sup>	Tumor tissue/ Formalin fixation and paraffin embedding Salivary rinse/ swishing with 15 mL of 0.9% normal saline, 15-30 s	MGMT methylation was detected in 29% of oral cancer/ dysplasia patient, showing MGMT gene is related to DNA repair and aberrant promoter hypermethylation.
Cyclin D1 gene amplification (CCND1, BCL1; PRAD1; U21B31; D11S287E) <sup>55,56</sup>	Tumor tissue/ Formalin fixation and paraffin embedding	14 of the 21 genes had copy number amplifications and losses, which included CCND1 (83.7%).
Maspin <sup>57,58</sup>	UWS/ samples were collected in the morning, without any oral stimulation for 90 minutes before collection	Downregulation of maspin; the functional importance of maspin includes the inhibition of tumor angiogenesis. The levels of maspin were reduced.

(NGS: next generation sequencing, DAPK: death-associated protein kinase, UWS: unstimulated whole saliva, TIMP3: tissue inhibitor of metalloproteinase-3, HNSCC: head and neck squamous cell carcinoma, MGMT: O-6-methylguanine-DNA methyltransferase, Maspin: mammary serine protease inhibitor)

Buyanbileg Sodnom-Ish et al: Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. *J Korean Assoc Oral Maxillofac Surg* 2022

differentiation biomarker found especially in the saliva of oral leukoplakia patients. IL-6 inactivated the p53 tumor suppressor gene by hypermethylation of its promoter region resulting in uncontrolled cell proliferation and suppression of programmed cell death<sup>33</sup>.

Researchers have also directed their research toward detection of human papilloma virus (HPV) and Epstein–Barr virus (EBV) in saliva; these viruses are etiological factors in cancer. The incidence for HPV positivity has been reported to be more than 45% in patients treated for oral cancer<sup>33</sup>.

#### IV. Current Cancer Biomarkers Using NGS-based Technique

Currently, several authors have reported data from the search for biomarkers using a NGS-based technique<sup>36,46-69</sup>. (Tables 3, 4) Most of these studies started from Gene Expression Omnibus datasets (<https://www.ncbi.nlm.nih.gov/gds>) in combination with differentially expressed genes, the protein-protein interaction network, gene ontology and the Kyoto Encyclopedia of Genes and Genomes pathway. Shanmugam et al.<sup>70</sup> have reported a customized NGS analysis of unique coding regions of seven mutated genes from OSCC saliva, and similar data from whole-exome sequencing were also

discovered in tumors from The Cancer Genome Atlas<sup>71</sup>, the International Cancer Genome Consortium gingiva-buccal cohort<sup>69</sup>, and MD Anderson Cancer Center OSCC cohort<sup>72</sup>.

From this review, NGS-based salivary biomarkers in OSCC could be categorized primarily as genomic (Table 3) and transcriptome markers.(Table 4) Within these main categories, we could present promoter hypermethylation of p16 and three miRNAs as the main candidate salivary biomarkers. Especially, miRNAs, endogenous, non-coding, single-stranded RNA molecules 22 nucleotides long, have tumor-controlling characteristics including tumor-suppression.

Hypermethylation of the p16 promoter is a useful serum biomarker for the early detection of alimentary tract cancer, especially gastric cancer. Zammit et al.<sup>73</sup> reported on the etiology of OSCC using NGS while focusing on smoking and HPV factors in a prospective observational study. This group showed that the most frequent mutations were found in TP53 and CDKN2A in the salivary specimen.(Table 3) Fadhil et al.<sup>74</sup> reported five miRNAs using NGS data in saliva of 12 HNSCC patients and 12 healthy controls. Among these, miR-let-7a-5p and miR-3928 were suggested to be NGS-based salivary biomarkers for early diagnosis of HNSCC when compared with other miRNAs such as miR-7703, miR-345-5p, and miR-1470.(Table 4) A recent study also showed the

**Table 4.** NGS-based salivary mRNA markers in oral cancer

Potential mRNA markers <sup>1</sup>	Sample/ Collection method	NGS-based studies
IL-8 (IL8) <sup>59,60</sup>	UWS/ samples were collected between 9 am and 10 am, following the standard protocol.	In oral, esophageal, lung, pancreatic, ovarian and breast cancers, certain salivary mRNA biomarkers have been proposed as a possible cancer biomarker, including IL-8.
IL-1β (IL-1) <sup>61,62</sup>	UWS/ 3-5 mL salivary specimen was collected into a tube containing 10 mL of RNAlater (Ambion, Austin, TX, USA), an aqueous tissue storage reagent that rapidly permeated tissue to stabilize and protect cellular RNA. The specimen was then placed on ice at 4°C; cell-free saliva supernatant was harvested.	IL-1β gene has been found to be an important biomarker for ovarian cancer.
S100P (S100 calcium binding protein P) <sup>63,64</sup>	UWS/ samples were collected between 6 am and 12 pm following standard protocol. A maximum of 8 mL of saliva were collected within 30 minutes.	Salivary S100 mRNA is a candidate biomarker for detecting OSCC development and in OLP patients determined by NGS.
SAT <sup>69</sup>	UWS/ Participants were asked to refrain from eating, drinking and any oral hygiene overnight and spit in 5-mL plastic vials used for biochemical examinations for 5 minutes. During the whole procedure and until centrifugation the vials were kept in ice.	The combination of SAT and IL-8 mRNA biomarkers are attractive candidates either for screening or for early diagnosis purposes. These exert a very good prediction ability together with a high sensitivity and specificity for screening oral squamous cell carcinoma.
miR 31 (MIRN31; miR-31; hsa-mir-31) <sup>65,66</sup>	UWS/ 3-5 mL saliva was collected from mouth floor after simple rinsing. Pre-treatment salivary sample was collected from 45 patients with OSCC and 10 patients with oral verrucous leukoplakia and 24 healthy participants.	A significantly high expression of miR-31 was found in the saliva sample of patients with OSCC at all clinical stages by RT-qPCR. More miR-31 levels were detected in the saliva than in plasma, suggesting salivary miR-31 to be a more sensitive.
miR 125 (MIRN125A; miR-125a; miRNA125A) <sup>36,67</sup>	UWS/ an aqueous tissue storage reagent was used to preserve UWS samples and SUPERase.In (Thermo Fisher Scientific, Waltham, MA, USA) was used for supernatant saliva preservation.	Significantly lower levels of miR-125a was found in saliva sample of OSCC patient than that of healthy controls.
miR 200a <sup>36,68</sup>	UWS/ an aqueous tissue storage reagent was used for the UWS samples and SUPERase.In (Thermo Fisher Scientific) was used for supernatant saliva preservation.	Significantly lower levels of miR 200a was found in saliva sample of OSCC patient than that of healthy controls.

(NGS: next generation sequencing, IL-8: interleukin 8, IL-1β: interleukin 1 beta, UWS: unstimulated whole saliva, OSCC: oral squamous cell carcinoma, OLP: oral lichen planus, miR: microRNA, RT-qPCR: quantitative real-time reverse transcriptase-polymerase chain reaction) Standard protocol was summarized as 1) no drinking neither using any oral hygiene care on day of saliva collection, 2) mouth rinsing with water on 5 minutes prior, and 3) upright sitting position and spitting into a 50-mL Falcon tube kept on ice.

Buyanbileg Sodnom-Ish et al: Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. J Korean Assoc Oral Maxillofac Surg 2022

**Table 5.** Known salivary biomarker and recommended NGS-based potential salivary biomarker

	Genomic	Transcriptome (mRNA)
Known salivary biomarkers	p53 FAT1 CASP8 PIK3CA HRAS NOTCH1 CDKN2A <sup>70</sup>	miRNA (saliva): miR-let-7a-5p and miR-3928 <sup>74</sup> Oncogenic (up-regulated, tissue): miR-21, miR-22, miR-26a, miR34c, miR-34b, miR-117, miR-118, miR-130b, miR-135, miR-142, miR-143, miR-148a, miR-150, miR-221, miR-222, miR-423, miR-542, miR-1269a <sup>81</sup>
Recommended NGS-based salivary biomarker	Mutation of p53 gene Promoter hypermethylation of DAPK, TIMP3, p16, and MGMT genes Cyclin D1 gene amplification Mammary serine protease inhibitor	Suppressive (down-regulated, tissue): miR-92b, miR-199, miR-214, miR-375, miR-486, miR-504, miR-499, miR-486 <sup>81</sup> miR 31 <sup>76</sup> , miR 125 <sup>77</sup> , miR 200 <sup>78,79</sup>

(NGS: next generation sequencing, FAT1: AT atypical cadherin 1, CASPS: caspase 8, PIK3CA: phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha, HRAS: Harvey rat sarcoma viral oncogene homolog, NOTCH1: Notch homolog 1, translocation-associated, CDKN2A: cyclin-dependent kinase inhibitor 2A, DAPK: death-associated protein kinase, TIMP3: tissue inhibitor of metalloproteinase-3, MGMT: O6-methylguanine-DNA methyltransferase, miR: microRNA)

Buyanbileg Sodnom-Ish et al: Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. J Korean Assoc Oral Maxillofac Surg 2022

potential benefits of NGS for mRNA expression profiling by using miR143-3p to detect chronic periodontitis<sup>75</sup>.

Lu et al.<sup>76</sup> reported that miR-31-5p may be an independent OSCC biomarker by showing its tumor growth inhibition capacity in oral cancer patient-derived xenograft models. Wang et al.<sup>77</sup> showed that mature miR-125b could control metabolism and immunity of cancer cells by regulating NF- $\kappa$ B or p53 signaling pathways through translation inhibition of 3' untranslated regions of target mRNAs. Korpala and Kang<sup>78</sup> also showed that miR-200 family members inhibit epithelial-mesenchymal transition (EMT) and metastasis. This occurs through a miRNA-mediated regulatory pathway by direct targeting of transcriptional repressors of E-cadherin, ZEB1, and ZEB2. The validity of this report was strengthened by Kabzinski et al.<sup>79</sup> who showed that DNA methylation of the miR-200c promoter in an epithelial originated tumor may occur during EMT. Majewska et al.<sup>80</sup> discovered a potentially targetable novel anaplastic lymphoma kinase fusion in an intraductal carcinoma of a minor salivary gland by NGS analysis.

Compared with known genomic salivary biomarkers, such as p53, FAT1, CASP8, PIK3CA, HRAS, NOTCH1, and CDKN2A, NGS-based genomics consider mutation of p53 and promoter hypermethylation of DAPK, TIMP3, p16, and MGMT, cyclone D1 and mammary serine protease inhibitors. Also, up-regulated oncogenic tissue miRNAs and down-regulated tissue suppressive marker miRNAs, including miR 31, miR 125, and miR 3928, are recommended as subjects for further research<sup>70,74,76-79,81</sup>.(Table 5)

## V. Discussion

This review article aimed to summarize current applications of NGS in cancer research and to propose potential genomic and proteomic saliva biomarkers for NGS-based study in OSCC screening and diagnosis programs. For the establishment of a standardized research and designed protocol for the detection of trace protein or nucleic acids biomarkers from saliva samples, efficient and stable collection, processing and preservation methods should first be confirmed. For the collection of saliva samples, confirmation of the patient's mouth cleanness is essential. This can be accomplished by rinsing the mouth with water to remove substances and by ensuring that the patient avoids any eating or drinking for at least 30 minutes prior to sampling. Approximately 2.5 mL of saliva in the buffered solution is recommended. Preparation should include addition of 2.5 mL of DNA stabilization buffer into a 10- or 15-mL conical tube. Excessive saliva might

to be degraded due to insufficient DNA stabilization buffer, and insufficient saliva collection may not provide adequate results. The cap needs to be replaced, mixing needs to occur by inversion not shaking, and the specimen needs to be stored at room temperature for immediate use or at 4°C for future use.

NGS-based research has been used to identify factors in metastasis of lung, prostate, ovarian, and bile duct cancer. NGS-based evaluation of candidate biomarkers in saliva could also be routinely used as a simple, non-invasive test in patients with OSCC. NGS has also been used as a valuable tool for detecting novel biomarkers in periodontal disease<sup>75</sup>. More recently, NGS-based research has led the ability to differentiate patients with primary Sjögren's syndrome from those with non-Sjögren's sicca<sup>82</sup>.

In various medical and non-medical institutions, prevention of unnecessary medical expenses may be possible by the quick and easy diagnosis of oral cancer in healthy or suspect patients. There would be no additional cost to the public health system, and provision of information on the potential of benefits to patients with OSCC by using NGS based salivary biomarkers can be effected. Since both proteins and genes can be applied as biomarkers, the accuracy of diagnosis can be increased. New biomarkers can be added quickly for expansion of a preventive platform for a variety of medical and dental diseases. Since both proteins and genes can be applied as biomarkers, the accuracy of diagnosis can be increased, and new biomarkers can be added quickly, so that it can be expanded to a preventive platform through early diagnosis of various diseases as well as various medical and dental treatments.

In conclusion, selection of potential OSCC biomarkers through NGS-based protein studies from epithelial tissue cells in collected saliva will be possible. These non-invasive methods could be a useful tool for the improvement of an immune biosensor for ensuring general public health and for diagnosis of OSCC before its metastasis or progressive infiltration to adjacent tissues. Based on NGS analysis, we will be able to verify the biomarkers of OSCC in saliva and to effect early diagnosis of oral cancer based on this.

## ORCID

Buyanbileg Sodnom-Ish, <https://orcid.org/0000-0002-4239-1420>

Mi Young Eo, <https://orcid.org/0000-0001-7055-9924>

Hoon Myoung, <https://orcid.org/0000-0002-9984-8479>

Jong Ho Lee, <https://orcid.org/0000-0002-8843-545X>

Soung Min Kim, <https://orcid.org/0000-0002-6916-0489>

## Authors' Contributions

B.S.I. and M.Y.E. participated in data collection and wrote the manuscript. H.M. and J.H.L. participated in the study design and helped to draft the manuscript. S.M.K. coordinated and approved the final manuscript.

## Acknowledgements

This study was supported by grant No. 03-2021-0045 from the SNUDH Research Fund.

## Conflict of Interest

No potential conflict of interest relevant to this article was reported.

## References

1. Nguyen TTH, Sodnom-Ish B, Choi SW, Jung HI, Cho J, Hwang I, et al. Salivary biomarkers in oral squamous cell carcinoma. *J Korean Assoc Oral Maxillofac Surg* 2020;46:301-12. <https://doi.org/10.5125/jkaoms.2020.46.5.301>
2. Salazar C, Nagadia R, Pandit P, Cooper-White J, Banerjee N, Dimitrova N, et al. A novel saliva-based microRNA biomarker panel to detect head and neck cancers. *Cell Oncol (Dordr)* 2014;37:331-8. <https://doi.org/10.1007/s13402-014-0188-2>
3. Pfaffe T, Cooper-White J, Beyerlein P, Kostner K, Punyadeera C. Diagnostic potential of saliva: current state and future applications. *Clin Chem* 2011;57:675-87. <https://doi.org/10.1373/clinchem.2010.153767>
4. Schulz BL, Cooper-White J, Punyadeera CK. Saliva proteome research: current status and future outlook. *Crit Rev Biotechnol* 2013;33:246-59. <https://doi.org/10.3109/07388551.2012.687361>
5. Genco RJ. Salivary diagnostic tests. *J Am Dent Assoc* 2012;143(10 Suppl):3S-5S. <https://doi.org/10.14219/jada.archive.2012.0340>
6. Fábryová H, Celec P. On the origin and diagnostic use of salivary RNA. *Oral Dis* 2014;20:146-52. <https://doi.org/10.1111/odi.12098>
7. Campuzano S, Yanez-Sedeno P, Pingarron JM. Electrochemical bioaffinity sensors for salivary biomarkers detection. *TrAC Trends Anal Chem* 2017;86:14-24. <https://doi.org/10.1016/j.trac.2016.10.002>
8. Malon RS, Sadir S, Balakrishnan M, Córcoles EP. Saliva-based biosensors: noninvasive monitoring tool for clinical diagnostics. *Biomed Res Int* 2014;2014:962903. <https://doi.org/10.1155/2014/962903>
9. Maxam AM, Gilbert W. A new method for sequencing DNA. *Proc Natl Acad Sci U S A* 1977;74:560-4. <https://doi.org/10.1073/pnas.74.2.560>
10. Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* 1977;74:5463-7. <https://doi.org/10.1073/pnas.74.12.5463>
11. Adams J. DNA sequencing technologies. *Nat Educ* 2008;1:193.
12. Sanger F, Coulson AR. A rapid method for determining sequences in DNA by primed synthesis with DNA polymerase. *J Mol Biol* 1975;94:441-8. [https://doi.org/10.1016/0022-2836\(75\)90213-2](https://doi.org/10.1016/0022-2836(75)90213-2)
13. Behjati S, Tarpey PS. What is next generation sequencing? *Arch Dis Child Educ Pract Ed* 2013;98:236-8. <https://doi.org/10.1136/archdischild-2013-304340>
14. Goodwin S, McPherson JD, McCombie WR. Coming of age: ten years of next-generation sequencing technologies. *Nat Rev Genet* 2016;17:333-51. <https://doi.org/10.1038/nrg.2016.49>
15. Gabusi A, Gissi DB, Tarsitano A, Asioli S, Marchetti C, Montebugnoli L, et al. Intratumoral heterogeneity in recurrent metastatic squamous cell carcinoma of the oral cavity: new perspectives afforded by multiregion DNA sequencing and mtDNA analysis. *J Oral Maxillofac Surg* 2019;77:440-55. <https://doi.org/10.1016/j.joms.2018.09.014>
16. Todorovic E, Dickson BC, Weinreb I. Salivary gland cancer in the era of routine next-generation sequencing. *Head Neck Pathol* 2020;14:311-20. <https://doi.org/10.1007/s12105-020-01140-4>
17. Massive parallel sequencing [Internet]. San Francisco (CA): Wikipedia [cited 2021 May 30]. Available from: [https://en.wikipedia.org/wiki/Massive\\_parallel\\_sequencing](https://en.wikipedia.org/wiki/Massive_parallel_sequencing)
18. Harrington CT, Lin EI, Olson MT, Eshleman JR. Fundamentals of pyrosequencing. *Arch Pathol Lab Med* 2013;137:1296-303. <https://doi.org/10.5858/arpa.2012-0463-RA>
19. The Next Generation Sequencing Platform of Roche 454 [Internet]. Shirley (NY): Biogene Blog [cited 2021 May 30]. Available from: <https://www.creative-biogene.com/blog/index.php/2017/02/02/the-next-generation-sequencing-platform-of-roche-454/#:~:text=Roche%20454%20sequencing%20system%20is,the%20next%20generation%20sequencing%20technology.&text=DNA%20Library%20construction%20in%20454,different%20adapters%20at%20both%20ends>
20. Gilles A, Meglécz E, Pech N, Ferreira S, Malausa T, Martin JF. Accuracy and quality assessment of 454 GS-FLX Titanium pyrosequencing. *BMC Genomics* 2011;12:245. <https://doi.org/10.1186/1471-2164-12-245>
21. 454 Life sciences [Internet]. San Francisco (CA): Wikipedia [cited 2021 May 30]. Available from: [https://en.wikipedia.org/wiki/454\\_Life\\_Sciences](https://en.wikipedia.org/wiki/454_Life_Sciences)
22. Ravi RK, Walton K, Khosroheidari M. MiSeq: a next generation sequencing platform for genomic analysis. *Methods Mol Biol* 2018;1706:223-32. [https://doi.org/10.1007/978-1-4939-7471-9\\_12](https://doi.org/10.1007/978-1-4939-7471-9_12)
23. Illumina. HiSeq™ sequencing systems: redefining the trajectory of sequencing [Internet]. San Diego (CA): Illumina [cited 2021 May 30]. Available from: [https://www.illumina.com/documents/products/datasheets/datasheet\\_hiseq\\_systems.pdf](https://www.illumina.com/documents/products/datasheets/datasheet_hiseq_systems.pdf)
24. Illumina. Genome Analyzer<sub>ix</sub> system: the most proven, widely adopted next-generation sequencing platform [Internet]. San Diego (CA): Illumina [cited 2021 May 30]. Available from: [https://support.illumina.com/content/dam/illumina-marketing/documents/products/datasheets/datasheet\\_genome\\_analyzerix.pdf](https://support.illumina.com/content/dam/illumina-marketing/documents/products/datasheets/datasheet_genome_analyzerix.pdf)
25. Castellana S, Romani M, Valente EM, Mazza T. A solid quality-control analysis of AB SOLiD short-read sequencing data. *Brief Bioinform* 2013;14:684-95. <https://doi.org/10.1093/bib/bbs048>
26. ABI Solid Sequencing [Internet]. San Francisco (CA): Wikipedia [cited 2021 May 30]. Available from: [https://en.wikipedia.org/wiki/ABI\\_Solid\\_Sequencing](https://en.wikipedia.org/wiki/ABI_Solid_Sequencing)
27. Thermo Fisher Scientific. Ion Proton™ System for next-generation sequencing [Internet]. Seoul: Thermo Fisher Scientific [cited 2021 May 30]. Available from: <https://www.thermofisher.com/kr/ko/home/life-science/sequencing/next-generation-sequencing/ion-torrent-next-generation-sequencing-workflow/ion-torrent-next-generation-sequencing-run-sequence/ion-proton-system-for-next-generation-sequencing.html>
28. Complete genomics [Internet]. San Francisco (CA): Wikipedia [cited 2021 May 30]. Available from: [https://en.wikipedia.org/wiki/Complete\\_Genomics](https://en.wikipedia.org/wiki/Complete_Genomics)
29. Thompson JF, Steinmann KE. Single molecule sequencing with a HeliScope genetic analysis system. *Curr Protoc Mol Biol* 2010;Chapter 7:Unit7.10. <https://doi.org/10.1002/0471142727.mb0710s92>



30. Helicos single molecule fluorescent sequencing [Internet]. San Francisco (CA): Wikipedia [cited 2021 May 30]. Available from: [https://en.wikipedia.org/wiki/Helicos\\_single\\_molecule\\_fluorescent\\_sequencing](https://en.wikipedia.org/wiki/Helicos_single_molecule_fluorescent_sequencing)
31. PacBio. SMRT sequencing [Internet]. Menlo Park (CA): PacBio [cited 2021 May 30]. Available from: <https://www.pacb.com/smrt-science/smrt-sequencing/>
32. Wong DT. Salivaomics. *J Am Dent Assoc* 2012;143(10 Suppl):19S-24S. <https://doi.org/10.14219/jada.archive.2012.0339>
33. Shah FD, Begum R, Vajaria BN, Patel KR, Patel JB, Shukla SN, et al. A review on salivary genomics and proteomics biomarkers in oral cancer. *Indian J Clin Biochem* 2011;26:326-34. <https://doi.org/10.1007/s12291-011-0149-8>
34. Singh P, Verma JK, Singh JK. Validation of salivary markers, IL-1 $\beta$ , IL-8 and Lgals3bp for detection of oral squamous cell carcinoma in an Indian population. *Sci Rep* 2020;10:7365. <https://doi.org/10.1038/s41598-020-64494-3>
35. Li Y, Zhou X, St John MA, Wong DT. RNA profiling of cell-free saliva using microarray technology. *J Dent Res* 2004;83:199-203. <https://doi.org/10.1177/154405910408300303>
36. Park NJ, Zhou H, Elashoff D, Henson BS, Kastratovic DA, Abemayor E, et al. Salivary microRNA: discovery, characterization, and clinical utility for oral cancer detection. *Clin Cancer Res* 2009;15:5473-7. <https://doi.org/10.1158/1078-0432.CCR-09-0736>
37. Patel RS, Jakymiw A, Yao B, Pauley BA, Carcamo WC, Katz J, et al. High resolution of microRNA signatures in human whole saliva. *Arch Oral Biol* 2011;56:1506-13. <https://doi.org/10.1016/j.archoralbio.2011.05.015>
38. Rapado-González Ó, López-Cedrún JL, López-López R, Rodríguez-Ces AM, Suárez-Cunqueiro MM. Saliva gene promoter hypermethylation as a biomarker in oral cancer. *J Clin Med* 2021;10:1931. <https://doi.org/10.3390/jcm10091931>
39. Viet CT, Schmidt BL. Methylation array analysis of preoperative and postoperative saliva DNA in oral cancer patients. *Cancer Epidemiol Biomarkers Prev* 2008;17:3603-11. <https://doi.org/10.1158/1055-9965.EPI-08-0507>
40. Nakahara Y, Shintani S, Mihara M, Hino S, Hamakawa H. Detection of p16 promoter methylation in the serum of oral cancer patients. *Int J Oral Maxillofac Surg* 2006;35:362-5. <https://doi.org/10.1016/j.ijom.2005.08.005>
41. Viet CT, Jordan RC, Schmidt BL. DNA promoter hypermethylation in saliva for the early diagnosis of oral cancer. *J Calif Dent Assoc* 2007;35:844-9.
42. Liao PH, Chang YC, Huang MF, Tai KW, Chou MY. Mutation of p53 gene codon 63 in saliva as a molecular marker for oral squamous cell carcinomas. *Oral Oncol* 2000;36:272-6. [https://doi.org/10.1016/s1368-8375\(00\)00005-1](https://doi.org/10.1016/s1368-8375(00)00005-1)
43. Shpitzer T, Hamzany Y, Bahar G, Feinmesser R, Savulescu D, Borovoi I, et al. Salivary analysis of oral cancer biomarkers. *Br J Cancer* 2009;101:1194-8. <https://doi.org/10.1038/sj.bjc.6605290>
44. Zimmermann BG, Wong DT. Salivary mRNA targets for cancer diagnostics. *Oral Oncol* 2008;44:425-9. <https://doi.org/10.1016/j.oraloncology.2007.09.009>
45. Franzmann EJ, Reategui EP, Carraway KL, Hamilton KL, Weed DT, Goodwin WJ. Salivary soluble CD44: a potential molecular marker for head and neck cancer. *Cancer Epidemiol Biomarkers Prev* 2005;14:735-9. <https://doi.org/10.1158/1055-9965.EPI-04-0546>
46. TP53 tumor protein p53 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/7157>
47. Grünewald I, Vollbrecht C, Meinrath J, Meyer MF, Heukamp LC, Drebber U, et al. Targeted next generation sequencing of parotid gland cancer uncovers genetic heterogeneity. *Oncotarget* 2015;6:18224-37. <https://doi.org/10.18632/oncotarget.4015>
48. DAPK1 death associated protein kinase 1 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/1612>
49. Rettori MM, de Carvalho AC, Bomfim Longo AL, de Oliveira CZ, Kowalski LP, Carvalho AL, et al. Prognostic significance of TIMP3 hypermethylation in post-treatment salivary rinse from head and neck squamous cell carcinoma patients. *Carcinogenesis* 2013;34:20-7. <https://doi.org/10.1093/carcin/bgs311>
50. TIMP3 TIMP metalloproteinase inhibitor 3 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/7078>
51. Sun W, Zaboli D, Wang H, Liu Y, Arnaoutakis D, Khan T, et al. Detection of TIMP3 promoter hypermethylation in salivary rinse as an independent predictor of local recurrence-free survival in head and neck cancer. *Clin Cancer Res* 2012;18:1082-91. <https://doi.org/10.1158/1078-0432.CCR-11-2392>
52. Cristaldi M, Maureri R, Di Fede O, Giuliana G, Campisi G, Panzarella V. Salivary biomarkers for oral squamous cell carcinoma diagnosis and follow-up: current status and perspectives. *Front Physiol* 2019;10:1476. <https://doi.org/10.3389/fphys.2019.01476>
53. CDKN2A cyclin dependent kinase inhibitor 2A [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/1029>
54. MGMT O-6-methylguanine-DNA methyltransferase [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/4255>
55. CCND1 cyclin D1 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/595>
56. Ku BM, Jung HA, Sun JM, Ko YH, Jeong HS, Son YI, et al. High-throughput profiling identifies clinically actionable mutations in salivary duct carcinoma. *J Transl Med* 2014;12:299. <https://doi.org/10.1186/s12967-014-0299-6>
57. SERPINB5 serpin family B member 5 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/5268>
58. Chattopadhyay I, Panda M. Recent trends of saliva omics biomarkers for the diagnosis and treatment of oral cancer. *J Oral Biosci* 2019;61:84-94. <https://doi.org/10.1016/j.job.2019.03.002>
59. CXCL8 C-X-C motif chemokine ligand 8 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/3576>
60. Cheng J, Nonaka T, Wong DTW. Salivary exosomes as nanocarriers for cancer biomarker delivery. *Materials (Basel)* 2019;12:654. <https://doi.org/10.3390/ma12040654>
61. IL1B interleukin 1 beta [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/3553>
62. Lee YH, Kim JH, Zhou H, Kim BW, Wong DT. Salivary transcriptomic biomarkers for detection of ovarian cancer: for serous papillary adenocarcinoma. *J Mol Med (Berl)* 2012;90:427-34. <https://doi.org/10.1007/s00109-011-0829-0>
63. S100P S100 calcium binding protein P [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/6286>
64. Cheng YS, Jordan L, Rees T, Chen HS, Oxford L, Brinkmann O, et al. Levels of potential oral cancer salivary mRNA biomarkers in oral cancer patients in remission and oral lichen planus patients. *Clin Oral Investig* 2014;18:985-93. <https://doi.org/10.1007/s00784-013-1041-0> Erratum in: *Clin Oral Investig* 2014;18:995. <https://doi.org/10.1007/s00784-013-1127-8>

65. MIR31 microRNA 31 [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/407035>
66. Liu CJ, Lin SC, Yang CC, Cheng HW, Chang KW. Exploiting salivary miR-31 as a clinical biomarker of oral squamous cell carcinoma. *Head Neck* 2012;34:219-24. <https://doi.org/10.1002/hed.21713>
67. MIR125A microRNA 125a [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/406910>
68. MIR200A microRNA 200a [*Homo sapiens* (human)] [Internet]. Bethesda (MD): National Center for Biotechnology Information [cited 2021 Feb 21]. Available from: <https://www.ncbi.nlm.nih.gov/gene/406983>
69. India Project Team of the International Cancer Genome Consortium. Mutational landscape of gingivo-buccal oral squamous cell carcinoma reveals new recurrently-mutated genes and molecular subgroups. *Nat Commun* 2013;4:2873. <https://doi.org/10.1038/ncomms3873>
70. Shanmugam A, Hariharan AK, Hasina R, Nair JR, Katragadda S, Irusappan S, et al. Ultrasensitive detection of tumor-specific mutations in saliva of patients with oral cavity squamous cell carcinoma. *Cancer* 2021;127:1576-89. <https://doi.org/10.1002/cncr.33393>
71. Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-origin patterns dominate the molecular classification of 10,000 tumors from 33 types of cancer. *Cell* 2018;173:291-304.e6. <https://doi.org/10.1016/j.cell.2018.03.022>
72. Pickering CR, Zhang J, Yoo SY, Bengtsson L, Moorthy S, Neskey DM, et al. Integrative genomic characterization of oral squamous cell carcinoma identifies frequent somatic drivers. *Cancer Discov* 2013;3:770-81. <https://doi.org/10.1158/2159-8290.CD-12-0537>
73. Zammit AP, Sinha R, Cooper CL, Perry CFL, Frazer IH, Tuong ZK. Examining the contribution of smoking and HPV towards the etiology of oral cavity squamous cell carcinoma using high-throughput sequencing: a prospective observational study. *PLoS One* 2018;13:e0205406. <https://doi.org/10.1371/journal.pone.0205406>
74. Fadhil RS, Wei MQ, Nikolarakos D, Good D, Nair RG. Salivary microRNA miR-let-7a-5p and miR-3928 could be used as potential diagnostic bio-markers for head and neck squamous cell carcinoma. *PLoS One* 2020;15:e0221779. <https://doi.org/10.1371/journal.pone.0221779>
75. Nisha KJ, Janam P, Harshakumar K. Identification of a novel salivary biomarker miR-143-3p for periodontal diagnosis: a proof of concept study. *J Periodontol* 2019;90:1149-59. <https://doi.org/10.1002/JPER.18-0729>
76. Lu Z, He Q, Liang J, Li W, Su Q, Chen Z, et al. miR-31-5p is a potential circulating biomarker and therapeutic target for oral cancer. *Mol Ther Nucleic Acids* 2019;16:471-80. <https://doi.org/10.1016/j.omtn.2019.03.012>
77. Wang Y, Zeng G, Jiang Y. The emerging roles of miR-125b in cancers. *Cancer Manag Res* 2020;12:1079-88. <https://doi.org/10.2147/CMAR.S232388>
78. Korpala M, Kang Y. The emerging role of miR-200 family of microRNAs in epithelial-mesenchymal transition and cancer metastasis. *RNA Biol* 2008;5:115-9. <https://doi.org/10.4161/rna.5.3.6558>
79. Kabzinski J, Maczynska M, Majsterek I. MicroRNA as a novel biomarker in the diagnosis of head and neck cancer. *Biomolecules* 2021;11:844. <https://doi.org/10.3390/biom11060844>
80. Majewska H, Gorczyński A, Czapiewski P, Menon R, Mueller J, Lakis S, et al. ALK alterations in salivary gland carcinomas. *Virchows Arch* 2021;478:933-41. <https://doi.org/10.1007/s00428-020-02971-w>
81. Kim S, Lee JW, Park YS. The application of next-generation sequencing to define factors related to oral cancer and discover novel biomarkers. *Life (Basel)* 2020;10:228. <https://doi.org/10.3390/life10100228>
82. Sembler-Møller ML, Belstrøm D, Loch H, Enevold C, Pedersen AML. Next-generation sequencing of whole saliva from patients with primary Sjögren's syndrome and non-Sjögren's sicca reveals comparable salivary microbiota. *J Oral Microbiol* 2019;11:1660566. <https://doi.org/10.1080/20002297.2019.1660566>

**How to cite this article:** Sodnom-Ish B, Eo MY, Myoung H, Lee JH, Kim SM. Next generation sequencing-based salivary biomarkers in oral squamous cell carcinoma. *J Korean Assoc Oral Maxillofac Surg* 2022;48:3-12. <https://doi.org/10.5125/jkaoms.2022.48.1.3>