



Original Article

Impact of the COVID-19 pandemic on oral healthcare among children and teachers in kindergarten and daycare centers

Myoung-Hee Kim¹ · Eun-Joo Hong² · Yu-Jin Kwon¹ · Young-Sun Hwang¹

¹Department of Dental Hygiene, College of Health Science, Eulji University

²Department of Early Childhood Education, Eulji University

Corresponding Author: Young-Sun Hwang, Department of Dental Hygiene, College of Health Science, Eulji University, 553 Sanseong-daero, Sejeong-gu, Seongnam-si, 13135, Korea. Tel: +82-31-740-7493, Fax: +82-31-740-7352, E-mail: kiteys@eulji.ac.kr

ABSTRACT

Objectives: The COVID-19 outbreak has resulted in changes in the instructional methods used in kindergartens and daycare centers to prevent viral infections. This study aimed to investigate the changes in oral health care for children before and during COVID-19 and the perceptions of kindergarten and daycare center teachers about oral health care. **Methods:** The study subjects were 189 teachers of kindergartens and daycare centers in charge of children aged 3 to 5. The data for the analysis were collected through an online survey. Frequency analysis and the chi-squared test were used for statistical analysis. **Results:** Compared to before COVID-19, the frequency of tooth brushing after lunch, the number of oral health education and regular oral examinations, and the rate of childcare teachers' experiences with oral health education had decreased during COVID-19. In addition, educators in kindergarten and daycare centers responded that an oral health officer at a public health center was the most desirable oral health educator. **Conclusions:** During the COVID-19 pandemic, the oral health care of children in kindergartens and daycare centers has declined. Efforts are needed to restore it by educating people about oral hygiene care and oral health education, both at home and in kindergartens and daycare centers.

Key Words: COVID-19, Kindergarten and daycare center, Oral healthcare, Teacher

Introduction

Deciduous teeth are important for chewing and pronunciation in children. They maintain the position for eruption of permanent teeth and help normal development of the jawbone. Early childhood caries (ECC) occurring in children under 5 years of age frequently occurs in maxillary primary incisors and primary first molars. Dental caries also increases according to the order of tooth eruption [1]. Therefore, even for deciduous teeth may need to be filled to reduce the risk of loss. Childhood caries can also impede the development of permanent teeth. Early loss of primary teeth due to caries can cause malocclusion by adversely affecting eruption of permanent teeth and growth of the maxilla. ECC also causes severe pain and infection. When it is associated with eating and sleep disturbances, ECC can negatively affect children's ability to focus on learning [2]. Therefore, thorough oral health care is required to prevent dental caries from eruption of primary teeth until they are replaced with permanent teeth. According to the Statistical Information Report on the Children's Oral Health Survey announced by the Ministry of Health and Welfare in 2019, the proportion of 5-year-olds who experienced deciduous teeth caries was as high as 68.5% and the average caries experience index per person was 3.43. This means that most children experience dental caries before complete eruption of their primary teeth (<https://www.narastat.kr/metasvc/index.do?confmNo=117051&inputYear=2018>, Statistical Information Report for Users_Children's Oral Health Survey_2019.pdf). The preventive oral health care effect of dental caries is significantly higher in children than in adults because primary caries affects permanent dental caries [3].

Received September 07, 2022

Revised October 10, 2022

Accepted October 13, 2022

Copyright © 2022 by Journal of Korean Society of Dental Hygiene. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License(<http://creativecommons.org/licenses/by-nc/4.0>)

Changes in family structure and an increase of working women have increased the demand for childcare facilities and early childhood education institutions. As the time spent in childcare facilities increases, the awareness and behavior of educators about oral care for young children have a great impact on children's oral health. Therefore, the duty and responsibility to guide children's oral health are being emphasized not only by family members, but also by teachers of childcare facilities and early childhood education institutions. Teachers with correct knowledge of dental practice had a lower average number of cavities in their infants than teachers without knowledge [4]. In addition, the average number of dental caries in infants was lower when childcare facilities and teachers provided correct oral care for infants. As a result of surveying the perception of oral health among preschool teachers in Seoul, 99.1% of respondents recognized the importance of oral healthcare for children. They also recognized their responsibility for providing children's brushing instruction and oral examinations [5]. Since childcare teachers' awareness and performance of children's oral health closely affect children's oral health, the need to develop an oral health education program for early childhood teachers was suggested.

To prevent the spread of COVID-19 through respiratory droplets transmission, from Feb. 29, 2020 until recently, a state-led quarantine system such as observing personal hygiene and social distancing has been maintained for a long time. Response guidelines for preventing infectious diseases were distributed to public and multi-purpose facilities. The Office of Education also distributed additional guidelines for preventing infectious diseases for kindergartens and daycare centers. Accordingly, over the past three years, kindergartens and daycare centers have been operating programs centered on individual play rather than group play while observing social distancing. In addition, for personal quarantine, continuous wearing of a mask is maintained except for eating and drinking. Various phenomena that have recently changed due to group quarantine have been reported, such as current state of the response system of daycare centers during the COVID-19 [6], the perception of kindergarten teachers on online non-face-to-face classes [7], and the phenomenological study of behavior changes in children after wearing masks in kindergarten [7]. The present study investigated changes in oral health care practices for children in kindergartens and daycare centers due to the COVID-19, trying to derive points that cause problems for children's oral health. Through this study, we intend to present a basis for establishing children oral care policies of childcare institutions in the future.

Methods

1. Subjects and data collection

This study conducted a survey of childcare teacher working in kindergartens and daycare centers from December 13, 2021 to Jan. 2, 2022 for about 3 weeks. This is COVID-19 special quarantine period corresponding to the level 3 and level 4 social distancing guideline in Korea. The period before Mar. 2020 was set as 'before COVID-19'. Through online community platform of childcare teachers, URL links and QR codes were provided. The survey was designed to be answered only by the teachers in charge of 3 to 5 years of age. After being provided the purpose of the study, only subjects who voluntarily agreed to the questionnaire participated in the study by checking the online consent question. A total of 193 questionnaires were returned and 189 questionnaires with responses to all questions were used as validated data, excluding 4 insincere responses.

The sample size was calculated using G*power sampling software (ver. 3.1.9.7 for Windows). The minimum required sample size was estimated to be 158 persons with an α error of 5%, power of 90%, degree of freedom of 3 and medium effect size of 0.3 by applying the chi-squared test as the main analysis method. Additional samples were collected in each group to compensate for processing errors.

Questionnaire contents used in two previous studies were partially modified and reflected to suit this study [9,10]. The questionnaire consisted of 32-item including general characteristics, oral hygiene practice, and oral education experience of children before and during COVID-19, teachers' oral education necessity, and teachers' oral health knowledge quiz related to childcare, dental caries, and fluoride application. The present study was approved by the Institutional Review Board of Eulji University (approval no. EU21-86).

2. Statistical analysis

The collected data were analyzed using SPSS Statistics program (ver. 28.0.1.1; IBM Corp., Armonk, NY, USA). All statistics were expressed as frequencies and percentages for categorical variables, and the chi-squared test was used to determine the statistical significance of associations between the variables. A $p \leq 0.05$ was considered statistically significant.

Results

Responses of 189 kindergarten and daycare center educator were analyzed for this study. Female educators accounted for 95.2% of respondents. Regarding ages of teachers, there were 90 in their 20s (47.6%), 72 in their 30s (38.1%), 21 in their 40s (11.2%), and 6 in their 50s (3.2%). As for the positions, there were 179 (94.7%) teachers and 10 (5.3%) directors. Regarding the length of their education career, there were 105 (55.6%) educators with less than 5 years of education carrier, 61 (32.3%) with 6-10 years of education career, 8 (4.2%) with 11-20 years of education career, and 15 (7.9%) with more than 20 years of education career. Regarding the age of the children that they were in charge of, there were 71 (37.6%) children aged 3 years, 58 (30.7%) children aged 4 years, and 60 (31.7%) children aged 5 years <Table 1>.

Table 1. General characteristics of respondents

(N=189)

Characteristics	Division	N(%)
Gender	Male	9(4.8)
	Female	180(95.2)
Age group	20's	90(47.6)
	30's	72(38.1)
	40's	21(11.2)
	≥50's	6(3.2)
Child age class in charge	3 years old	71(37.6)
	4 years old	58(30.7)
	5 years old	60(31.7)
Education carrier period (yr)	≤5	105(55.6)
	6-10	61(32.3)
	11-20	8(4.2)
	≥20	15(7.9)

Changes in the frequency of tooth brushing before and during the COVID-19 were investigated <Table 2>. Before the COVID-19 pandemic, 79.9% of children brushed their teeth daily after lunch. However, during the COVID-19, only 28% brushed their teeth daily (51.9% decrease) with statistically significant decrease ($p < 0.001$). On the other hand, the rate of not brushing their teeth during COVID-19 increased by 48.2%. The reason for the difficulty of brushing their teeth among respondents of 'not perform' and '1 or 2 times a week' was lack of time before the COVID-19 pandemic [9(39.1%), n=23]. However, during the COVID-19 pandemic, it was mostly due to the government's guidelines for COVID-19 prevention [110(87.3%), n=126] <Table 3>. The frequency of additional brushing of teeth other than brushing teeth after lunch before and during COVID-19 was also investigated <Table 2>. Before the COVID-19, 43.9% said they brushed teeth additionally occasionally or daily. However, during COVID-19, this percentage decreased to 23.8%. Additional toothbrushing frequency during COVID-19 was also significantly reduced than that before COVID-19 because of the government's guidelines for COVID-19 prevention [109(67.3%)] <Table 3>. This result meant that oral hygiene care of infant and children in kindergarten and daycare centers was neglected during the quarantine period.

Table 2. The rate of tooth brushing after lunch and additional brushing before and during COVID-19 (N=189)

Characteristics	Division	Before COVID-19	During COVID-19	<i>p</i> *
Tooth brushing after lunch	Not perform	4(2.1)	95(50.3)	<0.001
	Sometimes (1-2 times/week)	19(10.1)	31(16.4)	
	Sometimes (3-4 times/week)	15(7.9)	10(5.3)	
	Everyday	151(79.9)	53(28.0)	
Additional brushing	Not perform	106(56.1)	144(76.2)	0.001
	Sometimes (1-2 times/week)	29(15.3)	18(9.5)	
	Sometimes (3-4 times/week)	9(4.8)	5(2.6)	
	Everyday	45(23.8)	22(11.6)	

All values are presented as N(%) for categorical variables.

*by chi-squared test and percentages are expressed in columns to be 100%

Table 3. Affecting factors limiting post-meal brushing and additional brushing before and during COVID-19

Characteristics	Division	Before COVID-19 (N=23)	During COVID-19 (N=126)
Tooth brushing after lunch	Due to time constraint	9(39.1)	2(1.6)
	Lack of facilities for brushing teeth	7(30.4)	6(4.8)
	Unnecessary	1(4.3)	1(0.8)
	Understaffed	6(26.1)	5(4.0)
	Government guidelines for the prevention of infectious diseases	-	110(87.3)
	Others	0(0.0)	2(1.6)
Characteristics	Division	Before COVID-19 (N=135)	During COVID-19 (N=162)
Additional brushing	Due to time constraint	71(52.6)	22(13.6)
	Lack of facilities for brushing teeth	16(11.9)	8(4.9)
	Unnecessary	30(22.2)	15(9.3)
	Understaffed	16(11.9)	6(3.7)
	Government guidelines for the prevention of infectious diseases	-	109(67.3)
	Unnecessary	2(1.5)	2(1.2)

All values are presented as N(%) for categorical variables. Percentages are expressed in columns to be 100%

Changes in the toothbrushing instruction method between before and during COVID-19 were also investigated <Table 4>. Before COVID-19, 60% of children brush their teeth and then got checked by their teacher. This percentage decreased to 50.8% during COVID-19, although the number of children who brush their teeth was increased by 10.6% during COVID-19. The frequency of oral health education for infants and children was 88.4% before COVID-19. It decreased to 75.1% during COVID-19 <Table 5>. The results were statistically significant ($p=0.001$). The frequency of oral examination was 68.3% before COVID-19. It decreased to 58.2% during COVID-19 with statistical significance ($p=0.043$). This result means that oral education and oral examination of infants and children have decreased since group meetings and face-to-face meetings are limited to prevent COVID-19 infection. The frequency of oral health education experience for childcare teachers also decreased by 8.5% due to COVID-19. On the other hand, the demand for oral health education of childcare teachers during COVID-19 was very high at 95.2% <Table 6>. Of 180 childcare teachers who requested oral health education, the highest 'desirable oral health educator' was an oral health officer in public health center (52.8%), followed by a dentist (27.8%) and a dental hygienist (16.1%) <Table 7>.

Table 4. Tooth brushing behavior before and during COVID-19 (N=189)

Characteristics	Before COVID-19	During COVID-19	<i>p</i> *
Children do it themselves	56(29.6)	76(40.2)	0.096
After children brushing teeth, teacher examines them	114(60.3)	96(50.8)	
Teacher brushed the child's teeth	19(10.1)	17(9.0)	

All values are presented as N(%) for categorical variables.

*by chi-squared test and percentages are expressed in columns to be 100%

Table 5. Oral health education and oral examination for children before and during COVID-19 (N=189)

Characteristics	Division	Before COVID-19	During COVID-19	<i>p</i> *
Oral health education	Perform	167(88.4)	142(75.1)	0.001
	Not perform	22(11.6)	47(24.9)	
Oral examination	Perform	129(68.3)	110(58.2)	0.043
	Not perform	60(31.7)	79(41.8)	

All values are presented as N(%) for categorical variables.

*by chi-squared test and percentages are expressed in columns to be 100%

Table 6. Childhood oral health education experience for childcare teachers before and during COVID-19 (N=189)

Characteristics	Division	Before COVID-19	During COVID-19	<i>p</i> *
Childhood oral health education experience	Yes	144(76.2)	128(67.7)	0.067
	No	45(23.8)	61(32.3)	
The need for oral health education	Yes	-	180(95.2)	
	No	-	9(4.8)	

All values are presented as N(%) for categorical variables.

*by chi-squared test and percentages are expressed in columns to be 100%

Table 7. Desirable oral health educator for childcare teachers (N=90)

Characteristics	Before COVID-19
Dentist	50(27.8)
Dental hygienist	29(16.1)
Oral health officer in public health center	5(5.2.8)
Others	6(3.3)

All values are presented as N(%) for categorical variables.

Deciduous teeth, dental caries, and fluoride application related issues were tested to determine oral health knowledge levels of childcare teachers. There were 8.2 correct answers to 12 questions, representing an average of 68% <Table 8>. The correct answer for dental caries-related question was an average of 76% and the correct answer for fluoride application-related question was an average of 77%. On the other hand, correct answer for deciduous teeth related question had an average of 55%. Among items related to deciduous teeth, the correct answer to the effect of thumb sucking on deciduous teeth was high (79.4%), whereas correct answers to the timing and maintenance of eruption of deciduous teeth (46.6%) and the number of deciduous teeth (25.9%) were the lowest.

Table 8. Oral health knowledge of childcare teachers for children. (N=189)

Category	Questions	Correct answer rate (%)	Total rate
Deciduous teeth	The number of deciduous teeth is 18.	25.9	53.8
	Permanent teeth begin to appear at the age of six.	63.5	
	If the deciduous teeth fall out earlier than the normal time, it doesn't matter if you leave them.	46.6	
Dental caries	The habit of thumb sucking affects permanent teeth.	79.4	77.4
	Chewing xylitol gum eliminates the need to brush your teeth.	89.9	
	Toothpaste is more important than brushing teeth.	77.2	
	Sealants are effective in preventing tooth decay.	65.6	
	Dental caries in the deciduous teeth do not require treatment.	83.6	
	Uneven dentition more easily makes dental caries.	60.8	
	There is no need to treat dental caries if it doesn't hurt.	87.3	
Fluoride application	Fluoride application is more effective in infants than in adults.	63.0	70.6
	Fluoride application is a way to whitens teeth.	78.3	
Total average		68.3	

All values are presented as N(%) for categorical variables.

Discussion

The main goal of this study was to examine the changes in oral health care for children before and during COVID-19 in kindergartens and daycare centers. We found that the oral health care of children, including tooth brushing, oral examinations, and oral health education, in kindergartens and daycare centers has significantly declined during COVID-19.

According to the Korean Children's Oral Health Survey conducted every three years according to the Oral Health Act, the rate of experiencing dental caries among 5-year-olds is as high as 68.5%, suggesting that most children experience tooth decay before their primary teeth erupt completely (<https://www.narastat.kr/metascv/index.do?confmNo=117051&inputYear=2018>, Statistical Information Report for Users_Children's Oral Health Survey_2019.pdf). As social distancing to block the spread of COVID-19 and the obligation to wear a personal mask are maintained for a long time, factors that can harm oral health might have accumulated due to limited oral hygiene management in kindergartens and daycare centers that educate young children. Therefore, in this study, we investigated the status of oral health care for children before and during COVID-19 through a questionnaire targeting kindergarten and daycare center teachers and identified factors that could hinder promotion and maintenance of children's oral health.

Early childhood education institutions provide snacks and drinks in addition to lunch [5]. Fruit is served most frequently, followed by bread, vegetables and cereals. Milk is the most frequently served drink, followed by yogurt and fruit drinks. These are highly carbohydrate-containing products that are highly associated with dental caries. In this study, changes in brushing performance after lunch as well as after snack were investigated before and during COVID-19. As a result of the survey, the percentage of kindergartens and daycare centers that did not ask children to brush their teeth after eating increased from 2.1% to 50.3% during the COVID-19 pandemic. The percentage of institutions that do not ask children to perform additional brushing other than after meals also increased from 56.1% to 76.2% during the COVID-19 pandemic <Table 2>. Tooth brushing practice was dramatically reduced to prevention of droplet infections. Even in a report surveyed in 2020, the early days of the COVID-19 issue, the proportion of kindergartens and daycare centers that did not brush their teeth after meals increased significantly during the COVID-19 period, and the proportion of institutions that did not brush their teeth other than brushing after lunch also increased during the COVID-19 period [5]. In addition, during COVID-19, teacher's inspection of children's oral condition after teeth brushing decreased from 60% to 50.8% <Table 4>. This means that oral hygiene management for children who have to stay for a long time in the institution is

becoming poorer. The Korean Children's Oral Health Survey will be conducted in the next cycle. It will produce results reflecting the impact of COVID-19. It is predicted that oral hygiene practices that are continuously restricted under the influence of COVID-19 regulations will be harmful to the oral health of infants and children. Regular oral health education and oral examinations for infants and children have also been reduced by about 10-12% since COVID-19 <Table 5>. Due to social distancing caused by the COVID-19 epidemic, oral care for children at kindergarten and daycare centers was insufficient. Children who received oral examinations had more dental clinic visits. However, the cost of dental treatment was higher for children who did not receive oral examinations [11]. A decrease in the number of oral hygiene and oral examinations can cause not only harm to the oral health of growing children, but also can increase oral treatment costs in the future. Thus, prompt improvement is required. In a study comparing the oral hygiene behavior of pre-schoolers in Wuhan city during the sudden outbreak of COVID-19 with pre-schoolers residing in non-Wuhan, Wuhan children brushed their teeth more actively [13]. However, even if infectious diseases are adequately defended in the future, the intention of direct dental visits of Wuhan residents was lower than that of non-residents. These results suggest that the experience of being affected by a severe infectious disease may have long-term implications for future dental visits. Therefore, considering the long-term experience of infectious diseases, it is also necessary to provide the importance of oral healthcare through continuous counseling so that dental care is not restricted due to concerns about future infection.

Oral health education experience of childcare teachers has a significant impact on oral health education and practice for children. Thus, regular oral health education was necessary [10]. Childcare teachers in kindergarten and daycare centers recognize the importance of deciduous teeth and the necessity of brushing their teeth. They also recognize the need for oral education and oral examination. They acquire oral health knowledge mainly through online information, TV, and dentistry [14]. According to our survey results, the experience of oral health education of childcare teachers during COVID-19 decreased from 76.2% to 67.7%, although the demand for oral health education was very high at 95.2% <Table 6>.

In order to find out oral health knowledge of childcare teachers, a total of 12 questions related to deciduous teeth, dental caries, and fluoride application were used. The percentage of correct answers was 8.2 (68.3%) out of 12. Percentages of correct answers to questions related to deciduous teeth, dental caries, and fluoride application were 53.8, 77.4%, and 70.6%, respectively. Knowledge related to deciduous teeth was the least. In particular, there was a lack of knowledge about the number of deciduous teeth (percentage of correct answer: 25.9%) and problem of early loss of deciduous teeth (percentage of correct answer: 46.6%). Also, knowledge about the relationship between malocclusion and dental caries (percentage of correct answer: 65.6%), the relationship between dental caries and sealants (percentage of correct answer: 60.8%), and the effect of fluoride application to infants (percentage of correct answer: 63.0%) were relatively low <Table 8>. Therefore, oral health education for childcare teachers is required to include information related to them. In particular, the awareness level of the dental caries prevention effect of childcare teachers was significantly higher among those with more than 10 years of educational experience [15]. According to the 2021 basic statistics on education (<https://www.moe.go.kr/boardCnts/view.do?boardID=294&boardSeq=88238&dev=0&searchType=null&statusYN=W&page=1&s=moe&m=020402&opType=N>), the average age of kindergarten teachers registered with the Ministry of Education is 34.8 years old. Since the average age of daycare teachers managed by the Ministry of Health and Welfare is not reflected, the average age of daycare center and kindergarten teachers may be different. The age of daycare center and kindergarten teachers who participated in this study was 47.6% in their twenties. In addition, 55.6% of the participants had less than 5 years of educational experience. Therefore, considering age and education carrier period, it is necessary to provide an education program for improving oral health knowledge to prevent dental caries in children for a significant proportion of childcare teachers. In addition, in order to recover the lack of oral hygiene management for child that has persisted for the past 3 years, oral hygiene behaviors for children should be implemented through active oral health education for childcare teachers.

According to a previous report, professional personnel of dental hospitals are suitable for infant oral health education [14]. On the other hand, oral health officer in public health center (52.8%) was found to be the most suitable for oral health education for

childcare teachers, followed by dentistry (27.8%) and dental hygienist (16.1%) in this study <Table 7>. Before COVID-19, regular oral examinations at kindergartens and daycare centers were conducted mainly by professional personnel from dental clinics. The prolonged social distancing has caused a change in the perception of oral health educators. In particular, the suitability of dental hygienists as oral health educators as well as dentists was relatively low. Efforts are needed to improve the awareness of dental hygienists as oral health experts. Considering our results and the current public health care system in Korea, oral education by dental hygienists should be enhanced in the way of community oral health programs through cooperation with local dental hospitals. Active oral health activities through participation in education of dental hygienists for childcare teachers and community service are required.

Some limitations of our study should be noted. We did not consider area variables that may affect similarities and differences in oral health behaviors including affiliated institutions in the COVID-19 pandemic. In addition, this study did not overcome the issue of analytical biases by asking questions about the before COVID-19 situation, that is, recall bias.

However, this study has several strengths: (1) this is timely study providing the changes in oral health care for children before and during COVID-19 and (2) our study provided reliable information and results due to the Intensive survey period with same level of COVID-19 quarantine which affected people's behavioral patterns according to government's quarantine guidelines and social distancing levels

In this study, we confirmed that oral hygiene management for children was insufficient due to COVID-19. Efforts to recover it are urgently needed. Efforts should be made for oral hygiene management of children through close communication between childcare institutions and caregivers. In addition, continuous activities are needed to raise awareness of dental hygienists as suitable oral health care educators.

Conclusions

To investigate the changes in oral health care for children before and during COVID-19 and the perception of oral health care, a total of 189 teachers of kindergartens and daycare centers were analyzed.

1. Oral health care of children, including tooth brushing, oral examinations, and oral health education, has significantly declined during COVID-19.
2. Educators in kindergarten and daycare center responded that an oral health officer at a public health center was the most desirable oral health educator.

The oral health care behavior among children and teacher in kindergarten and day care center was significantly influenced by the COVID-19 epidemic, which might give rise to a long-term impact on oral healthcare in the future. The findings suggested that more attention to children as well as teacher among kindergarten and daycare center might be needed in order to maintain oral health.

Conflicts of Interest

The authors declared no conflicts of interest.

Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2022R1F1A1063204).

Authorship

Conceptualization: MH Kim, YS Hwang; Data collection: YJ Kwon; Formal analysis: MH Kim; Writing-original draft: YS Hwang; Writing-review&editing: MH Kim, HW Jeong, YS Hwang, EJ Hong

References

1. Fejerskov O, Nyvad B, Kidd E. Dental caries: the disease and its clinical management. 2nd ed. Victoria: Blackwell; 2009: 20-3.
2. Sheiham A. Dental caries affects body weight, growth and quality of life in preschool children. *Br Dent J* 2006;201(10):625-6. <https://doi.org/10.1038/sj.bdj.4814259>
3. Hong MH. Risk factors affecting dental caries in children. *JKAIS* 2020;21(5):320-6. <https://doi.org/10.5762/KAIS.2020.21.5.320>
4. Kim EH, Choi HS, Oh HS. The influence of the cognizance of childcare teachers about dental care on oral health of infants. *J Kor Dent Hyg Sci* 2006;8(1):17-28.
5. Lee SM, Lim MH, Jung JY, Hwang YS, Kim SH. Recognition and performance of preschool teachers on children's oral health in Seoul. *J Korean Soc Dent Hyg* 2010;10(3):449-57.
6. Korea Institute of Child Care and Education. Inspection and analysis of the response system for COVID-19 kindergartens and daycare centers [Internet]. KICCE[cited 2022 Aug 25]. Available from: <https://repo.kicce.re.kr/handle/2019.oak/5045>.
7. Lim EM. Early childhood teachers' perception and needs on distance education after COVID-19. *KALCI* 2021;21(24):631-45.
8. Kim YN. A phenomenological study on young children's face mask wearing in a kindergarten during the COVID-19 pandemic. *Early Childhood Education Research* 2021;41(6):107-31. <https://doi.org/10.18023/kjece.2021.41.6.005>
9. Jung YM. Oral care services for younger children by nursery school teachers in Daegu[Master's thesis]. Gyeongsan: Yeungnam University, 2012.
10. Han JY. A study on young child education practitioners' knowledge of oral health and its practice[Master's thesis]. Cheonan: Dankook University, 2008.
11. An ES, Shin HS. Effectiveness of oral examination for infants and toddlers: effects on subsequent utilization and costs. *J Korean Acad Oral Health* 2017;41(2):73-9. <https://doi.org/10.11149/jkaoh.2017.41.2.73>
12. Park IS. Oral health awareness and management of day care teachers in accordance with oral health education experience. *J Digit Converge* 2016;14(9):407-15. <https://doi.org/10.14400/JDC.2016.14.9.407>
13. Liu C, Zhang S, Zhang C, Tai B, Jiang H, Du M. The impact of coronavirus lockdown on oral healthcare and its associated issues of preschoolers in China: an online cross-sectional survey. *BMC Oral Health* 2021;21(54):1-6. <https://doi.org/10.1186/s12903-021-01410-9>
14. Lee HN, Shim HS. A study on preschool teachers' dental health recognition and behaviors about preschoolchildren dental health care. *J Dent Hyg Sci* 2008;8(4):283-9.
15. Yoon HS. Oral health knowledge and awareness of the prevention effects against dental caries among childcare teachers. *JKAIS* 2014;15(9):5668-74. <https://doi.org/10.5762/KAIS.2014.15.9.5668>

유치원과 어린이집 교사의 COVID-19 구강건강관리의 영향

초록

연구목적: COVID-19로 인해 유치원과 어린이집에서의 교육 방식이 변화되고 있다. 본 연구의 목적은 COVID-19 이전과 COVID-19 시기의 유아 구강건강관리 실태와 유치원 및 어린이집 교사들의 구강건강관리에 대한 인식을 조사하는 것이다. **연구방법:** 연구 대상은 만 3-5세 아동을 담당하는 유치원 및 어린이집 교사 189명으로 온라인 설문조사를 통해 자료를 수집하였다. 통계분석은 빈도분석과 카이제곱 검정을 이용하였다. **연구결과:** COVID-19 이전보다 COVID-19 이후 아동의 점심식사 후 칫솔 사용 횟수, 구강보건교육 및 정기 구강검진 횟수, 보육교사의 구강보건교육 경험률이 감소했다. 또한 유치원 및 어린이집 교육자들은 보건소 구강보건담당자가 가장 바람직한 구강보건교육자라고 응답하였다.

결론: COVID-19 팬데믹 이후 유치원과 어린이집에서 영유아의 구강건강관리가 취약해졌다. 구강위생과 구강건강교육을 통해 이를 회복시키려는 노력이 필요하다. 또한 구강건강교육자로서 치과위생사의 역할을 확대할 필요가 있다.

색인: COVID-19, 유치원 및 어린이집, 구강보건, 교사