

Morphology of First Zoea Stage of *Sphaerozius nitidus* (Decapoda: Eriphioidea: Menippidae) Reared in the Laboratory Material

Seok Hyun Lee*

Department of Life Science, Silla University, Busan 46958, Korea

ABSTRACT

Ovigerous crab of *Sphaerozius nitidus* of the family Menippidae was collected from Geojedo, Gyeongsangnamdo and hatched in the laboratory. Digital imaging of live zoeas of the first zoea stage of *S. nitidus* has been reported for the first time in the world, and its morphology has been redescribed and illustrated. This study is different from former study in that it has three unequal setae with exopod of antenna, 5 + 4 setae with basal endite of maxilla, and dorsomedial spine and lateral minute spine with fork of telson. The first zoea of *S. nitidus* has black pigments occurring behind the eyes, on the dorsal spine medially and on the basis of lateral carapace spines, on basal of basis of maxillipeds 1, 2, and posterior margins of somites 1–5, and yellowish green chromatophores on the dorsal spine.

Keywords: *Sphaerozius nitidus*, first zoea, Menippidae, Eriphioidea, Korea

INTRODUCTION

The order of Decapoda includes more than 15,000 species of shrimps, hermit crabs, lobsters, and crabs that belong to four of the 10 infraorders of malacostracan crustaceans (Cumberland et al., 2015). The larval stage of decapods consists of the nauplius, prezoa, zoea, and megalopa stages. However, most decapods larvae are hatched in the zoea stage because the nauplius stage is passed in the egg except for Dendrobranchiata, and prezoa stage is just hatched lava still covered by embryonic cuticle (Moore and McCormick, 1969), so, the zoea stage can be regarded as the first stage in the larval stage of decapods. The larval morphology of decapod is more useful in phylogenetic studies than adult morphology (Rice, 1980; Kornienko and Korn, 2009). The adult morphology of decapods has differences between individual variations because of convergent and divergent adaptations of the various life styles. However, larvae of decapods inhabit a planktonic environment with relatively uniform characteristics (Rice, 1980). And, larval morphology of decapods is more accordant with molecular data than with adult morphology (Hultgren et al., 2009). However, the number of studies of decapod larvae is inadequate and still unknown.

Crabs of the superfamily Eriphioidea comprise 68 species

21 genera six families (Ng et al., 2008; WoRMS, 2022) (Table 1). They are generally known as stone or rubble crabs and occupy a diverse range of habitats from intertidal rocky shores, mangrove swamps, coral reefs, to continental slopes about 800 meter in depth (Lai et al., 2013). In Korea, three species [*Eriphia smithii* MacLeay, 1838, *Hypothalassia armata* (De Haan, 1835), *Sphaerozius nitidus* Stimpson, 1858], in three families (Eriphiidae Macleay, 1838, Hypothalassidae Karasawa and Schweitzer, 2006, Menippidae Ortmann, 1893) are recorded. In the Eriphioidea, larval descriptions have been known for 20 species belonging to nine genera of four families in the world (Table 1), among them two species (*E. smithii* by Terada, 1982, and *S. nitidus* by Aikawa, 1933) have been reported from Korea and adjacent waters. However, the description and illustration of *S. nitidus* by Aikawa (1933) is not accurate in setal amature of appendages. So, the redescription of the first zoea stage of *S. nitidus* (Figs. 1B, 2) is needed.

In this study, the first zoea stage of *S. nitidus* is described and illustrated for the second time, and provided color digital image for the first time in the world. Its morphological characteristics are compared with those of other known belonging to the same superfamily of species in the Korea.

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

***To whom correspondence should be addressed**

Tel: 82-51-999-6931, Fax: 82-51-999-5471
E-mail: sky3284@silla.ac.kr

Table 1. List of species of the superfamily Eriphioidea and its known larval stages

Family	Genus	Species	Larval stages and Sources
Dairoiidae Štević, 2005	<i>Dairoides</i> Stebbing, 1920	<i>kusei</i> (T. Sakai, 1938) <i>margaritatus</i> Stebbing, 1920 <i>seafdeci</i> Takeda and Ananpongsuk, 1991 <i>ferox</i> Koh and PKL Ng, 2008 <i>gonagra</i> (JC Fabricius, 1781) <i>granulosa</i> A. Milne-Edwards, 1880 <i>scabricula</i> Dana, 1852 <i>sebana</i> (Shaw and Nodder, 1803) <i>smithii</i> MacLeay, 1838 <i>squamata</i> Stimpson, 1859 <i>verrucosa</i> (Forskål, 1775) <i>hispidula</i> (Stimpson, 1860) <i>palmeri</i> (Garth, 1986) <i>acerba</i> Koh and PKL Ng, 2000 <i>armata</i> (De Haan, 1835) <i>adina</i> AB Williams and Felder, 1986 <i>frontalis</i> A. Milne-Edwards, 1879 <i>hirtipes</i> (Jacquinot and Lucas, 1853) <i>mercenaria</i> (Say, 1818)	Z1-4, M: Fransozo (1987) Z1: Clark and Paula (2003) Z1: Ko (2005) Z1-4: Terada (1982) Z1-4, M: Lumare and Gozzo (1972)
Eriphiidae MacLeay, 1838	<i>Eriphia</i> Latreille, 1817	<i>Eriphides</i> Rathbun, 1897 <i>Garthopilumnus</i> Štević, 2011 <i>Hypothalassia</i> Gistel, 1848	
Hypothalassidae Karasawa and Schweitzer, 2006		<i>Menippe</i> De Haan, 1833	M: Martin et al. (1988)
Menippidae Ortman, 1893		<i>Myomenippe</i> Hilgendorf, 1879 <i>Ruppellioides</i> A. Milne-Edwards, 1867 <i>Sphaerozius</i> Stimpson, 1858	Z1: Hyman (1925) Z1-5, M: Kurata (1970) Z1-5, M: Scotto (1979) Z1-5, M: Kakati (1977)
Oziidae Dana, 1851		<i>Baptozius</i> Alcock, 1898 <i>Bountiana</i> Davie and PKL. Ng, 2000 <i>Epixanthoides</i> Balss, 1935 <i>Epixanthus</i> Heller, 1860	Z1: Aikawa (1933) Z1-4, M: Saba et al. (1978a) Z1, 2, M: Saba et al. (1978b) Z1: Clark and Paula (2003) Z1: Clark and Paula (2003) Z1-4, M: Aidaroos et al. (2014)

Continued

Table 1. Continued

Family	Genus	Species	Larval stages and Sources	
Oziidae Dana, 1851	<i>Eupilumnus</i> Kossmann, 1877	<i>hellerii</i> A. Milne-Edwards, 1867		
		<i>subcorrosus</i> de Man, 1891		
		<i>tenuidactylos</i> (Lockington, 1877)		
		<i>actumnoides</i> (A. Milne-Edwards, 1873)		
		<i>africanus</i> (A. Milne-Edwards, 1867)		
		<i>calmani</i> (Balss, 1933)		
		<i>ragaria</i> (SL Yang, Dai and PKL Ng, 1998)		
		<i>globosus</i> (Dana, 1852)		
		<i>kiensis</i> (Takeda and Nagai, 1983)		
		<i>laciniatus</i> (T. Sakai, 1980)		
		<i>stridulans</i> (Monod, 1956)		
		<i>xantusii</i> (Stimpson, 1860)		
		<i>annulipes</i> (H. Milne Edwards, 1834)		
		<i>granulosa</i> (A. Milne-Edwards, 1867)		
		<i>tenax</i> (Rüppell, 1830)		
		<i>truncata</i> (Strahl, 1862)		
		<i>deplanatus</i> (White, 1847)		
		<i>granulosus</i> de Man, 1879		
		<i>guttatus</i> H. Milne Edwards, 1834		
	<i>hawaiiensis</i> Rathbun, 1902			
	<i>lobatus</i> Heller, 1865			
	<i>perilatus</i> Stimpson, 1860			
	<i>reticulatus</i> (Desbonne and Schramm, 1867)			
<i>rugulosus</i> Stimpson, 1858				
<i>tricarinatus</i> Rathbun, 1907				
<i>truncatus</i> H. Milne Edwards, 1834				
<i>tuberculatus</i> H. Milne Edwards, 1834				
<i>verreauxii</i> de Saussure, 1853				
<i>crenulatus</i> (A. Milne-Edwards, 1879)				
			Z1-4, M: Kakati and Nayak (1977)	
			Z1-4, M: Wear (1968)	
			Z1-4, M: Dittel and Epifanio (1984)	
			Z1-4, M: Menu-Marque (1970)	
			Z1-4, M: Iorio and Boschi (1986)	
			Z1-4, M: Fagetti (1970)	
Platyxanthidae Guinot, 1977	<i>Danielethus</i> BP Thoma, PKL Ng and Felder, 2012			
	<i>Homalaspis</i> A. Milne-Edwards, 1863			
	<i>Otmroxanthus</i> Štević, 2011			
	<i>Peloeus</i> Eydoux and Souleyet, 1842			
	<i>Platyxanthus</i> A. Milne-Edwards, 1863			
				Z1: Clark and Paula (2003)

Z, zoea stage; M, megalopa stage.

MATERIALS AND METHODS

Ovigerous crab of *Sphaerozius nitidus* was collected from intertidal region in Geojedo, Gyeongsangnamdo, Korea on February 19, 2007 and its zoea was hatched in the laboratory on July 3, 2008. The first zoeas were preserved in 95% ethyl alcohol for examination. Digital photos with Leica EZ40 microscope (Wetzlar, Germany) and then processed in Photoshop were taken of living zoeas. Specimens of zoea were dissected using Leitz zoom stereomicroscope and appendages were examined under a Leitz Laborlux S microscope (Wetzlar, Germany). Appendages were mounted in ethylene glycol and drawings were made with the camera lucida. Setal counts on appendages and measurements were based on the mean of 10 specimens. The sequence of the description of zoea follows that of Clark et al., 1998. The long plumose natatory setae of the first and second maxillipeds were drawn truncated. The chromatophore patterns were observed with living zoeas. A micrometer was used for measurements of zoea: CL (carapace length) from the base of the rostral spine to the most posterior carapace margin and RDL (rostral and dorsal spine length) from the tip of rostral carapace spine to the tip of the dorsal carapace spine. Specimens examined and spent female were deposited at Silla University, Korea.

RESULTS

Sphaerozius nitidus Stimpson, 1858 (Fig. 1A)

First zoea (Figs. 1B, 2)

Size: CL 0.45 ± 0.05 mm; RDL 1.45 ± 0.05 mm.

Chromatophores (Fig. 1B): Black pigments which occurring behind and between eyes, on dorsal carapace spine medially, on basis of lateral carapace spines dorsally, on basal of basis

of maxillipeds 1, 2 and on posterior margins of somites 1–5; Yellowish green chromatophores on dorsal spine.

Carapace (Figs. 1B, 2A): Dorsal spine longer than CL, approximately same length as rostral spine, almost straight; rostral spine approximately 3 times longer than antennal protopod; lateral spines present, slightly shorter than rostral spine; all spines with blunt tip; 1 pair of posterodorsal setae present; each ventral margin unarmed; eyes sessile.

Antennule (Fig. 2B): Uniramous; endopod absent; exopod with 2 long, stout aesthetascs, 1 shorter, thinner aesthetasc, 2 simple (1 long, 1 shorter) setae, all terminal.

Antenna (Fig. 2C): Biramous; endopod bud absent; protopod 1/3 length of rostral spine, with 2 rows of spinules on distal half; exopod approximately 2/3 length of protopod, with 3 unequal (2 terminal, 1 subterminal) setae.

Mandibles (Fig. 2D): Asymmetrical; right molar with 3 teeth, left molar with 1 tooth, confluent with incisor process; palp absent.

Maxillule (Fig. 2E): Coxal endite with 7 setae; basial endite with 5 setae and 2 teeth; endopod 2-segmented, proximal segment with 1 seta, distal segment with 4 setae; exopod seta and epipod absent.

Maxilla (Fig. 2F): Coxal endite bilobed, with 5 + 4 setae; basial endite bilobed, with 5 + 4 setae; endopod bilobed, with 3 + 3 setae; exopod (scaphognathite) margin with 4 plumose setae, 1 distal process

First maxilliped (Fig. 2G): Basis with 10 setae, arranged 2 + 2 + 3 + 3; endopod 5-segmented, with 3, 2, 1, 2, 5 (1 subterminal, 4 terminal) setae; exopod 2-segmented, distal segment with 4 long terminal plumose natatory setae.

Second maxilliped (Fig. 2H): Basis with 4 setae arranged 1 + 1 + 1 + 1; endopod 3-segmented, with 0, 1, 5 setae; exopod 2-segmented, distal segment with 4 long terminal plumose natatory setae.




Fig. 1. Color photos of adult and the first zoeas of *Sphaerozius nitidus*. A, Adult of *S. nitidus*; B, First zoeas of *S. nitidus*.




Fig. 2. *Sphaerozius nitidus*, first zoea stage. A, Whole animal, lateral view; B, Antennule; C, Antenna, dorsal view; D, Mandibles, dorsal view; E, Maxillule, dorsal view; F, Maxilla, dorsal view; G, First maxilliped, lateral view; H, Second maxilliped, lateral view; I, Abdomen and telson, dorsal view; I', Fork of telson, dorsal view. Scale bars: A, I=0.3 mm, B-D, I'=0.05 mm, E-H=0.1 mm.

Table 2. Comparison of the characteristics of the first zoeas of Eriphioidea from Korea

	Eriphiidae <i>Eriphia smithii</i> Terada, 1982	Menippidae <i>Sphaerozius nitidus</i>	
		Aikawa, 1933	Present study
Carapace			
Dorsal, rostral, lateral spines	Long, long, short	Long, long, long	Long, long, long
Antennule	2 aesthetascs, 4 setae	2 aesthetascs, 2 setae	3 aesthetascs, 2 setae
Antenna			
Protopod	Spinulate	Spinulate	Spinulate
Exopod	1/2 protopod length, spinulate	1/2 protopod length, not spinulate	2/3 protopod length, not spinulate
	4 unequal terminal spines	2 unequal terminal setae	3 unequal terminal setae
Maxillule			
Coxal endite	7 setae	7 setae	7 setae
Basial endite	5 setae	5 setae	5 setae
Endopod	1, 2+4 setae	1, 4 setae	1, 4 setae
Maxilla			
Coxal endite	5+4 setae	5+3 setae	5+3 setae
Basial endite	6+4 setae	6+4 setae	5+4 setae
Endopod	3+5 setae	3+3 setae	3+3 setae
1st maxilliped			
Basis	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae
Endopod	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 5 setae
2nd maxilliped			
Basis	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae
Endopod	1, 1, 6 setae	0, 1, 5 setae	0, 1, 5 setae
Abdomen			
Lateral processes	Somites 2-5	Somites 2, 3	Somites 2, 3
Telson			
Fork	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	1 dorsomedial spine	1 lateral minute spine, 1 dorsomedial spine
	Not spinulate	Not spinulate	Not spinulate

Abdomen (Fig. 2I): With 5 somites; somite 2 with 1 pair of lateral processes directed anteriorly; somite 3 with 1 pair of lateral processes directed posteriorly; somites 2-5 each with 1 pair of posterodorsal setae, respectively; pleopod absent.

Telson (Fig. 2I'): Each fork long, with 1 minute lateral spine, and 1 dorsomedial spine; posterior margin with 3 pairs of setae.

DISCUSSION

The characteristics of the first zoea of *Sphaerozius nitidus* is redescribed in this study. So, zoea descriptions of Eriphioidea are available for two species, *Eriphia smithii* (Eriphiidae) and *S. nitidus* (Menippidae) in Korea.

After careful comparison of the first zoea stage of *S. nitidus* described by Aikawa (1933) with the present material, the author found some differences in three characters (Table 2).

Aikawa (1933) described that an exopod of antenna has two unequal terminal setae (vs. three unequal setae in this study), a basal endite of the maxilla has 6+4 setae (vs. 5+4 setae) and fork of telson has one dorsomedial spine (vs. a lateral minute spine and dorsomedial spine). These characteristics are considered as in significant and the above differences probably resulted from their mistakes in detailed examination.

Comparison of zoea can be made for only two species of two families, *E. smithii* (see Terada, 1982) of Eriphiidae and *S. nitidus* (present study) of Menippidae (Table 2). They are significantly different from each other in characters of lateral carapace spine, endopods of the maxillule, the maxilla, and the second maxilliped, and the abdomen. The first zoea of *E. smithii* of Eriphiidae has short lateral carapace spine, endopods of the maxillule and the maxilla with 1, 2+4 and 3+5 setae, respectively, an endopod of the second maxilliped with 1, 1, 6 setae, and abdominal somites 2-5 each with a pair of

lateral processes. These characteristics are similar to those of other pilumnoid and xanthoid zoeas except for abdominal somites. While, the first zoea of *S. nitidus* of Menippidae has long lateral carapace spine, endopods of the maxillule and the maxilla with 1, 4 and 3 + 3 setae, respectively, an endopod of the second maxilliped with 0, 1, 5 setae, and abdominal somites 2 and 3 each with a pair of lateral processes. A combination of characteristics of 1, 4 and 3, 3 setae in the endopods of the maxillule and the maxilla is reported for the first time in brachyuran zoeas of Korea and considered a diagnostic one of this species. Thus, the author considers that the Eriphioidea might be heterogeneous because *S. nitidus* is not related to *E. smithii* in characters of lateral carapace spine (long lateral carapace spine in *S. nitidus* vs. short lateral carapace spine in *E. smithii*), endopods of the maxillule (1, 4 setae in *S. nitidus* vs. 1, 2 + 4 setae in *E. smithii*) and the maxilla (3 + 3 setae in *S. nitidus* vs. 3 + 5 setae in *E. smithii*), and the second maxilliped (0, 1, 5 setae in *S. nitidus* vs. 1, 1, 6 setae in *E. smithii*) (Table 2).

ORCID

Seok-Hyun Lee: <https://orcid.org/0000-0003-4372-4909>

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This research was supported by grant (2021M00100) from the National Marine Biodiversity Institute of Korea (MABIK).

REFERENCES

- Aikawa H, 1933. On larval forms of some Brachyura, paper II; a note on indeterminable zoeas. Records of Oceanographic Work in Japan, 5:124-254.
- Al-Aidaros AM, Al-Haj AE, Kumar AAJ, 2014. Complete larval development of the Brachyuran crab (*Epixanthus frontalis* H. Milne Edwards, 1834) (Crustacea, Decapoda, Eriphioidea, Oziidae) under laboratory conditions. Zootaxa, 3815:29-50. <https://doi.org/10.11646/zootaxa.3815.1.2>
- Clark PF, Calazans D, Pohle GW, 1998. Accuracy and standardization of brachyuran larval descriptions. Invertebrate Reproduction and Development, 33:127-144. <https://doi.org/10.1080/07924259.1998.9652627>
- Clark PF, Paula J, 2003. Descriptions of ten xanthoidean (Crustacea: Decapoda: Brachyura) first stage zoeas from Inha-ca Island, Mozambique. The Raffles Bulletin of Zoology, 51:323-378.
- Cumberlidge N, Hobbs HH, Lodge DM, 2015. Class Malacostraca, Order Decapoda. In: Ecology and general biology: Thorp and Covich's freshwater invertebrates (Eds., Thorp J, Rogers DC). Academic Press, San Diego, CA, pp. 797-847. <https://doi.org/10.1016/B978-0-12-385026-3.00032-2>
- De Haan W, 1833-1850. Crustacea. In: Fauna Japonica sive Descriptio Animalium, quae in Itinere per Japoniam, Jussu et Auspiciis Superiorum, qui Summum in India Batava Imperium, Tenent, Wuscepto, Annis 1823-1830 Collegit, Notis Observationibus et Adumbrationibus Illustravit (Ed., von Siebold PF). Lugduni-Batavorum, Leiden, pp. 1-243.
- Dittel A, Epifanio CE, 1984. Desarrollo larval de *Ozius verreauxi* (Saussure) Brachyura: Xanthidae en el laboratorio. Revista de Biología Tropical, 32:171-172 (in Spanish).
- Fagetti E, 1970. Desarrollo larval en el laboratorio de *Homalaspis plana* (Milne-Edwards) (Crustacea Brachyura, Xanthidae). Revista de Biología Marina, 14:29-49.
- Fransozo A, 1987. Desenvolvimento larval de *Eriphia gonagra* (Fabricius, 1781) (Decapoda, Xanthidae), em laboratório. Revista Brasileira de Zoologia, 4:165-179. <https://doi.org/10.1590/S0101-81751987000300001>
- Hultgren KM, Guerao G, Marques FPL, Palero FP, 2009. Assessing the contribution of molecular and larval morphological characters in a combined phylogenetic analysis of the superfamily Majoidea. In: Decapod crustacean phylogenetics, crustacean issues (Eds., Martin JW, Crandall KA, Felder DL). CRC Press, Boca Raton, FL, 18:437-455.
- Hyman OW, 1925. Studies on the larvae of crabs of the family Xanthidae. Proceedings of the United States National Museum, 67:1-22. <https://doi.org/10.5479/si.00963801.67-2575.1>
- Iorio MI, Boschi EE, 1986. Studies on the larval stages of the crab *Platyxanthus patagonicus*, from laboratory rearing and plankton samples. Journal of Aquaculture Tropical, 1:7-24.
- Kakati VS, 1977. Larval development of the crab, *Menippe rumphii* (Fabricius) as observed in the laboratory (Crustacea: Brachyura). In: Proceedings Symposium of Warm Water Planktonic Species. Publishing National Institute of Oceanography, Goa, pp. 634-641.
- Kakati VS, Nayak VN, 1977. Larval development of the xanthid crab, *Ozius rugulosus* Stimpson (Decapoda, Brachyura) under laboratory conditions. Indian Journal of Marine Sciences, 6:26-30.
- Karasawa H, Schweizer CE, 2006. A new classification of the Xanthoidea sensu lato (Crustacea: Decapoda: Brachyura) based on phylogenetic analysis and traditional systematic and evaluation of all fossil Xanthoidea sensu lato. Contributions to Zoology, 75:23-73. <https://doi.org/10.1163/18759866-0750102002>
- Ko HS, 2005. First zoea of *Eriphia sebana* (Crustacea: Decapo-

- da: Xanthoidea: Menippidae) hatched in the laboratory. *Integrative Biosciences*, 9:135-138. <https://doi.org/10.1080/17386357.2005.9647263>
- Kornienko ES, Korn OM, 2009. Illustrated key for the identification of brachyuran zoeal stages (Crustacea: Decapoda) in the plankton of the Peter Great Bay (Sea of Japan). *Journal of the Marine Biological Association of the United Kingdom*, 89:379-386. <https://doi.org/10.1017/S0025315408002762>
- Kurata H, 1970. Studies on the life histories of decapod Crustacea of Georgia. Part III: Larvae of decapod Crustacea of Georgia. PhD dissertation, The University of Georgia, Athens, GA, pp. 1-192.
- Lai JCY, Thoma BP, Clark PF, Felder DL, Ng PKL, 2013. Phylogeny of eriphoid crabs (Brachyura, Eriphioidea) inferred from molecular and morphological studies. *Zoologica Scripta*, 43:52-64. <https://doi.org/10.1111/zsc.12030>
- Lumare F, Gozzo S, 1972. Sviluppo larvale del crostaceo Xanthidae *Eriphia verrucosa* (Forsk., 1775) in condizioni di laboratorio. *Bolletino di Pesca Piscicoltura e Idrobiologia*, 27:185-209.
- MacLeay WS, 1838. On the brachyurous decapod Crustacea. Brought from the cape by Dr. Smith. In: *Illustrations of the Zoology of South Africa; consisting chiefly of figures and descriptions of the objects of natural history collected during an expedition into the interior of South Africa, in the years 1834, 1835, and 1836; fitted out by "The Cape of Good Hope Association for Exploring Central Africa"* (Ed., Smith A). Published under the Authority of the Lords Commissioners of Her Majesty's Treasury, London, pp. 53-71.
- Martin JW, Truesdale FM, Felder DL, 1988. The megalopa state of the Gulf stone crab, *Menippe adina* Williams and Felder, 1986, with a comparison of megalopae in the genus *Menippe*. *Fishery Bulletin*, 86:289-297.
- Menu-Marque SA, 1970. Larval development of the crab *Platyxanthus crenulatus* (A. Milne Edwards, 1879) in the laboratory (Decapoda, Brachyura, Xanthidae). *Physis*, 29:477-494.
- Moore RC, McCormick L, 1969. General features of Crustacea. In: *Treatise on invertebrate paleontology, part R, Arthropoda 4, Vol. 1* (Ed., Moore RC). Geological Society of America and University of Kansas Press, Lawrence, KS, pp. 57-120.
- Ng PKL, Guinot D, Davie PJF, 2008. *Systema Brachyurorum: Part 1. An annotated checklist of extant brachyuran crabs of the world*. *The Raffles Bulletin of Zoology*, 17:1-286.
- Ortmann A, 1893. *Die Decapoden-Krebse des Strassburger Museums, mit besonderer Berücksichtigung der von Herrn Dr. Döderlein bei Japan und bei den Liu-Kiu-Inseln gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen*. VII. Theil. Abtheilung: Brachyura (Brachyura genuina Boas) II. Unterabtheilung: Cancroidea, 2. Section: Cancrinea, 1. Gruppe: Cyclometopa. *Zoologische Jahrbücher*, 7:411-495.
- Rice AL, 1980. Crab zoeal morphology and its bearing on the classification of the Brachyura. *Transactions of the Zoological Society of London*, 35:271-372. <https://doi.org/10.1111/j.1096-3642.1980.tb00060.x>
- Saba M, Takeda M, Nakasone Y, 1978a. Larval development of *Epixanthus dentatus* (White) (Brachyura, Xanthidae). *Bulletin of the National Science Museum, Series A (Zoology)*, 4:151-161.
- Saba M, Takeda M, Nakasone Y, 1978b. Larval development of *Baptozius vinosus* (H. Milne Edwards) (Crustacea, Brachyura, Xanthidae). *Proceedings of the Japanese Society of Systematic Zoology*, 14:29-42. https://doi.org/10.19004/pjssz.14.0_29
- Scotto LE, 1979. Larval development of the Cuban stone crab, *Menippe nodifrons* (Brachyura, Xanthidae), under laboratory conditions with notes on the status of the family Menippidae. *Fishery Bulletin*, 77:359-386.
- Stimpson W, 1858. *Prodromus descriptionis animalium evertebratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit W. Stimpson. Pars IV. Crustacea Cancroidea et Coryoidea*. *Proceedings of the Academy of Natural Sciences, Philadelphia*, 10:31-40.
- Terada M, 1982. Zoeal development of the menippid crab, *Eriphia smithi* MacLeay. *Zoological Magazine*, 91:255-262.
- Wear RG, 1968. Life-history studies on New Zealand Brachyura. *New Zealand Journal of Marine and Freshwater Research*, 2:698-707. <https://doi.org/10.1080/00288330.1968.9515267>
- WoRMS, 2022. *Eriphioidea* MacLeay, 1838. Accessed 30 Mar 2022, <<https://www.marinespecies.org/aphia.php?p=taxdetails&id=439029>>.

Received April 4, 2022
 Revised April 25, 2022
 Accepted April 25, 2022