

Posterior decentering of the humeral head in patients with arthroscopic rotator cuff repair

Hidehiro Nakamura¹, Masafumi Gotoh¹, Hirokazu Honda¹, Yasuhiro Mitsui¹, Hiroki Ohzono¹, Naoto Shiba², Shinichiro Kume¹, Takahiro Okawa¹

¹Department of Orthopedic Surgery, Kurume University Medical Center, Fukuoka, Japan

²Department of Orthopedic Surgery, Kurume University, Fukuoka, Japan

Background: In some patients with rotator cuff tear (RCT), the axial view of magnetic resonance imaging (MRI) shows subtle posterior decentering (PD) of the humeral head from the glenoid fossa. This is considered to result from a loss of centralization that is typically produced by rotator cuff function. There are few reports on PD in RCT despite the common occurrence of posterior subluxation in degenerative joint disease. In this study, we investigated the effect of PD in arthroscopic rotator cuff repair (ARCR).

Methods: We conducted a retrospective study of consecutive patients who underwent ARCR at our institute and were followed-up for at least 1 year. PD was identified as a 2-mm posterior shift of the humeral head relative to the glenoid fossa in the axial MRI view preoperatively. The tear size and fatty degeneration (FD, Goutallier classification) were also evaluated using preoperative MRI. Retears were evaluated through MRI at 1 year postoperatively.

Results: We included 135 shoulders in this study. Ten instances of PD (including seven retears) were observed preoperatively. Fifteen retears (three and 12 retears in the small/medium and large/massive tear groups, respectively) were observed postoperatively. PD was significantly correlated with tear size, FD, and re-tear occurrence ($p < 0.01$ each). The odds ratio for PD in retears was 34.1, which was greater than that for tear size ≥ 3 cm and FD grade ≥ 3 .

Conclusions: We concluded that large tear size and FD contribute to the occurrence of PD. Furthermore, PD could be a predictor of re-tear after ARCR.

Keywords: Rotator cuff tear; Humeral head; Shoulder

INTRODUCTION

In shoulders with rotator cuff deficiencies, humeral head (HH) decentralization appears as a superior migration [1,2]. There have been numerous reports on the clinical implications of superior HH migration in patients with rotator cuff tear (RCT) and its correlation with clinical outcomes following arthroscopic ro-

tator cuff repair (ARCR) [3-7]. Chalmers et al. [3] reported that proximal humeral migration occurred in patients with large RCT and infraspinatus involvement over a median follow-up period of 8 years. A study by Kuptniratsaikul et al. [4] found that superior HH migration correlated with arthroscopic reparability of large-to-massive RCTs.

Contrastingly, we occasionally observed subtle posterior HH

Received: September 21, 2021 Revised: November 22, 2021 Accepted: November 26, 2021

Correspondence to: Hidehiro Nakamura

Department of Orthopedic Surgery, Kurume University Medical Center, 155-1 Kokubu, Kurume, Fukuoka 839-0863, Japan

Tel: +81-942-22-6111, Fax: +81-942-22-6533, E-mail: nakamura_hidehiro@med.kurume-u.ac.jp, ORCID: <https://orcid.org/0000-0002-6269-9588>

Financial support: None.

Conflict of interest: None.

Copyright© 2022 Korean Shoulder and Elbow Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

migration in the glenoid fossa with RCT on axial magnetic resonance imaging (MRI) (Fig. 1). Considering the posterior HH subluxation and posterior glenoid wear that occurs in patients with cuff tear arthropathy, it is understandable that subtle posterior decentering (PD) of the HH can occur from rotator cuff deficiency. Although posterior subluxation is frequently reported in degenerative joint disease, there are few reports on PD in RCT without degenerative change [8,9]. Yun et al. [8] reported that PD detected on MRI arthrography was significantly associated with posterior synovial proliferation and rotator cuff atrophy. Kim and Seo [9] found that PD occurrence was related to tear size and fatty degeneration (FD) of the rotator cuff.

However, little is known about the effect of PD following RCT treatment. This study aimed to investigate the correlation between PD and retear after ARCR. We hypothesized that posterior HH decentering is correlated with retear after ARCR.

METHODS

This retrospective study was approved by Institutional Review Board of Kurume University (IRB No. 19072), and informed consent was obtained from patients. We identified all patients from a surgical registry with a minimum follow-up period of 12 months who underwent complete ARCR between January 2016 and December 2017 at our institution and who underwent both pre- and postoperative MRI evaluations. Patients who underwent partial repair or revision surgery were excluded.

Surgical Technique

The patients underwent ARCR in the beach-chair position under general anesthesia along with an interscalene block. Diagnostic arthroscopy was performed, and tear size was evaluated according to the DeOrio and Cofield classification [10]. Subsequently, the torn tendon was repaired using the single-row repair or double-row suture bridge techniques depending on tendon mobility and tear configuration. For single-row repairs, one row of an-

chors was placed on the lateral aspect of the rotator cuff footprint, and the torn cuff was fixed with an interrupted suture. For double-row suture bridge repairs, one row of anchors was placed on the medial aspect of the rotator cuff footprint with or without tying, and the torn cuff was transosseously fixed with a knotless anchor on the lateral aspect of the rotator cuff footprint. If needed, additional procedures including capsular release, tenotomy or tenodesis of the long head of the biceps tendon (LHB), distal clavicle excision, and arthroscopic subacromial decompression were performed. Particularly, partial repair was chosen when over-tensioning was required to move the tendon edge to a position just lateral to the articular margin, even after peritendinous adhesions and capsular contractures were released and any interval slides were performed [11,12].

Rehabilitation Protocol

Postoperatively, patients were immobilized using a sling with an abduction pillow, with the shoulder internally rotated at 30°–40° and abducted at 20°. Passive shoulder range of motion (ROM) exercises were commenced on postoperative day 4, and active ROM exercises were allowed at postoperative week 6. Isotonic muscle strengthening exercises were allowed at postoperative week 12 [11,12].

Clinical Assessment

Clinical outcomes were assessed using the Japanese Orthopaedic Association (JOA) and University of California, Los Angeles (UCLA) scores. Physical examinations were performed by an independent physical therapist who was blinded to the study. The outcome measures were assessed preoperatively and at final follow-up.

Structural Assessment

According to the arthroscopic findings, tear morphology (anterosuperior tear/posterosuperior tear/others) and LHB pathology (intact LHB/injured LHB/absent LHB) were analyzed [13]. The radiographic outcome was evaluated according to the Hamada classification on standard anteroposterior shoulder radiographs prior to surgical treatment [14].

All patients underwent MRI preoperatively and at final follow-up. All imaging was performed on a 1.5-Tesla scanner (Excelart; Toshiba Medical Systems, Tokyo, Japan) using routine pulse sequences with the patient in the supine position and the arm in a neutral position. The scans were reviewed by experienced orthopedic doctors who were blinded to this study.

When a fluid-equivalent signal was found or when there was non-visualization of the rotator cuff tendon on more than one




Fig. 1. Axial T2-weighted magnetic resonance imaging showing posterior decentering of the humeral head in patients with rotator cuff tear. (A) A 70-year-old man. (B) A 67-year-old man.

T2-weighted image, a retear was diagnosed at final follow-up [15]. FD of the subscapularis (SSc), supraspinatus (SSp), and infraspinatus (ISp) muscles was preoperatively evaluated on the most lateral oblique sagittal T2-weighted MRI that visualized the scapular spine in contact with the scapular body (the “Y-view”), using the Goutallier classification system [16].

PD was evaluated on pre- and postoperative MRI scans. We used T2-weighted axial images at the mid-glenoid level. The transverse axis of the scapula was detected as a line drawn from the medial end of the scapula to the mid-point of the glenoid fossa. PD was confirmed as present if the center of the HH had shifted more than 2 mm posterior to the scapula (Fig. 2) [8,9]. To assess the reproducibility, a random sample of 30 axial MRI slices from 30 individuals was reviewed twice by two observers (KE and YN) who were blinded to the study.

Statistical Analysis

Statistical analysis was performed using JMP 14 (SAS Institute, Cary, NC, USA). For all calculations, the level of significance was defined as $p < 0.05$. Data are expressed as mean value with standard deviation. The Mann-Whitney U-test was used to compare the pre- and postoperative clinical outcomes. The chi-square test was used to compare the nominal variables (retear, DeOrio and Cofield classification, Goutallier classification, and PD) in the structural outcomes. Intra- and interobserver agreement for the PD analysis were evaluated using the Cohen κ statistic and divided into the following categories: very good (0.81–1.00), good (0.61–0.80), moderate (0.41–0.60), fair (0.21–0.40),




Fig. 2. Posterior decentering. Yellow line: transverse axis of the scapula, red dot: center of the humeral head. If the red dot lays ≥ 2 mm posterior to the yellow line, the humeral head position is defined as decentered posteriorly.

or poor (0.00–0.20) [17].

RESULTS

There were 162 RCT patients who underwent ARCR at our institution from January 2016 to December 2017. Among these patients, 135 shoulders of 132 patients were enrolled in this study. The remaining 30 patients were excluded, including 2, 11, and 17 who underwent revision surgeries, partial repairs, or who were not followed up for at least 1 year, respectively.

Among the 135 shoulders of the 132 patients included, the mean age at the time of surgery was 63.7 ± 10.3 years (range, 30–88 years). The mean symptom duration preoperatively was 8.4 ± 7.9 months (range, 1–48 months). Single-row repairs were performed in 22 shoulders (16.3%), and the double-row suture bridge technique were performed in 113 shoulders (83.7%). All the patients’ JOA and UCLA scores improved significantly from 69.0 ± 11.8 and 15.9 ± 5.0 points preoperatively to 88.0 ± 11.2 and 29.0 ± 5.9 points at 1 year postoperatively ($p = 0.049$ and $p = 0.006$, respectively). In this study, postoperative retears were noted in 15 of the 135 shoulders (11.1%) at 1 year postoperatively (Table 1).

PD was observed in 10 patients (7.4%) preoperatively (Fig. 1) but not (0%) postoperatively. Among these 10 patients with preoperative PD, seven retears were observed. We noted a significant correlation between PD and re-tear ($p < 0.001$; odds ratio, 34.1; 95% confidence interval [CI], 7.4–157.7). In all preoperative PD patients, the postoperative MRI scans showed PD improvement and HH centralization (Fig. 3). Regarding PD analysis reproducibility, Cohen’s κ values were 1.0 and 0.92 (95% CI, 1.0–1.0 and 0.81–1.0, respectively) for intra-observer and inter-observer agreement, respectively.

According to the DeOrio and Cofield classification, small/medium and large/massive tears (< 3.0 cm and ≥ 3.0 cm, respectively) were observed in 98 and 37 shoulders, respectively (72.6% and 27.4%, respectively). The retear rate in the patients who had large/massive tears (12 retears/37 shoulders, 32.4%) was significantly higher than that in those who had small/medium tears (three retears/98 shoulders, 3.1%; $p < 0.001$; odds ratio, 15.2; 95% CI, 4.0–58.0). The preoperative PD rate in the large/massive tear group (8 PD/37 shoulders, 21.6%) was significantly higher than that in the small/medium tear group (2 PD/98 shoulders, 2.04%) ($p = 0.001$).

According to tear morphology, 53 anterosuperior tears (39.3%) and 43 posterosuperior tears were observed in this study. There was no significant correlation between preoperative PD and tear morphology ($p = 0.34$). According to intraoperative findings, the LHB was intact in 28 shoulders (20.7%), injured in 92 shoulders

Table 1. Patients' demographic data

Variable	Value (n = 135)
Age (yr)	63.4 ± 10.2 (30–85)
Sex	
Male	73 (54.1)
Female	62 (45.9)
Arm dominance	87 (64.9)
Symptom duration (wk)	8.4 ± 7.9 (1–48)
Traumatic onset	55 (40.7)
DeOrio and Cofield's classification	
Small	44 (32.6)
Medium	54 (40)
Large	33 (24.4)
Massive	4 (3)
Large/massive tear size	37 (27.4)
Goutallier classification	
SSc	
Stage 0–2	129 (95.6)
Stage 3–4	6 (4.4)
SSp	
Stage 0–2	118 (87.4)
Stage 3–4	17 (12.6)
ISp	
Stage 0–2	133 (98.5)
Stage 3–4	2 (1.5)
Goutallier stage 3–4 in any off the rotator cuff muscles	23 (17.0)
Preoperative PD	10 (7.4)
Postoperative PD	0
Repair technique	
Single low	37 (27.4)
Suture bridge	98 (62.6)
Retear	15 (11.1)

Values are presented as mean ± standard deviation (range) or number (%).

SSc: subscapularis, SSp: supraspinatus, ISp: infraspinatus, PD: posterior decentering.




Fig. 3. Preoperative (A) and postoperative (B) T2-weighted magnetic resonance imaging axial views in the patient who underwent arthroscopic rotator cuff repair. The posterior decentering improved and centralization of the humeral head was observed postoperatively.

(68.1%), and absent in 15 shoulders (4.3%). Surgical procedures were performed as below: preservation of LHB (n = 41, 30.4%),

LHB tenotomy (n = 82, 60.7%), and LHB tenodesis (n = 12, 8.9%). No significant correlation was observed between PD and LHB morphology/surgical procedures. ($p = 0.74$ and $p = 0.42$, respectively). No significant correlation was also detected between PD and the Hamada classification ($p = 0.23$). Eight PD (6.5%) were observed in Hamada grade 1 (n = 124), two PD (20%) in grade 2 (n = 10), and none in grade 3 (n = 1).

Low-grade Goutallier stages (stages 0–2) were seen in the SSCs, SSp, and ISp of 129, 118, and 133 shoulders, respectively (95.6%, 87.4%, and 98.5%). Preoperatively, high-grade Goutallier stages (stages 3 and 4) were seen in the SSCs, SSp, and ISp of 6, 17, and 2 shoulders, respectively, (4.4%, 12.6%, and 1.5%). The retear rate in the patients who had a high-grade Goutallier stage in any of the rotator cuffs (8 retears/23 shoulders, 34.8%) was significantly higher than that in those who did not (7 retears/112 shoulders, 6.3%; $p < 0.001$; odds ratio, 8.0; 95% CI, 2.5–25.3). Significant correlations were observed between PD and the Goutallier classification of SSp and ISp but not with that of SSC ($p = 0.002$, $p < 0.001$, and $p = 0.311$, respectively).

DISCUSSION

This study showed that 7.4% of RCT patients had PD preoperatively; as a result, PD could be recognized as a common MRI finding of RCT. PD had significant correlations with tear size and FD of SSp/ISp. Hence, it is understandable that these patients with PD had significantly higher postoperative retear rate in this study (70%; odds ratio, 34.1). The severity of RCT could be a reason for PD occurrence, and this could be a reason why PD is not found in all RCT patients. Although LHB is considered as a stabilizer of the shoulder joint, this study did not detect a significant correlation between arthroscopic LHB findings and preoperative PD. Particularly, ISp contributes to posterior stability of shoulder joints, and loss of this contribution could lead HH posterior shift in the supine position.

There are no specific methods needed to recognize PD; it can easily be seen on the axial view of a regular MRI scan in a clinical setting, and PD analysis reproducibility was excellent in this study. Therefore, we consider PD as a predictor of retear after ARCR, as well as tear size and FD of the torn rotator cuffs [13, 18,19].

Typically, HH decentralization appears as antero-superior migration in patients with rotator cuff dysfunction [2,3]. The Hamada classification describes the progression of cuff tear arthropathy as a gradual decrease in acromiohumeral interval. In this study, no significant relationship between PD and the Hamada classification was detected. Contrastingly, the Walch classi-

cation describes posterior HH subluxation and posterior glenoid fossa wear as the result of HH decentralization in degenerative disease [20,21]. In patients with rotator cuff dysfunction, PD can occur as a pre-stage pathology as a result of a similar mechanism to that of posterior HH subluxation in degenerative disease. Although posterior HH subluxation in degenerative disease is currently described as a percentage change from the midline of the scapula [22,23], PD in RCTs is a subtle change. Therefore, we defined PD as a greater than 2 mm posterior shift of the HH to the scapula, based on past studies [8,9].

Interestingly, all the patients with preoperative PD improved postoperatively. Thus, we recognized that PD has no direct influence on postoperative retear after ARCR. Here, as both PD and retear were significantly correlated with preoperative tear size (≥ 3.0 cm) and with high-grade FD (Goutallier classifications 3 and 4), we propose that a spurious correlation exists between PD and retears after ARCR. Considering the high retear rate, PD can present a difficult situation to arthroscopic rotator cuff treatment. Other procedures such as tendon transfer, superior capsular reconstruction, or reverse shoulder arthroplasty are alternative treatment options for RCT with PD [24,25].

Our study had some limitations. First, because of the small number of patients with PD, it lacked adequate statistical power required to evaluate the outcomes. Second, the nature of this retrospective cohort resulted in inherent biases related to follow-up and measurement. Third, we did not evaluate the correlation between superior migration amount and PD. Resolution of these issues will provide further clarification as to the outcomes obtained.

PD is a common MRI finding in patients with RCT and is significantly correlated with tear size, FD, and postoperative retears after ARCR. PD might be a predictor of retear after ARCR. Moreover, PD can complicate arthroscopic repair, and an appropriate treatment for PD should be established.

ACKNOWLEDGMENTS

We would like to thank for Kenichiro Eshima and Yousuke Nakamura who gave us clinical support.

ORCID

Hidehiro Nakamura <https://orcid.org/0000-0002-6269-9588>
 Masafumi Gotoh <https://orcid.org/0000-0002-1079-3425>
 Hirokazu Honda <https://orcid.org/0000-0002-6742-0208>
 Yasuhiro Mitsui <https://orcid.org/0000-0001-5689-438X>
 Hiroki Ohzono <https://orcid.org/0000-0001-7170-5401>

Naoto Shiba <https://orcid.org/0000-0002-9307-0280>
 Shinichiro Kume <https://orcid.org/0000-0001-8150-1357>
 Takahiro Okawa <https://orcid.org/0000-0003-2601-6177>

REFERENCES

1. Macaulay AA, Greiwe RM, Bigliani LU. Rotator cuff deficient arthritis of the glenohumeral joint. *Clin Orthop Surg* 2010;2: 196-202.
2. Hamada K, Yamanaka K, Uchiyama Y, Mikasa T, Mikasa M. A radiographic classification of massive rotator cuff tear arthritis. *Clin Orthop Relat Res* 2011;469:2452-60.
3. Chalmers PN, Salazar DH, Steger-May K, et al. Radiographic progression of arthritic changes in shoulders with degenerative rotator cuff tears. *J Shoulder Elbow Surg* 2016;25:1749-55.
4. Kuptniratsaikul V, Laohathaimongkol T, Umprai V, Yeekian C, Prasathaporn N. Pre-operative factors correlated with arthroscopic reparability of large-to-massive rotator cuff tears. *BMC Musculoskelet Disord* 2019;20:111.
5. Keener JD, Wei AS, Kim HM, Steger-May K, Yamaguchi K. Proximal humeral migration in shoulders with symptomatic and asymptomatic rotator cuff tears. *J Bone Joint Surg Am* 2009;91:1405-13.
6. Park SH, Choi CH, Yoon HK, Ha JW, Lee C, Chung K. What can the radiological parameters of superior migration of the humeral head tell us about the reparability of massive rotator cuff tears. *PLoS One* 2020;15:e0231843.
7. Taniguchi N, D'Lima DD, Suenaga N, et al. Translation of the humeral head scale is associated with success of rotator cuff repair for large-massive tears. *BMC Musculoskelet Disord* 2017; 18:511.
8. Yun G, Kang Y, Ahn JM, et al. Posterior decentering of the humeral head on shoulder MR arthrography: significant association with posterior synovial proliferation. *AJR Am J Roentgenol* 2017;208:1297-303.
9. Kim JH, Seo HW. Relationship of posterior decentering of the humeral head with tear size and fatty degeneration in rotator cuff tear. *Clin Shoulder Elb* 2019;22:121-7.
10. DeOrto JK, Cofield RH. Results of a second attempt at surgical repair of a failed initial rotator-cuff repair. *J Bone Joint Surg Am* 1984;66:563-7.
11. Honda H, Gotoh M, Mitsui Y, et al. Clinical and structural outcomes after arthroscopic rotator cuff repair: a comparison between suture bridge techniques with or without medial knot tying. *J Orthop Surg Res* 2018;13:297.
12. Nakamura H, Gotoh M, Mitsui Y, et al. Factors affecting clinical outcome in patients with structural failure after arthroscopic

- rotator cuff repair. *Arthroscopy* 2016;32:732-9.
13. Shimokobe H, Gotoh M, Honda H, et al. Risk factors for retear of large/massive rotator cuff tears after arthroscopic surgery: an analysis of tearing patterns. *J Orthop Surg Res* 2017;12:140.
 14. Hamada K, Fukuda H, Mikasa M, Kobayashi Y. Roentgenographic findings in massive rotator cuff tears: a long-term observation. *Clin Orthop Relat Res* 1990;(254):92-6.
 15. Davidson JF, Burkhart SS, Richards DP, Campbell SE. Use of preoperative magnetic resonance imaging to predict rotator cuff tear pattern and method of repair. *Arthroscopy* 2005; 21:1428.
 16. Goutallier D, Postel JM, Bernageau J, Lavau L, Voisin MC. Fatty muscle degeneration in cuff ruptures: pre- and postoperative evaluation by CT scan. *Clin Orthop Relat Res* 1994;(304):78-83.
 17. Cheung S, Dillon E, Tham SC, et al. The presence of fatty infiltration in the infraspinatus: its relation with the condition of the supraspinatus tendon. *Arthroscopy* 2011;27:463-70.
 18. Kim JR, Cho YS, Ryu KJ, Kim JH. Clinical and radiographic outcomes after arthroscopic repair of massive rotator cuff tears using a suture bridge technique: assessment of repair integrity on magnetic resonance imaging. *Am J Sports Med* 2012;40:786-93.
 19. Oh JH, Kim SH, Ji HM, Jo KH, Bin SW, Gong HS. Prognostic factors affecting anatomic outcome of rotator cuff repair and correlation with functional outcome. *Arthroscopy* 2009;25:30-9.
 20. Walch G, Badet R, Boulahia A, Khoury A. Morphologic study of the glenoid in primary glenohumeral osteoarthritis. *J Arthroplasty* 1999;14:756-60.
 21. Bercik MJ, Kruse K 2nd, Yalozis M, Gauci MO, Chaoui J, Walch G. A modification to the Walch classification of the glenoid in primary glenohumeral osteoarthritis using three-dimensional imaging. *J Shoulder Elbow Surg* 2016;25:1601-6.
 22. Ma CB, Xiao W, Salesky M, et al. Do glenoid retroversion and humeral subluxation affect outcomes following total shoulder arthroplasty. *JSES Int* 2020;4:649-56.
 23. Walch G, Moraga C, Young A, Castellanos-Rosas J. Results of anatomic nonconstrained prosthesis in primary osteoarthritis with biconcave glenoid. *J Shoulder Elbow Surg* 2012;21:1526-33.
 24. Juhan T, Stone M, Jalali O, et al. Irreparable rotator cuff tears: current treatment options. *Orthop Rev (Pavia)* 2019;11:8146.
 25. Sellers TR, Abdelfattah A, Frankle MA. Massive rotator cuff tear: when to consider reverse shoulder arthroplasty. *Curr Rev Musculoskelet Med* 2018;11:131-40.