

RESEARCH NOTE


Development of HRM Markers for Discrimination of Pyogo (*Lentinula edodes*) Cultivars Sanjo 701 and Chamaram

Suyun Moon¹, and Hojin Ryu^{1,2,*}¹Department of Biological Sciences and Biotechnology, Chungbuk National University, Cheongju 28644, Korea²Department of Biology, Chungbuk National University, Cheongju 28644, Korea*Corresponding author: hjryu96@chungbuk.ac.kr

ABSTRACT

Pyogo (Shiitake, *Lentinula edodes*) is one of the most important edible mushrooms because of its outstanding nutritive and medicinal value. In the registration and protection procedure for newly developed mushroom cultivars, the application of molecular markers that can supplement the morphological characteristic-based distinction has been strongly requested. Sanjo 701 and Chamaram, newly developed at the Federation Forest Mushroom Research Center of Korea, have been characterized as innovative cultivars suitable for customer demands because of their high yields and cultivation rates. However, no technical tools can protect the rights to these important cultivars. In this study, using comparative genomic information from 23 commercially available pyogo cultivars, we identified single nucleotide polymorphisms (SNPs) that accurately differentiated Sanjo701 and Chamaram from the other cultivars. We also developed high-resolution melting analysis (HRM)-based SNP markers that discriminate among the tested 23 pyogo cultivars. The developed SNP markers can be utilized for rapid, accurate identification of pyogo cultivars with low genetic diversity and to prevent cultivar contamination caused by illegally distributed inocula. In addition, these markers can serve as a crucial scientific basis for securing the right to conserve new cultivars in international markets.

Keywords: Cultivar discrimination, HRM (high-resolution melting), *Lentinula edodes*, Molecular marker, SNP (single nucleotide polymorphism)



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249Kor. J. Mycol. 2022 September, 50(3): 225-233
<https://doi.org/10.4489/KJM.20220023>**Received:** June 17, 2022**Revised:** August 15, 2022**Accepted:** August 19, 2022

© 2022 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pyogo (*Lentinula edodes*) is a basidiomycete of the order Agaricales that causes white rot in dead hardwoods [1]. This mushroom is rich in nutrients and a good source of provitamin D2. Pyogo is also known for its high pharmaceutical value, owing to the presence of polysaccharides and many compounds with antiviral, anticancer, antioxidant, and immunomodulatory activities [2-4]. Because of these positive attributes, pyogo is one of the most cultivated edible mushrooms, accounting for approximately 22% of global mushroom production, and is commercially cultivated in many countries, including by major producers in Korea, China, Japan, Australia, the United States, and Canada [5,6]. In 2020, pyogo production in Korea was approximately 20,207 tons, making it one of the most important forest products, accounting for approximately 98% of total forest mushroom production [7]. Because pyogo has a high economic value, individual breeders, the seed industry, and national institutions are developing new cultivars [8].

Plant variety protection is a system that protects new cultivars from unauthorized use and encourages breeding [9]. A specific cultivation test is required to confirm the differentiation, uniformity, and stability of cultivars for registration and protection. This cultivation test is a time-consuming, labor-intensive process because it is mainly based on morphological characteristics and is vulnerable to environmental factors [10,11]. For these reasons, the International Union for the Protection of New Cultivars (UPOV) suggests using sequence-based markers, such as reproducible simple sequence repeats (SSR), single nucleotide polymorphisms (SNP), and cleavage amplification polymorphisms (CAPS) [12,13].

Single nucleotide polymorphisms (SNPs) are mutations in a single nucleotide sequence that are caused by a transition between purines (A, G) or pyrimidines (C, T). SNP is the most widely used molecular marker because of its broad-spectrum distribution and lower analysis cost than other markers [14-16]. SNP discovery via Sanger sequencing technology is based on PCR and has low throughput. However, with the development of next-generation sequencing (NGS), high-throughput SNPs in short time navigation became possible [17]. Electrophoresis has been used in traditional SNP genotyping methods, such as cleaved amplified polymorphic sequence (CAPS), dCAPs (derived CAPS), and AS-PCR (allele-specific PCR) [18]. Because of technical advances, high-resolution melting (HRM)-based SNP marker analysis can efficiently use a fluorescent dye interposed between DNA double strands to measure the rate of DNA dissociation from double to single strands. It has been reported as one of the most powerful methods for analyzing genetic variation [19-21]. HRM analysis has been successfully used to identify crop cultivars, such as cherries, tangerines, peppers, and pomelo, as well as to ensure the authenticity of olive oil and wine [22-26]. In the case of mushrooms, HRM analysis has been used to identify hallucinogenic mushrooms and enoki mushrooms, as well as the pyogo cultivar Gyunheung No. 115 [27-29].

The most popular pyogo mushrooms cultivated on sawdust medium in Korea are Sanjo 701 and Chamaram, two new cultivars developed in the Federation Forest Mushroom Research Center of Korea. In 2020, the preference for the two cultivars in Korea was 45.2 percent and 25.2 percent, respectively [30]. Sanjo 701 is characterized by its bright color, firm flesh, and heft. Chamaram has a quantitative trait similar to that of Sanjo 701, with the additional advantage of being easily diversified in commercial distribution because of its excellent growth properties over a wide temperature spectrum [30]. However, no scientific or technical approach has been developed to adequately protect these two important cultivars from illegal distribution in domestic and global markets. In this study, we developed SNP markers that differentiated Chamaram and Sanjo 701 from 23 commercialized pyogo cultivars and then applied HRM analysis to secure a scientific method for distinguishing the two major pyogo cultivars.

For developing SNP markers that differentiate Sanjo 701 and Chamaram from one another, 22 locally developed strains and one imported strain were used [31], with B17 serving as a standard whole genome of *L. edodes* (Table 1). Mycelia of the tested strains were cultivated in dark culture at 25°C in a medium containing potato dextrose agar.

Table 1. List of strains analyzed in this study.

No.	Cultivars	Origin
1	Sanlim10 ^a	Korea
2	Gaeulhyang ^a	Korea
3	Chunjang2 ^a	Korea
4	Sanmaru1 ^a	Korea
5	Sanmaru2 ^a	Korea
6	Sanbaekhyang ^a	Korea
7	Suhyanggo ^a	Korea
8	KFRI 31 ^a	Japan
9	Sanjo101 ^b	Korea
10	Sanjo102 ^b	Korea
11	Sanjo103 ^b	Korea
12	Sanjo109 ^b	Korea
13	Sanjo111 ^b	Korea
14	Sanjo502 ^b	Korea
15	Sanjo701 ^b	Korea
16	Sanjo704 ^b	Korea
17	Sanjo705 ^b	Korea
18	Sanjo706 ^b	Korea
19	Sanjo707 ^b	Korea
20	Sanjo708 ^b	Korea
21	Sanjo709 ^b	Korea
22	Sanjo710 ^b	Korea
23	Chamaram ^b	Korea

^a Cultivars provided by National Institute of Forest Science in Korea Forest Service

^b Cultivars provided by Forest Mushroom Research Center.

Genomic DNA was extracted from mycelium cultured by shaking at 25°C and 110 rpm under dark conditions for approximately 2 weeks. The cultured mycelium was washed with PBS buffer (NaCl 135 mM, KCl 2.7 mM, Na₂HPO₄ 4.3 mM, KH₂PO₄ 1.4 mM) and dried with a kitchen towel. Approximately 100 mg of dried mycelium was frozen in liquid nitrogen and finely ground with a mortar, and genomic DNA was extracted using the GenEx™ Plant kit (GeneAll, Seoul, Korea). The extracted DNA was quantified with a micro-spectrophotometer K5600 (BioFuture Inc., China), diluted to 20 ng/μL and used in this study.

The method developed by Hillier et al. [32] was used to identify SNPs found in Sanjo 701 and Chamaram. Genomic DNA extracted from 23 cultivars was resequenced on the HiSeq 2500 platform, and SNPs found specifically in the genomes of Sanjo 701 and Chamaram were selected (Tables 2 and 3). One homozygous SNP, three heterozygous SNPs from Sanjo 701, and three heterozygous SNPs from Chamaram were chosen because only missense SNPs in the coding region of specific genes were selected (Tables 2 and 3).

Table 2. List of specific SNPs presented in Sanjo701.

#Chr	POS	REF	ALT	Genotype	Codon	Gene	Description
Scaffold2	817310	C	T	Homozygous	c.283G>A	GENE10946	-
Scaffold7	4123084	G	A	Heterozygous	c.700G>A	GENE07542	Hypothetical protein Moror_8994
Scaffold15	668068	A	G	Heterozygous	c.1166A>G	rbm8a (RNA binding motif protein 8A)	RNA-binding protein 8A
Scaffold24	187022	C	T	Heterozygous	c.-1G>A	SEC65 (signal recognition particle subunit SEC65)	Signal recognition particle subunit SEC65

The SNP used for designing the HRM marker is in red.

SNP: Single nucleotide polymorphism; POS: Position ; REF: References; ALT: Alternative nucleotides; HRM: High-resolution melting.

Table 3. List of specific SNPs presented in Chamaram.

#Chr	POS	REF	ALT	Genotype	Codon	Gene	Description
Scaffold7	1072836	T	G	Heterozygous	c.*3510T>G	NST1 (negatively-affecting salt tolerance protein 1)	Stress response protein NST1
Scaffold14	823041	A	C	Heterozygous	c.659A>C	INT6 (integration site 6)	Eukaryotic translation initiation factor 3 subunit E
Scaffold20	243980	C	G	Heterozygous	c.-1G>C	URE1 (urease)	Urease

The SNP used for designing the HRM marker is in red.

SNP: Single nucleotide polymorphism; POS: Position ; REF: References; ALT: Alternative nucleotides; HRM: High-resolution melting.

Table 4. Characteristics of primer sets for discrimination of Sanjo701 and Chamaram.

Primer name	Primer sequences (5'-3')	Tm (°C) ^a	Amplicon size (bp) ^b
RL-LE-202 F	GCTTGATCAATTCGTTGGAA	54.9	104
RL-LE-202 R	GCAATGGAAAGTTGGACATC	54.2	
RL-LE-203 F	ACTCGTTCGTGAAGCCATAG	54.2	112
RL-LE-203 R	TCGGGAATGAAGCTGACTTA	54.7	

^aTm: Annealing temperature;

^bAmplicon size: Size of the fragment amplified using high-resolution melting (HRM) primers.

Primer Express software (Applied Biosystems, Foster City, CA, USA) was used to extract the flanking sequence from the selected SNP and amplify the marker sequence. PCR amplification of the marker regions was performed by 35 cycles at 95°C for 3 min, 95°C for 30 s, 55°C for 30 s, and 72°C for 20 s, followed by a 5 min reaction at 72°C.

PCR for HRM analysis was performed using 20 ng genomic DNA 1 µL, 2X MeltDoctor™ HRM Master Mix (Applied Biosystems, Foster City, CA, USA), 10 µL 10 pmole/µL primers F and R (1 µL each), and 7 µL DW. After denaturation at 95°C for 10 min using QuantStudio 3 (Applied Biosystems, Foster City, CA, USA), denaturation at 95°C for 15 s, annealing at 54°C for 30 s, and extension at 72°C for 30 s were performed on a mixture of 20 µL. After 40 repetitions and 10 s of heating at 95°C, the melt curve was determined by gradually increasing the temperature from 60°C to 95°C at a rate of 0.025°C/s. High-resolution melt software version 3.2 was used to analyze the melting curve (Applied Biosystems, Foster City, CA, USA).




Fig. 1. Sanger sequencing validation of SNPs in Sanjo 701 and Chamaram. (A) In Sanjo701, heterozygous G to A transition was observed. (B) In Chamaram, heterozygous A to C transition was confirmed.




Fig. 2. Discrimination of Sanjo701 and Chamaram by high-resolution melting curve analysis among 23 pyogo cultivars. Aligned melting curve analysis (A) and differential plot (B) of RL-LE-002 using ‘Sanjo101’ as a reference. Aligned melting curve analysis (C) and differential plot (D) of RL-LE-203 using ‘Chunjang2’ as a reference. Results for Sanjo701 and Chamaram are in blue, and those for the other 22 strains are in red.

Using Sanger sequencing, we confirmed the SNPs on 4,123,084 bp in Scaffold 7 in the genome of Sanjo 701, which changed the nucleotide sequence from G to A (Fig.1A; Table 2), and 823,041bp in Scaffold 14, which changed the nucleotide sequence from A to C in the Chamaram genome (Fig.1B; Table 3). However, Sanger sequencing further proved that additional non-specific SNPs existed in genomic regions other than the selected SNPs (highlighted in red in Tables 2 and 3). On the basis of these results, two specific SNPs, Chamaram and Sanjo 701, were selected as the most suitable targets for HRM marker development. The RL-LE-202 and RL-LE-203 primer sets were designed to produce 104 and 112 bp amplicons for each SNP-containing sequence, respectively, by PCR. (Table 4).

HRM-PCR analysis of 23 strains, including Sanjo 701 and Chamaram, revealed that the RL-LE-202 marker clearly produced a heterotype curve for Sanjo 701 and homotype curves for the other strains (Figs. 2A and B). Chamaram was also distinguished by a specific heterotype curve by using the RL-LE-203 primer set (Figs. 2C and D). These findings indicate that the cultivars Sanjo 701 and Chamaram can be precisely differentiated using the SNP-based HRM markers developed in this study.

SNP and indels are the two most common types of nucleotide sequence polymorphism. Most of them are biallelic and found in large quantities across the genome [33]. SNPs found in the coding area of genes can be employed more effectively than randomly selected SNPs to discover functionally significant variations [34]. However, because the amount of genetic information per marker is modest, the duration of the marker development selection process is long and is restricted because it needs to evaluate many loci [35,36]. NGS technology delivers large-scale sequence datasets that can be used for genetic mapping, genetic diversity analysis, gene identification, and molecular breeding. These large-scale sequence datasets can also be used to define sequence diversity and create polymorphism and genotyping data [37-39]. Other applications of this technology include digital molecular breeding. This instrument is excellent for performing such tasks [40]. Additionally, it allows for the discovery of new or reference-based genomic variations, even in organisms with little or no genetic information, and provides a quick, easy method to obtain the location and genotype information of the related genome [41,42].

This study demonstrates that SNP identification, marker development, and HRM analysis can be successfully applied to the discrimination of pyogo cultivars with low genetic diversity by using comparative genomics with the whole genomic information of the B17 *L. edodes* strain. The developed SNP marker can accurately identify shiitake varieties even at the mycelial growth stage; thus, we expect it to prevent the mixing of varieties. In addition, the SNP marker development method presented in this study can be used to develop markers that can discriminate among other edible mushrooms.

CONFLICT OF INTERESTS

No conflict of interest was reported by the author(s).

ACKNOWLEDGEMENT

This research was supported by the Chungbuk National University Korea National University Development Project (2022).

REFERENCES

1. Lee KW, Jeon JO, Kim MJ, Kim IJ, Jang MJ, Park HS. Effects of difference in medium composition on the growth of *Lentinula edodes*. *J Mushrooms* 2018;16:267-71.
2. Bisen P, Baghel RK, Sanodiya BS, Thakur GS, Prasad G. *Lentinus edodes*: a macrofungus with pharmacological activities. *Curr Med Chem* 2010;17:2419-30.
3. Finimundy TC, Dillon AJP, Henriques JAP, Ely MR. A review on general nutritional compounds and pharmacological properties of the *Lentinula edodes* mushroom. *Food Nutr Sci* 2014;5:12.
4. Wasser SP. Shiitake (*Lentinus edodes*). New York: Marcel Dekker Inc.; 2004. p. 653-64.
5. Mata G, Gaitán-Hernández R. Cultivation of the edible mushroom *Lentinula edodes* (Shiitake) in pasteurized wheat straw—alternative use of geothermal energy in Mexico. *Eng Life Sci* 2004;4:363-7.
6. Royse DJ, Baars J, Tan Q. Current overview of mushroom production in the world. Hoboken: John Wiley & Sons Ltd.; 2017. p. 5-13.
7. Horticulture Business Division. Minister of Agriculture Food and Rural Affairs. 2019 The production record of a special crop. Sejong: Minister of Agriculture, Food and Rural Affairs; 2020. p. 64-7.
8. Jang YS, Kwon YR, Kim TH. Prospect and status of plant variety protection (PVP) of forest-sector in Korea. *Korean J Breed Sci* 2020;52:31-9.
9. Nguyen NN, Kim M, Jung JK, Shim EJ, Chung SM, Park Y, Lee GP, Sim SC. Genome-wide SNP discovery and core marker sets for assessment of genetic variations in cultivated pumpkin (*Cucurbita* spp.). *Hortic Res* 2020;7:121.
10. Bonow S, Von Pinho EV, Vieira MG, Vosman B. Microsatellite markers in and around rice genes: applications in variety identification and DUS testing. *Crop Sci* 2009;49:880-6.
11. Jamali SH, Cockram J, Hickey LT. Insights into deployment of DNA markers in plant variety protection and registration. *Theor Appl Genet* 2019;132:1911-29.
12. Hong JH, Chae CW, Choi KJ, Kwon YS. A database of simple sequence repeat (SSR) marker-based DNA profiles of citrus and related cultivars and germplasm. *Korean J Hortic Sci Technol* 2016;34:142-53.
13. International Union for the Protection of New Varieties of Plants (UPOV). Guidelines for DNA-Profiling: Molecular Marker Selection and Database Construction ("BMT Guideline") [Internet]. Geneva: UPOV; 2010 [cited 2022 Aug 13]. Available from https://www.upov.int/edocs/infdocs/en/upov_inf_17_1.pdf.
14. Cheng A, Chai HH, Ho WK, Bamba ASA, Feldman A, Kendabie P, Halim RA, Tanzi A, Mayes S, Massawe F. Molecular marker technology for genetic improvement of underutilised crops. In: Abdullah SNA, Chai-Ling H, Wagstaff C, editors. *Crop Improvement: Sustainability Through Leading-Edge Technology*. Cham: Springer International Publishing; 2017. p. 47-70.

15. Schlötterer C. The evolution of molecular markers—just a matter of fashion? *Nat Rev Genet* 2004;5:63-9.
16. Trick M, Long Y, Meng J, Bancroft I. Single nucleotide polymorphism (SNP) discovery in the polyploid *Brassica napus* using Solexa transcriptome sequencing. *Plant Biotechnol J* 2009;7:334-46.
17. Zhang S, Li B, Chen Y, Shaibu AS, Zheng H, Sun J. Molecular-assisted distinctness and uniformity testing using SLAF-sequencing approach in soybean. *Genes* 2020;11:175.
18. Zhang J, Yang J, Zhang L, Luo J, Zhao H, Zhang J, Wen C. A new SNP genotyping technology Target SNP-seq and its application in genetic analysis of cucumber varieties. *Sci Rep* 2020;10:1-11.
19. Herrmann MG, Durtschi JD, Wittwer CT, Voelkerding KV. Expanded instrument comparison of amplicon DNA melting analysis for mutation scanning and genotyping. *Clin Chem* 2007;53:1544-8.
20. Reed GH, Wittwer CT. Sensitivity and specificity of single-nucleotide polymorphism scanning by high-resolution melting analysis. *Clin Chem* 2004;50:1748-54.
21. White HE, Hall VJ, Cross NC. Methylation-sensitive high-resolution melting-curve analysis of the SNRPN gene as a diagnostic screen for Prader-Willi and Angelman syndromes. *Clin Chem* 2007;53:1960-2.
22. Ganopoulos I, Tsaballa A, Xanthopoulou A, Madesis P, Tsaftaris A. Sweet cherry cultivar identification by high-resolution-melting (HRM) analysis using gene-based SNP markers. *Plant Mol Biol Report* 2013;31:763-8.
23. Jeong HJ, Jo YD, Park SW, Kang BC. Identification of *Capsicum* species using SNP markers based on high resolution melting analysis. *Genome* 2010;53:1029-40.
24. Jin SB, Kim HB, Park S, Kim MJ, Choi CW, Yun SH. Identification of the ‘Haryejaeang’ mandarin cultivar by multiplex PCR-based SNP genotyping. *Mol Biol Rep* 2020;47:8385-95.
25. Pereira L, Gomes S, Barrias S, Fernandes JR, Martins-Lopes P. Applying high-resolution melting (HRM) technology to olive oil and wine authenticity. *Food Res Int* 2018;103:170-81.
26. Run-ting Y, Bo W, Chong L, Pei Z, Ji-wu Z, Yun Z, Bo J, Bi-rong Z, Guang-yan Z. Comparison of allele-specific PCR and high resolution melting analysis in SNP genotyping and their application in pummelo cultivar identification. *Acta Horticulturae Sinica* 2013;40:1061.
27. Im JH, Oh MJ, Oh YL, Raman J, Jang KY, Kong WS. Development of SNP markers for discriminating color of *Flammulina velutipes*. *J Mushrooms*; 2019. p. 42.
28. Maeda A, Terashima K, Hasebe K. Development of a method for rapid strain-typing of a shiitake cultivar, Kinko 115, by high-resolution melting (HRM) analysis. *Mushroom Science and Biotechnology* 2015;23:114-9.
29. Zhang X, Yu H, Yang Q, Wang Z, Xia R, Chen C, Qu Y, Tan R, Shi Y, Xiang P. A forensic detection method for hallucinogenic mushrooms via high-resolution melting (HRM) analysis. *Genes* 2021;12:199.
30. Korea Rural Economic Institute. Forestry observation for *Lentinula edodes*, November. Naju: Korea Rural Economic Institute; 2020. p. 1-6.
31. Shim D, Park SG, Kim K, Bae W, Lee GW, Ha BS, Ro HS, Kim M, Ryoo R, Rhee SK. Whole genome de novo sequencing and genome annotation of the world popular cultivated edible mushroom, *Lentinula edodes*. *J Biotechnol* 2016;223:24-5.
32. Hillier LW, Marth GT, Quinlan AR, Dooling D, Fewell G, Barnett D, Fox P, Glasscock JJ,

- Hickenbotham M, Huang W. Whole-genome sequencing and variant discovery in *C. elegans*. *Nat Methods* 2008;5:183-8.
33. Cabezas JA, Ibáñez J, Lijavetzky D, Vélez D, Bravo G, Rodríguez V, Carreño I, Jermakow AM, Carreño J, Ruiz-García L. A 48 SNP set for grapevine cultivar identification. *BMC Plant Biol* 2011;11:1-12.
 34. Luikart G, England PR, Tallmon D, Jordan S, Taberlet P. The power and promise of population genomics: from genotyping to genome typing. *Nat Rev Genet* 2003;4:981-94.
 35. Glaubitz JC, Rhodes Jr OE, DeWoody JA. Prospects for inferring pairwise relationships with single nucleotide polymorphisms. *Mol Ecol* 2003;12:1039-47.
 36. Oh YL, Choi IG, Kong WS, Jang KY, Oh MJ, Im JH. Evaluating genetic diversity of *Agaricus bisporus* accessions through phylogenetic analysis using single-nucleotide polymorphism (SNP) markers. *Mycobiology* 2021;49:61-8.
 37. Oh YL, Choi IG, Jang KY, Kim MS, Oh MJ, Im JH. SNP-based genetic linkage map and quantitative trait locus mapping associated with the agronomically important traits of *Hypsizygus marmoreus*. *Mycobiology* 2021;49:589-98.
 38. Woo SI, Kim ES, Han JG, Jang KY, Shin PG, Oh YL, Oh MJ, Jo SH, Lee JH, Kim KS. Genome-wide single nucleotide polymorphism-based assay for phylogenetic relationship of the *Flammulina velutipes*. *Kor J Mycol* 2015;43:231-8.
 39. Wu J, Choi J, Asiegbu FO, Lee YH. Comparative genomics platform and phylogenetic analysis of fungal laccases and multi-copper oxidases. *Mycobiology* 2020;48:373-82.
 40. Li D, Zeng R, Li Y, Zhao M, Chao J, Li Y, Wang K, Zhu L, Tian W-M, Liang C. Gene expression analysis and SNP/InDel discovery to investigate yield heterosis of two rubber tree F1 hybrids. *Sci Rep* 2016;6:1-12.
 41. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, Del Angel G, Rivas MA, Hanna M. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 2011;43:491-8.
 42. Kumar S, Banks TW, Cloutier S. SNP discovery through next-generation sequencing and its applications. *Int J Plant Genomics* 2012;2012:831460.