



## Research Article

Syringaresinol derived from *Panax ginseng* berry attenuates oxidative stress-induced skin aging via autophagyWooram Choi<sup>a,1</sup>, Hyun Soo Kim<sup>b,1</sup>, Sang Hee Park<sup>c</sup>, Donghyun Kim<sup>b</sup>, Yong Deog Hong<sup>b,\*\*\*</sup>, Ji Hye Kim<sup>a,\*\*</sup>, Jae Youl Cho<sup>a,c,\*</sup><sup>a</sup> Department of Integrative Biotechnology, And Biomedical Institute for Convergence at SKKU (BICS), Sungkyunkwan University, Suwon, Republic of Korea<sup>b</sup> Research and Innovation Center, AMOREPACIFIC, Yongin, Republic of Korea<sup>c</sup> Department of Biocosmetics, Sungkyunkwan University, Suwon, Republic of Korea

## ARTICLE INFO

## Article history:

Received 28 June 2021

Received in revised form

15 August 2021

Accepted 17 August 2021

Available online 21 August 2021

## Keywords:

*Panax ginseng* berry

Syringaresinol

Autophagy

Antioxidant

Anti-aging

## ABSTRACT

**Background:** In aged skin, reactive oxygen species (ROS) induces degradation of the extracellular matrix (ECM), leading to visible aging signs. Collagens in the ECM are cleaved by matrix metalloproteinases (MMPs). Syringaresinol (SYR), isolated from *Panax ginseng* berry, has various physiological activities, including anti-inflammatory action. However, the anti-aging effects of SYR via antioxidant and autophagy regulation have not been elucidated.

**Methods:** The preventive effect of SYR on skin aging was investigated in human HaCaT keratinocytes in the presence of H<sub>2</sub>O<sub>2</sub>, and the keratinocyte cells were treated with SYR (0–200 µg/mL). mRNA and protein levels of MMP-2 and -9 were determined by real-time PCR and Western blotting, respectively. Radical scavenging activity was researched by 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) assays. LC3B level was assessed by Western blotting and confocal microscopy.

**Results:** SYR significantly reduced gene expression and protein levels of MMP-9 and -2 in both H<sub>2</sub>O<sub>2</sub>-treated and untreated HaCaT cells. SYR did not show cytotoxicity to HaCaT cells. SYR exhibited DPPH and ABTS radical scavenging activities with an EC<sub>50</sub> value of 10.77 and 10.35 µg/mL, respectively. SYR elevated total levels of endogenous and exogenous LC3B in H<sub>2</sub>O<sub>2</sub>-stimulated HaCaT cells. 3-Methyladenine (3-MA), an autophagy inhibitor, counteracted the inhibitory effect of SYR on MMP-2 expression.

**Conclusion:** SYR showed antioxidant activity and up-regulated autophagy activity in H<sub>2</sub>O<sub>2</sub>-stimulated HaCaT cells, lowering the expression of MMP-2 and MMP-9 associated with skin aging. Our results suggest that SYR has potential value as a cosmetic additive for prevention of skin aging.

© 2022 The Korean Society of Ginseng. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Abbreviations:** ROS, reactive oxygen species; PCR, polymerase chain reaction; ECM, extracellular matrix; MMP, matrix metalloproteinase; SYR, Syringaresinol; DPPH, 2,2-diphenyl-1-picrylhydrazyl; ABTS, 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid; 3-MA, 3-Methyladenine; UV, ultraviolet; CMA, chaperone-mediated autophagy; FoxO3a, Forkhead box O3; MAPK, mitogen-activated protein kinase; DMEM, Dulbecco's Modified Eagle's medium; FBS, fetal bovine serum; PBS, phosphate-buffered saline; DMSO, dimethyl sulfoxide.

\* Corresponding author. Department of Integrative Biotechnology, and Biomedical Institute for Convergence at SKKU (BICS), Sungkyunkwan University, Suwon, 16419, Republic of Korea.

\*\* Corresponding author.

\*\*\* Corresponding author.

E-mail addresses: [chwoo1028@naver.com](mailto:chwoo1028@naver.com) (W. Choi), [hsookim@amorepacific.com](mailto:hsookim@amorepacific.com) (H.S. Kim), [84701@naver.com](mailto:84701@naver.com) (S.H. Park), [hydhong@amorepacific.com](mailto:hydhong@amorepacific.com) (Y.D. Hong), [kjhkhmlml@skku.edu](mailto:kjhkhmlml@skku.edu) (J.H. Kim), [jaecho@skku.edu](mailto:jaecho@skku.edu) (J.Y. Cho).

<sup>1</sup> These authors equally contributed to this work.

<https://doi.org/10.1016/j.jgr.2021.08.003>

1226-8453/© 2022 The Korean Society of Ginseng. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Skin consists of two main layers, the epidermis and dermis. During skin aging processes, degenerative changes of the dermis are more pronounced than those of the epidermis. The dermis is composed of keratinocytes and extracellular matrix (ECM) such as collagen, elastin, and proteoglycans [1]. With the aging of skin, the dermis loses ECMs, particularly collagen [2–4]. One of the main mechanisms for reducing ECM levels in the dermis is activation of matrix metalloproteinases (MMPs). MMPs are zinc-dependent endopeptidases composed of four subgroups (collagenases, gelatinases, stromelysins, and membrane-type MMPs (MT-MMPs)), cleaving ECM proteins and are functionally classified into [5]. MMP-2 and -9 belong to the gelatinase group and are involved in the breakdown of collagen and elastic fibers [6,7]. MMPs are increased

in both intrinsically aged skin and in photoaged skin caused by prolonged exposure to ultraviolet (UV) light [8,9]. Therefore, lowering MMP levels might be a promising approach for anti-aging.

Autophagy, derived from the Greek word “self-eating,” is a set of highly conserved cellular mechanisms by which unnecessary or dysfunctional organelles and cellular components are broken down and recycled in response to nutrient or stress conditions [10]. Macroautophagy, microautophagy, and chaperone-mediated autophagy (CMA) are classified types of autophagy process [11]. Macroautophagy forms autophagosomes with double layered-membrane structures that engulf cellular cargo [12,13]. Formation of autophagolysosomes is generated by fusion of autophagosomes with lysosomes [14,15]. Autophagy plays a role in various normal cellular functions, including cellular homeostasis maintenance, differentiation [16], and immune responses [17]. Dysregulation of autophagy can contribute to diverse pathologies, including cancer, neurodegeneration, aging, and heart disease [18]. Defects of autophagy also are associated with skin aging. The role of autophagy in skin aging is understood mainly in terms of regulating proteostasis, but its precise function remains to be elucidated [19]. Nevertheless, attempts to increase autophagy in senescent skin are warranted for anti-aging.

*Panax ginseng* has numerous therapeutic biochemical and pharmacological effects such as immunomodulatory [20], anti-inflammatory [21,22], antioxidative [23], anti-cancer [24,25], anti-fatigue [23], and anti-diabetic effects [26]. Based on these therapeutic effects, various parts and active pharmacological components of ginseng, including root, berry, and ginsenosides, have been used as functional foods [27]. Syringaresinol (SYR) (Fig. 1A Left panel), a plant lignan, is one of the biologically active ingredients present in ginseng berry [28]. SYR has anti-inflammatory [29], anti-fungal, and anti-cancer activities [30]. In addition, in skin-related studies, SYR has been reported to decrease the activity of Forkhead box O3 (FoxO3a) and mitogen-activated protein kinase (MAPK), which are aging-related factors [31,32]. However, the regulatory roles of SYR in aging process of skin are not fully elucidated. Therefore, in the present study, the beneficial effects of SYR derived from *Panax ginseng* on skin aging and the underlying biological mechanism for these effects, particularly those on antioxidant and autophagy regulation were looked over.

## 2. Materials and methods

### 2.1. Materials

HaCaT skin keratinocyte cells (American Type Culture Collection (ATCC) (Rockville, MD, USA) was used. Antibiotics (penicillin-streptomycin solution) was purchased from Hyclone (Logan, UT, USA). 3-Methyladenine (3-MA, Fig. 1A Left panel), 3-(4-(5-dimethylthiazol-2-yl)-2-5-diphenyltetrazolium bromide (MTT) was obtained from Sigma-Aldrich (St. Louis, MO, USA). The cDNA synthesis kit was purchased from Thermo Fisher Scientific (Waltham, MA, USA). Primers for determining the mRNA levels of MMP-2 and -9 were from Bioneer (Seoul, Korea). Antibodies against MMP-2, -9, LC-3B, and  $\beta$ -actin were employed, as reported previously [33]. SYR (Fig. 1A Left panel) was purified from *Panax ginseng* berry as reported previously [34]. The purity of the compound was >80% (Fig. 1A Right panel). A stock solution of SYR was prepared with 100% DMSO.

### 2.2. Cell culture

Keratinocyte HaCaT cells were cultured in DMEM with 10% FBS and 1% antibiotics (penicillin and streptomycin) in a 5% CO<sub>2</sub> incubator at 37 °C.

### 2.3. RNA extraction and quantitative polymerase chain reaction

HaCaT cells ( $2 \times 10^5$  cells/ml) plated for 18 h were exposed with H<sub>2</sub>O<sub>2</sub> (500  $\mu$ M) or SYR (100 or 200  $\mu$ g/mL). After 24 h, total RNA was prepared by general extraction method with TRIzol reagent as carried out before [35]. For synthesizing complementary DNA, a cDNA synthesis kit was used, and quantitative polymerase chain reaction was performed with Pcrbio's qPCR BIO SyGreen mix [36]. Primer sequences of MMP-2 and -9 were summarized in Table 1.

### 2.4. Preparation of cell lysates and immunoblotting analysis

Total lysates of HaCaT cells were obtained using lysis buffer as reported previously [37]. The prepared whole cell lysates were tested after clarification by centrifugation at 12,000 rpm for 1 min at 4 °C. Immunoblotting analysis was used to detect levels of MMP2, MMP9, LC-3B, and  $\beta$ -actin. Each antibody was treated to the PVDF membrane in 3% BSA in TBST at a ratio of 1:2500 at 4 °C. After 18 h, the second antibodies (anti-mouse or anti-rabbit antibodies) were reacted for 1 h at a ratio of 1:2500 at 20 °C. Immunoreactive bands were detected with enhanced peroxidase detection (EPD) of ELPIS-BIOTECH in Chemidoc of ATTO.

### 2.5. Cell viability assay

Whether SYR induces death of HaCaT cells was elucidated by a conventional MTT assay [38].


In  $2 \times 10^5$  of HaCaT cells, SYR (12.5–200  $\mu$ g/mL) was treated for 24 h.

### 2.6. DPPH and ABTS colorimetric assays

Either SYR (12.5–200  $\mu$ g/mL) or ascorbic acid (250  $\mu$ M) was reacted with 300  $\mu$ M DPPH or 740  $\mu$ M ABTS for 15 min at 37 °C. Then, the absorbance of each mixture was observed at 517 nm or 730 nm with spectrophotometer. The radical scavenging activity was presented as percent inhibition using the formula: Radical scavenging activity (%) = [(A0-A1)/A0]  $\times$  100, where A0 is the absorbance of DPPH or ABTS alone, and A1 is the absorbance of the sample.

### 2.7. Confocal microscopy

HaCaT cells ( $5 \times 10^4$  cells/mL) were transfected with 1  $\mu$ g/mL of a GFP-LC3B plasmid using transfection reagent, lipofectamine. After 24 h incubation, the cells were exposed to H<sub>2</sub>O<sub>2</sub> (500  $\mu$ M) or SYR (200  $\mu$ g/mL) for 12 h. Then, the cells were fixed with 4% paraformaldehyde (PFA) in PBS for 10 min, followed by treating Triton X-100 (0.5%) in PBS for 10 min for permeabilization of the cell membrane. The cells were continuously treated with PBS containing 1% BSA for 1 h at 20 °C for blocking. DNA staining was carried out by adding PBS with 10  $\mu$ g/mL Hoechst 33342 for 10 min. After mounting on glass slides, images of the cells were observed using a laser-scanning confocal microscope (Zeiss LSM 710 META, Oberkochen, Germany) with a 63x oil-immersion objective lens.



**Fig. 1.** Anti-aging activity of SYR in HaCaT cells. (A) Chemical structure of syringaresinol and 3-Methyladenin (Left panel) and analysis profile of syringaresinol by HPLC (Right panel). (B, C, and D) Following SYR (100 and 200 μg/mL) treatment, HaCaT cells were incubated with H<sub>2</sub>O<sub>2</sub> (500 μM) for 24 h mRNA levels of MMP-2 (B) and MMP-9 (C) were determined by real-time PCR. Protein levels of MMP-2 and MMP-9 were measured by Western blotting (D). (E) HaCaT cells were treated with various concentrations (0–200 μg/mL) of SYR and then incubated for 24 h. Protein levels of MMP-2 and MMP-9 were determined by immunoblotting analysis. (F) Viability of SYR-treated HaCaT cells was measured by MTT assay. #*p* < 0.05 and ##*p* < 0.01 compared with the normal group; \**p* < 0.05 and \*\**p* < 0.01 compared with the control group.

**Table 1**  
Primer list used in this study.

Gene	Direction	Sequences
MMP-2	F	CAAGTGGAGAGCAGTTGAGGACATC
	R	TGAGGACATCTCCACGTCAA
MMP-9	F	GCCACTGTGCGGCGATAAGG
	R	CACTGTCCACCCCTCAGAGC
GAPDH	F	CACTCACGGCAAAATTCACCGCA
	R	GACTCCACGACATACTCAGCAC

F: Forward, R: Reverse.

2.8. Statistical analysis


PCR and Western blot analyses were performed three independent experiments with each experimental group. Band intensity was determined by ImageJ. For MTT, DPPH, and ABTS assays, two separate experiments were carried out. All data in this study are presented as mean ± standard deviation (SD) obtained from each experiment. Mann-Whitney test was used to evaluate

significance of each data. Statistical significance was judged with a *p* value less than 0.05.


3. Results

3.1. Anti-aging activity of SYR in HaCaT cells

Regardless of intrinsic or extrinsic aging, ROS functions a significant role in the skin aging process, and the generated ROS activate MMPs to accelerate skin aging [39–41]. Therefore, anti-aging activity of SYR was investigated by measuring changes of MMP-2 and -9 in hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)-treated HaCaT cells. Gene expression and protein levels of MMP-2 were found to be meaningfully enhanced in response to H<sub>2</sub>O<sub>2</sub> (Fig. 1B and D). MMP-9 expression also was increased in H<sub>2</sub>O<sub>2</sub>-stimulated cells, although to a lesser extent than MMP-2 (Fig. 1C and D). SYR (100 and 200 μg/mL) reduced the MMP-2 and MMP-9 mRNA levels that were elevated by H<sub>2</sub>O<sub>2</sub> (Fig. 1B and C). Consistently, protein levels of MMP-2 and MMP-9 was downregulated in H<sub>2</sub>O<sub>2</sub>-treated cells



**Fig. 2.** Antioxidant activity of SYR. (A) DPPH and SYR (0–200 µg/mL) were incubated together at 37 °C for 30 min. Absorbance at 517 nm was measured by spectrophotometry. (B) SYR (0–200 µg/mL) was mixed with ABTS in the dark at 37 °C for 30 min. Absorbance at 730 nm was observed by spectrophotometry. \**p* < 0.05 and \*\**p* < 0.01 compared with the control group.



**Fig. 3.** Effect of SYR on autophagy activation. (A) SYR (100 and 200 µg/mL)-treated HaCaT cells were incubated with H<sub>2</sub>O<sub>2</sub> for 12 h. Then, the total level of LC3B was determined by Western blotting. β-actin was used as a loading control. (B) HaCaT cells were treated with SYR (50, 100, and 200 µg/mL) for 24 h, and then the total levels of LC3B and β-actin were determined by immunoblotting analysis. (C and D) HaCaT cells were transfected with a plasmid expressing GFP-LC3B and incubated with 200 µg/mL SYR and H<sub>2</sub>O<sub>2</sub> for 12 h (C). HaCaT cells were transfected with plasmid GFP-LC3B and incubated with 200 µg/mL SYR for 24 h (D). Confocal microscopy images were observed using a laser-scanning confocal microscope (Zeiss LSM 710 META). #*p* < 0.05 and ##*p* < 0.01 compared with the normal group; \**p* < 0.05 and \*\**p* < 0.01 compared with the control group.

(Fig. 1D). SYR at a concentration of 200  $\mu\text{g}/\text{mL}$  reduced MMP-2 and MMP-9 expression even in the absence of  $\text{H}_2\text{O}_2$  stimulation (Fig. 1E). Cell viability was not affected by SYR (Fig. 1F), indicating that inhibition of MMP-2 and MMP-9 was not due to cytotoxicity.

### 3.2. In vitro antioxidant activity of SYR

Antioxidant activity of SYR was analyzed by DPPH and ABTS assays, which are broadly used methods for determining antioxidant activity of a wide range of plant components. The DPPH radical scavenging activity was dose-dependently enhanced with treatment of SYR compared with the blank group, and the lowest half-maximal effective concentration ( $\text{EC}_{50}$ ) was 10.77  $\mu\text{g}/\text{mL}$  (Fig. 2A). SYR also significantly increased the ABTS radical scavenging activity, and the  $\text{EC}_{50}$  against ABTS radicals was 10.35  $\mu\text{g}/\text{mL}$  (Fig. 2B). In addition, SYR had a radical scavenging activity similar to that of ascorbic acid (250  $\mu\text{M}$ ), a strong antioxidant, indicating that SYR has potent antioxidant activity.

### 3.3. Activation of autophagy by SYR in HaCaT cells

To investigate whether SYR is participated in autophagy regulation, we detected the autophagy marker LC3B using Western blotting and confocal microscopy. SYR (100 and 200  $\mu\text{g}/\text{mL}$ ) significantly increased LC3B expression in  $\text{H}_2\text{O}_2$ -treated HaCaT cells, and SYR (50, 100, and 200  $\mu\text{g}/\text{mL}$ ) did not affect LC3B expression in HaCaT cells without  $\text{H}_2\text{O}_2$  (Fig. 3A and B). In parallel, exogenous LC3B expression was upregulated by SYR (200  $\mu\text{g}/\text{mL}$ ) in  $\text{H}_2\text{O}_2$ -stimulated HaCaT cells but not in  $\text{H}_2\text{O}_2$ -free HaCaT cells (Fig. 3C and D).

### 3.4. The association between anti-aging activity of SYR and autophagy activation

The relevance of autophagy activation to anti-aging was assessed in recovery experiments with 3-MA, an autophagy


inhibitor. SYR (200  $\mu\text{g}/\text{mL}$ ) inhibited the MMP-2 expression elevated by  $\text{H}_2\text{O}_2$ , but the decreased MMP-2 level was restored in the 3-MA-treated group (Fig. 4A). However, the decreased MMP-2 expression by SYR was not affected by 3-MA in HaCaT cells without  $\text{H}_2\text{O}_2$  (Fig. 4B).

## 4. Discussion

As the amount of UV light increases due to the recent deterioration of the ecological environment, methods to prevent or delay skin aging are receiving increased attention [42]. In addition, there is growing demand for natural, herbal cosmetics because of their weaker or absent side effects [43]. In this study, we investigated the feasibility of using SYR as an anti-aging material. Since degradation of the epidermal and dermal extracellular matrix during skin aging promotes visible signs such as wrinkle formation, the anti-aging activity of SYR was assessed in terms of MMP-2 and MMP-9 regulation. SYR (200  $\mu\text{g}/\text{mL}$ ) inhibited the expression of MMP-2 and -9 at mRNA and protein levels regardless of the presence of  $\text{H}_2\text{O}_2$  in human keratinocyte HaCaT cells (Fig. 1B–E), which indicates SYR has skin anti-aging activity.

Antioxidants such as ascorbic acid, tocopherols, and polyphenols increase resistance to oxidative stress, preventing and improving skin aging [44]. Most plant compounds have antioxidant activity as part of their physiological adaptation to a highly oxidized environment [45], and these compounds exert anti-aging activity via antioxidant properties. SYR also exhibited strong antioxidant activity similar in potency to that of ascorbic acid (Fig. 2A and B). This suggests the antioxidant activity of SYR is an effective mechanism for preventing aging.

In addition, SYR significantly activated autophagy in  $\text{H}_2\text{O}_2$ -treated HaCaT cells (Fig. 3A and C). While the relevance of autophagy to the aging process is not yet fully understood, we directly identified the relationship between the anti-aging activity of SYR and autophagy. The inhibitory activity of SYR on MMP-2 was reduced under conditions in which autophagy activity was



**Fig. 4.** Relevance of autophagy activation to the anti-aging activity of SYR. (A and B) HaCaT cells were incubated with  $\text{H}_2\text{O}_2$  (500  $\mu\text{M}$ ) in the presence or absence of SYR (200  $\mu\text{g}/\text{mL}$ ) or 3-MA (5 mM) for 24 h (A). HaCaT cells were treated with SYR (200  $\mu\text{g}/\text{mL}$ ) in the presence or absence of 3-MA (5 mM) for 24 h (B). The expression of MMP-2 and  $\beta$ -actin was determined by Western blotting. # $p < 0.05$  and ## $p < 0.01$  compared with the normal group; \* $p < 0.05$  and \*\* $p < 0.01$  compared with the control group; ns, not significant.




Fig. 5. Schematic summary of the anti-aging effects of SYR and its mechanisms.

suppressed by 3-MA (Fig. 4A), suggesting that autophagy activation by SYR is another important mechanism for its anti-aging activity. Autophagy generally is thought to delay aging by maintaining homeostasis through recycling mechanisms, but the role of autophagy has been expanding. For example, autophagy suppresses NF- $\kappa$ B signaling in the inflammatory response [46]. Given that SYR inhibits MMP-2 and -9 expression at the transcription level, and that NF- $\kappa$ B is an essential transcription factor for MMP expression (Fig. 1B–D) [47,48], autophagy might exert anti-aging effects by regulating transcription factors such as NF- $\kappa$ B. Interestingly, SYR did not activate autophagy in cells without H<sub>2</sub>O<sub>2</sub>, and the inhibitory effect of SYR against MMP-2 was not affected by 3-MA. These results imply that the anti-aging mechanism of SYR could vary depending on the stimulant.

## 5. Conclusion

Collectively, in this study, we looked over the preventive activity of SYR on skin aging. SYR showed antioxidant activity and up-regulated autophagy activity in H<sub>2</sub>O<sub>2</sub>-stimulated HaCaT cells, thereby decreasing the mRNA expression of MMP-9 and MMP-2 in relation to skin aging (Fig. 5). Therefore, we propose SYR as a novel active ingredient for anti-aging cosmetics.

## Declaration of competing interest

The authors have no conflicts of interest to declare.

## Acknowledgments

This research was funded by the Basic Science Research Program through the National Research Foundation of Korea (NRF), the Ministry of Science and ICT, Republic of Korea (2017R1A6A1A03015642) and by AmorePacific Co. (Yongin, Republic of Korea).

## References

- [1] Mecham RP. Overview of extracellular matrix. *Curr Protoc Cell Biol* 2012. <https://doi.org/10.1002/0471143030.cb1001s00> [Chapter 10]:Unit 10 11.
- [2] Craven NM, Watson RE, Jones CJ, Shuttleworth CA, KIELTY CM, Griffiths CE. Clinical features of photodamaged human skin are associated with a reduction in collagen VII. *Br J Dermatol* 1997;137:344–50.
- [3] Talwar HS, Griffiths CE, Fisher GJ, Hamilton TA, Voorhees JJ. Reduced type I and type III procollagens in photodamaged adult human skin. *J Invest Dermatol* 1995;105:285–90.
- [4] Kohl E, Steinbauer J, Landthaler M, Szeimies RM. Skin ageing. *J Eur Acad Dermatol Venereol* 2011;25:873–84.
- [5] Pittayapruek P, Meephanan J, Prapapan O, Komine M, Ohtsuki M. Role of matrix metalloproteinases in photoaging and photocarcinogenesis. *Int J Mol Sci* 2016;17.
- [6] Salo T, Turpeenniemi-Hujanen T, Tryggvason K. Tumor-promoting phorbol esters and cell proliferation stimulate secretion of basement membrane (type IV) collagen-degrading metalloproteinase by human fibroblasts. *J Biol Chem* 1985;260:8526–31.
- [7] Aimes RT, Quigley JP. Matrix metalloproteinase-2 is an interstitial collagenase. Inhibitor-free enzyme catalyzes the cleavage of collagen fibrils and soluble native type I collagen generating the specific 3/4- and 1/4-length fragments. *J Biol Chem* 1995;270:5872–6.
- [8] Quan T, Little E, Quan H, Qin Z, Voorhees JJ, Fisher GJ. Elevated matrix metalloproteinases and collagen fragmentation in photodamaged human skin: impact of altered extracellular matrix microenvironment on dermal fibroblast function. *J Invest Dermatol* 2013;133:1362–6.
- [9] Qin Z, Balimunkwe RM, Quan T. Age-related reduction of dermal fibroblast size upregulates multiple matrix metalloproteinases as observed in aged human skin *in vivo*. *Br J Dermatol* 2017;177:1337–48.
- [10] Mizushima N, Komatsu M. Autophagy: renovation of cells and tissues. *Cell* 2011;147:728–41.
- [11] Eckhart L, Tschachler E, Gruber F. Autophagic control of skin aging. *Front Cell Dev Biol* 2019;7:143.
- [12] Galluzzi L, Green DR. Autophagy-independent functions of the autophagy machinery. *Cell* 2019;177:1682–99.
- [13] Klionsky DJ, Emr SD. Autophagy as a regulated pathway of cellular degradation. *Science* 2000;290:1717–21.
- [14] Berg TO, Fengsrud M, Stromhaug PE, Berg T, Seglen PO. Isolation and characterization of rat liver amphisomes. Evidence for fusion of autophagosomes with both early and late endosomes. *J Biol Chem* 1998;273:21883–92.
- [15] Fader CM, Sanchez D, Furlan M, Colombo MI. Induction of autophagy promotes fusion of multivesicular bodies with autophagic vacuoles in k562 cells. *Traffic* 2008;9:230–50.
- [16] Goldman SJ, Zhang Y, Jin S. Autophagic degradation of mitochondria in white adipose tissue differentiation. *Antioxidants Redox Signal* 2011;14:1971–8.

- [17] Levine B, Deretic V. Unveiling the roles of autophagy in innate and adaptive immunity. *Nat Rev Immunol* 2007;7:767–77.
- [18] Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008;132:27–42.
- [19] Murase D, Kusaka-Kikushima A, Hachiya A, Fullenkamp R, Stepp A, Imai A, Ueno M, Kawabata K, Takahashi Y, Hase T, Ohuchi A, Nakamura S, Yoshimori T. Autophagy declines with premature skin aging resulting in dynamic alterations in skin pigmentation and epidermal differentiation. *Int J Mol Sci* 2020;21.
- [20] Ratan ZA, Youn SH, Kwak YS, Han CK, Haidere MF, Kim JK, Min H, Jung YJ, Hosseinzadeh H, Hyun SH, Cho JY. Adaptogenic effects of *Panax ginseng* on modulation of immune functions. *J Ginseng Res* 2021;45:32–40.
- [21] Cabral de Oliveira AC, Perez AC, Merino G, Prieto JG, Alvarez AI. Protective effects of *Panax ginseng* on muscle injury and inflammation after eccentric exercise. *Comp Biochem Physiol C Toxicol Pharmacol* 2001;130:369–77.
- [22] Kim JH, Yi YS, Kim MY, Cho JY. Role of ginsenosides, the main active components of *Panax ginseng*, in inflammatory responses and diseases. *J Ginseng Res* 2017;41:435–43.
- [23] Kim HG, Cho JH, Yoo SR, Lee JS, Han JM, Lee NH, Ahn YC, Son CG. Antifatigue effects of *Panax ginseng* C.A. Meyer: a randomised, double-blind, placebo-controlled trial. *PLoS One* 2013;8:e61271.
- [24] Shin HR, Kim JY, Yun TK, Morgan G, Vainio H. The cancer-preventive potential of *Panax ginseng*: a review of human and experimental evidence. *Cancer Causes Control* 2000;11:565–76.
- [25] Li C, Cai J, Geng J, Li Y, Wang Z, Li R. Purification, characterization and anticancer activity of a polysaccharide from *Panax ginseng*. *Int J Biol Macromol* 2012;51:968–73.
- [26] Vuksan V, Sung MK, Sievenpiper JL, Stavro PM, Jenkins AL, Di Buono M, Lee KS, Leiter LA, Nam KY, Arnason JT, Choi M, Naeem A. Korean red ginseng (*Panax ginseng*) improves glucose and insulin regulation in well-controlled, type 2 diabetes: results of a randomized, double-blind, placebo-controlled study of efficacy and safety. *Nutr Metabol Cardiovasc Dis* 2008;18:46–56.
- [27] Baeg IH, So SH. The world ginseng market and the ginseng (Korea). *J Ginseng Res* 2013;37:1–7.
- [28] Choi SR, Lee MY, Reddy CK, Lee SJ, Lee CH. Evaluation of metabolite profiles of ginseng berry pomace obtained after different pressure treatments and their correlation with the antioxidant activity. *Molecules* 2021;26.
- [29] Bajpai VK, Alam MB, Quan KT, Ju MK, Majumder R, Shukla S, Huh YS, Na M, Lee SH, Han YK. Attenuation of inflammatory responses by (+)-syringaresinol via MAP-Kinase-mediated suppression of NF-kappaB signaling *in vitro* and *in vivo*. *Sci Rep* 2018;8:9216.
- [30] Park BY, Oh SR, Ahn KS, Kwon OK, Lee HK. (-)-Syringaresinol inhibits proliferation of human promyelocytic HL-60 leukemia cells via G1 arrest and apoptosis. *Int Immunopharmacol* 2008;8:967–73.
- [31] Kim J, Toda T, Watanabe K, Shibuya S, Ozawa Y, Izuo N, Cho S, Seo DB, Yokote K, Shimizu T. Syringaresinol reverses age-related skin atrophy by suppressing FoxO3a-mediated matrix metalloproteinase-2 activation in Copper/Zinc superoxide dismutase-deficient Mice. *J Invest Dermatol* 2019;139:648–55.
- [32] Oh JH, Joo YH, Karadeniz F, Ko J, Kong CS. Syringaresinol inhibits UVA-induced MMP-1 expression by suppression of MAPK/AP-1 signaling in HaCaT keratinocytes and human dermal fibroblasts. *Int J Mol Sci* 2020;21.
- [33] Jeong D, Park SH, Kim MH, Lee S, Cho YK, Kim YA, Park BJ, Lee J, Kang H, Cho JY. Anti-melanogenic effects of ethanol extracts of the leaves and roots of *Patrinia villosa* (Thunb.) Juss through their inhibition of CREB and induction of ERK and autophagy. *Molecules* 2020;25.
- [34] Park HW, Cho SY, Kim HH, Yun BS, Kim JU, Lee SJ, Park J. Enantioselective induction of SIRT1 gene by syringaresinol from *Panax ginseng* berry and *Acanthopanax senticosus* Harms stem. *Bioorg Med Chem Lett* 2015;25:307–9.
- [35] Lorz LR, Kim D, Kim MY, Cho JY. *Panax ginseng*-derived fraction BIOGF1K reduces atopic dermatitis responses via suppression of mitogen-activated protein kinase signaling pathway. *J Ginseng Res* 2020;44:453–60.
- [36] Lee JO, Kim JH, Kim S, Kim MY, Hong YH, Kim HG, Cho JY. Gastroprotective effects of the nonsaponin fraction of Korean Red Ginseng through cyclooxygenase-1 upregulation. *J Ginseng Res* 2020;44:655–63.
- [37] Lee JO, Hwang SH, Shen T, Kim JH, You L, Hu W, Cho JY. Enhancement of skin barrier and hydration-related molecules by protopanaxatriol in human keratinocytes. *J Ginseng Res* 2021;45:354–60.
- [38] Hong YH, Kim D, Nam G, Yoo S, Han SY, Jeong SG, Kim E, Jeong D, Yoon K, Kim S, Park J, Cho JY. Photoaging protective effects of BIOGF1K, a compound-K-rich fraction prepared from *Panax ginseng*. *J Ginseng Res* 2018;42:81–9.
- [39] Pillai S, Oresajo C, Hayward J. Ultraviolet radiation and skin aging: roles of reactive oxygen species, inflammation and protease activation, and strategies for prevention of inflammation-induced matrix degradation - a review. *Int J Cosmet Sci* 2005;27:17–34.
- [40] Poljsak B, Dahmane RG, Godic A. Intrinsic skin aging: the role of oxidative stress. *Acta Dermatovenerol Alpina Pannonica Adriatica* 2012;21:33–6.
- [41] Kammeyer A, Luiten RM. Oxidation events and skin aging. *Ageing Res Rev* 2015;21:16–29.
- [42] Gu Y, Han J, Jiang C, Zhang Y. Biomarkers, oxidative stress and autophagy in skin aging. *Ageing Res Rev* 2020;59:101036.
- [43] Shital GS, Aishwarya JJ, Amar ZG, Ravindra SB. A Review on oils used in herbal cosmetics. *Res J Top Cosmet Sci* 2013;4:61.
- [44] Masaki H. Role of antioxidants in the skin: anti-aging effects. *J Dermatol Sci* 2010;58:85–90.
- [45] Cavinato M, Waltnerberger B, Baraldo G, Grade CVC, Stuppner H, Jansen-Durr P. Plant extracts and natural compounds used against UVB-induced photoaging. *Biogerontology* 2017;18:499–516.
- [46] Peng X, Wang Y, Li H, Fan J, Shen J, Yu X, Zhou Y, Mao H. ATG5-mediated autophagy suppresses NF-kappaB signaling to limit epithelial inflammatory response to kidney injury. *Cell Death Dis* 2019;10:253.
- [47] Ukaji T, Lin Y, Okada S, Umezawa K. Inhibition of MMP-2-mediated cellular invasion by NF-kappaB inhibitor DHMEQ in 3D culture of breast carcinoma MDA-MB-231 cells: a model for early phase of metastasis. *Biochem Biophys Res Commun* 2017;485:76–81.
- [48] Bond M, Chase AJ, Baker AH, Newby AC. Inhibition of transcription factor NF-kappaB reduces matrix metalloproteinase-1, -3 and -9 production by vascular smooth muscle cells. *Cardiovasc Res* 2001;50:556–65.