

Editorial



Magnetic Resonance Imaging-based Right Ventricular Strain Evaluation

Jongmin Lee , MD, PhD

Department of Radiology, Kyungpook National University, School of Medicine, Daegu, Korea



► See the article “Right Ventricular Strain Is Associated With Increased Length of Stay After Tetralogy of Fallot Repair” in volume 30 on page 50.

Received: Aug 17, 2021
Revised: Oct 4, 2021
Accepted: Oct 14, 2021
Published online: Nov 4, 2021

Address for Correspondence:

Jongmin Lee, MD, PhD

Department of Radiology, Kyungpook National University, School of Medicine, 130, Dongdeok-ro, Jung-gu, Daegu 41944, Korea.
Email: jonglee@knu.ac.kr

Copyright © 2022 Korean Society of Echocardiography

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Jongmin Lee 
<https://orcid.org/0000-0002-4163-913X>

Conflict of Interest

The author has no financial conflicts of interest.

Cardiac integrity comprises normal morphology and pumping function of the heart's 4 chambers. Evaluation of the heart aims to check the chambers' shape and function. Since the major role of the heart is to pump blood in the circulatory system, cardiac function is typically prioritized over morphology. Echocardiography is an important modality for assessing cardiac function and morphology, although it is more proficient for hemodynamic evaluation. Computed tomography (CT) and magnetic resonance imaging (MRI) joining cardiac imaging has strengthened the evaluation of cardiac morphology and facilitated the objective assessment of cardiac chamber function.

The function of each chamber is influenced by preload, afterload, and its own wall function. If the conditions in and around the chambers provoke dysfunction, the cardiac chambers will begin to adjust their morphology and motion, resulting in remodeling. Chamber volumes and their variations along a cardiac cycle can be measured so that the global function of a single chamber can be estimated, even during the remodeling process. For regional function evaluation, chamber wall motion can be evaluated qualitatively in routine practice. However, detecting chamber dysfunction as early as possible is important, and more sensitive imaging markers have been reported.

Unlike the left ventricle (LV), the right ventricle (RV) has a more complex shape without fixed reference points since it partly wraps around the LV and has a thinner and more compliant myocardium. In healthy subjects, the wall thickness of the RV varies between 2 and 5 mm.¹⁾ To evaluate RV volume and its variability, complete three-dimensional measurements along a compact timeline are critical. The magnetic resonance (MR) cine image could provide a solution in this context. As comparing with MR, the echocardiography could evaluate right ventricle in a simpler and more convenient way. Already several echocardiographic imaging markers for right ventricular function are popular. The tricuspid annular plane systolic excursion, the time-velocity integral, the fractional area change, and the Tei index of the RV feasibly measure global RV function.²⁾ However, these markers cannot provide regional RV function, and they cannot differentiate active RV motion from passive entrainment by the neighboring LV.

The global and regional strains along promised planes were introduced to evaluate active myocardial motion from the standpoint of the myocardium itself. Since echocardiography

depicts myocardial motion in real-time and speckle tracking is relatively easier, the echocardiography is an essential modality for strain imaging.³⁾ Strain presents as myocardial deformation during the chamber motion. Despite the handicaps of RV imaging, echocardiographic strain has been successfully and widely applied on RV function evaluation in diverse pathological situations, including heart failure, ischemic heart disease, valvular heart disease, congenital heart disease, cardiomyopathy, and pulmonary hypertension.²⁾

As a gold standard modality for the evaluation of chamber function, cardiac MRI provides three-dimensional information on all 4 chambers without any imaging window limitations. Several techniques were introduced for strain evaluation based on cine MRI data. Initially, a conventional tagged MRI technique was used for strain imaging despite insufficient spatial resolution. Soon, a technical advancement occurred by introducing feature tracking imaging (FTI) operable in an independent post-processing console. FTI has been the most popular option for strain MRI thus far, although features at the endocardial and epicardial borders are the main targets. Actually, measuring strain in the mid-myocardial layer by MR FTI is challenging due to a homogenous mid-myocardial texture without a target feature.⁴⁾ Recently an improved tagged MRI technique, strain-encoded (SENC) MRI, was introduced, which showed superior performance to the MR FTI technique.⁵⁾

Since strain MRI is a relatively younger methodology than echocardiography, its clinical application is not as wide as other methodologies. In acute myocardial infarction cases, the longitudinal strain evaluated by MR FTI reliably predicted clinical outcomes during follow-up.⁶⁾ Measuring the difference between epicardial and endocardial circumferential strain based on MR FTI permitted early detection of systolic dysfunction in patients with hypertrophic cardiomyopathy, even at the pre-clinical and pre-dysmorphic stages.⁴⁾ In cases of arrhythmogenic right ventricular dystrophy, MR FTI successfully and reproducibly evaluated regional strain in the right ventricular myocardium.⁷⁾ In patients with pulmonary hypertension, the SENC MR technique revealed decreased longitudinal and circumferential strains compared with those in healthy subjects.⁸⁾ However, due to the limited MRI quality of the RV in pediatric patients, strain MRI for congenital heart diseases is still unrecognized.

Tetralogy of Fallot (TOF) is well known as the most common cyanotic congenital heart disease. Due to the high incidence of TOF, initial palliative and corrective surgical treatments are well established. In parallel, postoperative follow-up and prognosis prediction becomes more interesting and meticulous. By correcting the TOF, the RV is exposed to a new environment, from high afterload pressure to volume overload by inevitable pulmonary regurgitation. From the moment of correction, the RV remodels itself over time.⁹⁾ Together with the possible improvement of RV hypertrophy, RV dilatation with hindered systolic function is a common postoperative feature. During this remodeling process, medical surveillance is required for pulmonary valvular replacement at a suitable timing. The decision regarding the appropriate time for this correction depends on the patient's clinical expression, RV volume and function, and systemic venous congestion. Meanwhile, RV strain provides more sensitive and accurate information for detection of noticeable stress of the RV as early as possible. Echocardiographic strain evaluation was successfully applied on postoperative TOF cases despite a limited imaging window.¹⁰⁾ Strain MRI will allow more feasible evaluation of RV strain for patients with surgically corrected TOF. In this context, the report by Srinivasan et al.¹¹⁾ sought to improve the role of strain MRI for pediatric patients with congenital heart disease, including surgically corrected TOF.

Using MRI, Srinivasan et al.¹¹⁾ focused on RV myocardial strain immediately after TOF repair for early detection of adverse effects on future RV function. In addition, based on their previous work, they performed strain MRI exams without anesthesia in infants. In this study, early postoperative RV deformation was observed as a decrease in mean RV peak longitudinal strain (median [interquartile range], 16.51 [-19.49, -9.95]). In addition, RV peak radial strain was always available and was significantly associated with an extended hospital stay. These findings were detected prior to the expression of conventional markers, such as changes in RV ejection fraction. A series of postoperative RV strain adaptation processes could not be disclosed in this study; however, MR-based RV strain was adequate as an early prognostic indicator for surgically-corrected TOF patients.

REFERENCES

1. Haddad F, Hunt SA, Rosenthal DN, Murphy DJ. Right ventricular function in cardiovascular disease, part I: anatomy, physiology, aging, and functional assessment of the right ventricle. *Circulation* 2008;117:1436-48. [PUBMED](#) | [CROSSREF](#)
2. Ayach B, Fine NM, Rudski LG. Right ventricular strain: measurement and clinical application. *Curr Opin Cardiol* 2018;33:486-92. [PUBMED](#) | [CROSSREF](#)
3. Aneq MÁ, Engvall J, Brudin L, Nylander E. Evaluation of right and left ventricular function using speckle tracking echocardiography in patients with arrhythmogenic right ventricular cardiomyopathy and their first degree relatives. *Cardiovasc Ultrasound* 2012;10:37. [PUBMED](#) | [CROSSREF](#)
4. Vigneault DM, Yang E, Jensen PJ, et al. Left ventricular strain is abnormal in preclinical and overt hypertrophic cardiomyopathy: cardiac MR feature tracking. *Radiology* 2019;290:640-8. [PUBMED](#) | [CROSSREF](#)
5. Ohyama Y, Ambale-Venkatesh B, Chamera E, et al. Comparison of strain measurement from multimodality tissue tracking with strain-encoding MRI and harmonic phase MRI in pulmonary hypertension. *Int J Cardiol* 2015;182:342-8. [PUBMED](#) | [CROSSREF](#)
6. Gavara J, Rodriguez-Palomares JF, Valente F, et al. Prognostic value of strain by tissue tracking cardiac magnetic resonance after ST-segment elevation myocardial infarction. *JACC Cardiovasc Imaging* 2018;11:1448-57. [PUBMED](#) | [CROSSREF](#)
7. Vigneault DM, te Riele AS, James CA, et al. Right ventricular strain by MR quantitatively identifies regional dysfunction in patients with arrhythmogenic right ventricular cardiomyopathy. *J Magn Reson Imaging* 2016;43:1132-9. [PUBMED](#) | [CROSSREF](#)
8. Shehata ML, Basha TA, Tantawy WH, et al. Real-time single-heartbeat fast strain-encoded imaging of right ventricular regional function: normal versus chronic pulmonary hypertension. *Magn Reson Med* 2010;64:98-106. [PUBMED](#) | [CROSSREF](#)
9. DiLorenzo MP, Elci OU, Wang Y, et al. Longitudinal changes in right ventricular function in tetralogy of Fallot in the initial years after surgical repair. *J Am Soc Echocardiogr* 2018;31:816-21. [PUBMED](#) | [CROSSREF](#)
10. Sabate Rotes A, Bonnicksen CR, Reece CL, et al. Long-term follow-up in repaired tetralogy of Fallot: can deformation imaging help identify optimal timing of pulmonary valve replacement? *J Am Soc Echocardiogr* 2014;27:1305-10. [PUBMED](#) | [CROSSREF](#)
11. Srinivasan R, Faerber JA, DeCost G, et al. Right ventricular strain is associated with increased length of stay after tetralogy of Fallot repair. *J Cardiovasc Imaging* 2022;30:50-8. [CROSSREF](#)