



Corrigendum: Marginal bone loss around crestal or subcrestal dental implants: prospective clinical study

Naser Sargolzaie¹, Hosein Hoseini Zarch², Hamidreza Arab¹, Tahereh Koohestani³, Mahdiye Fasihi Ramandi⁴

¹Dental Research Center, School of Dentistry, Mashhad University of Medical Sciences,

²Department of Oral and Maxillofacial Radiology, School of Dentistry and Dental Research Center, Mashhad University of Medical Sciences,

³School of Dentistry, Mashhad University of Medical Sciences,

⁴Department of Periodontics, School of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran

J Korean Assoc Oral Maxillofac Surg 2022;48:159-166.

<https://doi.org/10.5125/jkaoms.2022.48.3.159>

This correction is being published to correct the column headings of Table 6.

The authors apologize for any inconvenience this may have caused.

Before correction

Table 6. Association of age and bone level changes

	Mesial marginal bone loss		Distal marginal bone loss	
	3 mo	6 mo	3 mo	6 mo
Spearman correlation coefficient	-0.087	-0.321	-0.051	-0.431
P-value	0.552	0.025	0.729	0.002

Naser Sargolzaie et al: Marginal bone loss around crestal or subcrestal dental implants: prospective clinical study. J Korean Assoc Oral Maxillofac Surg 2022

After correction

Table 6. Association of age and bone level changes

	3 mo (mesial side)	3 mo (distal side)	6 mo (mesial side)	6 mo (distal side)
Spearman correlation coefficient	-0.087	-0.321	-0.051	-0.431
P-value	0.552	0.025	0.729	0.002

Naser Sargolzaie et al: Marginal bone loss around crestal or subcrestal dental implants: prospective clinical study. J Korean Assoc Oral Maxillofac Surg 2022