



Understanding of oral potentially malignant disorders and epithelial dysplasia among oral and maxillofacial surgeons

So-Young Choi, DDS, PhD

Section Editor of JKAOMS

Department of Oral and Maxillofacial Surgery, School of Dentistry, Kyungpook National University, Daegu, Korea

Oral mucosal diseases have various etiologic factors; however, the characteristics of each disease are similar. Moreover, even the same disease can present as various clinical symptoms as it progresses, making treatment difficult. Oral mucosal diseases can be diagnosed and treated not only in dentistry but also in otolaryngology, internal medicine, and pediatrics, which can confuse patients. Oral and maxillofacial surgeons tend to be interested in diseases that require surgical treatment and less interested in oral mucosal diseases as these are often managed at the oral medicine department through biopsies or medications.

However, as society ages, the number of patients visiting hospitals for oral mucosal diseases has been increasing recently. Furthermore, the interest in precancerous lesions increasing along with as interest in oral cancer. The term “potentially malignant diseases of the oral cavity” has been recently used to integrate precancerous lesions and precancerous conditions with a risk of malignant transformation.

As stated by the World Health Organization, oral potentially malignant disorders (OPMDs) represent “a heterogeneous group of clinically defined conditions associated with a variable risk of progression to oral squamous cell carcinoma”^{1,2}. OPMDs include leukoplakia, proliferative verrucous leukoplakia, erythroplakia, erythroleukoplakia, oral submucous fibrosis, actinic cheilitis, palatal lesions associated with reverse smoking, smokeless tobacco keratosis, oral lichenoid patho-

ses, and heritable conditions with cancer predisposition, such as Fanconi anemia, congenital dyskeratosis, and xeroderma pigmentosum³. A meta-analysis of the literature revealed that the overall malignant transformation rate of OPMDs is approximately 8%⁴.

Leukoplakia, erythroplakia, and erythroleukoplakia are clinical terms. These disorders of the oral mucosa are closely associated with the degree of epithelial dysplasia. Dysplasia is observed in some cases of leukoplakia but is consistently observed in cases with erythroplakia and erythroleukoplakia.

The cytological changes in dysplastic epithelial cells include abnormal variations in nuclear size, nuclear shape, cell size, and cell shape; increased nuclear-to-cytoplasmic ratio, nuclear size, and nucleoli number and size; and hyperchromasia. Architectural alterations include irregular epithelial stratification, loss of polarity of basal cells, drop-shaped rete ridges, increased number of mitotic figures, abnormal superficial mitotic figures, premature keratinization in single cells, keratin pearls within rete ridges, and loss of epithelial cell cohesion¹⁻³.

Despite recent advances in biomarker research and molecular profiling for predicting the risk for malignant transformation, investigation on biomarker is still lacking, and further prospective clinical studies are warranted. The standard method for predicting the risk for malignant transformation is histopathologic grading of oral epithelial dysplasia and consideration of clinical factors.

Traditionally, oral epithelial dysplasia can be classified into three grades of severity: mild epithelial dysplasia with alterations limited mainly to the basal and parabasal layers, moderate epithelial dysplasia with dysplastic changes extending to the midportion of the epithelium, and severe epithelial dysplasia with alterations mostly in epithelial thickness³. Carcinoma *in situ* in the oral cavity is considered synonymous with severe dysplasia¹. However, this traditional definition

So-Young Choi

Department of Oral and Maxillofacial Surgery, School of Dentistry, Kyungpook National University, 2177 Dalgubeol-daero, Jung-gu, Daegu 41940, Korea

TEL: +82-53-600-7576

E-mail: dentalchoi@knu.ac.kr

ORCID: <https://orcid.org/0000-0002-2563-3539>

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2023 The Korean Association of Oral and Maxillofacial Surgeons.

does not fully reflect the complexity of oral epithelial dysplasia grading. Alternatively, some authorities advocate the use of a binary grading system, i.e., low-grade versus high-grade dysplasia. This binary system categorizes oral epithelial dysplasia as having a low or high risk for malignant transformation⁵. The cutoff points between low- and high-grade dysplasia are four architectural and five cytological changes, irrespective of the level of dysplasia within the epithelium⁵. Such a two-tier system may have improved reproducibility than the three-tier system. However, outcome studies are still warranted to validate the efficacy of binary scoring before it can be routinely applied clinically¹.

Mehanna et al.⁶ reported that the mean malignant transformation rate for mild/moderate dysplasia was 10% compared with 24% for severe dysplasia/carcinoma *in situ*. Furthermore, the risk of progression to oral cancer significantly increases for all grades of dysplasia⁶. A population-based cohort study reported an overall hazard ratio of 4.9 for mild dysplasia, 6.8 for moderate dysplasia, and 15.8 for severe dysplasia⁷. A recent meta-analysis also revealed a six-fold increase in the odds of malignant transformation in high-risk lesions compared with low-risk lesions in the binary grading of oral epithelial dysplasia⁸.

Oral mucosa with no mild dysplasia and adverse clinical outcomes is often conservatively managed with risk factor modification and periodic clinical reevaluation rather than surgical excision. Some oral mucosa with no or minimal dysplasia may disappear or diminish in size within a few months after the modification. However, if the lesion increases in size over time, additional biopsies may be performed. Moreover, for lesions with mild dysplasia and adverse clinical outcomes, surgical excision and/or short-interval clinical follow-up may be considered^{6,9}. Moderate or severe epithelial dysplasia in the oral mucosa is treated with complete excision and close clinical follow-up. Complete removal can be achieved with equal efficacy by surgical excision, electrocautery, cryosurgery, or laser ablation^{6,9}.

Even after removal or obvious resolution of oral leukoplakia, the reported overall recurrence rates range from 7% to 38%, and the development of additional lesions is common¹⁰. Therefore, long-term postoperative follow-up is crucial.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

References

1. Reibel J, Tilakaratne W, Gale N, Westra W, Hille J, Williams M, et al. Oral potentially malignant disorders & oral epithelial dysplasia. In: El-Naggar AK, Chan JKC, Grandis JR, Takata T, Slootweg PJ, eds. WHO classification of head and neck tumours. 4th ed. International Agency for Research on Cancer; 2017:112-5.
2. Muller S, Tilakaratne WM. Update from the 5th edition of the World Health Organization classification of head and neck tumors: tumours of the oral cavity and mobile tongue. *Head Neck Pathol* 2022;16:54-62. <https://doi.org/10.1007/s12105-021-01402-9>
3. Neville BW, Damm DD, Allen CM, Chi AC. Oral and maxillofacial pathology. 5th ed. Elsevier; 2023:380-1.
4. Iocca O, Sollecito TP, Alawi F, Weinstein GS, Newman JG, De Virgilio A, et al. Potentially malignant disorders of the oral cavity and oral dysplasia: a systematic review and meta-analysis of malignant transformation rate by subtype. *Head Neck* 2020;42:539-55. <https://doi.org/10.1002/hed.26006>
5. Kujan O, Oliver RJ, Khattab A, Roberts SA, Thakker N, Sloan P. Evaluation of a new binary system of grading oral epithelial dysplasia for prediction of malignant transformation. *Oral Oncol* 2006;42:987-93. <https://doi.org/10.1016/j.oraloncology.2005.12.014>
6. Mehanna HM, Rattay T, Smith J, McConkey CC. Treatment and follow-up of oral dysplasia - a systematic review and meta-analysis. *Head Neck* 2009;31:1600-9. <https://doi.org/10.1002/hed.21131>
7. Chaturvedi AK, Udaltsova N, Engels EA, Katz JA, Yanik EL, Katki HA, et al. Oral leukoplakia and risk of progression to oral cancer: a population-based cohort study. *J Natl Cancer Inst* 2020;112:1047-54. <https://doi.org/10.1093/jnci/djz238>
8. Yan F, Reddy PD, Nguyen SA, Chi AC, Neville BW, Day TA. Grading systems of oral cavity pre-malignancy: a systematic review and meta-analysis. *Eur Arch Otorhinolaryngol* 2020;277:2967-76. <https://doi.org/10.1007/s00405-020-06036-1>
9. Awadallah M, Idle M, Patel K, Kademani D. Management update of potentially premalignant oral epithelial lesions. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2018;125:628-36. <https://doi.org/10.1016/j.oooo.2018.03.010>
10. Kuribayashi Y, Tsushima F, Sato M, Morita K, Omura K. Recurrence patterns of oral leukoplakia after curative surgical resection: important factors that predict the risk of recurrence and malignancy. *J Oral Pathol Med* 2012;41:682-8. <https://doi.org/10.1111/j.1600-0714.2012.01167.x>

How to cite this article: Choi SY. Understanding of oral potentially malignant disorders and epithelial dysplasia among oral and maxillofacial surgeons. *J Korean Assoc Oral Maxillofac Surg* 2023;49:309-310. <https://doi.org/10.5125/jkaoms.2023.49.6.309>