



Research Article

Antiviral effects of Korean Red Ginseng on human coronavirus OC43

Chi Hwan Jeong, Jisu Kim, Bo Kyeong Kim, Kang Bin Dan, Hyeyoung Min*

College of Pharmacy, Chung-Ang University, Seoul, Republic of Korea



ARTICLE INFO

Article history:

Received 3 August 2022

Received in revised form

15 September 2022

Accepted 29 September 2022

Available online 5 October 2022

Keywords:

aAntiviral
coronavirus
ginsenoside
Korea Red Ginseng

ABSTRACT

Background: *Panax ginseng Meyer* is a medicinal plant well-known for its antiviral activities against various viruses, but its antiviral effect on coronavirus has not yet been studied thoroughly. The antiviral activity of Korean Red Ginseng (KRG) and ten ginsenosides against Human coronavirus OC43 (HCoV-OC43) was investigated in vitro.

Methods: The antiviral response and mechanism of action of KRG extract and ginsenoside Rc, Re, Rf, Rg1, Rg2-20 (R) and -20 (S), Rg3-20 (R) and -20 (S), and Rh2-20 (R) and -20 (S), against the human coronavirus strain OC43 were investigated by using plaque assay, time of addition assay, real-time PCR, and FACS analysis.

Results: Virus plaque formation was reduced in KRG extract-treated and HCoV-OC43-infected HCT-8 cells. KRG extract decreased the viral proteins (Nucleocapsid protein and Spike protein) and mRNA (N and M gene) expression, while increased the expression of interferon genes.

Conclusion: KRG extract exhibits antiviral activity by enhancing the expression of interferons and can be used in treating infections caused by HCoV-OC43.

© 2022 The Korean Society of Ginseng. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Severe acute respiratory syndrome-associated coronavirus-2 (SARS-CoV-2) is a highly pathogenic coronavirus and was responsible for causing the coronavirus disease in 2019 (COVID-19) which has been declared a pandemic worldwide. A few therapies, including vaccines, are available for COVID-19 treatment. Considering the outbreak of SARS-CoV-2, developing an effective treatment for the highly pathogenic virus is urgent. It is also essential to find broad-spectrum antiviral agents that effectively prevent or cure viral infection caused by the current reigning strain of coronavirus (CoV) and future-emerging strains of CoVs. However, manipulating highly pathogenic viruses, including SARS-CoV-2, for research must be performed under high biosafety level conditions. Accordingly, it is crucial to find alternatives to overcome the limitations of research accessibility [1].

Coronaviruses are enveloped viruses with a diameter of 120–160 nm, and the envelope derived from intracellular membranes contains multiple club-shaped spikes of a crown [2,3]. They have a capped, non-segmented, and linear, positive-sense single-stranded

RNA genome of approximately 30 kb [2–4]. The genus Coronavirus belongs to the family Coronaviridae [5]. Based on genetic similarities, the coronavirinae subfamily comprises four genera: alpha, beta, gamma, and delta coronaviruses. Human coronaviruses (HCoV) are limited to the alpha and beta genera [6]. Alphacoronaviruses include HCoV-NL63 and HCoV-229E, while Betacoronaviruses contain HCoV-HKU1, SARS-CoV, MERS-CoV, and HCoV-OC43 [4]. HCoVs cause multiple respiratory diseases, and their severity varies from the mild common cold to life-threatening pneumonia [4,7].

HCoV-OC43 belongs to the same viral genus as SARS-CoV and SARS-CoV-2 [8]. HCoV-OC43 has been associated with common cold [9] and mild upper respiratory tract infections [10], and occasionally can cause a severe form of the disease in people with underlying respiratory conditions, infants, and the elderly [11,12]. The genome of HCoV-OC43 encodes several well-conserved motifs that match with SARS-CoV-2, and cross-reactivity between HCoV-OC43 and SARS-CoV-2 is reported, indicating that HCoV-OC43 might be used as a substitute for the study of SARS-CoV-2, which can only be conducted in a BSL-4 facility [13–16].

Panax ginseng Meyer has been shown to possess diverse physiological activities [17,18], and the beneficial effects of Korean Red Ginseng (KRG) in various diseases such as cancer, immune diseases, and neuronal diseases are well known [19–22]. The antiviral properties of *P. ginseng* have also been reported [23], and KRG

* Corresponding author. 84 Heukseokro, Dongjakgu, Seoul, 06974, Republic of Korea.

E-mail address: hymin@cau.ac.kr (H. Min).

extract and ginsenosides have been shown to exert antiviral activity against influenza virus, human immunodeficiency virus, norovirus, herpes simplex virus, and hepatitis B virus [24–29].

Most medicinal effects of ginseng are derived from ginsenosides, which are triterpene glycosides [30]. Ginsenosides can be divided based on the chemical structure of the aglycone part: the protopanaxadiol group (e.g., Rc, Rg3, Rh2) and the protopanaxatriol group (e.g., Re, Rf, Rg1, Rg2) [31]. Although it has been reported that *P. ginseng* possesses antiviral activities, the antiviral property of KRG and ginsenosides against coronavirus has not been elucidated [23–25,27]. Therefore, in this study, the antiviral activity of KRG and ginsenosides against HCoV-OC43 was investigated in vitro.

2. Materials and methods

2.1. Cell line, virus, Korean Red Ginseng extract, and ginsenosides

The human colon cell-8 (HCT-8) cell line was obtained from Korean Cell Line Bank (Seoul, Korea). HCT-8 cells were cultured in RPMI 1640 supplemented with 10% fetal bovine serum (Welgene, Gyeongsan, Gyeong-buk, Korea), 25 mM HEPES (Gibco, Waltham, MA, USA), 100 U/mL penicillin/streptomycin (Welgene). The cells were incubated at 37°C in a 5% CO₂ humidified air environment. HCoV-OC43 was obtained from Korea Bank for Pathogenic Viruses (Seoul, Korea). KRG extract was provided by the Korea Ginseng Corporation (Buyeo, Chung-nam, Korea). For KRG extract preparation, fresh ginseng roots were prepared and processed by steaming and drying to make red ginseng in red ginseng manufacturing factory of Korea Ginseng Corporation. Washed fresh ginseng roots were steamed for 4 hours while slowly raising its temperature from 50 °C to 98 °C and then firstly dried at 60–70 °C for 15 hours. Thereafter, secondary drying process was performed in a closed chamber at 50°C for 5 days to resulting the red ginseng roots. To prepare red ginseng extract, the root was sequentially extracted 7 times at 87°C for 12 hours with distilled water. The extracted water was combined followed by filtering and concentrating process. The KRG extract contains ginsenoside Rg1 (1.74 mg/g), Re (1.88 mg/g), Rf (1.33 mg/g), Rg2 (S) (1.21 mg/g), Rb1 (7.9 mg/g), Rc (2.83 mg/g), Rb2 (2.58 mg/g), Rd (1.04 mg/g), Rg3 (S) (2.06 mg/g), Rg3 (R) (0.93 mg/g), and Rh1 (0.89 mg/g). Ginsenosides Rc, Re, Rf, Rg1, Rg2 -20 (R), and -20 (S), Rg3 -20 (R) and -20 (S), and Rh2 -20 (R) and -20 (S) were purchased from Ambo Institute (Daejeon, Korea). -20 (R) and -20 (S) represent the isomer designation for carbon 20 of each ginsenoside.

2.2. Cell viability assay

HCT-8 cells were seeded in 96-well plates (5×10^4 cells) and treated with 100 µg/mL KRG or 100 µM ginsenoside for 7 days. The cells were then treated with 10 µL MTT solution (Sigma-Aldrich, St. Louis, MO, USA) and incubated at 37°C in 5% CO₂ for 4 h. This step was followed by adding 100 µL of 0.04 N HCl-isopropanol to dissolve the formazan crystals. The cell viability was quantified at an absorbance of 540 nm using an E_{max} microplate reader (Molecular Devices, Sunnyvale, CA, USA)

2.3. Plaque assays

HCT-8 cells were seeded at 5×10^5 cells/well in 12-well plates. After 24 h of culture, virus stock solutions were added to the cells and allowed to incubate for 60 min at 37°C, rocking every 15 min for 1 h. HCT-8 cells were washed once in PBS and overlaid with overlay medium (complete DMEM containing 0.6% methylcellulose). KRG or ginsenoside was added into the overlay medium at 100 µg/mL and 100 µmol/mL, respectively. After 7 days of culture,

the cells were fixed and stained with 2% crystal violet (Sigma-Aldrich) in 20% ethanol overnight. The plaques were counted to determine the antiviral effect of KRG.

2.4. Time-of-addition assays

HCT-8 cells (5×10^5 cells) were seeded in 12-well plates. KRG (25, 50, 100 µg/mL) was added to the HCT-8 cells at 24 h before HCoV-OC43 infection (pre-treatment), during infection (co-treatment), and after removal of the virus (post-treatment). The infected cells were cultured for 7 days at 37°C in 5% CO₂, and a plaque assay was performed.

2.5. Quantitative real-time PCR (qRT-PCR)

HCT-8 cells (1×10^6 cells) were seeded in 6-well plates. They were treated with 25, 50, and 100 µg/mL of KRG or 5 µM of remdesivir (Hunan Hua Teng Pharmaceutical Co., Ltd, Changsha, Hunan, China) for 1–4 days. After washing, the cells were treated with RNAiso (Takara Bio Inc., Kusatsu, Shiga, Japan), and total RNA was obtained according to the manufacturer's protocol. The cDNA was synthesized and reverse-transcribed from 3 µg of total RNA. qRT-PCR was performed with a CFX Connect™ RT-PCR Detection System (Bio-Rad Laboratories, Hercules, CA, USA). The amplification condition was an initial denaturation (95°C, 10 min), followed by amplification cycles (40 cycles of 95°C, 10 s; 65°C, 15 s; 72°C, 30 s).

2.6. Quantification of viral RNA copy number

According to the manufacturer's instructions, the viral RNA was isolated using a QiaAMP viral RNA purification kit (Qiagen, Hilden, Germany). qRT-PCR was performed using TOPreal™ qPCR 2X Pre-MIX (Enzymomics, Daejeon, Korea). For RNA quantification of HCoV-OC43 nucleoprotein (N protein) and matrix protein (M protein), N and M protein genes were amplified by PCR, digested with BamHI and EcoRI restriction enzymes (Enzymomics), and inserted into the MDH1-PGK-GFP_2.0 vector (Addgene, Watertown, MA, USA). The HCoV-OC43 RNA copy number was calculated by a standard curve of viral RNA concentrations. The primer sequences are listed in Table 1.

Table 1
Primers Sequences

Primer	Direction	Sequence (5' – 3')
GAPDH	Fwd	AATGGTGAAGGTCGGTGTGAAC
	Rev	GAAGATGGTGATGGGCTTCC
N	Fwd	AGCAACCAGGCTGATGTCAATACC
	Rev	AGCAGACCTTCCTGAGCCTTCAAT
N (for cloning)	Fwd	AATTATGGATCCAGCAACCAGGCTGATGTCAATACC
	Rev	ATAGTCGAATTCAGCAGACCTTCTGAGCCTTCAAT
M	Fwd	GGCTTATGTGGCCCTTACT
	Rev	GGCAAACTGCCAAGAATA
M (for cloning)	Fwd	ATATCTGGATCCCGCTTATGTGGCCCTTACT
	Rev	GCGCCTGAATTCGGCAAATCTGCCAAGAATA
IFN-α	Fwd	GTGCTCAGCTGCAAGTCAAG
	Rev	TTATCCAGGCTGTGGGTCTC
IFN-β	Fwd	ACCAACAAGTGTCTCTCTCCA
	Rev	GTAGTGGAGAAGCACAACAGG
IFN-γ	Fwd	TCTTGGCTGTACT GCCAGGACCCA
	Rev	TCAGCCACTTCGTGCCG GTCTTC
MxA	Fwd	CAACTGTGACGACGATATG
	Rev	GTCTGTCCACACTAGAG




Fig. 1. Cytotoxicity and antiviral effect of KRG and ginsenosides. For cytotoxicity, HCT-8 cells were treated with KRG (100 µg/mL) or each of the 10 different ginsenosides (100 µM) for 7 days, and cell viability was measured by the MTT assay (A). For testing antiviral effect, plaque formation assay was performed (B). Cell viability is shown as relative values in each treatment compared to the untreated control group (N). HCoV-OC43 plaque formation levels were represented as the relative number of plaques compared with that obtained from DMSO-treated control (B). The data are represented as mean ± SEM of three independent experiments with similar results. One-way ANOVA plus Tukey’s multiple comparison test was applied.

2.7. Fluorescence-activated cell sorting (FACS) analysis

HCT-8 cells (1 × 10⁶ cells) were seeded in a 6-well plate. After 24 h of culture, HCT-8 cells were infected by HCoV-OC43 for 1 h. After infection, HCT-8 cells were washed once in PBS to remove the uninfected virus. The cells were cultured with 25, 50, 100 µg/mL of KRG and 5 µM of remdesivir for 4 days. After 4 days of infection, HCT-8 cells were harvested by trypsinization and washed in PBS. The cells were fixed and permeabilized for 15 min at 4°C with fixation/permeabilization working solution diluted from fixation/permeabilization concentrate (eBioscience, San Diego, CA, USA) with fixation/permeabilization diluent (eBioscience). Afterward, samples were incubated for 2 h at 4°C with primary anti-HCoV-OC43 antibody (Merck, Rahway, NJ, USA) or anti-HCoV-OC43 S antibody (Cusabio, Houston, TX, USA). Subsequently, samples were incubated for 1 h with Alexa Fluor 488 labeled goat Anti-Mouse IgG (Cell Signaling, Danvers, MA, USA) for anti-HCoV-OC43 antibody or Alexa Fluor 488-conjugated goat Anti-rabbit IgG (H + L), F(ab’)2 Fragment (Cell Signaling) for anti-HCoV-OC43 S antibody. Cells were resuspended in PBS and analyzed by FACS Calibur (Becton Dickinson, San Diego, CA, USA). Data were analyzed by CellQuest Pro software (Becton Dickinson).

2.8. Statistics

All experiments were repeated at least three times, and data are presented as mean ± standard deviation. For statistical analysis, one-way ANOVA plus Tukey’s multiple comparison test was

applied using Prism 5.0 (GraphPad Software, San Diego, CA, USA). *p* < 0.05 was considered statistically significant. **p* < 0.05; ***p* < 0.01; and ****p* < 0.001.

3. Results

3.1. Cytotoxic effects of KRG extract and ginsenosides

To evaluate the cytotoxicity of KRG extract and ginsenosides in HCT-8 cells, KRG and ginsenosides were added to HCT-8 cells for 7 days, and cytotoxicity was measured using an MTT assay. KRG and all ginsenosides except Rg3 (S) and Rh2 (S) showed no significant cytotoxic effects (Fig. 1A). Therefore, subsequent experiments were conducted with KRG and the eight ginsenosides devoid of cytotoxicity.

3.2. Antiviral activity of KRG and ginsenosides against HCoV-OC43 in HCT-8 cells

To test the antiviral activity of KRG and ginsenosides, the number of plaques formed in HCT-8 cells infected with HCoV-OC43 was counted following treatment with KRG or the eight ginsenosides. Compared with the DMSO control, the plaque formation in HCT-8 was significantly decreased in KRG-treated cells at 25, 50, and 100 µg/mL. However, none of the ginsenoside-treated groups had antiviral activity at 100 µM against HCoV-OC43 infection (Fig. 1B).




Fig. 2. Time of addition assay to determine the antiviral activities of KRG. HCT-8 cells were infected with HCoV-OC43, and KRG was added to the cells at 24 h before, during, or after HCoV-OC43 infection. The overall scheme of the time-of-addition assay (A). Plaque assay image of HCoV-OC43-infected HCT-8 cells with or without KRG treatment (B). The plaques of pre-, co-, and post-KRG-treated cells were quantified at 7 dpi (Fig. C–E). The data are representative of three experiments with similar results.




Fig. 3. Expression of HCoV-OC43 nucleoprotein (N protein) and spike protein (S protein) in HCT-8 cells. The cells were infected with HCoV-OC43 followed by KRG treatment, and the expression of N protein (A) and S protein (B) was analyzed by flow cytometry at 4 dpi. The FACS plot is representative of three experiments with similar results.

3.3. Identification of the step of virus life cycle affected by KRG

A time-of-addition assay was implemented to identify the step of the virus life cycle at which KRG exerts its antiviral activity (Fig. 2A). When KRG was added to the cells during virus infection (co-treatment), no protection against viral replication was observed. However, the addition of KRG at 24 h before (pre-treatment) and after (post-treatment) HCoV-OC43 infection significantly reduced the number of viral plaques dose-dependently (Fig. 2B and C). The reduction of viral plaques against HCoV-OC43 infection was more prominent upon exposure of cells to KRG post-treatment than pre-treatment (Fig. 2B and C).

3.4. Inhibition of HCoV-OC43 replication by KRG

To assess the inhibitory effects of KRG on HCoV-OC43 replication, the presence of viral nucleocapsid (N) protein and spike (S) protein in HCT-8 cells was measured by flow cytometry. The cells were infected with HCoV-OC43, and the culture medium was replaced with a fresh medium containing KRG to remove unbound virus particles. After 7 days post-infection (dpi), the expression of N protein was analyzed by flow cytometry. KRG treatment decreased the expression of N and S proteins in a dose-dependent manner (Fig. 3), indicating that KRG significantly affects virus replication in HCT-8 cells.

In addition, to confirm the antiviral effect of KRG on HCoV-OC43 replication, HCT-8 cells were treated with KRG after viral infection, and the culture supernatants were harvested at 3 and 4 dpi. The abundance of HCoV-OC43 N and M protein mRNA in the supernatant was then measured by qRT-PCR. KRG treatment significantly reduced the expression of viral N protein mRNA (Fig. 4A and B), confirming that KRG blocks the replication of HCoV-OC43 in HCT-8

cells. In addition, viral matrix (M) protein mRNA expression was also decreased by KRG treatment (Fig. 4C and D).

3.5. Induction of antiviral gene expression by KRG treatment

To confirm the antiviral activity of KRG, the levels of IFN genes (IFN- α , IFN- β , and IFN- γ) and MxA were evaluated in HCT-8 cell lysates 4 dpi. KRG treatment after virus infection augmented the mRNA levels of IFN- α , IFN- β , and IFN- γ (Fig. 5A-C). MxA mRNA expression was not increased by both virus infection and KRG treatment (Fig. 5D). These data suggest that KRG suppresses viral infection by inducing the expression of antiviral cytokines such as interferons in HCoV-OC43-infected HCT-8 cells.

4. Discussion

Antiviral effects of ginsenosides have been demonstrated in previous studies. Ginsenosides Re, Rf, and Rg2 were found to protect the host from coxsackievirus and rhinovirus 3 infections [27], while ginsenoside Rg3 significantly inhibited hepatitis B virus secretion [32]. In addition, ginsenoside Rb1 suppressed the replication of various viruses including SARS-CoV [24,33–35]. However, no study reports the antiviral activity of ginsenosides against HCoV-OC43. Thus, this study assessed the antiviral effect of KRG extract and ginsenosides against HCoV-OC43. The data showed that KRG extract considerably reduced the number of HCoV-OC43 plaques dose-dependently (Fig. 1B), but none of the ginsenosides tested (ginsenosides Rc, Re, Rf, Rg1, and each isoform of Rg2, Rg3, and Rh2) showed antiviral activities against HCoV-OC43.

To find the steps of the HCoV-OC43 life cycle that could be affected by KRG extract, KRG extract was added to HCT-8 cells at three time points: after (post-treatment), during (co-treatment),




Fig. 4. Decrease in nucleoprotein (N) and membrane protein (M) mRNA levels in KRG-treated HCT-8 cells. HCT-8 cells were treated with 25, 50, and 100 µg/mL KRG for 4 days after infection with HCoV-OC43, and the culture supernatant was harvested at 3 dpi and 4 dpi. The viral RNA was obtained from the culture supernatant of HCoV-OC43-infected cells, and the mRNA level of HCoV-OC43 N protein (A, B) and M protein (C, D) was examined by qRT-PCR.

and 24 h before (pre-treatment) virus infection. Our data demonstrated that KRG extract did not directly interact with virus particle and block the viral entry, but inhibited viral replication, as shown by the decrease in plaque numbers in the pre- and post-treatment groups (Fig. 2). Considering the anti-viral effect of pre-treated KRG extract, KRG can be expected to have preventive effect on HCoV-OC43 infection as well as inhibition of virus replication.

In addition, to confirm the antiviral effect of KRG extracts, the mRNA and protein levels of HCoV-OC43 N gene were examined in HCoV-OC4-infected HCT-8 cells. The nucleocapsid N protein is a structural protein forming a helical capsid while binding to genomic RNA, and plays a critical role in CoV replication [36,37]. KRG extract treatment reduced N protein expression in HCT-8 cells and decreased mRNA expression of N protein at 3 and 4 dpi. The membrane protein (M) is a type III transmembrane glycoprotein and most abundantly present in the CoV particle. It plays a central role in virus assembly. The data demonstrated that KRG extract significantly also decreased the mRNA level of M protein. Furthermore, the expression of spike protein was also reduced by KRG extract treatment.

To investigate the effects of KRG on the host antiviral response, the production of IFN genes was measured in HCoV-OC43-infected HCT-8 cells. Type I and II IFNs are important cytokines induced

upon viral infections [38,39], and they promote an antiviral state that resists viral spreading in uninfected cells and replication in infected cells [40]. The mRNA levels of IFN-α, IFN-β, and IFN-γ were increased in the lysates of virus-infected cells at 4 dpi, and the increase in IFN was more prominent in the KRG-treated groups (Fig. 5). However, the mRNA level of MxA was not improved at 4 dpi. MxA is an IFN-inducible gene product, and MxA induced by type I IFNs (IFN-α and IFN-β) can block viral replication. Beidas and Chehadah have reported that M and N proteins of HCoV-OC43 downregulated the expression of antiviral genes associated with the type I IFN and NF-κB signaling pathways [41]. In addition to the direct increase in IFN response, KRG treatment may promote IFN response by decreasing M and N proteins.

Although ginsenosides are well known active ingredients of KRG, ten individual ginsenoside tested in this study did not show any antiviral effect on HCoV-OC43. Since KRG extract consists of numerous compounds, the combination of multiple ingredients present in KRG extract may exert synergistic effect on HCoV-OC43. In addition, further research will be required to investigate the antiviral effects of other active constituents of KRG such as polyacetylenes, phenolic compounds, and polysaccharides, and ginsenosides on HCoV-OC43 [42].




Fig. 5. Changes in the expression of IFN- α , IFN- β , IFN- γ , and MxA in KRG-treated, HCoV-OC43-infected HCT-8 cells. Total RNA was extracted from HCT-8 cells at 4 dpi, and the mRNA expression levels of IFN- α (A), IFN- β (B), IFN- γ (C), and MxA (D) were measured by qRT-PCR. GAPDH was used for data normalization.

In conclusion, this study indicates that KRG extract can impede the progression of CoV infection by enhancing virus-induced IFN responses. Further investigation would require evaluating the antiviral effects of KRG on SARS-CoV-2.

Declaration of competing interest

All authors have no conflicts of interest to declare.

Acknowledgement

This research was supported by the Chung-Ang University Research Scholarship Grants in 2022, and by a 2021 grant from the Korean Society of Ginseng.

References

[1] Mourya DT, Sapkal G, Yadav PD, Sk MB, Shete A, Gupta N. Biorisk assessment for infrastructure & biosafety requirements for the laboratories providing coronavirus SARS-CoV-2/(COVID-19) diagnosis. *Indian J Med Res* 2020;151:172–6.
 [2] Woo PC, Huang Y, Lau SK, Yuen KY. Coronavirus genomics and bioinformatics analysis. *Viruses* 2010;2:1804–20.
 [3] Vijgen L, Keyaerts E, Moes E, Thoelen I, Wollants E, Lemey P, Vandamme AM, Van Ranst M. Complete genomic sequence of human coronavirus OC43:

molecular clock analysis suggests a relatively recent zoonotic coronavirus transmission event. *J Virol* 2005;79:1595–604.
 [4] Lim YX, Ng YL, Tam JP, Liu DX. Human coronaviruses: a review of virus-host interactions. *Diseases* 2016;4.
 [5] Cabeca TK, Granato C, Bellei N. Epidemiological and clinical features of human coronavirus infections among different subsets of patients. *Influenza Other Respir Viruses* 2013;7:1040–7.
 [6] Brown AJ, Won JJ, Graham RL, Dinnon 3rd KH, Sims AC, Feng JY, Cihlar T, Denison MR, Baric RS, Sheahan TP. Broad spectrum antiviral remdesivir inhibits human endemic and zoonotic deltacoronaviruses with a highly divergent RNA dependent RNA polymerase. *Antiviral Res* 2019;169:104541.
 [7] van der Hoek L. Human coronaviruses: what do they cause? *Antivir Ther* 2007;12:651–8.
 [8] Yang C-W, Peng T-T, Hsu H-Y, Lee Y-Z, Wu S-H, Lin W-H, Ke Y-Y, Hsu T-A, Yeh T-K, Huang W-Z. Repurposing old drugs as antiviral agents for coronaviruses. *Biomedical Journal* 2020;43:368–74.
 [9] Hu Y, Meng X, Zhang F, Xiang Y, Wang J. The in vitro antiviral activity of lactoferrin against common human coronaviruses and SARS-CoV-2 is mediated by targeting the heparan sulfate co-receptor. *Emerg Microbes Infect* 2021;10:317–30.
 [10] Larson HE, Reed SE, Tyrrell DA. Isolation of rhinoviruses and coronaviruses from 38 colds in adults. *J Med Virol* 1980;5:221–9.
 [11] Zhao X, Guo F, Liu F, Cuconati A, Chang J, Block TM, Guo JT. Interferon induction of IFITM proteins promotes infection by human coronavirus OC43. *Proc Natl Acad Sci U S A* 2014;111:6756–61.
 [12] Gaunt ER, Hardie A, Claas EC, Simmonds P, Templeton KE. Epidemiology and clinical presentations of the four human coronaviruses 229E, HKU1, NL63, and OC43 detected over 3 years using a novel multiplex real-time PCR method. *J Clin Microbiol* 2010;48:2940–7.

- [13] St-Jean JR, Jacomy H, Desforges M, Vabret A, Freymuth F, Talbot PJ. Human respiratory coronavirus OC43: genetic stability and neuroinvasion. *J Virol* 2004;78:8824–34.
- [14] Beidas M, Chehadah W. PCR array profiling of antiviral genes in human embryonic kidney cells expressing human coronavirus OC43 structural and accessory proteins. *Arch Virol* 2018;163:2065–72.
- [15] Tamminen K, Salminen M, Blazevic V. Seroprevalence and SARS-CoV-2 cross-reactivity of endemic coronavirus OC43 and 229E antibodies in Finnish children and adults. *Clin Immunol* 2021;229:108782.
- [16] Yamaguchi T, Shinagawa T, Kobata H, Nakagawa H. Immunity against seasonal human coronavirus OC43 mitigates fatal deterioration of COVID-19. *Int J Infect Dis* 2021;109:261–8.
- [17] Scaglione F, Ferrara F, Dugnani S, Falchi M, Santoro G, Fraschini F. Immunomodulatory effects of two extracts of *Panax ginseng* C.A. Meyer. *Drugs Exp Clin Res* 1990;16:537–42.
- [18] See DM, Broumand N, Sahl L, Tilles JG. In vitro effects of echinacea and ginseng on natural killer and antibody-dependent cell cytotoxicity in healthy subjects and chronic fatigue syndrome or acquired immunodeficiency syndrome patients. *Immunopharmacology* 1997;35:229–35.
- [19] Helms S. Cancer prevention and therapeutics: *Panax ginseng*. *Altern Med Rev* 2004;9:259–74.
- [20] Kim SK, Park JH. Trends in ginseng research in 2010. *J Ginseng Res* 2011;35:389–98.
- [21] Kim S, Lee Y, Cho J. Korean red ginseng extract exhibits neuroprotective effects through inhibition of apoptotic cell death. *Biol Pharm Bull* 2014;37:938–46.
- [22] Sung WS, Lee DG. The combination effect of Korean red ginseng saponins with kanamycin and cefotaxime against methicillin-resistant *Staphylococcus aureus*. *Biol Pharm Bull* 2008;31:1614–7.
- [23] Im K, Kim J, Min H. Ginseng, the natural effectual antiviral: protective effects of Korean Red Ginseng against viral infection. *J Ginseng Res* 2016;40:309–14.
- [24] Lee MH, Lee BH, Lee S, Choi C. Reduction of hepatitis A virus on FRhK-4 cells treated with Korean red ginseng extract and ginsenosides. *J Food Sci* 2013;78:M1412–5.
- [25] Sung H, Kang SM, Lee MS, Kim TG, Cho YK. Korean red ginseng slows depletion of CD4 T cells in human immunodeficiency virus type 1-infected patients. *Clin Diagn Lab Immunol* 2005;12:497–501.
- [26] Kang S, Im K, Kim G, Min H. Antiviral activity of 20(R)-ginsenoside Rh2 against murine gammaherpesvirus. *J Ginseng Res* 2017;41:496–502.
- [27] Song JH, Choi HJ, Song HH, Hong EH, Lee BR, Oh SR, Choi K, Yeo SG, Lee YP, Cho S, et al. Antiviral activity of ginsenosides against coxsackievirus B3, enterovirus 71, and human rhinovirus 3. *J Ginseng Res* 2014;38:173–9.
- [28] Cho YK, Kim JE. Effect of Korean Red Ginseng intake on the survival duration of human immunodeficiency virus type 1 patients. *J Ginseng Res* 2017;41:222–6.
- [29] Lee MH, Lee BH, Jung JY, Cheon DS, Kim KT, Choi C. Antiviral effect of Korean red ginseng extract and ginsenosides on murine norovirus and feline calicivirus as surrogates for human norovirus. *J Ginseng Res* 2011;35:429–35.
- [30] Park JD, Rhee DK, Lee YH. Biological activities and chemistry of saponins from *Panax ginseng* CA Meyer. *Phytochemistry Reviews* 2005;4:159–75.
- [31] Kang S, Min H. Ginseng, the 'immunity boost': the effects of *Panax ginseng* on immune system. *J Ginseng Res* 2012;36:354–68.
- [32] Kang LJ, Choi YJ, Lee SG. Stimulation of TRAF6/TAK1 degradation and inhibition of JNK/AP-1 signalling by ginsenoside Rg3 attenuates hepatitis B virus replication. *Int J Biochem Cell Biol* 2013;45:2612–21.
- [33] Lee MH, Seo DJ, Kang JH, Oh SH, Choi C. Expression of antiviral cytokines in Crandell-Reese feline kidney cells pretreated with Korean red ginseng extract or ginsenosides. *Food Chem Toxicol* 2014;70:19–25.
- [34] Liang YY, Wang B, Qian DM, Li L, Wang ZH, Hu M, Song XX. Inhibitory effects of Ginsenoside Rb1 on apoptosis caused by HSV-1 in human glioma cells. *Virology* 2012;27:19–25.
- [35] Wu CY, Jan JT, Ma SH, Kuo CJ, Juan HF, Cheng YS, Hsu HH, Huang HC, Wu D, Brik A, et al. Small molecules targeting severe acute respiratory syndrome human coronavirus. *Proc Natl Acad Sci U S A* 2004;101:10012–7.
- [36] Parker MM, Masters PS. Sequence comparison of the N genes of five strains of the coronavirus mouse hepatitis virus suggests a three domain structure for the nucleocapsid protein. *Virology* 1990;179:463–8.
- [37] Kuo L, Masters PS. Genetic evidence for a structural interaction between the carboxy termini of the membrane and nucleocapsid proteins of mouse hepatitis virus. *J Virol* 2002;76:4987–99.
- [38] Samuel CE. Antiviral actions of interferons. *Clin Microbiol Rev* 2001;14:778–809. table of contents.
- [39] McNab F, Mayer-Barber K, Sher A, Wack A, O'Garra A. Type I interferons in infectious disease. *Nat Rev Immunol* 2015;15:87–103.
- [40] Fritsch SD, Weichhart T. Effects of interferons and viruses on metabolism. *Front Immunol* 2016;7:630.
- [41] Beidas M, Chehadah W. Effect of human coronavirus OC43 structural and accessory proteins on the transcriptional activation of antiviral response elements. *Intervirology* 2018;61:30–5.
- [42] Hyun SH, Kim SW, Seo HW, Youn SH, Kyung JS, Lee YY, In G, Park CK, Han CK. Physiological and pharmacological features of the non-saponin components in Korean Red Ginseng. *J Ginseng Res* 2020;44:527–37.