

Pre-pregnancy Diet to Maternal and Child Health Outcome: A Scoping Review of Current Evidence

Fadila Wirawan¹, Desak Gede Arie Yudhantari², Aghnaa Gayatri³

¹Department of Public Health Nutrition, Faculty of Public Health, Universitas Indonesia, Depok, Indonesia; ²Child Health Division, Leuwiliang General District Hospital, Bogor District, Indonesia; ³Department of Family and Community Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

Objectives: Pre-pregnancy diet has an important role in preparing for healthy generation. However, evidence on this issue has been scarce. A scoping review synthesising current evidence will support the demand to map ‘what has been researched’ on pre-pregnancy diet and maternal and child health.

Methods: Systematic search was performed using PICOS (Population, Intervention, Comparison, Outcomes, and Study design) framework in electronic databases. Articles were screened for eligibility, summarized, and the quality was assessed using the National Institute of Health assessment tool. The review structure complies with Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews guide.

Results: Forty-two articles were included after full-text screening. Twenty-five studies were in high-income countries (HICs), six in each upper-middle income, five in lower-middle income countries (LMICs), and one in low-income countries (LIC). Based on the regions: North America (n=16), Europe (n=5), South America (n=4), Australia (n=4), Asia (n=5), Middle East (n=2), and sub-Saharan Africa (n=1). The two-most observed diet-related exposures were dietary pattern (n=17) and dietary quality (n=12). The most assessed outcome was gestational diabetes mellitus (n=28) and fetal and newborn anthropometry (n=7). The average quality score \pm standard deviation was $70 \pm 18\%$.

Conclusions: Research related to pre-pregnancy diet is still concentrated in HICs. The context of diet may vary; therefore, future research is encouraged in LMICs and LICs context, and Mediterranean, South-East Asia, Pacific, and African regions. Some maternal and child nutrition-related morbidity, such as anemia and micronutrient deficiencies, have not been discussed. Research on these aspects will benefit to fill in the gaps related to pre-pregnancy diet and maternal and child health.

Key words: Preconception care, Preconception, Maternal health, Child health

INTRODUCTION

Pre-pregnancy or preconception health, not until recently, has come to attention as a window of opportunity to prepare for a healthy pregnancy. Pregnancy itself is not always a prepared event. In the United Kingdom, 45% of pregnancies were unplanned at the time of conception [1], and the rate may be even higher in developing countries with less access to family planning. Preconception health includes a wide range of nutrition and lifestyle aspects, one of which is diet. Healthy diets

Received: November 14, 2022 Accepted: February 20, 2023

Corresponding author: Fadila Wirawan
Department of Public Health Nutrition, Faculty of Public Health,
Universitas Indonesia, Depok 16424, Indonesia

E-mail: fadila.wirawan@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

contribute to preventing malnutrition, and in the long term, reducing the risk of diet-related non-communicable diseases (NCDs) [2]. Studies have observed the associations between pre-pregnancy lifestyle and maternal and child health outcomes [3]. Evidence also suggests that women's nutritional status before pregnancy is associated with maternal and child outcomes [4]. However, the role of diet before pregnancy in the development of maternal and child outcomes is not much discussed.

Understanding the role of pre-pregnancy diet in maternal and child health will contribute to maternal and child morbidity and mortality as well as NCDs prevention. Nutritional status is directly influenced by dietary intake, whereas, a healthy diet contributes to macronutrient and micronutrient adequacy and balanced energy [5]. In high-income countries (HICs), child-bearing age women's diet typically had a high intake of refined sugar and a high-fat diet, but low intakes of fruits, vegetables, and protein source food [3]. The national surveys in some HICs also revealed that the young adult age group have also been reported to have lower than recommended intakes for iodine, iron, and folate, which are important for pregnancy [3,6]. Therefore, it raises concerns about nutritional preparedness in pregnancy.

Research related to the pre-pregnancy diet are highly diverse. The context of diet is widely affected by socio-cultural, geographical, and economy. The maternal and child health constraints also vary depending on the region and country's socioeconomic level, leading to the possible differences in the evidence-based application based on regional context. Information related to what evidence is available, what outcomes already measured, and in what context, will support the demand for more evidence on pre-pregnancy diet's role in future health. This scoping review aims to explore the current evidence of diet in the pre-pregnancy stage to the maternal and child health outcome. A preliminary search of PROSPERO, MEDLINE, the Cochrane Database of Systematic Reviews, and *JBI* Evidence Synthesis was conducted and no current or in-progress scoping reviews or systematic reviews on the topic were identified. To answer the question of: What have been done in the existing research of pre-pregnancy diet as exposure to maternal and child health outcome?, this scoping review systematically maps the available evidence and identify the gaps for further research.

METHODS

This scoping review was structured according to the Preferred Reporting Items for Systematic Reviews and Meta-Anal-

Table 1. Population, intervention, comparison, outcomes, and study design (PICOS) framework

PICOS	Keywords
Population	"preconception"[All Fields] OR "preconceptional"[All Fields] OR "preconceptions"[All Fields] OR "pre-pregnancy"[All Fields]
Intervention	"diet"[MeSH Terms] OR "diet"[All Fields]
Comparison	Not available
Outcomes	"health"[MeSH Terms] OR "health"[All Fields] OR "healthful"[All Fields] OR "healthfulness"[All Fields] OR "healths"[All Fields]
Study design	Filters: Full text, case reports, clinical study, clinical trial, comparative study, controlled clinical trial, meta-analysis, multicenter study, observational study, randomized controlled trial, systematic review, humans, English, Indonesian

yses Extension for Scoping Reviews guide for scoping review. The literature search was performed using PICOS (Population, Intervention, Comparison, Outcomes, and Study design) framework (Table 1). The use of PICOS was chosen as the framework includes the necessary observation and more sensitivity for a scoping review related to quantitative health research compared to other frameworks, such as PEOS, SPICE, SPIDER, and others. PICOS framework also provides a comprehensive search that benefits research with limited time and resources [7]. The topic to adhere to was pre-pregnancy or preconception diet and its relation to maternal and/or child health-related outcomes. Pre-pregnancy was defined as the time before conception or pregnancy, which includes the phase in preparing for pregnancy as well as the adolescent period. The diet variable may include nutritional content, food-based supplementation, dietary quality, dietary pattern, and other dietary-related variables, however, did not include non-food-based supplementation and nutritional status. Maternal health was defined as the health-related condition of the mother during the pregnancy, childbirth, and post-natal period [8], therefore did not include the fertility outcome variable. Child health was considered as all aspects of childhood illness starting from birth. Maternal and child health search were combined as a 'health' term due to the wider scope of search compared to the use of separated 'maternal' and 'child' terms. Further selection was conducted in the article identification process.

The Article Identification

We conducted a search through some potential bibliographic databases in July 2022. The systematic search was conducted in PubMed, the largest electronic medical bibliography.

The search strategy based on the PICOS framework resulted in the use of the following concept filters: (1) pre-pregnancy or preconception; (2) diet; and (3) health. The final search strategy is available in Supplemental Material 1. An additional systematic search using keywords combination of "Pre-pregnancy diet", "Pregpregnancy diet", "Pre-conception diet", "Preconception diet", and "maternal health" and "child health" was also conducted through PubMed to identify more related articles. Articles were filtered for publication in full text in Indonesian or English languages, involving human participants, and article type based on PubMed NCBI filters: reviews, case reports, clinical study, clinical trial, comparative study, controlled clinical trial, multicentre study, observational study, randomized controlled trial (RCT), systematic review, and meta-analysis. Another search attempt was conducted through Indonesia research bibliography; SINTA; Indonesian Scientific Journal Database; and Google Scholar using local keywords for pre-pregnancy diet; "*Pola makan prahamil*"; "*Pola makan prakonsepsi*"; and "*Pola makan prakehamilan*". Only peer-reviewed articles were included.

To be included, the article needed to observe pre-pregnancy diet as exposure and maternal and/or child health as the outcome. The article should be peer-reviewed original article, involving human participants, and available in full-text. Articles that are methodological paper, pre-preprints, conference article, and in language that was not understood by the authors (aside from English and Bahasa Indonesia) were excluded. The list of inclusion and exclusion criteria can also be found in Supplemental Material 2.

The Selection of Relevant Articles

After the articles were cleaned for duplicates, two authors (FW and DGAY), with sufficient capacity to understand English scientific article, each screened for the article separately. The articles were screened by title and abstract for adherence to the topic. The authors then converged their selection of articles to be assessed further for eligibility. Full-text articles were assessed separately by the authors. The final inclusion of the articles in the scoping review was based on the authors' mutual agreement on the article's adherence to the topic. In the condition when the authors did not reach a consensus, AG was consulted. The article selection flow was as presented in Figure 1.

Data Extraction and Quality Appraisal

After the final article selection, the studies were grouped into primary research (case reports, clinical study, clinical trial, comparative study, controlled clinical trial, multicentre study, observational study, and RCT) and secondary research (review, systematic review, and meta-analysis). The following information was recorded using a Microsoft Excel spreadsheet for primary research; author(s); year of publication; year of data collection; study design; study group (if any); population; sample size; subject's age group; country; country's income level; observed or intervention group; outcome variable; time of the assessed diet; dietary assessment method (if any); maternal health finding(s); and child health finding(s). The time of the assessed diet represents the time frame in which the participant is required to recall or report. As for secondary researchs, the following information were recorded; author(s); year of publication; study design; sample size; exposure variable; outcome variable; maternal health finding(s); and child health finding(s). We also summarized the observed maternal and child health outcome for each diet-related variable.

To provide better insight in the currently available evidence, the quality of the articles were appraised based on the National Institute of Health study quality assessment tool [9] with the FW and DY as the Raters. The overall article's quality was decided by the average percentage of 'Yes' answers on the study quality assessment tools given by the Raters. When no consensus reached by the Raters, AG was consulted. We grouped the article's quality ratings into three groups; poor (fulfilling less than 33.3% of the criteria); fair (fulfilling 33.3 to 66.6% of the criteria); good (fulfilling more than 66.6% of the criteria). Quality assessment of articles were conducted to evaluate the current quality of evidence available in this review to map the currently available study's quality, not as a basis of inclusion.

Protocol Registration

This scoping review protocol has been registered in the Open Science Framework Registries: <https://doi.org/10.17605/OSF.IO/7F9C2>.

Ethics Statement

This study does not include human subject involvement, therefore exempted for institutional review board approval.




Figure 1. Article selection flow. PICO, population, intervention, comparison, outcomes, and study design.

RESULTS

The article identification and selection are presented in Figure 1. A search using the PICOS framework in the PubMed database was conducted and yielded 267 articles after filtering for full-text, language, and types of articles. An additional 29 articles were obtained from linked research, database, and the web. After cleaning for duplicates, 286 articles were screened for adherence to the topic. A total of 42 articles were included after the full-text screening, consisting of 37 primary research articles and five secondary research articles to be included in the current scoping review.

The Characteristics of the Studies

The extracted findings from the included primary research were as shown in Table 2 [10-46]. One study is cross-sectional, three are case-control, one is a post-hoc observational study, 27 are cohort, and five studies are RCT. All of the primary studies were quantitative. Out of 42 included studies, 27 of them

are based on big projects, mostly based on the Nurses' Health Study II cohort (13 studies). This scoping review did not use a limiter of date of publication in the search process. However, studies included in the final selection were published in 2009 to 2022.

Of the 42 studies included, 25 studies were conducted in high-income countries (HICs), six in upper-middle-income countries (UMICs), five in lower-middle-income countries (LMICs), and only one was conducted in low-income country (LIC). Based on the geographical areas, 16 were conducted in North America, five were in Europe, four were in South America, four were in Australia, four were in East and South Asia, two were in the Middle East, one in South-East Asia, and one was in sub-Saharan Africa. The proportion of observed pre-pregnancy diet variables based on region was presented in Figure 2A. The pre-pregnancy dietary patterns had been observed in almost all regions but sub-Saharan Africa and East and South Asia. Meanwhile, pre-pregnancy dietary quality research had only been brought up in regions with UMICs and HICs, without any known stud-

Table 2. Primary studies included in the scoping review

Study	Year of data collection	Study design	Study group	Population	Sample size	Subjects' age (y)	Country	Country's income level'	Exposure/ intervention variable	Outcome variable	Time of the assessed diet	Dietary assessment method	Maternal health finding(s)	Child health finding(s)
Alves-Santos et al., 2019 [16]	2009-2012	Cohort	NA	Pregnant women	193	20-40	Brazil	Upper-middle	Pre-pregnancy dietary pattern	LGA, birth length, Apgar score, and preterm birth	Recall 6 mo before pregnancy	Semi-quantitative FFQ	NA	Fast food and candies were associated with higher LGA; "beans, bread, and fat" was associated with lower Apgar score; and "vegetables and dairy" was associated with lower preterm birth
Asadi et al., 2019 [11]	2014-2015	Case-control	NA	Pregnant women	296	19-40	Iran	Lower-middle	Pre-pregnancy dietary pattern history	GDM	Recall of diet in the previous year	FFQ	Prudent diet was associated with lower risk of GDM	NA
Bao et al., 2013 [35]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	15 294	25-44	USA	High	Protein intake	GDM	Before pregnant	Semi-quantitative FFQ	Red meat consumption was associated with GDM; Nut intake was associated with a lower risk of GDM	NA
Bao et al., 2014 [31]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	15 027	25-44	USA	High	Fried-food consumption	GDM	Before pregnant	FFQ	Fried food consumption was associated with GDM risk	NA
Bao et al., 2014 [12]	1999-2001	Cohort	Nurses' Health Study II	Non-pregnant women	21 411	25-44	USA	High	Low-carbohydrate diet	GDM	Before pregnant	Semi-quantitative FFQ	Low-carbohydrate, high animal protein, and animal fat diet were associated with increased GDM risk	NA
Bao et al., 2016 [38]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	15 632	25-44	USA	High	Potato intake	GDM	Before pregnant	Semi-quantitative FFQ	Higher potato consumption was significantly associated with an increased risk of GDM	NA
Bao et al., 2018 [43]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	15 225	24-44	USA	High	Dietary vitamin D intake	GDM	Before pregnant	FFQ	Dietary and total vitamin D intakes were inversely associated with GDM risk	NA
Beraim et al., 2018 [19]	2009-2011	Cohort	NA	Pregnant women	154	22-31	Brazil	Upper-middle	Food intake 6 mo before pregnant	Changes of serum EPA and DHA, total PUFA, total n-3 and omega-6/omega-3 ratio during pregnancy	Recall of diet 6 mo before pregnant	Semi-quantitative FFQ	Healthy dietary pattern was associated with serum fatty acids, however the effect was modified by BMI	NA

(Continued to the next page)

Table 2. Continued from the previous page

Study	Year of data collection	Study design	Study group	Population	Sample size	Subjects' age (y)	Country	Country's income level ¹	Exposure/ intervention variable	Outcome variable	Time of the assessed diet	Dietary assessment method	Maternal health finding(s)	Child health finding(s)
Bowers et al., 2012 [33]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	13 475	22-44	USA	High	Fat intake	GDM	Before pregnant	Semi-quantitative FFQ	Higher consumption of dietary cholesterol and animal fat was associated with GDM	NA
Chen et al., 2009 [41]	1992-2001	Cohort	Nurses' Health Study II	Non-pregnant women	13 475	24-44	USA	High	Sugar-sweetened beverage intake	GDM	Before pregnant	Semi-quantitative FFQ	Sugar-sweetened cola was associated with the risk of GDM	NA
Chen et al., 2014 [37]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	13 475	24-44	USA	High	Fruit and fruit juice intake	GDM	Before pregnant	Semi-quantitative FFQ	Apple intake was associated with lower risk of GDM	NA
Chen et al., 2019 [39]	2018	Case-control	NA	Pregnant women	260	Child-bearing age	China	Upper-middle	Pre-pregnancy probiotic yogurt intake	GDM	Recall	Structured questionnaire	No association	NA
Dhana et al., 2018 [28]	1989-2013	Cohort	Nurses' Health Study II	Non-pregnant women	4698	25-42	USA	High	Alternate HEI 2010	BMI of offspring	Before pregnant	FFQ	NA	No association
Gaskins et al., 2014 [22]	1992-2009	Cohort	Nurses' Health Study II	Non-pregnant women	15 950	24-44	USA	High	Alternate HEI 2010, alternate MeD, and fertility diet	Pregnancy loss	Before pregnant	Semi-quantitative FFQ	NA	No association
Gaskins et al., 2018 [40]	1991-2009	Cohort	Nurses' Health Study II	Non-pregnant women	15 590	24-44	USA	High	Coffee intake	Spontaneous abortion	Before pregnant	FFQ	NA	High coffee consumption was associated with the increased risk of spontaneous abortion
Gete et al., 2021 [30]	2003-2016	Cohort	Australian Longitudinal Study in Women's Health (ALSWH) and Mother and Their Children's Health Study (MatCh)	Non-pregnant women	1554	25-30	Australia	High	HEI 2015	Child behavioral problems using Strengths and Difficulties Questionnaire (SDQ)	Before pregnant	Semi-quantitative FFQ	NA	Better diet quality was associated with lower risk of offspring total behavioral difficulties, externalizing score, hyperactivity, and peer problems
Gete et al., 2021 [29]	1996-2015	Cohort	ALSWH and MatCh	Non-pregnant women	1936	18-23	Australia	High	HEI 2015	Childhood BMI of the offspring	Before pregnant	Semi-quantitative FFQ	NA	High diet quality was associated with reduced risk of underweight and obesity

(Continued to the next page)

Table 2. Continued from the previous page

Study	Year of data collection	Study design	Study group	Population	Sample size	Subjects' age (y)	Country	Country income level ¹	Exposure/intervention variable	Outcome variable	Time of the assessed diet	Dietary assessment method	Maternal health finding(s)	Child health finding(s)
Gievic et al., 2018 [25]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	41 229	24-44	USA	High	Dietary diversity (FGI and MDD-W) and dietary and dietary quality (alternate HEI 2010 and PDIOS)	GDM and hypertensive disorder in pregnancy	Before pregnant	Semi-quantitative FFQ	Alternate HEI-2010 and PDIOS were associated with a lower risk of GDM	NA
Grieger et al., 2016 [17]	2009-2013	Cross-sectional	NA	Pregnant asthmatic women	309	>18	Australia	High	Pre-pregnancy protein; fat and sugar; and vegetarian diet	Asthma control	Recall 12 mo before pregnancy	FFQ	High fat and sugar diet was associated with uncontrolled asthma	NA
Grieger et al., 2019 [32]	2009-2013	Cohort	NA	Pregnant women	234	>18	Australia	High	Food intake 12 mo before pregnant	Allergies in children	Recall of diet 12 mo before pregnant	FFQ	NA	Dairy, fresh fruit, unsaturated and saturated spreads, takeaway foods, and non-oily fish were protective against some types of allergies; Poultry and fruit juice were adversely associated with allergy
Hillesund et al., 2018 [23]	2009-2013	Post-hoc observational study	Norwegian Fit for Delivery (NFFD)	Pregnant women	591	>18	Norway	High	NFFD diet score pre-pregnancy recall	Gestational weight gain, preterm delivery, birth anthropometry, GDM, preeclampsia	Recall	FFQ	A higher diet score was associated with a lower risk of excessive weight gain	A higher diet score was associated with a lower risk of preterm delivery
Jarman et al., 2018 [24]	2009-2012	Cohort	Alberta Pregnancy Outcomes and Nutrition (APrON) study	Pregnant women	1598	>16	Canada	High	Pre-pregnancy healthy eating pattern score	Gestational hypertension and GDM	Recall of diet 12 mo before pregnancy	FFQ	A higher healthy eating pattern score was associated with lower odds of gestational hypertension	NA
Lamyian et al., 2017 [34]	2010-2011	Cohort	NA	Pregnant women in prenatal clinic	1054	18-45	Iran	Lower-middle	Fast food consumption in the previous year	GDM	Recall of diet in previous year	Semi-quantitative FFQ	Fast food intake was significantly associated with increased risk of developing GDM	NA
Lawande et al., 2018 [44]	2006-2012	RCT	Mumbai Maternal Nutrition Project	Non-pregnant women	6513	22-28	India	Lower-middle	Food-based micronutrient-rich snack	Fetal size and growth	Before pregnant	NA	NA	No association

(Continued to the next page)

Table 2. Continued from the previous page

Study	Year of data collection	Study design	Study group	Population	Sample size	Subjects' age (y)	Country	Country's income level	Exposure/intervention variable	Outcome variable	Time of the assessed diet	Dietary assessment method	Maternal health finding(s)	Child health finding(s)
Van Lippevelde et al., 2021 [20]	1995-2008	Cohort	Young-HUNT study	Adolescent	8980	<19	Norway	High	Dietary and meal patterns	Birth anthropometry	Before pregnant	Self-reported questionnaire	NA	Higher energy-dense food consumption was associated with the risk of LBW
Mari-Sanchis et al., 2018 [36]	1999-2012	Cohort	The SUN Project	Non-pregnant women	3298	Child-bearing age	Spain	High	Meat consumption and iron intake	GDM	Up to 2 y before pregnant	Semi-quantitative FFQ	Higher meat consumption associated with increased risk of GDM	NA
Mohammed et al., 2020 [42]	2011-2014	RCT	NA	Non-pregnant women	1220	≥18	Ethiopia	Low	Iodized salt intervention	Cognition of offspring	Before pregnant	24-hr food frequency interview	NA	Offspring's cognitive scores were better in intervention group compared to control
Olmedo-Requena et al., 2019 [13]	n/a	Case-control	NA	Pregnant women with GDM	1466	>18	Spain	High	Pregnancy with GDM	Pre-pregnancy adherence to MeD	Recall of diet in previous year before pregnant	FFQ	Very high adherence to MeD was associated with reduced GDM case; Low consumption of meat and its derivatives were protective of GDM	NA
Potdar et al., 2014 [45]	2006-2012	RCT	Mumbai Maternal Nutrition Project	Non-pregnant women	6513	<40	India	Lower-middle	Micronutrient-rich vegetarian snack	Birth weight	≥90 day before last menstrual period	NA	NA	Micronutrient-rich snack was associated higher birth weight and lower low-birth weight outcome
Sahariah et al., 2022 [46]	2006-2011	RCT	Mumbai Maternal Nutrition Project	Non-pregnant women	1255	Child-bearing age	India	Lower-middle	Micronutrient-rich vegetarian snack	Cardiometabolic risk markers and body composition in offspring (5-10 y follow up after intervention)	Before pregnant	NA	NA	No association
Siega-Riz et al., 2021 [27]	2008-2011	Cohort	Hispanic Community Health Study	Non-pregnant women	497	18-44	USA	High	HEI 2010	infant birth weight	Mean of 3.2 y before pregnancy	24-hr food recall	NA	High dietary quality was associated with infant birth weight
Teixeira et al., 2021 [21]	2011-2013	Cohort	ProciAr study	Pregnant women	299	Child-bearing age	Brazil	Upper-middle	Pre-pregnancy dietary pattern	Newborn anthropometry	Recall of diet in previous 12 mo	FFQ	NA	Energy-dense nutrient-poor pre-pregnancy diet was a risk factor for SGA newborn

(Continued to the next page)

Table 2. Continued from the previous page

Study	Year of data collection	Study design	Study group	Population	Sample size	Subjects' age (y)	Country	Country's income level ¹	Exposure/intervention variable	Outcome variable	Time of the assessed diet	Dietary assessment method	Maternal health finding(s)	Child health finding(s)
Tobias et al., 2012 [14]	1991-2001	Cohort	Nurses' Health Study II	Non-pregnant women	15 254	24-44	USA	High	Med, DASH, and alternate HEI scores	GDM	Before pregnant	Semi-quantitative FFQ	Adherence to alternate Med, DASH, and alternate HEI dietary patterns was associated with lower GDM risk	NA
Valkama et al., 2018 [10]	2008-2014	RCT	RADIEL Study	Non-pregnant women	75	≥18	Finland	High	Dietary counselling and diet	GDM	Before pregnant	FFQ	Low-fat cheese consumption was associated with lower GDM risk	NA
Vilela et al., 2014 [18]	2009-2011	Cohort	NA	Pregnant women	248	20-40	Brazil	Upper-middle	Pre-pregnancy dietary pattern	Depressive symptoms during pregnancy	Recall 6 mo before pregnancy	Semi-quantitative FFQ	Health dietary pattern was associated with lower risk of depressive symptoms	NA
Yee et al., 2020 [26]	2010-2013	Cohort	The Nulliparous Pregnancy Outcomes Study: Monitoring Mothers-To-Be (nuMoM2b)	Pregnant women	8259	>13	USA	High	Preconception HEI 2010	Maternal pregnancy morbidities and child birth outcomes	Recall of diet 3 mo before conception	Semi-quantitative FFQ	Poor dietary quality was associated with higher preterm birth, NICU admission, SGA, and LBW, however lower risk of macrosomia	NA
Yong et al., 2020 [15]	2008-2014	Cohort	The Seremban Cohort Study (SECOST)	Pregnant women	737	Child-bearing age	Malaysia	Upper-middle	Pre-pregnancy dietary pattern	GDM	Recall of diet before pregnancy in the first trimester	Semi-quantitative FFQ	High sugar and fat diet reduced the risk of GDM	NA

RCT, randomized controlled trial; FFQ, Food-frequency questionnaire; HEI, Healthy Eating Index; Med, Mediterranean diet; DASH, dietary approach to stop hypertension; NFFD, Norwegian fit for delivery diet; FGI, Food Groups Index; MDD-W, minimum dietary diversity for women; PDOS, Prime Diet Quality Score; GDM, gestational diabetes mellitus; NICU, neonatal intensive care unit; LGA, large-for-gestational age; SGA, small-for-gestational age; LBW, low birth weight; EPA, eicosapentaenoic acid; PUFA, polyunsaturated fatty acid; BMI, body mass index; NA, not available.




Figure 2. (A) Pre-pregnancy diet research variable theme based on regions; (B) Pre-pregnancy diet research variable theme based on observed outcomes; (C) Timeline of the available studies.

ies of pre-pregnancy dietary quality from Asia, Middle East, and sub-Saharan Africa. The diet variable as the exposure was mostly (30 studies) measured by food frequency questionnaire (FFQ), other methods used were 24-hour dietary recall, questionnaire, and interview. A total of 13 studies were conducted in the already pregnant population and rely on recall of the diet during the pre-pregnancy phase. Figure 2B shows the type of diet-related variables that were considered as exposures

based on the included studies and its maternal and child health outcome variable. The duration of observation from pre-pregnancy to outcome were varied as seen in Figure 2C. The shortest and the most common observation, 15 studies were started no more than six months before pregnancy to observe the outcomes during pregnancy. The longest observation was until adolescent by only one study.

Table 3 [47-51] presents the findings of secondary studies

Table 3. Secondary studies included in the scoping review

Study	Study design	Sample size (study)	Exposure variable	Outcome variable	Maternal health finding(s)	Child health finding(s)
Cui et al., 2021 [47]	Meta-analysis	38	Pre-pregnancy diet	GDM	Higher pre-pregnancy consumption of fried food, fast food, red and processed meat, heme iron, and low-carbohydrate diet pattern were associated with increased risk of gestational diabetes mellitus; Higher intake of dietary fiber was associated with lower risk of gestational diabetes mellitus	High pre-pregnancy caffeine intake was associated with increased risk of small-for-gestational age.
Tieu et al., 2017 [48]	Systematic review	0	Interconception intervention	Maternal and child morbidity	NA	NA
Stephenson et al., 2018 [49]	Review	NA	Preconception diet	Maternal and child morbidity	Diet before pregnancy may modify maternal outcomes through BMI status and micronutrient deficiency	Diet before pregnancy may modify prenatal outcomes through BMI status and micronutrient deficiency
Temel et al., 2014 [50]	Systematic review	44	Preconceptional nutrition support	Maternal and child health	NA	Long-term nutritional support was associated with a positive effect on birth weight
Hanson et al., 2012 [51]	Review	NA	Pre-pregnancy diet	NCD	NA	Maternal diet before pregnancy are related to phenotypic characteristics of the child, such as adiposity at birth and in childhood, and markers of cardiovascular risk

GDM, gestational diabetes mellitus; NCD, non-communicable disease; BMI, body mass index; NA, not available.

that are included in the current scoping review. One is meta-analysis, two are systematic reviews, and the other two are reviews. The publication year ranges from 2012 to 2022. One of the systematic reviews is without results because did not find any relevant intervention study related to preconception diet.

Quality of the Studies

The quality of the included studies was appraised (Supplemental Material 3) with an average quality score of $70 \pm 18\%$. Most of the observational studies were based on secondary data from a large cohort project. However, the quality assessment was made solely on how the authors' reporting in the article. Three articles are considered poor, ten articles are fair, and 29 are good (marked in Table 4) [10-46]. Articles with poor quality were the two reviews and one RCT that did not provide enough information related to the study methods. Most of the studies that have fair quality did not mention the justification of the sample size, their study power and effect. The studies with fair quality also did not clearly explain whether the assessors were blinded to the exposure status, therefore, assumed as 'No' or 'Not Clear'. The outcome in the fair studies was self-reported, and therefore considered to have a high risk of bias. Regardless of the rank of quality, most studies in use recall and only measured the exposure once.

Assessed Diet-related Exposure and Outcomes

We found 13 observations of dietary patterns and 16 observations of dietary quality (Figure 2B). The discussion of dietary pattern includes various known dietary patterns; Western diet; Prudent diet; Mediterranean diet; alternate Mediterranean diet; Nordic diet; and Diet to Stop Hypertension, and other defined dietary patterns; low caloric diet with controlled micro-nutrient; energy-dense poor-nutrient diet; and pre-defined healthy diet. Dietary quality includes observation for Healthy Eating Index, alternate Healthy Eating Index, and Prime Diet Quality Score. The mention of dietary pattern or quality score can be found in Table 1.

Maternal gestational diabetes mellitus (GDM) was the most discussed topic for the maternal outcome, with 28 observations for different diet-related exposures. Hypertension disorder in pregnancy (preeclampsia) and asthma were the second observed maternal outcome ($n=4$). Meanwhile, fetal and newborn anthropometry was the most observed child health outcome to different diet-related exposures ($n=7$) followed by preterm birth and neonatal morbidity ($n=3$).

DISCUSSION

The current scoping review observed that, to date, research related to pre-pregnancy diet was mostly conducted in the

Table 4. Synthesis of diet variable and observed outcomes from primary studies

Pre-pregnancy diet variables	Observed outcome														
	Maternal outcome					Child outcome									
	GDM	Hypertension in pregnancy	Nutritional status	Type of delivery	Asthma	Depression	Cardio-metabolic serum marker	Fetal and newborn anthropometry	Preterm birth and neonatal morbidity	Spontaneous abortion and infant mortality	Child nutrition status	Allergy and eczema	Cognitive and behaviour	Cardio-metabolic risk	Total (n)
Diet counselling	[10]														1
Dietary pattern	[11], [12], [13], [14], [15]		[16]		[17]	[18]	[19]	[20], [21]	[16]	[22]					13
Dietary quality	[23], [24], [25]	[23], [25], [26]	[23]	[26]				[23], [26], [27]	[23], [26]		[28], [29]		[30]		16
Dietary diversity	[25]	[25]													2
Food component															
Fried food	[31]											[32]			1
General food															1
Dietary fat	[33]				[17]										2
Fast food	[10], [34]														2
Dairy	[10]														1
Protein intake	[10], [35]				[17]										3
Meat	[36]														1
Fish	[10]														1
Fruit	[10], [37]														2
Vegetables and legumes	[10]														1
Potato	[38]														1
Bread, cereal, grains	[10]														1
Probiotic yogurt	[39]														1
Caffeine										[40]					1
Sugar-sweetened beverage and sugary snacks	[10], [41]				[17]										3
Food-based micronutrient															
Iodized salt intake													[42]		1
Dietary vitamin D	[43]														1
Food-based micronutrient-rich snack								[44], [45]						[46]	3
Total (n)	28	4	1	2	4	1	1	7	3	2	2	1	2	1	

GDM, gestational diabetes mellitus. ¹Study with good quality.

HIC population. The most observed area was North America (the United States and Canada) and the least observed based on the area was the African region. The dietary pattern and dietary quality were the two-most observed diet-related exposures. The most assessed outcome was GDM, hypertension disorder in pregnancy, and fetal and newborn anthropometry. FFQ was the most common dietary assessment tool used in research related to the pre-pregnancy diet.

Pregnancy is, most of the time, an unpredictable event. Even in the case of intended pregnancy, the starting point of early pregnancy can be unknown. Therefore, research related to pre-pregnancy diet often requires a long longitudinal observation from exposure to outcome. The time constraint explains the high number of studies that extracted secondary data from big longitudinal study. However, in studies using such method, the exposure and outcome assessments were sometimes cannot be implemented consistently.

The types of maternal morbidity are varying based on the region and types of economy. Hemorrhage, hypertension and preeclampsia, and sepsis were recognized as the leading cause of maternal mortality in high income countries. Hemorrhage, hypertension, dystocia, and sepsis in sub-Saharan Africa. Hypertension and hemorrhage in North Africa and Middle East. Hemorrhage, hypertension, anemia, and sepsis in Asia. Hypertension, hemorrhage, and infection in Latin America [10]. Hypertension, hemorrhage, and sepsis or infection are problems in all regions and economy, however, only three articles observed four dietary parameters to hypertensive disorder in pregnancy. Pre-pregnancy diet exposure to hemorrhage and infection were not found, but the nature of both etiologies was not directly related to diet and nutrition.

Anemia is one of the most prevalent maternal morbidities in Asia [52] and globally in LMICs and LICs with global prevalence of 42% in pregnant women [53]. Anemia is highly affected by nutrition [54]. Iron, folic acid, vitamin B12, and vitamin A deficiency, as well as protein energy malnutrition are among the contributing factors of nutritional anemia [53]. A population study in Indonesia shows that pre-pregnancy anemia increased the risk of child anemia in under-5 years old [55]. However, we did not identify any study of pre-pregnancy diet and anemia.

Malnourished mother's nutrition is unlikely to suddenly improved during pregnancy due to even increasing nutritional demand. There are risk of insufficient weight gain and chronic energy and micronutrient deficiencies, that contribute to

poor birth outcomes, neonatal mortality, and subsequent childhood malnutrition [56]. Newborn mortality incidence was reported the highest in sub-Saharan Africa, followed by Central and South Asia. Most neonatal deaths were related to preterm birth, childbirth-related complications, infections, and birth defects [57]. Malnutrition in children also contributes to under-5 death globally. Stunting and chronic protein-energy malnutrition are prevalent in Africa and Asia, while micronutrient deficiencies, mainly iron, vitamin A, iodine, and zinc, are prevalent in developing countries [56,58,59]. Referring to the burden of child mortality and morbidity, regions with high prevalence of child mortality still had the least number of available research in preconception diet and child health outcome. Micronutrient deficiencies in children are connected to maternal nutrition, such as iron storage in newborns that are affected by maternal iron [60]. In addition, evidence already supports the prevention of birth defects through adequate maternal nutrition, for example folate for neural tube defect prevention [61,62]. Maternal protein malnutrition also linked to hippocampal formation and neurobehavioral development [63,64]. But research related to preconception diet and child micronutrient deficiency, birth defect, and development were still scarce.

Although the evidence level was still weak, maternal nutrition is associated with intergenerational effects on NCDs risk in adult offspring [65]. Study in epigenetics suggests the role of nutrition in the early phase of life in the development of allergy, metabolic disorders risk, and cancer in the future [66-69]. However, further study is still required, including maternal pre-pregnancy diet's role in the future NCDs risk.

Globally, there has been a shift in the diet not only in HICs but also in developing countries. The plant-based diet has been shifted to high-fat, energy-dense diet [70]. Adolescent diet practice is crucial that is likely to extend to adulthood and representative of pre-pregnancy diet [71]. Several research in adolescent dietary quality report the poor dietary habit in adolescent girls in LMICs [71,72]. Pre-pregnancy diet has not been attention in LMICs and LICs, although is recognized as a critical base for birth preparedness and health in the lower economy countries. Some experts agreed that pre-pregnancy care priority of urgency is high due to the remaining-high maternal and child mortality and morbidity in some regions. Research related to pre-pregnancy context also needs to consider different local circumstances and context [73]. The currently available studies were mostly in HICs, which may not be

completely suitable for implementation in LMICs and LICs context.

The current study used systematic search method and additional hand search to obtain more articles that engaged with the topic. The quality of included articles was also assessed by its methodology. The approach also conducted by two assessors, which, lowers the risk of bias. However, some studies related to pre-pregnancy may be missed during the search process. Studies in other language from English and Bahasa were also not included and may cause bias in selection process. The current scoping review also only included primary original research and did not include review and grey literature.

CONCLUSION

Pre-pregnancy diet is a potential opportunity to prepare for the healthier next generation and stop intergenerational cycle of malnutrition. The current evidence related to pre-pregnancy diet is still limited. Future research is encouraged in LMICs and LICs contexts, as well as South-East Asia, Pacific, and African regions. Some maternal and child nutrition-related morbidity, such as anemia, micronutrient deficiencies, and birth defects have not much or at all discussed yet. More research on these aspects and the regional dietary context will benefit to fill in the gaps related to pre-pregnancy diet and maternal and child health.

SUPPLEMENTAL MATERIALS

Supplemental materials are available at <https://doi.org/10.3961/jpmph.22.472>.

CONFLICT OF INTEREST

The authors have no conflicts of interest associated with the material presented in this paper.

FUNDING

None.

ACKNOWLEDGEMENTS

None.

AUTHOR CONTRIBUTIONS

Conceptualization: Wirawan F, Gayatri A. Data curation: Wirawan F, Yudhantari DGA, Gayatri A. Formal analysis: Wirawan F, Yudhantari DGA, Gayatri A. Funding acquisition: None. Methodology: Wirawan F, Yudhantari DGA, Gayatri A. Project administration: Wirawan F. Visualization: Wirawan F. Writing – original draft: Wirawan F, Yudhantari DGA, Gayatri A. Writing – review & editing: Wirawan F, Yudhantari DGA, Gayatri A.

ORCID

Fadila Wirawan <https://orcid.org/0000-0003-1621-1470>

Desak Gede Arie Yudhantari

<https://orcid.org/0009-0000-3728-0526>

Aghnaa Gayatri

<https://orcid.org/0000-0001-5760-8488>

REFERENCES

1. Public Health England. Health matters: reproductive health and pregnancy planning; 2018 [cited 2022 Jul 23]. Available from: <https://www.gov.uk/government/publications/health-matters-reproductive-health-and-pregnancy-planning/health-matters-reproductive-health-and-pregnancy-planning>.
2. World Health Organization. Sustainable healthy diets: guiding principles; 2019 [cited 2022 Jul 23]. Available from: <https://www.who.int/publications/i/item/9789241516648>.
3. Stephenson J, Heslehurst N, Hall J, Schoenaker DA, Hutchinson J, Cade JE, et al. Before the beginning: nutrition and lifestyle in the preconception period and its importance for future health. *Lancet* 2018;391(10132):1830-1841.
4. Li C, Zhu N, Zeng L, Dang S, Zhou J, Pei L, et al. Effect of maternal pre-pregnancy underweight and average gestational weight gain on physical growth and intellectual development of early school-aged children. *Sci Rep* 2018;8(1):12014.
5. National Research Council (US) Committee on Diet and Health. Diet and health: implications for reducing chronic disease risk. Washington, D.C.: National Academies Press; 1989, p. 1-3.
6. United Nations Children's Fund (UNICEF). Maternal nutrition: preventing malnutrition in pregnant and breastfeeding women [cited 2022 Jul 23]. Available from: <https://www.unicef.org/nutrition/maternal>.
7. Methley AM, Campbell S, Chew-Graham C, McNally R, Cheraghi-Sohi S. PICO, PICOS and SPIDER: a comparison study of specificity and sensitivity in three search tools for qualitative

- systematic reviews. *BMC Health Serv Res* 2014;14:579.
8. World Health Organization. Maternal health [cited 2022 Jul 30]. Available from: https://www.who.int/health-topics/maternal-health#tab=tab_1.
 9. National Heart, Lung, and Blood Institute. Study quality assessment tools; 2021 [cited 2022 Jul 30]. Available from: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>.
 10. Valkama AJ, Meinilä J, Koivusalo S, Lindström J, Rönö K, Stach-Lempinen B, et al. The effect of pre-pregnancy lifestyle counselling on food intakes and association between food intakes and gestational diabetes in high-risk women: results from a randomised controlled trial. *J Hum Nutr Diet* 2018;31(3):301-305.
 11. Asadi M, Shahzeidi M, Nadjarzadeh A, Hashemi Yusefabad H, Mansoori A. The relationship between pre-pregnancy dietary patterns adherence and risk of gestational diabetes mellitus in Iran: a case-control study. *Nutr Diet* 2019;76(5):597-603.
 12. Bao W, Bowers K, Tobias DK, Olsen SF, Chavarro J, Vaag A, et al. Prepregnancy low-carbohydrate dietary pattern and risk of gestational diabetes mellitus: a prospective cohort study. *Am J Clin Nutr* 2014;99(6):1378-1384.
 13. Olmedo-Requena R, Gómez-Fernández J, Amezcua-Prieto C, Mozas-Moreno J, Khan KS, Jiménez-Moleón JJ. Pre-pregnancy adherence to the mediterranean diet and gestational diabetes mellitus: a case-control study. *Nutrients* 2019;11(5):1003.
 14. Tobias DK, Zhang C, Chavarro J, Bowers K, Rich-Edwards J, Rosner B, et al. Prepregnancy adherence to dietary patterns and lower risk of gestational diabetes mellitus. *Am J Clin Nutr* 2012;96(2):289-295.
 15. Yong HY, Mohd Shariff Z, Mohd Yusof BN, Rejali Z, Appannah G, Bindels J, et al. The association between dietary patterns before and in early pregnancy and the risk of gestational diabetes mellitus (GDM): data from the Malaysian SECOST cohort. *PLoS One* 2020;15(1):e0227246.
 16. Alves-Santos NH, Cocate PG, Benaim C, Farias DR, Emmett PM, Kac G. Prepregnancy dietary patterns and their association with perinatal outcomes: a prospective cohort study. *J Acad Nutr Diet* 2019;119(9):1439-1451.
 17. Grieger JA, Grzeskowiak LE, Wood LG, Clifton VL. Asthma control in pregnancy is associated with pre-conception dietary patterns. *Public Health Nutr* 2016;19(2):332-338.
 18. Vilela AA, Farias DR, Eshriqui I, Vaz Jdos S, Franco-Sena AB, Castro MB, et al. Prepregnancy healthy dietary pattern is inversely associated with depressive symptoms among pregnant Brazilian women. *J Nutr* 2014;144(10):1612-1618.
 19. Benaim C, Freitas-Vilela AA, Pinto TJ, Lepsch J, Farias DR, Dos Santos Vaz J, et al. Early pregnancy body mass index modifies the association of pre-pregnancy dietary patterns with serum polyunsaturated fatty acid concentrations throughout pregnancy in Brazilian women. *Matern Child Nutr* 2018;14(1):e12480.
 20. Van Lippevelde W, Vik FN, Wills AK, Strömmer ST, Barker ME, Skreden M, et al. The impact of diet during adolescence on the neonatal health of offspring: evidence on the importance of preconception diet. The HUNT study. *J Dev Orig Health Dis* 2021;12(5):798-810.
 21. Teixeira JA, Hoffman DJ, Castro TG, Saldiva SR, Francisco RP, Vieira SE, et al. Pre-pregnancy dietary pattern is associated with newborn size: results from ProcriAr study. *Br J Nutr* 2021;126(6):903-912.
 22. Gaskins AJ, Rich-Edwards JW, Hauser R, Williams PL, Gillman MW, Penzias A, et al. Prepregnancy dietary patterns and risk of pregnancy loss. *Am J Clin Nutr* 2014;100(4):1166-1172.
 23. Hillesund ER, Bere E, Sagedal LR, Vistad I, Seiler HL, Torstveit MK, et al. Pre-pregnancy and early pregnancy dietary behavior in relation to maternal and newborn health in the Norwegian Fit for Delivery study - a post hoc observational analysis. *Food Nutr Res* 2018. doi: <https://doi.org/10.29219/fnr.v62.1273>.
 24. Jarman M, Mathe N, Ramazani F, Pakseresht M, Robson PJ, Johnson ST, et al. Dietary patterns prior to pregnancy and associations with pregnancy complications. *Nutrients* 2018;10(7):914.
 25. Gicevic S, Gaskins AJ, Fung TT, Rosner B, Tobias DK, Isanaka S, et al. Evaluating pre-pregnancy dietary diversity vs. dietary quality scores as predictors of gestational diabetes and hypertensive disorders of pregnancy. *PLoS One* 2018;13(4):e0195103.
 26. Yee LM, Silver RM, Haas DM, Parry S, Mercer BM, Iams J, et al. Quality of periconceptional dietary intake and maternal and neonatal outcomes. *Am J Obstet Gynecol* 2020;223(1):121.e1-121.e8.
 27. Siega-Riz AM, Vladutiu CJ, Butera NM, Daviglius M, Gellman M, Isasi CR, et al. Preconception diet quality is associated with birth weight for gestational age among women in the Hispanic community health study/study of Latinos. *J Acad Nutr Diet* 2021;121(3):458-466.
 28. Dhana K, Zong G, Yuan C, Schernhammer E, Zhang C, Wang X, et al. Lifestyle of women before pregnancy and the risk of offspring obesity during childhood through early adulthood. *Int*

- J Obes (Lond) 2018;42(7):1275-1284.
29. Gete DG, Waller M, Mishra GD. Pre-pregnancy diet quality is associated with lowering the risk of offspring obesity and underweight: finding from a prospective cohort study. *Nutrients* 2021;13(4):1044.
 30. Gete DG, Waller M, Mishra GD. Pre-pregnancy diet quality and its association with offspring behavioral problems. *Eur J Nutr* 2021;60(1):503-515.
 31. Bao W, Tobias DK, Olsen SF, Zhang C. Pre-pregnancy fried food consumption and the risk of gestational diabetes mellitus: a prospective cohort study. *Diabetologia* 2014;57(12):2485-2491.
 32. Grieger JA, Pelecanos AM, Hurst C, Tai A, Clifton VL. Pre-conception maternal food intake and the association with childhood allergies. *Nutrients* 2019;11(8):1851.
 33. Bowers K, Tobias DK, Yeung E, Hu FB, Zhang C. A prospective study of prepregnancy dietary fat intake and risk of gestational diabetes. *Am J Clin Nutr* 2012;95(2):446-453.
 34. Lamyian M, Hosseinpour-Niazi S, Mirmiran P, Moghaddam Banaem L, Goshtasebi A, Azizi F. Pre-pregnancy fast food consumption is associated with gestational diabetes mellitus among Tehranian women. *Nutrients* 2017;9(3):216.
 35. Bao W, Bowers K, Tobias DK, Hu FB, Zhang C. Prepregnancy dietary protein intake, major dietary protein sources, and the risk of gestational diabetes mellitus: a prospective cohort study. *Diabetes Care* 2013;36(7):2001-2008.
 36. Mari-Sanchis A, Díaz-Jurado G, Basterra-Gortari FJ, de la Fuente-Arrillaga C, Martínez-González MA, Bes-Rastrollo M. Association between pre-pregnancy consumption of meat, iron intake, and the risk of gestational diabetes: the SUN project. *Eur J Nutr* 2018;57(3):939-949.
 37. Chen L, Hu FB, Yeung E, Tobias DK, Willett WC, Zhang C. Pre-pregnancy consumption of fruits and fruit juices and the risk of gestational diabetes mellitus: a prospective cohort study. *Diabetes Care* 2012;35(5):1079-1082.
 38. Bao W, Tobias DK, Hu FB, Chavarro JE, Zhang C. Pre-pregnancy potato consumption and risk of gestational diabetes mellitus: prospective cohort study. *BMJ* 2016;352:h6898.
 39. Chen X, Jiang X, Huang X, He H, Zheng J. Association between probiotic yogurt intake and gestational diabetes mellitus: a case-control study. *Iran J Public Health* 2019;48(7):1248-1256.
 40. Gaskins AJ, Rich-Edwards JW, Williams PL, Toth TL, Missmer SA, Chavarro JE. Pre-pregnancy caffeine and caffeinated beverage intake and risk of spontaneous abortion. *Eur J Nutr* 2018;57(1):107-117.
 41. Chen L, Hu FB, Yeung E, Willett W, Zhang C. Prospective study of pre-gravid sugar-sweetened beverage consumption and the risk of gestational diabetes mellitus. *Diabetes Care* 2009;32(12):2236-2241.
 42. Mohammed H, Marquis GS, Aboud F, Bougma K, Samuel A. Pre-pregnancy iodized salt improved children's cognitive development in randomized trial in Ethiopia. *Matern Child Nutr* 2020;16(3):e12943.
 43. Bao W, Song Y, Bertrand KA, Tobias DK, Olsen SF, Chavarro JE, et al. Prepregnancy habitual intake of vitamin D from diet and supplements in relation to risk of gestational diabetes mellitus: a prospective cohort study. *J Diabetes* 2018;10(5):373-379.
 44. Lawande A, Di Gravio C, Potdar RD, Sahariah SA, Gandhi M, Chopra H, et al. Effect of a micronutrient-rich snack taken pre-conceptionally and throughout pregnancy on ultrasound measures of fetal growth: the Mumbai Maternal Nutrition Project (MMNP). *Matern Child Nutr* 2018;14(1):e12441.
 45. Potdar RD, Sahariah SA, Gandhi M, Kehoe SH, Brown N, Sane H, et al. Improving women's diet quality preconceptionally and during gestation: effects on birth weight and prevalence of low birth weight—a randomized controlled efficacy trial in India (Mumbai Maternal Nutrition Project). *Am J Clin Nutr* 2014;100(5):1257-1268.
 46. Sahariah SA, Gandhi M, Chopra H, Kehoe SH, Johnson MJ, di Gravio C, et al. Body composition and cardiometabolic risk markers in children of women who took part in a randomized controlled trial of a preconceptional nutritional intervention in Mumbai, India. *J Nutr* 2022;152(4):1070-1081.
 47. Cui Y, Liao M, Xu A, Chen G, Liu J, Yu X, et al. Association of maternal pre-pregnancy dietary intake with adverse maternal and neonatal outcomes: a systematic review and meta-analysis of prospective studies. *Crit Rev Food Sci Nutr* 2021. doi: <https://doi.org/10.1080/10408398.2021.1989658>.
 48. Tieu J, Shepherd E, Middleton P, Crowther CA. Dietary advice interventions in pregnancy for preventing gestational diabetes mellitus. *Cochrane Database Syst Rev* 2017;1(1):CD006674.
 49. Stephenson J, Heslehurst N, Hall J, Schoenaker DA, Hutchinson J, Cade JE, et al. Before the beginning: nutrition and lifestyle in the preconception period and its importance for future health. *Lancet* 2018;391(10132):1830-1841.
 50. Temel S, van Voorst SF, Jack BW, Denктаş S, Steegers EA. Evidence-based preconceptional lifestyle interventions. *Epidemiol Rev* 2014;36:19-30.
 51. Hanson MA, Gluckman PD, Ma RC, Matzen P, Biesma RG. Early life opportunities for prevention of diabetes in low and mid-

- dle income countries. *BMC Public Health* 2012;12:1025.
52. Geller SE, Koch AR, Garland CE, MacDonald EJ, Storey F, Lawton B. A global view of severe maternal morbidity: moving beyond maternal mortality. *Reprod Health* 2018;15(Suppl 1):98.
 53. Balarajan Y, Ramakrishnan U, Ozaltin E, Shankar AH, Subramanian SV. Anaemia in low-income and middle-income countries. *Lancet* 2011;378(9809):2123-2135.
 54. Ryan LM, Mahmood MA, Laurence CO. Incidence of concomitant illnesses in pregnancy in Indonesia: estimates from 1990-2019, with projections to 2030. *Lancet Reg Health West Pac* 2021;10:100139.
 55. Wirawan F, Nurrika D. Maternal pre-pregnancy anemia and childhood anemia in Indonesia: a risk assessment using a population-based prospective longitudinal study. *Epidemiol Health* 2022;44:e2022100.
 56. Ahmed T, Hossain M, Sanin KI. Global burden of maternal and child undernutrition and micronutrient deficiencies. *Ann Nutr Metab* 2012;61 Suppl 1:8-17.
 57. World Health Organization. Infant mortality [cited 2022 Sep 22]. Available from: <https://www.who.int/data/gho/data/themes/topics/indicator-groups/indicator-group-details/GHO/infant-mortality>.
 58. Dhaded SM, Hambidge KM, Ali SA, Somannavar M, Saleem S, Pasha O, et al. Preconception nutrition intervention improved birth length and reduced stunting and wasting in newborns in South Asia: the Women First Randomized Controlled Trial. *PLoS One* 2020;15(1):e0218960.
 59. Young MF, Ramakrishnan U. Maternal undernutrition before and during pregnancy and offspring health and development. *Ann Nutr Metab* 2020;76(suppl 3):41-53.
 60. Shukla AK, Srivastava S, Verma G. Effect of maternal anemia on the status of iron stores in infants: a cohort study. *J Family Community Med* 2019;26(2):118-122.
 61. Yang J, Cheng Y, Zeng L, Dang S, Yan H. Maternal dietary diversity during pregnancy and congenital heart defects: a case-control study. *Eur J Clin Nutr* 2021;75(2):355-363.
 62. Dean SV, Lassi ZS, Imam AM, Bhutta ZA. Preconception care: nutritional risks and interventions. *Reprod Health* 2014;11(Suppl 3):S3.
 63. Morgane PJ, Mokler DJ, Galler JR. Effects of prenatal protein malnutrition on the hippocampal formation. *Neurosci Biobehav Rev* 2002;26(4):471-483.
 64. Rizzo TA, Metzger BE, Dooley SL, Cho NH. Early malnutrition and child neurobehavioral development: insights from the study of children of diabetic mothers. *Child Dev* 1997;68(1):26-38.
 65. Wilkins E, Wickramasinghe K, Pullar J, Demaio AR, Roberts N, Perez-Blanco KM, et al. Maternal nutrition and its intergenerational links to non-communicable disease metabolic risk factors: a systematic review and narrative synthesis. *J Health Popul Nutr* 2021;40(1):20.
 66. Greco EA, Lenzi A, Migliaccio S, Gessani S. Epigenetic modifications induced by nutrients in early life phases: gender differences in metabolic alteration in adulthood. *Front Genet* 2019;10:795.
 67. Hardy TM, Tollefsbol TO. Epigenetic diet: impact on the epigenome and cancer. *Epigenomics* 2011;3(4):503-518.
 68. Peral-Sanchez I, Hojeij B, Ojeda DA, Steegers-Theunissen RP, Willaime-Morawek S. Epigenetics in the uterine environment: how maternal diet and ART may influence the epigenome in the offspring with long-term health consequences. *Genes (Basel)* 2021;13(1):31.
 69. Acevedo N, Alashkar Alhamwe B, Caraballo L, Ding M, Ferrante A, Garn H, et al. Perinatal and early-life nutrition, epigenetics, and allergy. *Nutrients* 2021;13(3):724.
 70. World Health Organization. Diet, nutrition and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation: WHO technical report series 916; 2003 [cited 2022 Sep 22]. Available from: <https://www.who.int/publications/i/item/924120916X>.
 71. Keats EC, Rappaport AI, Shah S, Oh C, Jain R, Bhutta ZA. The dietary intake and practices of adolescent girls in low- and middle-income countries: a systematic review. *Nutrients* 2018;10(12):1978.
 72. Agustina R, Nadiya K, Andini EA, Setianingsih AA, Sadariskar AA, Prafantini E, et al. Associations of meal patterning, dietary quality and diversity with anemia and overweight-obesity among Indonesian school-going adolescent girls in West Java. *PLoS One* 2020;15(4):e0231519.
 73. Dean S, Rudan I, Althabe F, Webb Girard A, Howson C, Langer A, et al. Setting research priorities for preconception care in low- and middle-income countries: aiming to reduce maternal and child mortality and morbidity. *PLoS Med* 2013;10(9):e1001508.