



Expression of cytokines and co-stimulatory molecules in the *Toxoplasma gondii*-infected dendritic cells of C57BL/6 and BALB/c mice



Jae-Hyung Lee , Jae-Min Yuk, Guang-Ho Cha, Young-Ha Lee*

Department of Medical Science and Department of Infection Biology, Chungnam National University College of Medicine, Daejeon 35015, Korea

Abstract

Received: 30 October 2022
Accepted: 14 March 2023

*Correspondence
(yhalee@cnu.ac.kr)

Citation

Lee JH, Yuk JM, Cha GH, Lee YH. Expression of cytokines and co-stimulatory molecules in the *Toxoplasma gondii*-infected dendritic cells of C57BL/6 and BALB/c mice. Parasites Hosts Dis 2023;61(2):138-146.

Toxoplasma gondii is an intracellular protozoan parasite which can infect most warm-blooded animals and humans. Among the different mouse models, C57BL/6 mice are more susceptible to *T. gondii* infection compared to BALB/c mice, and this increased susceptibility has been attributed to various factors, including T-cell responses. Dendritic cells (DCs) are the most prominent type of antigen-presenting cells and regulate the host immune response, including the response of T-cells. However, differences in the DC responses of these mouse strains to *T. gondii* infection have yet to be characterized. In this study, we cultured bone marrow-derived DCs (BMDCs) from BALB/c and C57BL/6 mice. These cells were infected with *T. gondii*. The activation of the BMDCs was assessed based on the expression of cell surface markers and cytokines. In the BMDCs of both mouse strains, we detected significant increases in the expression of cell surface T-cell co-stimulatory molecules (major histocompatibility complex (MHC) II, CD40, CD80, and CD86) and cytokines (tumor necrosis factor (TNF)- α , interferon (IFN)- γ , interleukin (IL)-12p40, IL-1 β , and IL-10) from 3 h post-*T. gondii* infection. The expression of MHC II, CD40, CD80, CD86, IFN- γ , IL-12p40, and IL-1 β was significantly higher in the *T. gondii*-infected BMDCs obtained from the C57BL/6 mice than in those from the BALB/c mice. These findings indicate that differences in the activation status of the BMDCs in the BALB/c and C57BL/6 mice may account for their differential susceptibility to *T. gondii*.

Keywords: *Toxoplasma gondii*, dendritic cell, co-stimulatory molecule, cytokine, mouse

Introduction

Toxoplasma gondii is an obligate intracellular protozoan parasite which can infect most warm-blooded animals and humans [1]. Although to date, only a single species of this parasite has been identified, there is a marked variation in susceptibility according to its host [2-5]. Among the different mouse models, C57BL/6 mice have been determined to be more susceptible than BALB/c mice to *T. gondii* infection [2-5]. This difference in susceptibility has been ascribed to multiple factors, including genes associated with the major histocompatibility complex (MHC) [4] and the cytokine storm associated with CD4+ T-cell dependent interferon- γ (IFN- γ) [5]. However, the precise cause of the difference in susceptibility of the different mouse strains to *T. gondii* infection remains unclear, including the associated cellular immunity control.

Antigen-presenting cells (APCs) play an essential role in the process of cellular immuni-

© 2023 The Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author contributions

Conceptualization: Lee JH, Lee YH
 Data curation: Lee JH, Lee YH
 Formal analysis: Lee JH, Lee YH
 Funding acquisition: Lee YH
 Investigation: Lee JH, Yuk JM, Cha GH, Lee YH
 Methodology: Lee JH, Lee YH
 Project administration: Lee YH
 Resources: Lee YH
 Software: Lee JH
 Supervision: Yuk JM, Cha GH
 Validation: Lee JH, Yuk JM, Cha GH, Lee YH
 Visualization: Lee JH
 Writing – original draft: Lee YH

Conflict of interest

The authors declare no conflict of interest related to this study.

ORCID

Jae-Hyung Lee
 (<https://orcid.org/0009-0002-0703-9330>)
 Young-Ha Lee
 (<https://orcid.org/0000-0002-5795-8960>)

ty, among which dendritic cells (DCs), which can induce innate and adaptive immune responses, have been identified as the most important APC type in the immune system [6]. The maturation of DCs occurs when antigen fragments are presented on the cell surface of immature DCs using MHC molecules, and these cells upregulate co-stimulatory molecules, such as CD40, CD80, and CD86, thereby markedly enhancing their ability to activate T-cells [6]. *T. gondii* exploits DCs to disseminate throughout the body, produce pro-inflammatory cytokines, and presenting its antigens to the immune system, and in doing so, it is able to subvert their signaling pathways and evade detection [7]. DCs play a vital role in the resistance of murine hosts to *T. gondii*, and mice with depleted DCs have been found to be more susceptible to *T. gondii* infection [8,9]. The differences in the DC responses are attributable to the genetic susceptibility or resistance of the mouse strains to *Trypanosoma cruzi* [10], *Leishmania major* [11], and *Coccidioides posadasii* [12]. However, no information is currently regarding the activation status or immune response against *T. gondii* among the DCs derived from susceptible and resistant mouse strains.

Thus, to investigate the activation status of the DCs derived from genetically susceptible or resistant mouse strains in response to *T. gondii* infection, we cultured bone marrow-derived DCs (BMDCs) from BALB/c and C57BL/6 mice, infected these with *T. gondii*, and evaluated the expression of the cell surface T-cell co-stimulatory molecules (MHC II, CD40, CD80, and CD86) and cytokines (tumor necrosis factor (TNF)- α , IFN- γ , interleukin (IL)-12p40, IL-1 β , IL-10, and transforming growth factor (TGF)- β) based on flow cytometry and quantitative real-time polymerase chain reaction (qRT-PCR) analysis.

Materials and Methods

Survival monitoring of *T. gondii*-infected mice

C57BL/6 and BALB/c mice ($n = 5$ per group) were obtained from Damool Science, Daejeon, Korea. Each mouse was orally infected with 100 cysts of the *T. gondii* ME49 strain, and survival was monitored every day for 30 days after infection. All animal-related procedures were reviewed and approved by the Institutional Animal Care and Use Committee of Chungnam National University College of Medicine (Approval no. 202012A-CNU-200). All procedures were conducted in accordance with the relevant guidelines.

Culture and isolation of BMDCs

BMDCs were generated as previously described [9]. Briefly, the tibias and femurs of the mice were disinfected with 70% ethanol for 5 min and washed with phosphate-buffered saline (PBS). Both ends of the bone were cut off with scissors, and the bone marrow cavity was rinsed with sterile PBS using a 1 ml syringe. On day 0, the bone marrow cells were seeded at a density of 4×10^6 cells per 90×15 mm Petri dish with 10 ml of complete medium (RPMI 1640 medium supplemented with 100 U/ml penicillin and 100 μ g/ml streptomycin; WelGENE, Gyeongsan, Gyeongsangbuk-do, Korea), 2-mercaptoethanol (Gibco BRL, Grand Island, NY, USA), 10% fetal bovine serum (FBS), and granulocyte-macrophage colony stimulating factor (GM-CSF; 20 ng/ml; PeproTech, Rocky Hill, NJ, USA) in combination with IL-4 (1 ng/ml; PeproTech). On day 3, 10 ml of fresh complete medium containing GM-CSF and IL-4 was added to the Petri dishes. On day 6, non-adherent

and loosely adherent cells were collected by gently pipetting the medium against the dish plate. The collected cells were seeded at a density of 2×10^6 cells per 24-well plate and were used for subsequent experiments.

***In vitro T. gondii* infection of BMDCs**

The BMDCs were suspended in RPMI 1640 medium containing 10% FBS and antibiotics at 37°C in 5% CO₂, and infected with freshly harvested live *T. gondii* tachyzoites at a multiplicity of infection (MOI) of 1. For comparison, BMDCs were also incubated with 100 ng/ml lipopolysaccharide (LPS; Sigma, St. Louis, MO, USA). The BMDCs were treated with live *T. gondii* tachyzoites for 3, 6, 18, and 24 h, and LPS for 3 and 24 h.

Phenotypic analysis of BMDCs by flow cytometry

The BMDCs were stained with LIVE/DEAD Fixable Dead Cell Staining dye (Life Technologies, Carlsbad, CA, USA) to discriminate between the live and dead cells. The BMDCs were stained with anti-CD40 fluorescein isothiocyanate (FITC), anti-CD86 PE, anti-MHC II PerCP-Cy5.5 (BD Biosciences, San Jose, CA, USA), anti-CD80 PE-Cy7 (eBioscience, Santa Clara, CA, USA), anti-CD11c APC (Tonbo Biosciences, San Diego, CA, USA), and anti-CD11b APC-H7 (BioLegend, San Diego, CA, USA). The samples were run on a BD Canto II flow cytometer (BD Biosciences), and the data were analyzed using FlowJo software (Treestar, Ashland, OR, USA).

Gene expression of cytokines by qRT-PCR

Total RNA was extracted from the BMDCs using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) and reverse transcribed to cDNA using the ReverTra Ace RT Kit (Toyobo, Osaka, Japan). PCR was performed in triplicate on a Step One Plus instrument (Applied Biosystems, Waltham, MA, USA). The PCR reagents were obtained from Applied Biosystems, and each sample was run in triplicate in a 20 µl reaction. The PCR conditions were as follows: initial denaturation and enzyme activation at 95°C for 10 min and 40 cycles of 95°C for 15 sec, 60°C for 60 sec, and 72°C for 30 sec. The expression level of the target gene was normalized to that of the housekeeping gene (β -actin) for each sample. The PCR primers used are shown in Supplementary Table S1.

Statistical analysis

The results are expressed as the mean \pm standard deviation (SD) of at least 3 independent experiments. Significant differences between groups were determined using the two-tailed unpaired Student's *t*-test. Statistical significance was set at $P < 0.05$.

Results

Survival of *T. gondii*-infected BALB/c and C57BL/6 mice

The survival of the C57BL/6 and BALB/c mice infected with *T. gondii* was monitored over a 30 day post-infection period. All of the C57BL/6 mice had died within 9 day post-infection, whereas all of the BALB/c mice were still alive at the end of the 30 day monitoring period (Fig. 1).




Fig. 1. Survival rates of the C57BL/6 and BALB/c mice orally infected with *Toxoplasma gondii*. Each C57BL/6 or BALB/c mouse ($n = 5$ mice per group) was orally infected with 100 *T. gondii* cysts (ME49 strain), and their survival was monitored for 30 days post-infection.





Fig. 2. Gating strategy for the bone marrow-derived dendritic cell (BMDCs) analysis. Singlet cells were gated along the diagonal when the forward scatter height (FSC-H) versus forward scatter area (FSC-A) was plotted. Cells were gated when the side scatter area (SSC-A) versus the FSC-A was plotted. Live cells were live/dead Amcyan negative when the SSC-A was plotted versus the live/dead Amcyan. The CD11c⁺CD11b⁺ cells were discriminated by plotting CD11b versus CD11c. The BMDCs were defined as CD11c⁺CD11b⁺.

MHC II and co-stimulatory molecule expression was higher in the *T. gondii*-infected BMDCs from the C57BL/6 mice than from the BALB/c mice

We obtained the BMDCs from uninfected normal C57BL/6 and BALB/c mice. The BMDCs were defined as CD11c⁺CD11b⁺ cells (Fig. 2). To compare the activation status of the BMDCs from both mouse strains in response to live *T. gondii*, we initially investigated the expression of MHC II and co-stimulatory molecules (CD40, CD80, and CD86) via flow cytometry.

Although the expression of MHC II in the BMDCs derived from the BALB/c mice did not significantly differ following *T. gondii* infection, we detected an increase in MHC II expression in the C57BL/6 BMDCs in response to infection ($P=0.067$). Moreover, during the entire experimental period, the MHC II expression in *T. gondii*-infected BMDCs from the C57BL/6 mice was significantly higher than that in the BMDCs from the BALB/c mice (Fig. 3A).

In the BMDCs of both mouse strains, we detected a gradual increase in the expression of CD40, CD80, and CD86 following *T. gondii* infection, which peaked at 24 h post-infection. The expression of CD40 in the *T. gondii*-infected BMDCs from the C57BL/6 mice was found to be significantly higher than that in the *T. gondii*-infected BMDCs from the




BALB/c mice from 6 h post-infection (Fig. 3B) ($P < 0.01$). In contrast, the CD80 and CD86 expression in the *T. gondii*-infected BMDCs from the C57BL/6 mice was observed to be significantly higher than those in the infected BMDCs from the BALB/c mice from 18 h post-infection (Fig. 3C, D).

LPS promotes an increase in the expression of co-stimulatory proteins and induces the production of polarizing Th1 and Th2 cytokines/chemokines in DCs [13], thus we used LPS as a positive stimulant of the BMDCs from both mouse strains. The expression of MHC II, CD40, CD80, and CD86 in the *T. gondii*-infected BMDCs from the C57BL/6 mice was found to be significantly higher than that in the *T. gondii*-infected BMDCs from the BALB/c mice at 3 and 24 h post-infection. In the LPS-treated cells, we detected a significantly higher expression of CD40 and CD80 than in the *T. gondii*-infected cells, although this was not observed for MHC II and CD86 (Fig. 3A-D).

Cytokine expression was higher in the *T. gondii*-infected BMDCs from the C57BL/6 mice than in the BALB/c mice

We subsequently investigated the cytokine production in the *T. gondii*-infected BMDCs derived from the C57BL/6 and BALB/c mice based on qRT-PCR analysis (Fig. 4). In both the mouse strains, compared to the uninfected BMDCs, we detected a significant increase in the mRNA expression of TNF- α , IFN- γ , IL-12p40, IL-1 β , and IL-10 in the BMDCs from 3 h post-infection, with the mRNA expression of IFN- γ , IL-12p40, IL-1 β , and TGF- β in the C57BL/6 mice being significantly higher than that of the BALB/c mice (Fig. 4A-F). In



both strains, the mRNA expression of TNF- α peaked at 3 h post-infection and thereafter declined (Fig. 4A). In contrast, we detected a reduction in the TGF- β mRNA expression level in the BALB/c BMDCs in response to *T. gondii* infection, whereas a slight increase was observed in C57BL/6 BMDCs (Fig. 4F).

In the LPS-treated BMDCs, significantly higher mRNA expression levels of TNF- α , IFN- γ , IL-12p40, IL-1 β , IL-10, and TGF- β were found in the C57BL/6 mice compared to the BALB/c mice. In the case of TNF- α , IFN- γ , and TGF- β the mRNA expression levels were significantly higher in the LPS-treated cells than in the *T. gondii*-infected cells, although this was not observed for the mRNA expression levels IL-12p40, IL-1 β , or IL-10 (Fig. 4A-F).

Discussion

In this study, we investigated the activation status of the BMDCs derived from the BALB/c and C57BL/6 mice in response to *T. gondii* infection. Our findings revealed that the different responses of the BMDCs to *T. gondii* infection in these 2 inbred mouse strains which differ with respect to their susceptibility to *T. gondii* infection. We found that the survival of the C57BL/6 mice was considerably shorter than that of the BALB/c mice following infection with *T. gondii* Me49 cysts. This is consistent with previous reports indicating that C57BL/6 mice are more susceptible to *T. gondii* infection than BALB/c mice (Fig. 1) [2-5]. Having established this different response between the 2 strains, we subsequently evaluated the expression of the cell surface T-cell co-stimulatory molecules and cytokines in the BMDCs derived from the 2 mouse strains.

DCs are specialized APCs characterized by their unique capacity to initiate and modulate immune responses via their priming of naïve T-cells, the activation of which is dependent on antigens, co-stimulatory molecules, and inflammatory cytokines [6]. Upon stimulation, DCs undergo several morphological, phenotypic, and functional changes during their maturation process [6]. In this regard, a previous transcriptomic analysis of *T. gondii*-infected BMDCs revealed significant alterations in the transcripts associated with cellular metabolism, activation of T-cells, and inflammation-mediated chemokine and cytokine signaling pathways [14]. In addition, *T. gondii*-induced activation of human DCs is associated with the T-cell production of IFN- γ mediated via the CD40-CD40L-dependent release of IL-12 and through the CD80/CD86-CD28 interaction in the absence of bioactive IL-12 [15]. The findings of these studies indicate that in response to *T. gondii* infection, DCs may play a key role in the association between the innate immune response and adaptive immunity.

In this study, we analyzed the response of the BMDC cell surface markers, including MHC II and the co-stimulatory molecules CD40, CD80, and CD86 to *T. gondii* infection. We detected significant increases in the expression of these markers in the BMDCs of the C57BL/6 and BALB/c mice following *T. gondii* infection (Fig. 3). Interestingly, the expression of the cell surface markers was found to be significantly higher in the *T. gondii*-infected BMDCs of the C57BL/6 mice than in the BALB/c mice. These observations indicate that whereas *T. gondii* induces the activation of BMDCs in both mouse strains, but compared with the BALB/c BMDCs, those in the C57BL/6 mice were more effective in capturing and processing the antigens to elicit specific T-cell effector responses following *T. gondii*

infection.

Another hallmark of DC activation is a shift in the profile of the secreted cytokines. Mature DCs secrete inflammatory cytokines (IL-12, IL-1 β , and TNF- α) and anti-inflammatory cytokines (IL-10 and TGF- β) and present antigens to naïve T-cells for activation [6,16]. In this regard, as previously reported, the production of pro-inflammatory cytokines, such as IL-1 α , IL-1 β , IL-6, IL-10, and TNF- α , was significantly increased in the BALB/c mouse splenic CD11c⁺ DCs at week 1 post-infection [17]. It has been shown that in response to stimulation with *Toxoplasma* antigens or live parasites, the T-cells derived from the C57BL/6 mice preferentially produced Th1 cytokines with high IFN- γ and low IL-4 levels, whereas those from the BALB/c mice typically produced Th2 cytokines with low IFN- γ and high IL-4 levels [3,18]. Similar T-cell responses have been observed when these mouse strains were infected with *Leishmania major* [19]. Consistent with the findings of Watanabe et al. [3], we observed that the IFN- γ , IL-12p40, IL-1 β , and TGF- β mRNA expression levels were significantly higher in the BMDCs obtained from the C57BL/6 mice than in the BALB/c mice from 3 h post-infection (Fig. 4). Our findings indicate that *T. gondii* infection induces cytokine profiles polarized to Th1 and Th2 immune responses in the BMDCs of the C57BL/6 and BALB/c mice, respectively.

In conclusion, our findings in this study revealed that the differences in the activation status of the BMDCs in the BALB/c and C57BL/6 mice could be a factor contributing to the differences in the susceptibility of these 2 mouse strains to *T. gondii* infection.

Acknowledgment

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MIST) (2022R1A2B5B01001641, 2022K2A9A2A06037100).

References

- Montoya JG, Liesenfeld O. Toxoplasmosis. *Lancet* 2004;363(9425):1965-1976. [https://doi.org/10.1016/S0140-6736\(04\)16412-X](https://doi.org/10.1016/S0140-6736(04)16412-X)
- Lee YH, Kasper LH. Immune responses of different mouse strains after challenge with equivalent lethal doses of *Toxoplasma gondii*. *Parasite* 2004;11(1):89-97. <https://doi.org/10.1051/parasite/200411189>
- Watanabe H, Numata K, Ito T, Takagi K, Matsukawa A. Innate immune response in Th1- and Th2-dominant mouse strains. *Shock* 2004;22(5):460-466. <https://doi.org/10.1097/01.shk.0000142249.08135.e9>
- Johnson J, Suzuki Y, Mack D, Mui E, Estes R, et al. Genetic analysis of influences on survival following *Toxoplasma gondii* infection. *Int J Parasitol* 2002;32(2):179-185. [https://doi.org/10.1016/s0020-7519\(01\)00321-6](https://doi.org/10.1016/s0020-7519(01)00321-6)
- Liesenfeld O, Kosek J, Remington JS, Suzuki Y. Association of CD4+ T cell-dependent, interferon-gamma-mediated necrosis of the small intestine with genetic susceptibility of mice to peroral infection with *Toxoplasma gondii*. *J Exp Med* 1996;184(2):597-607. <https://doi.org/10.1084/jem.184.2.597>
- Liu K, Nussenzweig MC. Origin and development of dendritic cells. *Immunol Rev* 2010;234(1):45-54. <https://doi.org/10.1111/j.0105-2896.2009.00879.x>
- Sanecka A, Frickel EM. Use and abuse of dendritic cells by *Toxoplasma gondii*. *Virulence* 2012;3(7):678-869. <https://doi.org/10.4161/viru.22833>
- Liu CH, Fan YT, Dias A, Esper L, Corn RA, et al. Cutting edge: dendritic cells are essential for in vivo IL-12 production and development of resistance against *Toxoplasma gondii* infection in mice. *J Immunol* 2006;177(1): 31-35. <https://doi.org/10.4049/jimmunol.177.1.31>
- Mashayekhi M, Sandau MM, Dunay IR, Frickel EM, Khan A, et al. CD8 α (+) dendritic cells are the critical source of interleukin-12 that controls acute infection by *Toxoplasma gondii* tachyzoites. *Immunity* 2011;35(2):249-259. <https://doi.org/10.1016/j.immuni.2011.08.008>
- Planelles L, Thomas MC, Marañón C, Morell M, López MC. Differential CD86 and CD40 co-stimulatory molecules and cytokine expression pattern induced by *Trypanosoma cruzi* in

- APCs from resistant or susceptible mice. *Clin Exp Immunol* 2003;131(1):41-47. <https://doi.org/10.1046/j.1365-2249.2003.02022.x>
11. von Stebut E, Ehrchen JM, Belkaid Y, Kostka SL, Molle K, et al. Interleukin 1alpha promotes Th1 differentiation and inhibits disease progression in *Leishmania major*-susceptible BALB/c mice. *J Exp Med* 2003;198(2):191-199. <https://doi.org/10.1084/jem.20030159>
 12. Awasthi S, Magee DM. Differences in expression of cell surface co-stimulatory molecules, Toll-like receptor genes and secretion of IL-12 by bone marrow-derived dendritic cells from susceptible and resistant mouse strains in response to *Coccidioides posadasii*. *Cell Immunol* 2004;231(1-2):49-55. <https://doi.org/10.1016/j.cellimm.2004.11.006>
 13. Neves BM, Cruz MT, Francisco V, Garcia-Rodriguez C, Silvestre R, et al. Differential roles of PI3-Kinase, MAPKs and NF-kappaB on the manipulation of dendritic cell T(h)1/T(h)2 cytokine/chemokine polarizing profile. *Mol Immunol* 2009;46(13):2481-2492. <https://doi.org/10.1016/j.molimm.2009.05.021>
 14. Hargrave KE, Woods S, Millington O, Chalmers S, Westrop GD, et al. Multi-omics studies demonstrate *Toxoplasma gondii*-induced metabolic reprogramming of murine dendritic cells. *Front Cell Infect Microbiol* 2019;9:309. <https://doi.org/10.3389/fcimb.2019.00309>
 15. Subauste CS, Wessendarp M. Human dendritic cells discriminate between viable and killed *Toxoplasma gondii* tachyzoites: dendritic cell activation after infection with viable parasites results in CD28 and CD40 ligand signaling that controls IL-12-dependent and -independent T cell production of IFN-gamma. *J Immunol* 2000;165(3):1498-1505. <https://doi.org/10.4049/jimmunol.165.3.1498>
 16. Bazan SB, Walch-Rückheim B, Schmitt MJ, Breinig F. Maturation and cytokine pattern of human dendritic cells in response to different yeasts. *Med Microbiol Immunol* 2018;207(1):75-81. <https://doi.org/10.1007/s00430-017-0528-8>
 17. Nam HW, Ahn HJ, Yang HJ. Pro-inflammatory cytokine expression of spleen dendritic cells in mouse toxoplasmosis. *Korean J Parasitol* 2011;49(2):109-114. <https://doi.org/10.3347/kjp.2011.49.2.109>
 18. Mukhopadhyay D, Arranz-Solís D, Saeij JPJ. Influence of the host and parasite strain on the immune response during *Toxoplasma* infection. *Front Cell Infect Microbiol* 2020;10:580425. <https://doi.org/10.3389/fcimb.2020.580425>
 19. Iborra S, Abánades DR, Parody N, Carrión J, Risueño RM, et al. The immunodominant T helper 2 (Th2) response elicited in BALB/c mice by the *Leishmania* LiP2a and LiP2b acidic ribosomal proteins cannot be reverted by strong Th1 inducers. *Clin Exp Immunol* 2007;150(2):375-385. <https://doi.org/10.1111/j.1365-2249.2007.03501.x>