



ARTICLE

Preparation of Hypoallergenic Whey Protein Hydrolysate by a Mixture of Alcalase and Prozyme and Evaluation of Its Digestibility and Immunoregulatory Properties

Jiyeon Yang^{1,2,†}, Se Kyung Lee^{1,†}, Young Suk Kim³, Hyung Joo Suh^{1,4}, and Yejin Ahn^{1,*}

OPEN ACCESS

Received May 3, 2023

Revised May 15, 2023

Accepted May 16, 2023

*Corresponding author : Yejin Ahn
Department of Integrated Biomedical and Life Science, Graduate School, Korea University, Seoul 02841, Korea
Tel: +82-2-940-2764
Fax: +82-2-940-2850
E-mail: ahnyj708@gmail.com

*ORCID
Jiyeon Yang
<https://orcid.org/0000-0003-1842-1631>
Se Kyung Lee
<https://orcid.org/0009-0001-5394-9653>
Young Suk Kim
<https://orcid.org/0009-0007-1821-3695>
Hyung Joo Suh
<https://orcid.org/0000-0001-8869-3929>
Yejin Ahn
<https://orcid.org/0000-0002-7016-9798>

† These authors contributed equally to this work.

¹Department of Integrated Biomedical and Life Science, Graduate School, Korea University, Seoul 02841, Korea

²Maeil Innovation Center, Maeil Dairies Co., Ltd., Pyeongtaek 17714, Korea

³Department of Food and Nutrition, Ansan University, Ansan 15328, Korea

⁴Transdisciplinary Major in Learning Health Systems, Department of Healthcare Sciences, Graduate School, Korea University, Seoul 02841, Korea

Abstract Whey protein (WP) has nutritional value, but the presence of β -lactoglobulin (β -LG) and α -lactalbumin (α -LA) cause allergic reactions. In this study, hypoallergenic whey protein hydrolysate (HWPH) was prepared by decomposing β -LG and α -LA of WP using exo- and endo-type proteases. The enzyme mixing ratio and reaction conditions were optimized using response surface methodology (RSM). Degradation of α -LA and β -LG was confirmed through gel electrophoresis, and digestion, and absorption rate, and immunostimulatory response were measured using *in vitro* and *in vivo* systems. Through RSM analysis, the optimal hydrolysis conditions for degradation of α -LA and β -LG included a 1:1 mixture of Alcalase and Prozyme reacted for 10 h at a 1.0% enzyme concentration relative to substrate. The molecular weight of HWPH was <5 kDa, and leucine was the prominent free amino acid. Both *in vitro* and *in vivo* tests showed that digestibility and intestinal permeability were higher in HWPH than in WP. In BALB/c mice, as compared to WP, HWPH reduced allergic reactions by inducing elevated Type 1/Type 2 helper T cell ratio in the blood, splenocytes, and small intestine. Thus, HWPH may be utilized in a variety of low allergenicity products intended for infants, adults, and the elderly.

Keywords hydrolysate, whey protein, response surface methodology, hypoallergenic, infant nutrition

Introduction

Whey, a by-product resulting from the production of cheese, is a generic term for the remaining water-soluble components other than cheese (milk concentrate curd). With

the exception of casein, which is involved in cheese production, whey is valued highly nutritionally and physiologically because of the presence of active ingredients including proteins, lactose, minerals, vitamins, and inorganic components (Boscaini et al., 2023; Yiğit et al., 2023).

Whey protein (WP) not only boosts immunity due to the presence of β -lactoglobulin (β -LG), α -lactalbumin (α LA), bovine serum albumin, lactoferrin (LF), immunoglobulin (Ig), enzymes, and glycomacropeptide, and is also a source of essential amino acids (Marshall, 2004). Though it is obvious that WP is an excellent nutritive food, some of its components have been linked to allergic conditions in infants and young children (Jaiswal and Worku, 2022; Wright et al., 2022). Among the above mentioned proteins, the major antigens include casein, β -LG, and α LA, and it has been reported that most patients allergic to milk allergy have specific IgE antibodies against at least two antigens (Cohen et al., 2022; Savilahti and Kuitunen, 1992; Sicherer and Sampson, 1999). Among WPs, β -LG is known to be more allergenic than α LA. In human milk, in particular, β -LG is present in smaller amounts ranging from 1–150 $\mu\text{g}/\text{mL}$ (Exl and Fritsché, 2001), and its content is lower than that of casein, another allergen of milk, which comprises approximately 7% to 12% of milk protein.

To lessen milk allergies, β -LG is hydrolyzed using digestive enzymes, and hypoallergenic milk powder prepared using proteolytic enzymes is commercially available (Lynch and Buckin, 2022). To reduce milk allergic reactions, it is necessary to provide a diet that restricts the causative milk or provided in a form that lowers the antigenicity of milk. Different treatment methods, such as heat, ultra-high pressure (Landim et al., 2023), and enzymes, can be used to reduce the antigenicity of proteins. Because one of the main allergens in milk, casein, is heat-resistant, the reduction in antigenicity by heat treatment is insignificant (Bu et al., 2013; Vandenplas et al., 2022). Thus, milk proteins containing both structural and sequential epitopes are mainly hydrolyzed by enzymes. Although it doesn't exist in human milk and has a lower content than α LA, the primary allergen of WP, β -LG, shows resistance during digestion in the stomach and penetrates the intestines to cause milk allergies.

Therefore, this study aimed to prepare WP hydrolysates (WPH) with reduced antigenicity of α -LA and β -LG allergens as target substrates. Optimal hydrolysis conditions for degrading antigenic WPs using a mixture of endo- and exo-type enzymes were selected using response surface methodology (RSM). By preparing a hypoallergenic whey protein hydrolysate (HWPH), we developed a milk protein manufacturing technology that can be fed to milk-allergic infants. The digestibility and immune stimulation of the prepared hydrolysate were also evaluated.

Materials and Methods

Materials

In this study, whey protein concentrate (WPC; Agri-mark, Coral Gables, FL, USA) with a protein content of 80% (dry basis) was used as the raw WP. A 75% α -LA and a 95% β -LG from bovine milk (Samyoung Innovation, Pyeongtaek, Korea) were used as the allergen standards. Alcalase (Novozymes, Copenhagen, Denmark), protamex (Novozymes), flavourzyme (Novozymes), and collupulin (DSM, Heerlen, Netherlands) were employed as endo-type proteolytic enzymes, whereas prozyme (DuPont, Wilmington, DE, USA) was used as an exo-type enzyme. Pepsin and pancreatin (Sigma-Aldrich, St. Louis, MO, USA) were used as the artificial digestive enzymes. L-leucine was purchased from Sigma-Aldrich.

Preparation of whey protein hydrolysate

To prepare the whey protein hydrolysate, a mixture of endo- and exo-type protease enzymes was added to a 10% suspension of WP at a concentration of 1.0% (v/w) relative to the substrate solution. This was then hydrolyzed for 8 h at 50°C

at pH 7.8 with shaking at 70×g. Each reaction solution was treated at 90°C for 10 min to inactivate protease and then centrifuged at 5,500×g for 20 min to separate the supernatant. A WP solution without protease treatment was used as a control. The reaction conditions and enzyme characteristics used for WP hydrolysis are listed in Supplementary Table S1.

Identification of β -lactoglobulin (β -LG) and α -lactalbumin (α -LA) in whey protein hydrolysate using sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE)

HWP (10 mg) was dissolved in 1 mL of sample buffer [0.5 M trihydrochloric acid (HCl), glycerol, 10% SDS, 0.1% bromophenol blue, pH 6.8], heat-treated at 100°C for 5 min, centrifuged, and the supernatant separated. A 20 μ L of this was run on a 12% polyacrylamide gel at 135 V for approximately 90 min. To confirm the presence of α -LA and β -LG proteins, the gel was stained with Coomassie Brilliant Blue R-250 for 6 h, and then destained using a solution containing 10% methanol and 10% acetic acid until the protein bands were clearly visible.

Amino-nitrogen (A-N) contents and degree of hydrolysis (DH) calculation

The A-N content of HWP was measured using the 2,4,6-trinitrobenzenesulfonic acid (TNBS) method (Nielsen et al., 2001). Briefly, 125 μ L of the sample, 400 μ L of 0.212 M phosphate buffer (pH 8.2), and 0.4 mL of 0.1% TNBS were mixed and incubated in the dark at 50°C for 60 min. Next, 0.8 mL of 100 mM HCl was added and left for 20 min. After another 10 min, 0.8 mL of distilled water was added, and the absorbance was measured at 340 nm. The amount of A-N was calculated using L-leucine as a standard.

The DH of proteins following enzyme treatment was calculated using the following equation by estimated the A-N content of whey protein hydrolysates (Benjakul and Morrissey, 1997). The total A-N content was measured by neutralizing WP with sodium hydroxide (NaOH) after hydrolysis at 100°C for 24 h using 6 N HCl.

$$DH = (L_t - L_0) / (L_{\max} - L_0) \times 100 \quad (1)$$

Where L_t is the A-N content after hydrolysis at time t , while L_0 is the A-N content before hydrolysis, and L_{\max} is the A-N content after acid hydrolysis with 6 N HCl.

Establishment of optimal mixing ratio of Alcalase and Prozyme and of optimal reaction conditions using response surface methodology (RSM)

The RSM was used to select the optimal mixing ratio of Alcalase and Prozyme. A total of 15 combinations were composed of three repetitions of the experiment at 10% substrate, 50°C, pH 7.8, with the mixing ratio of the two enzymes, Alcalase and Prozyme, as an independent variable (Supplementary Table S2). The optimal mixing ratio of the two enzymes was selected by measuring the A-N of the hydrolysate corresponding to the experimental combination. RSM was used to analyze the experimental results, and the regression equation representing the optimal mixing ratio of Alcalase and Prozyme was as follows:

$$aX + bY + cXY = 32.29x + 68.87y + 96.58xy \quad (2)$$

where X is Alcalase (%) and Y is Prozyme (%).

A total of 39 combinations were composed of three repetitions of the experiment at 10% substrate, 50°C, pH 7.8, with the amount of enzyme added and hydrolysis time by the mixed enzyme of Alcalase and Prozyme as independent variables (Supplementary Table S3). The A-N of the hydrolysate corresponding to the experimental combination was measured to select the optimal enzyme addition amount and hydrolysis time. RSM was used to analyze the experimental results, and the regression equation representing the optimal conditions for the amount of enzyme added and hydrolysis time was as follows:

$$A + aX + bY + cX^2 + dY^2 + eXY = 29.6286 - 0.9220 X + 4.5520 Y + 0.4706 X^2 + 8.3643 Y^2 - 2.0389 XY \quad (3)$$

where X is the hydrolysis time and Y is the amount of the enzyme added.

Amino acid composition by high-performance liquid chromatography (HPLC) analysis

Amino acids were derivatized using AccQ-Tag reagent (Waters, Milford, MA, USA) according to the manufacturer's instructions and then analyzed using an HPLC system (Waters). An AccQ-Tag column (3.9 mm×150 mm, Waters) was used with the following parameters: temperature, 37°C; flow rate of the mobile phase [Water AccQ-Tag Eluent A (acetate-phosphate buffer), acetonitrile, and Milli-Q Water], 1.0 mL/min, and a fluorescence detector (250 nm of excitation and 395 nm of emission) was used to analyze the data.

The analysis for total amino acids was performed after acid hydrolysis. Briefly, 10 mL of 6 N HCl solution was added to 0.5 g of the samples taken in a test tube, which was sealed and hydrolyzed at 110°C for 24 h. The filtrate obtained was centrifuged, and the supernatant concentrated at 50°C to completely evaporate the acid and water, and then diluted to 5 mL using 20 mM HCl (pH 2.2). The dissolved solution was filtered through a 0.45 µm membrane, and the filtrate was used as a sample for HPLC analysis.

***In vitro* digestibility and intestinal permeability assay**

In vitro digestibility was determined as described by Garrett et al. (1999) using pepsin and pancreatin. After dissolving 1,000 mg of the sample in distilled water, the pH was adjusted to 2.0. Pepsin (2.5% of the substrate) was added, reacted at 37°C for 1 h, and the pH was then adjusted to 5.3 with sodium bicarbonate and further adjusted to 7.5 with 5 N NaOH. Pancreatin was added at 4% per substrate, reacted at 37°C for 2 h, and was then heated at 90°C for 10 min to inactivate the digestive enzyme. A-N (mg) was measured in the supernatant obtained by centrifugation (14,000×g, 10 min), and digestibility (%) was calculated using the following formula:

$$\text{Digestibility (\%)} = \frac{[\text{A-N in digested sample (mg)} - \text{A-N in sample before digestion (mg)}]}{[\text{Total A-N in sample before digestion (mg)} - \text{A-N in sample before digestion (mg)}]} \times 100 \quad (4)$$

Total A-N in sample before digestion: A-N content after acid hydrolysis.

A Caco-2 Ready 24-well plate (Komabiotec, Seoul, Korea) was used for the permeability analysis of WP and HWP (van Breemen and Li, 2005). The plate was incubated in a 5% CO₂ incubator (37°C) for 4 h, thawed, and then replaced with Dulbecco's Modified Eagle's Medium containing 1 g/L glucose, 10% fetal bovine serum, and 1% glutamine. The assay was conducted 48 h after the medium was replaced. For the intestinal permeability analysis, WP and HWP prepared using

digestive enzymes (pepsin and pancreatin) were diluted 10-fold with HBSS buffer. After washing the Caco-2 plate with a complete monolayered film three times using HBSS buffer, 250 and 750 μL of the diluted sample and HBSS buffer were dispensed into the insert and receiver plates, respectively. The buffer on the receiver plate was recovered over time (20, 40, 60, and 120 min). After analyzing the protein content of the sample initially distributed in the insert and the sample collected over time using the bicinchoninic acid assay, the permeability of the sample was analyzed using the following formula (Park et al., 2021):

$$\text{Permeability (\%)} = [\text{Total protein (mg) in the receiver plate}] / [\text{Initial protein (mg) added to the insert}] \times 100 \quad (5)$$

Evaluation of absorption rate by high-dose single oral administration

Sprague-Dawley (SD) rats (6-week-old, male) were purchased from Oriental Bio (Seongnam, Korea). The animals were housed in a breeding room at $21 \pm 1^\circ\text{C}$, a relative humidity of 50%–55%, and a light-dark cycle of 12 h. Drinking water and feed were provided *ad libitum*. After acclimation for a week, four SD-rats were randomly assigned per experimental group. WP and HWPB were orally administered at 500 mg/kg, sacrificed according to the sample administration times (20, 40, 60, and 120 min), and blood was collected. Blood samples were collected from the control group at 0 min. Following centrifugation ($2,000 \times g$, 15 min, 4°C), serum was collected to measure the total A-N content. All animal experiments were approved by the Korea University Animal Experiment Ethics Committee (KUIACUC-2022-0095).

Molecular weight distribution of hydrolysates

To measure the molecular weight distribution of the hydrolysate using HPLC, the sample was filtered (microfilter, 0.45 μm) and 20 μL was injected into a Superdex G-75 column (10×300 mm, GE Healthcare, Anaheim, CA, USA). Subsequently, absorbance was measured at 220 nm, and the column was eluted with 50 mM ammonium formate buffer (pH 5.5; 0.5 mL/min). The molecular weight distribution of the hydrolysate was measured according to the molecular weight distribution curve prepared using glutathione (MW 307), aprotinin (MW 6,512), cytochrome c (MW 12,384), enolase (MW 67,000), lactate dehydrogenase (MW 142,000), and glutamate dehydrogenase (MW 290,000) as standard proteins, which were purchased from Sigma-Aldrich.

Evaluation of immunological responses of hypoallergenic hydrolysates

A specific pathogen free male BALB/c mice aged 8 weeks were purchased from Daehan Biolink (Eumseong, Korea) and experimented under aseptic management (breeding room temperature, 25°C ; humidity, 55%; and sterile distilled water supply). Mice were divided into groups administered 250 and 500 mg/kg of WP and HWPB, respectively, and a control group that was administered isotonic sodium chloride solution. Eight mice were included in each group. Gastric intubation was performed five times per week for 4 weeks. Body weight was measured twice a week at 3-day intervals. After administration, blood was collected and centrifuged at $2,000 \times g$ for 10 min to separate the serum.

After sacrifice, aseptically excised splenocytes for cell culture were dispensed into 24 well-culture plates and cultured in a CO_2 incubator for 48 h. Subsequently, phytohemagglutinin (Sigma-Aldrich, $5 \mu\text{g/L} \times 10^6$ cells) was added as a polyclonal stimulator for *in vitro* activation. The concentration of cytokines (INF- γ and IL4), antibodies (IgG2a and IgG1) and IgE in the cell culture medium were measured using an ELISA kit purchased from BD Bioscience (San Diego, CA, USA). For the small

intestine cytokines, 2 mL of PBS was added to 0.5 g of duodenum aseptically extracted after sacrifice, vortexed, and left at 4°C for 2 days. Thereafter, the supernatant obtained by centrifugation was analyzed for cytokines and IgE using the ELISA kits.

Statistical analysis

Data were statistically analyzed using SPSS (ver. 12.0; SPSS, Chicago, IL, USA), and the mean and SD were calculated for all measured values. The significance between the experimental groups was tested by ANOVA, followed by Tukey's multiple range test at $p < 0.05$ level. RSM analysis was performed using Minitab 17 software (Minitab, State College, PA, USA) to confirm the suitability of the model to understand the relationship between the independent and dependent variables.

Results and Discussion

Selection of enzyme combinations for the preparation of hypoallergenic whey protein hydrolysate (HWPH)

WP accounts for approximately 20% of the total milk protein. β -LG and α -LA are the major allergens, and serum albumin, Ig, and LF also act as allergens (Hochwallner et al., 2014). α -LA and β -LG are more effective in reducing allergic reactions using proteolytic enzymes due to their acid-resistant and heat-resistant structures (Stănciuc and Rapeanu, 2010; Yuan et al., 2018). Proteolytic enzymes under mild conditions convert proteins to shorter chain peptides and amino acids, yielding minimal byproducts. It is documented that enzymatic protein hydrolysis can alter the protein structure without compromising its nutritional value, thereby reducing the functional properties or allergens (Cui et al., 2023; Kheroufi et al., 2022).

To select enzymes that could degrade α -LA and β -LG, endo-type enzymes (Alcalase, Collupulin, and Protamex) and exo-type enzymes (Prozyme) were mixed, and hydrolysis was performed for 8 h. A combination of enzymes in which α -LA and β -LG were degraded was selected by SDS-PAGE. Following electrophoresis, the α -LA and β -LG bands were not seen in the hydrolysate prepared using the mixture of Alcalase and Prozyme, revealing that the two allergens were decomposed (Supplementary Fig. S1). Fig. 1 shows the A-N and DH in the hydrolysate obtained after treatment with the enzyme mixture. The levels of A-N and DH in the HWPHs obtained from by mixing Alcalase and Prozyme were significantly higher than those in the other hydrolysates. Thus, this mixture was the most suitable enzyme combination for the hydrolysis of α -LA and




Fig. 1. Amino-nitrogen (A-N) (A) and degree of hydrolysis (DH) (B) of whey protein hydrolysate by mixing endo- and exo-types protease. Values are mean \pm SD for each group (n=3). ^{a-d} Different letters represent significant difference at $p < 0.05$ based on Tukey's multiple range test.

β -LG.

Proteolytic enzymes have different properties depending on the order of hydrolysis and the combination of endo-type enzymes that hydrolyze proteins roughly and exo-type enzymes that degrade proteins from the end (Villanueva et al., 1999). Alcalase, a serine protease (endo-type), cleaves the internal chains of proteins and peptides as well as peptide bonds following Glu, Met, Leu, Tyr, Lys, and Gln (Adamson and Reynolds, 1996). Prozymes are fungal exo-proteases from *Aspergillus oryzae* suitable for producing HWP (Suh et al., 2017). Among the hydrolysates prepared using exo- and endo-type enzymes, α -LA and β -LG were not detected in the hydrolysates prepared using Alcalase and Prozyme, and their A-N content and DH were the highest (Fig. 1 and Supplementary Fig. S1).

Mixing ratio of Alcalase and Prozyme for the preparation of hypoallergenic whey protein hydrolysate (HWP)

According to the 15 combinations recommended by the RSM, hydrolysates were prepared to choose the appropriate mixing ratio of Alcalase and Prozyme for the production of HWP (Supplementary Table S2). DH was calculated by measuring the A-N content of the hydrolysate (Fig. 2). The A-N content and DH of the hydrolysate obtained by mixing Alcalase and Prozyme at 1:1 and 1:3, respectively, were significantly higher than those of the hydrolysate obtained at other mixing ratios ($p < 0.05$). The regression equation for the A-N content obtained from the multiple regression analysis is as follows:




Fig. 2. A-N content (A, C) and DH (B, D) of whey protein hydrolysate optimized using response surface methodology (RSM). (A, B) RSM conditions according to the mixing ratio of Alcalase and Prozyme. Whey protein hydrolysis was performed using a mixture of Alcalase and Prozyme in 10% substrate for 8 h at 50°C and pH 7.8. (C, D) RSM conditions according to the addition amount and reaction time of the mixed enzymes. Values are mean \pm SD for each group ($n=3$). ^{a-d} Other letters are significantly different at $p < 0.05$ by Tukey's multiple range test. A-N, amino-nitrogen; DH, degree of hydrolysis.

$$\text{A-N (mg/mL)} = 32.29 x + 68.87 y + 96.58 xy \quad (6)$$

x: Alcalase (%), y: Prozyme (%), $r^2 = 97.54\%$.

As a result of predicting the optimal ratio of an enzyme mixture using RSM, when Alcalase and Prozyme were mixed 1:1, the A-N content was expected to be the highest at 50.42 mg/mL. The A-N content of the hydrolysate produced by Alcalase and Prozyme mixed in a 1:1 ratio, as predicted by RSM, was 51.52 mg/mL, and the actual measured value (50.42 mg/mL) was close to this predicted value. Thus, for the preparation of HWPH, the combination of Alcalase and Prozyme at 0.5/0.5 or 0.25/0.75 was most appropriate (Fig. 2).

Hydrolysis conditions such as temperature, pH, and enzyme–substrate ratio also affect protein hydrolysis (Diniz and Martin, 1997; Vieira et al., 1995). When enzymes are to be mixed, the mixing ratio plays an important role in determining the hydrolysate's properties. The hydrolysate produced by the mixture of Alcalase and Prozyme at a ratio of 1:1 showed the highest A-N and DH (Fig. 2).

Optimization of enzyme amount and hydrolysis time for the preparation of hypoallergenic whey protein hydrolysate (HWPH) by Alcalase-Prozyme mixture

To select the optimal enzyme concentration and the hydrolysis time for HWPH production by the Alcalase-Prozyme mixture, the A-N content of the hydrolysate was measured according to the 39 combinations suggested by the RSM (Supplementary Table S3). The regression equation for the A-N content obtained from the multiple regression analysis is as follows:

$$\text{A-N (mg/mL)} = 0.471 X^2 + 8.364 Y^2 - 2.039 XY - 0.922 X + 4.552 Y + 29.62 \quad (7)$$

X: hydrolysis time (h), Y: enzyme addition amount (%), $r^2 = 96.80\%$.

With an increase in the amount of enzyme and the hydrolysis time, the A-N content also increased (Fig. 2). The highest amount of A-N was observed when hydrolysis was performed for 10 h with the addition of 1.0% enzyme compared to the substrate. The expected amount of A-N at the optimal enzyme concentration and hydrolysis time predicted by RSM was 60.00 mg/mL, and the actual measured value of 59.58 mg/mL was close to this, with the r^2 value being 0.968. To produce a HWPH by the Alcalase-Prozyme mixture, hydrolysis with 1.5% enzyme addition was found to be most suitable compared to the substrate for 8 h.

Under the optimal reaction conditions of the WPH to produce low allergens, when WP and allergen-inducing substances α -LA and β -LG were hydrolyzed as substrates, the decomposition of these substances was confirmed through gel electrophoresis (Fig. 3). The bands for these allergens appeared dark before hydrolysis (0 h), and faded after 4 h of hydrolysis. After an 8 h hydrolysis, almost no bands were visible, confirming that α -LA and β -LG were decomposed by the mixed enzyme treatment.

Amino acids composition of hypoallergenic whey protein hydrolysate (HWPH)

The amino acid composition of HWPH treated with Alcalase and Prozyme enzymes (Table 1) revealed the composition of




Fig. 3. Bands of α -lactalbumin (α -LA) and β -lactoglobulin (β -LG) as seen in gel electrophoresis according to hydrolysis time of whey protein (WP).

Table 1. Composition of amino acids in hypoallergenic whey protein hydrolysates

Contents (μ mole/g)	Free amino acids	Total amino acids
Aspartic acid (Asp)	46.96 \pm 0.33	516.35 \pm 4.63
Serine (Ser)	151.70 \pm 0.86	308.41 \pm 2.31
Glutamine (Glu)	169.00 \pm 2.30	720.73 \pm 9.91
Glycine (Gly)	25.27 \pm 0.90	182.34 \pm 3.57
Histidine (His)	61.55 \pm 4.06	95.91 \pm 3.63
Arginine (Arg)	79.38 \pm 0.91	394.24 \pm 21.16
Threonine (Thr)	57.53 \pm 28.23	72.17 \pm 13.46
Alanine (Ala)	48.65 \pm 35.25	288.05 \pm 5.65
Proline (Pro)	100.55 \pm 2.67	388.43 \pm 6.84
Cysteine (Cys)	10.91 \pm 1.14	282.89 \pm 8.71
Tyrosine (Tyr)	73.76 \pm 0.93	93.46 \pm 2.77
Valine (Val)	288.57 \pm 6.28	396.82 \pm 4.46
Methionine (Met)	66.35 \pm 0.28	78.32 \pm 2.07
Lysine (Lys)	234.71 \pm 1.66	308.88 \pm 5.78
Isoleucine (Ile)	182.55 \pm 3.47	249.27 \pm 5.98
Leucine (Leu)	351.73 \pm 4.62	469.35 \pm 4.30
Phenylalanine (Phe)	140.56 \pm 3.47	141.59 \pm 0.99
Total	2,089.72 \pm 76.18	4,987.21 \pm 93.44

Values are mean \pm SD for each group (n=3).

free amino acids as $2,089.72 \pm 76.18$ $\mu\text{mole/g}$ and that of total amino acids as $4,987.21 \pm 93.44$ $\mu\text{mole/g}$. In all, 17 types of free amino acids were detected, with leucine being the most prominent (16.8%, 351.73 ± 4.62 $\mu\text{mole/g}$), followed by valine (13.8%), lysine (11.2%), and isoleucine (8.7%). As high levels of these amino acids cause bitterness, the bitterness in the sample was consistent with their presence.

Digestibility and intestinal permeability by Caco-2 cell

Changes in A-N content and digestibility were measured when WP and HWPH were treated with the artificial digestive enzymes pepsin and pancreatin (Figs. 4A and B). As the digestion of pepsin and pancreatin progressed, the A-N content and digestibility tended to increase. WP and HWPH were more hydrolyzed by pancreatin treatment, and the increase in A-N content and digestibility was significantly higher than that by pepsin treatment. The amount of A-N at each stage of the digestive enzyme treatment was significantly higher in HWPH than in WP (Fig. 4A). As for the change in digestibility, HWPH, which had a high A-N content before hydrolysis, showed lower digestibility than WP in the pepsin treatment stage but a higher level of digestibility increase than WP in the pancreatin treatment stage was noted (Fig. 4B).

By measuring the intestinal permeability of Caco-2 cells (Figs. 4C, D, E, and F), it was confirmed that the permeability of HWPH was significantly higher than that of the WP sample at the indicated treatment time points of 20, 40, 60, and 120 min. The intestinal permeability of Caco-2 cells at each stage of treatment with pepsin and pancreatin was also significantly higher in HWPH than in WP. Thus, it was confirmed that the digestibility and intestinal permeability of HWPH determined by Alcalase and Prozyme were higher than those of WP.

The size of the peptides constituting the protein also affects the digestibility and absorption. Peptides of ≤ 500 Da have higher bioavailability than peptides of $\geq 2,000$ Da (Feng and Betti, 2017; Wang and Li, 2017). Hydrolysates composed of small peptides are absorbed into the bloodstream through the intestinal wall and confer their effects (Chatterjee et al., 2018). WP and HWPH are converted into smaller peptides by pepsin and pancreatin and absorbed through the intestinal mucosa. During *in vitro* digestion, pancreatin treatment resulted in higher A-N content and digestibility than pepsin treatment (Fig. 4).

Evaluation of absorption rate of whey protein (WP) and hypoallergenic whey protein hydrolysate (HWPH) in Sprague-Dawley (SD) rats

Changes in the amount of A-N in the blood were measured after the oral administration of WP and HWPH to SD rats (Fig. 5). Twenty min after oral administration, the highest amount of A-N was observed in the blood, which decreased gradually thereafter. After 20 and 40 min, as compared to WP, the administration of HWPH resulted in a significantly higher amount of A-N in the blood ($p < 0.001$). HWPH, a HWPH produced by Alcalase-Prozyme, is composed of smaller peptides than WP and is easily digested and absorbed; therefore, the amount of A-N in the blood may have been high at the beginning of oral administration.

Pancreatin contains endopeptidases (trypsin and chymotrypsin) and exopeptidases (carboxypeptidase; Andriamihaja et al., 2013), which can hydrolyze WPs into smaller peptides than pepsin. Since β -LG is resistant to digestive enzymes (Villas-Boas et al., 2015), HWPH, in which β -LG is decomposed by Alcalase/Prozyme treatment, showed higher digestibility than WP. The measurement of the absorption rate through Caco-2 cells and animal experiments (Figs. 4 and 5) showed that the hydrolysate (HWPH) had better permeability than before hydrolysis (WP). Animal experiments showed a difference in the absorption rate within 40 min of oral administration.




Fig. 4. Changes in (A) A-N content, (B) digestibility, and (C, D, E, F) intestinal permeability according to time in Caco-2 cells exposed to WP and hypoallergenic whey protein hydrolysate (HWP) treated with *in vitro* digestive enzymes. Values are mean±SD for each group (n=3). ^{a-c} Different letters represent significant differences at p<0.05 based on Tukey’s multiple range test. *** p<0.001 for HWP as compared to WP. WP, whey protein; ns, not significant; A-N, amino-nitrogen.

Molecular weight distribution of hypoallergenic whey protein hydrolysate (HWP)

The molecular weight distributions of the WP and HWP were determined using a Superdex G-75 column (Fig. 6). WP was noted to contain several peptides with molecular weights of 18 and 5.8 kDa. In contrast, HWP produced by Alcalase




Fig. 5. A-N changes in serum of Sprague-Dawley (SD) rats following oral administration of WP and HWPH before and after hydrolysis of WP with mixed enzymes. Values are mean \pm SD for each group (n=4). ^{a-d} Different letters represent significant difference at p<0.05 based on Tukey's multiple range test. *** p<0.001 for HWPH compared to WP. WP, whey protein; HWPH, hypoallergenic whey protein hydrolysate; ns, not significant; A-N, amino-nitrogen.




Fig. 6. Molecular weight distribution of WP (A, B, C) and HWPH (D, E, F). Absorbance was measured at 220 nm while eluting with 50 mM ammonium formate buffer (pH 5.5). Glutathione (MW 307), aprotinin (MW 6,512), cytochrome c (MW 12,384), enolase (MW 67,000), lactate dehydrogenase (MW 142,000), and glutamate dehydrogenase (290,000) were used as molecular weight standard proteins. WP, whey protein; HWPH, hypoallergenic whey protein hydrolysate.

and Prozyme mainly consists of peptides of 5 kDa or less, and the main peaks were approximately 384 Da and 214 Da, respectively. WP was degraded into small peptides of 2.6, 1.3, and 0.491 kDa when treated with pepsin, and following pancreatin treatment, it was hydrolyzed into peptides with smaller molecular weights than when treated with pepsin (Figs. 6B and C). The molecular weight distribution of HWPH was similar after pepsin and pancreatin treatments. Since the HWPH had already been hydrolyzed by the enzyme (Figs. 6E and F), the effect of pepsin and pancreatin treatments on the molecular

weight distribution seems to be minimal.

Allergens can also be identified by their molecular weights. Enzymatic hydrolysis of milk proteins can prevent allergic sensitization by minimizing the number of sensitizing protein epitopes through the decomposition of allergenic sites or reduction in protein size (<1,500 Da; Crittenden and Bennett, 2005; Fiocchi et al., 2003). WP is mainly composed of 1,800 and 5,800 Da peptides, but Alcalase/Prozyme hydrolysate (HWPH) was composed of short peptides of 384 and 214 Da (Fig. 6). Therefore, the peptides constituting HWPH are small in size, which increases the absorption and bioavailability, but will be limited in antibody production due to their size.

Effect of hypoallergenic whey protein hydrolysate (HWPH) administration on cellular immunity of mice

The content and ratio of cytokines (INF- γ and IL-4) and antibodies (IgG2a and IgG1) corresponding to Type 1 helper T (Th1) and Type 2 helper T (Th2) effector cells in the blood, spleen cells, and small intestine are closely related to the immune response caused by allergens. INF- γ and IL-4 in the blood of mice administered with WP and HWPH tended to marginally increase as compared to mice in the control group, but this difference was not significant (Table 2). Although not statistically significant, the ratio of Th1/Th2 (INF- γ /IL-4) also showed an increase with HWPH treatment. The ratio of IgG2a/IgG1 between WP and HWPH was significantly increased by the oral administration of 250 and 500 mg/kg HWPH.

Table 2. Changes in cytokines and antibodies in the blood, spleen cells and small intestine after oral administration of WP and HWPH

Sample	(mg/kg)	Serum					
		INF- γ (pg/mL)	IL-4 (pg/mL)	INF- γ /IL-4	IgG2a (ng/mL)	IgG1 (ng/mL)	IgG2a/IgG1
CON		8.87 \pm 1.50 ^{ns}	5.30 \pm 0.21 ^{ns}	1.65 \pm 0.24 ^{ns}	52.62 \pm 2.18	26.32 \pm 0.65	2.01 \pm 0.13
WP	250	8.82 \pm 2.07	7.73 \pm 0.72	1.12 \pm 0.32	136.70 \pm 1.76 ^{***}	69.29 \pm 1.57	1.97 \pm 0.02
	500	9.95 \pm 1.71	10.53 \pm 2.90	1.16 \pm 0.35	103.27 \pm 2.98 ^{**}	500.69 \pm 36.01 ^{***}	0.21 \pm 0.01 ^{***}
HWPH	250	10.70 \pm 1.89	7.64 \pm 1.16	1.55 \pm 0.47	155.63 \pm 5.22 ^{***,#}	64.15 \pm 3.33	2.43 \pm 0.12 [#]
	500	11.66 \pm 0.81	8.94 \pm 2.61	1.63 \pm 0.33	339.82 \pm 13.33 ^{***,###}	63.96 \pm 3.20 ^{###}	5.35 \pm 0.44 ^{***,###}
Sample	(mg/kg)	Spleen					
		INF- γ (μ g/g)	IL-4 (μ g/g)	INF- γ /IL-4	IgG2a (μ g/g)	IgG1 (μ g/g)	IgG2a/IgG1
CON		45.37 \pm 0.89	2.85 \pm 0.28	16.74 \pm 1.72	0.36 \pm 0.08	0.32 \pm 0.04	1.18 \pm 0.30
WP	250	44.30 \pm 0.85	3.68 \pm 0.28	12.37 \pm 0.87	0.40 \pm 0.05	0.82 \pm 0.04 ^{***}	0.50 \pm 0.08
	500	45.13 \pm 0.63	4.57 \pm 0.51 [*]	10.50 \pm 1.15	0.48 \pm 0.05	1.89 \pm 0.08 ^{***}	0.26 \pm 0.03
HWPH	250	44.30 \pm 0.62	3.51 \pm 0.98	13.28 \pm 1.21 [*]	0.70 \pm 0.08 ^{*,#}	0.53 \pm 0.02 ^{*,###}	1.35 \pm 0.16 ^{*,###}
	500	52.19 \pm 0.92 ^{***,###}	3.85 \pm 0.16	13.67 \pm 1.71	0.77 \pm 0.08 ^{*,#}	0.72 \pm 0.01 ^{***,###}	1.08 \pm 0.12 ^{###}
Sample	(mg/kg)	Small intestine					
		INF- γ (μ g/g)	IL-4 (μ g/g)	INF- γ /IL-4	IgG2a (μ g/g)	IgG1 (μ g/g)	IgG2a/IgG1
CON		1.92 \pm 0.34	3.17 \pm 0.003	0.61 \pm 0.11	0.28 \pm 0.01	6.08 \pm 0.25	0.05 \pm 0.002
WP	250	1.91 \pm 0.79	3.97 \pm 0.01 ^{***}	0.48 \pm 0.20	0.39 \pm 0.01 ^{***}	11.18 \pm 0.18 ^{***}	0.04 \pm 0.001
	500	3.77 \pm 0.17	5.37 \pm 0.02 ^{***}	0.70 \pm 0.03	0.73 \pm 0.03 ^{***}	14.83 \pm 0.13 ^{***}	0.05 \pm 0.002
HWPH	250	4.54 \pm 1.04	3.54 \pm 0.02 ^{***,###}	1.281 \pm 0.30	0.94 \pm 0.04 ^{***,###}	7.19 \pm 0.06 ^{*,###}	0.13 \pm 0.01 ^{###}
	500	6.85 \pm 0.62 [*]	4.58 \pm 0.01 ^{***,###}	1.49 \pm 0.14 ^{*,###}	1.68 \pm 0.03 ^{***,###}	6.81 \pm 0.22 ^{###}	0.25 \pm 0.01 ^{###}

Values are presented as the mean \pm SEM for each group (n=8).

* p<0.05, ** p<0.01, and *** p<0.001 indicate significant difference from the control group (CON) based on Tukey's multiple test.

p<0.05, ## p<0.01, and ### p<0.001 indicate significant differences between groups (WP vs. HWPH) at the same concentration by Student's t-test.

WP, whey protein before hydrolysis; HWPH, hypoallergenic whey protein hydrolysate; ns, no significant difference between the groups.

CD4⁺ helper T (Th) cells play a central role in the immune response and secrete various cytokines that affect the function of B cells and the immune system. Depending on the secreted cytokines, they are divided into Th1 and Th2 effector cells (Herzyk et al., 2001; Lavigne et al., 1998) which exhibit characteristic cytokine profiles (MacLeod et al., 2008). Cytokines (IFN- γ) and antibodies (IgG2a) secreted from Th1 cells have an antagonistic action with cytokines (IL-4) and antibodies (IgG1) secreted from Th2 cells.

The ratio of INF- γ /IL-4 in the splenocytes decreased significantly in the group administered with 500 mg/kg of WP compared to that in the control group ($p < 0.01$). The ratio of IgG2a/IgG1 (Th1:Th2) was similar or higher in the HWPH-administered groups than in the control group, whereas the WP orally administered group showed a lower ratio, but the difference was not significant. There was a significant difference in the IgG2a/IgG1 ratios between the HWPH and WP groups.

Cytokines and antibodies specific to Th1 and Th2 cells in the small intestine of mice orally administered WP and HWPH were analyzed (Table 2). The IL-4 content in the groups orally administered with WP and HWPH was significantly higher than that in the control group ($p < 0.05$). The ratio of INF- γ /IL-4, corresponding to the Th1/Th2 ratio, tended to increase as the dose of HWPH increased, and the 500 mg/kg oral administration group showed a significantly higher level than the control group ($p < 0.05$). IgG2a levels in the small intestine in the two groups were significantly higher than in the control group. The ratio of IgG2a/IgG1 and the ratio of Th1 to Th2 cells were significantly higher in the HWPH-administered group than in the control group ($p < 0.001$). There was a significant difference in the IgG2a/IgG1 ratio between the WP and HWPH groups in terms of the dose ($p < 0.001$).

Abnormalities in the maintenance of Th1 and Th2 cell homeostasis are known to lead to various immunopathological abnormalities. When Th2 is relatively dominant, it promotes decreased defense against microbial infections, such as bacteria and viruses, allergic development, and some autoimmune diseases (Prigent et al., 1995; Umetsu and DeKruyff, 1997), and excessive immune responses (cell-mediated immunity) of Th1 cells are associated with auto-immune diseases (Gans et al., 2003; Margalit et al., 2005). As depicted in Table 2, the ratio of INF- γ /IL-4 was higher for HWPH than for WP, indicating that the allergic reaction induced by Th2 was somewhat suppressed. This is due to the fact that IgG2a isotype switching is induced by IFN- γ , and isotype switching of IgG1 or IgE is induced by IL-4, which inhibits the function or production of IFN- γ (Kühn et al., 1991; Snapper and Paul, 1987). The IgG2a/IgG1 ratio was higher in the HWPH than in the WP, suggesting that the Th1 response was relatively stimulated rather than the Th2 response, and thus, antibody conversion was induced, leading to a significant suppression of the allergic reaction (Table 2).

Oral administration of HWPH increased the levels of Th1 cytokines and antibodies in the blood, spleen, and small intestine, whereas the levels of Th2 cytokines and antibodies were lower than those in the WP group. In addition, the IgE content in the blood, splenocytes, and small intestine was significantly different between the two groups (Fig. 7). HWPH, in which α -LA and β -LG had been decomposed, had a lower level of allergen-related intracellular immune response than WP, and thus presumed to have a lower allergenic potential due to the hydrolysis of the allergen-inducing substances. The allergic response to WP is representative of type I immune hypersensitivity involving IgE (Martorell-Aragonés et al., 2015). Oral administration of HWPH resulted in a significantly lower IgE production than that of WP (Fig. 7). HWPH, a hydrolysate produced by Alcalase and Prozyme, seems to reduce allergic reactions as it hydrolyzes LG and LA, the two allergens in WP.

Conclusion

The mixing ratio of Alcalase and Prozyme, and the enzyme treatment conditions for HWPH production were optimized




Fig. 7. Changes in immunoglobulin (Ig) E in the blood, spleen cells, and small intestine in mice after oral administration of WP and HWPH. Values are presented as the mean±SEM for each group (n=8). * p<0.05, ** p<0.01, and *** p<0.001 as compared to the control group based on Tukey's multiple test. # p<0.05, ## p<0.01, and ### p<0.001 between groups of the same concentration using student t-test. WP, whey protein; HWPH, hypoallergenic whey protein hydrolysate; ns, no significant difference between the groups.

using RSM. The mixed treatment of Alcalase and Prozyme effectively degraded α -LA and β -LG, the major allergens of WP. Because HWPH is converted into small peptides by enzyme treatment and has a low molecular weight, its digestibility and intestinal permeability are higher than those of WP. In addition, it was confirmed that HWPH suppressed the allergic immune response by stimulating the Th1 response, but not the Th2 response, and by inducing antibody conversion. Thus, HWPH can be expected to be useful in a variety of products with low allergenicity targeting infants, adults, and the elderly.

Supplementary Materials

Supplementary materials are only available online from: <https://doi.org/10.5851/kosfa.2023.e21>.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through High Value-added Food Technology Development Program, funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (No. 322011-4).

Author Contributions

Conceptualization: Ahn Y. Data curation: Yang J, Lee SK. Formal analysis: Yang J, Lee SK. Methodology: Kim YS, Suh HJ. Software: Kim YS, Suh HJ. Validation: Kim YS, Suh HJ, Ahn Y. Investigation: Yang J, Lee SK. Writing - original draft: Yang J, Lee SK, Ahn Y. Writing - review & editing: Yang J, Lee SK, Kim YS, Suh HJ, Ahn Y.

Ethics Approval

All animal experiments were approved by the Korea University Animal Experiment Ethics Committee (KUIACUC-2022-0095).

References

- Adamson NJ, Reynolds EC. 1996. Characterization of casein phosphopeptides prepared using alcalase: Determination of enzyme specificity. *Enzyme Microb Technol* 19:202-207.
- Andriamihaja M, Guillot A, Svendsen A, Hagedorn J, Rakotondratohanina S, Tomé D, Blachier F. 2013. Comparative efficiency of microbial enzyme preparations versus pancreatin for *in vitro* alimentary protein digestion. *Amino Acids* 44:563-572.
- Benjakul S, Morrissey MT. 1997. Protein hydrolysates from pacific whiting solid wastes. *J Agric Food Chem* 45:3423-3430.
- Boscaini S, Skuse P, Nilaweera KN, Cryan JF, Cotter PD. 2023. The ‘whey’ to good health: Whey protein and its beneficial effect on metabolism, gut microbiota and mental health. *Trends Food Sci Technol* 133:1-14.
- Bu G, Luo Y, Chen F, Liu K, Zhu T. 2013. Milk processing as a tool to reduce cow’s milk allergenicity: A mini-review. *Dairy Sci Technol* 93:211-223.
- Chatterjee C, Gleddie S, Xiao CW. 2018. Soybean bioactive peptides and their functional properties. *Nutrients* 10:1211.
- Cohen CG, Zhao WW, Ke D, Beaudette L, Lejtenyi D, McCusker C, Zhang X, Chan ES, Upton JEM, Grunebaum E, Clarke AE, Mazer BD, Ben-Shoshan M. 2022. Elevated cow’s milk-specific IgE levels prior to oral immunotherapy decrease the likelihood of reaching the maintenance dose. *J Allergy Clin Immunol Pract* 10:215-221.E2.
- Crittenden RG, Bennett LE. 2005. Cow’s milk allergy: A complex disorder. *J Am Coll Nutr* 24:582S-591S.
- Cui Q, Zhang Z, Li M, Zhou M, Sun X. 2023. Peptide profiles and allergy-reactivity of extensive hydrolysates of milk protein. *Food Chem* 411:135544.
- Diniz FM, Martin AM. 1997. Optimization of nitrogen recovery in the enzymatic hydrolysis of dogfish (*Squalus acanthias*) protein. Composition of the hydrolysates. *Int J Food Sci Nutr* 48:191-200.
- Exl BM, Fritsché R. 2001. Cow’s milk protein allergy and possible means for its prevention. *Nutrition* 17:642-651.
- Feng M, Betti M. 2017. Transepithelial transport efficiency of bovine collagen hydrolysates in a human Caco-2 cell line model. *Food Chem* 224:242-250.
- Fiocchi A, Martelli A, De Chiara A, Moro G, Warm A, Terracciano L. 2003. Primary dietary prevention of food allergy. *Ann Allergy Asthma Immunol* 91:3-12.
- Gans H, DeHovitz R, Forghani B, Beeler J, Maldonado Y, Arvin AM. 2003. Measles and mumps vaccination as a model to investigate the developing immune system: Passive and active immunity during the first year of life. *Vaccine* 21:3398-3405.
- Garrett DA, Failla ML, Sarama RJ. 1999. Development of an *in vitro* digestion method to assess carotenoid bioavailability from meals. *J Agric Food Chem* 47:4301-4309.
- Herzyk DJ, Gore ER, Polsky R, Nadwodny KL, Maier CC, Liu S, Hart TK, Harmsen AG, Bugelski PJ. 2001. Immunomodulatory effects of anti-CD4 antibody in host resistance against infections and tumors in human CD4 transgenic mice. *Infect Immun* 69:1032-1043.
- Hochwallner H, Schulmeister U, Swoboda I, Spitzauer S, Valenta R. 2014. Cow’s milk allergy: From allergens to new forms of diagnosis, therapy and prevention. *Methods* 66:22-33.

- Jaiswal L, Worku M. 2022. Recent perspective on cow's milk allergy and dairy nutrition. *Crit Rev Food Sci Nutr* 62:7503-7517.
- Kheroufi A, Brassesco ME, Campos DA, Boughellout H, Pintado ME. 2022. Functional properties of peptides obtained from whey proteins by ficin extract hydrolysis. *Food Biosci* 47:101707.
- Kühn R, Rajewsky K, Müller W. 1991. Generation and analysis of interleukin-4 deficient mice. *Science* 254:707-710.
- Landim APM, Tiburski JH, Mellinger CG, Juliano P, Rosenthal A. 2023. Potential application of high hydrostatic pressure on the production of hydrolyzed proteins with antioxidant and antihypertensive properties and low allergenicity: A review. *Foods* 12:630.
- Lavigne LM, Schopf LR, Chung CL, Maylor R, Sypek JP. 1998. The role of recombinant murine IL-12 and IFN- γ in the pathogenesis of a murine systemic *Candida albicans* infection. *J Immunol* 160:284-292.
- Lynch R, Buckin V. 2022. Ultrasonic analysis of effects of varying temperature, pH, and proteolytic enzymes on hydrolysis of lactose by neutral lactase formulations in infant milk and in the infant digestive system. *Food Res Int* 157:111004.
- MacLeod MKL, McKee A, Crawford F, White J, Kappler J, Marrack P. 2008. CD4 memory T cells divide poorly in response to antigen because of their cytokine profile. *Proc Natl Acad Sci USA* 105:14521-14526.
- Margalit M, Shibolet O, Klein A, Elinav E, Alper R, Thalenfeld B, Engelhardt D, Rabbani E, Ilan Y. 2005. Suppression of hepatocellular carcinoma by transplantation of *ex-vivo* immune-modulated NKT lymphocytes. *Int J Cancer* 115:443-449.
- Marshall K. 2004. Therapeutic applications of whey protein. *Altern Med Rev* 9:136-156.
- Martorell-Aragonés A, Echeverría-Zudaire L, Alonso-Lebrero E, Boné-Calvo J, Martín-Muñoz MF, Nevot-Falcó S, Piquer-Gibert M, Valdesoiro-Navarrete L, Food Allergy Committee of SEICAP (Spanish Society of Pediatric Allergy, Asthma and Clinical Immunology). 2015. Position document: IgE-mediated cow's milk allergy. *Allergol Immunopathol* 43:507-526.
- Nielsen PM, Petersen D, Dambmann C. 2001. Improved method for determining food protein degree of hydrolysis. *J Food Sci* 66:642-646.
- Park WS, Koo KA, Kim HJ, Kwon JM, Kang DM, Chung HJ, Lee SW, Kim HS, Kwak SS, Ahn MJ. 2021. Intestinal epithelial monolayer permeability of sweet potato-derived anthocyanin and carotenoid extracts in Caco-2 cells. *J Korean Soc Food Sci Nutr* 50:765-773.
- Prigent P, Saoudi A, Pannetier C, Graber P, Bonnefoy JY, Druet P, Hirsch F. 1995. Mercuric chloride, a chemical responsible for T helper cell (Th)2-mediated autoimmunity in brown Norway rats, directly triggers T cells to produce interleukin-4. *J Clin Invest* 96:1484-1489.
- Savilahti E, Kuitunen M. 1992. Allergenicity of cow milk proteins. *J Pediatr* 121:S12-S20.
- Sicherer SH, Sampson HA. 1999. Cow's milk protein-specific IgE concentrations in two age groups of milk-allergic children and in children achieving clinical tolerance. *Clin Exp Allergy* 29:507-512.
- Snapper CM, Paul WE. 1987. Interferon- γ and B cell stimulatory factor-1 reciprocally regulate Ig isotype production. *Science* 236:944-947.
- Stănciuc N, Rapeanu G. 2010. An overview of bovine α -lactalbumin structure and functionality. *Ann Univ Dunarea Jos Galati Fascicle VI Food Technol* 34:82-93.
- Suh HJ, Shin JC, Kim JH, Jang JH, Han SH. 2017. Optimal enzyme selection for organic whey protein hydrolysis. *Korean J Food Nutr* 30:1359-1363.
- Umetsu DT, Dekruyff RH. 1997. T_{H1} and T_{H2} CD4⁺ cells in human allergic diseases. *J Allergy Clin Immunol* 100:1-6.

- van Breemen RB, Li Y. 2005. Caco-2 cell permeability assays to measure drug absorption. *Expert Opin Drug Metab Toxicol* 1:175-185.
- Vandenplas Y, Ksiazyk J, Luna MS, Migacheva N, Picaud JC, Ramenghi LA, Singhal A, Wabitsch M. 2022. Partial hydrolyzed protein as a protein source for infant feeding: Do or don't? *Nutrients* 14:1720.
- Vieira GHF, Martin AM, Saker-Sampaiao S, Omar S, Goncalves RCF. 1995. Studies on the enzymatic hydrolysis of Brazilian lobster (*Panulirus* spp.) processing wastes. *J Sci Food Agric* 69:61-65.
- Villas-Boas MB, Benedé S, de Lima Zollner R, Netto FM, Molina E. 2015. Epitopes resistance to the simulated gastrointestinal digestion of β -lactoglobulin submitted to two-step enzymatic modification. *Food Res Int* 72:191-197.
- Villanueva A, Vioque J, Sánchez-Vioque R, Clemente A, Bautista J, Millán F. 1999. Production of an extensive sunflower protein hydrolysate by sequential hydrolysis with endo- and exo-proteases. *Grasas y Aceites* 50:472-476.
- Wang B, Li B. 2017. Effect of molecular weight on the transepithelial transport and peptidase degradation of casein-derived peptides by using Caco-2 cell model. *Food Chem* 218:1-8.
- Wright K, Feeney M, Yerlett N, Meyer R. 2022. Nutritional management of children with food allergies. *Curr Treat Options Allergy* 9:375-393.
- Yiğit A, Bielska P, Cais-Sokolińska D, Samur G. 2023. Whey proteins as a functional food: Health effects, functional properties, and applications in food. *J Am Nutr Assoc* (in press). doi: 10.1080/27697061.2023.2169208.
- Yuan F, Ahmed I, Lv L, Li Z, Li Z, Lin H, Lin H, Zhao J, Tian S, Ma J. 2018. Impacts of glycation and transglutaminase-catalyzed glycosylation with glucosamine on the conformational structure and allergenicity of bovine β -lactoglobulin. *Food Funct* 9:3944-3955.