

Unraveling the Contributing Factors of Sarcopenia in Young Korean Male Adults: A Study of Occurrence, Somatometric, Biochemical, and Behavioral Characteristics

Jongseok Hwang, PT, Ph.D.[†]
Institute of Human Ecology, Yeungnam University

Received: July 28 2023 / Revised: July 28 2023 / Accepted: August 3 2023

© 2023 J Korean Soc Phys Med

| Abstract |

PURPOSE: The present study evaluated the occurrence, somatometric, biochemical, and behavioral characteristics of young Korean sarcopenic males aged between 30 to 39 years.

METHODS: The study involved a total of 1,546 participants. These individuals were divided into two distinct groups based on their skeletal muscle mass index. Of these, 49 participants were categorized into the sarcopenia group, while the remaining 1,497 individuals were placed in the normal group. The researchers analyzed the occurrence of sarcopenia, somatometric, biochemical, and behavioral characteristics such as height, weight, body mass index (BMI), waist circumference (WC), skeletal muscle mass index, fasting glucose (FG), triglyceride and total cholesterol (TC) levels, systolic (SBP) and diastolic blood pressure (DBP), and the drinking and smoking status. A complex sampling data analysis was performed.

RESULTS: The weighted occurrence of sarcopenia was

found to be 3.24%, (95% confidence interval (CI) of 2.39% to 4.36%). The somatometric contributing factors were height, BMI, and WC ($p < .05$). Triglyceride and TC levels exhibited statistically significant differences in the biochemical variables ($p < .05$). The behavioral characteristics, including the drinking and smoking status, had no statistical significance ($p > .05$).

CONCLUSION: This study identified specific occurrences of sarcopenia and contributing factors in young Korean men.

Key Words: Contributing factors, Occurrence, Sarcopenia, Younger adult

I. Introduction

Global advances in applied science and healthcare technology, along with improvements in socioeconomic conditions, have led to a notable increase in the human lifespan. According to estimates from 2019, the worldwide population of individuals aged 65 years and older was approximately 703 million, and this number is projected to reach 1.5 billion by 2050 [1]. Life expectancy has risen in all countries, resulting in a significant increase in the proportion of the elderly in the population [2]. Consequently, healthcare services are facing a strain due to the growing

[†]Corresponding Author : Jongseok Hwang
sfscfc44@naver.com, <https://orcid.org/0000-0003-3376-5619>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

burden of caring for the elderly. To address age-related chronic illnesses and alleviate the pressure on healthcare systems, it is imperative to gain a better understanding of these conditions and develop innovative approaches to address aging-related diseases.

Sarcopenia is a condition characterized by the gradual and widespread decline in muscle mass and function as people age [3]. It is closely associated with frailty, a state in which elderly individuals become more susceptible to stressors that can lead to adverse health outcomes, and increase the chances of disability and mortality [4]. Understanding and addressing sarcopenia is crucial in improving the health and well-being of aging individuals, as these conditions can put a significant strain on healthcare systems and affect the quality of life of older adults. Developing interventions to prevent or manage sarcopenia and frailty can help alleviate the burden on healthcare services and promote healthy aging.

Sarcopenia exhibits a greater prevalence in men when compared to women, as evidenced by several studies. For instance, Brown et al. [5] conducted a study involving 4,425 adults in the US, revealing that the prevalence of sarcopenia was higher in males (44.8%) than in females (30.24%). Similarly, Liu et al. [6] investigated 4,500 Chinese individuals in an urban community and reported a prevalence of 22.1% in men and 17.8% in women. In addition, Hai et al. [7], in their assessment of 834 community-dwelling Chinese individuals, found that the incidence of sarcopenia was 11.3% in men and 9.8% in women. Likewise, Chan et al. [8] observed a prevalence of 9.30% in males and 5.30% in females among a group of 3,957 people in Hong Kong. These collective findings consistently suggest that sarcopenia is more commonly found in males than females. Notwithstanding the notable prevalence of this condition among males, identifying contributing factors and effectively addressing sarcopenia in this specific population pose considerable challenges, particularly when compared to the well-explored research

on sarcopenia in females [9-12].

Most existing research on sarcopenia has focused on individuals aged 40 years and older [20-28]. Nonetheless, emerging evidence suggests that age-related muscle decline may commence in younger adult males as early as in the 30s [29-31]. Sarcopenia in young adults can be attributed to a combination of multiple lifestyle-related and other physiological factors [13,14]. Among these, lifestyle choices play a pivotal role—sedentary habits, insufficient physical activity, and suboptimal dietary patterns can all precipitate muscle weakening and atrophy even in individuals of a younger age [15]. The hormonal decline after the 20s constitutes another significant facet; disruptions in the growth hormone, testosterone, and insulin-like growth factor-1 (IGF-1) levels can have an intricate effect on the maintenance and development of muscle mass [16,17]. Furthermore, young adults prefer instant foods that offer only limited nutrition [18]. The deficiency of crucial nutrients such as proteins, vitamins, and essential minerals [19], which are fundamental for nurturing muscle health, also significantly contributes to the early onset of sarcopenia among young adults.

It is important to evaluate the factors contributing to sarcopenia in young men between 30 and 39 years old, as this can lead to the development of early prevention strategies against age-related muscle loss. Consequently, the present study aimed to investigate the occurrence, somatometric, biochemical, and behavioral characteristics of young Korean sarcopenic men aged 30 to 39 years.

II. Methods

1. Data Sampling

The current investigation utilized data from the Korea National Health and Nutrition Examination Survey (KNHANES), an initiative by the Centers for Disease Control and Prevention aimed at monitoring the health-contributing behaviors of the population. The survey

Table 1. General characteristics of participants in the study

	Sarcopenia group (n =49)	Normal group (n = 1,497)	p (N = 1,546)
Age (years)	35.49 ± 2.9	35.00 ± 2.78	.235
Height (cm)	164.29 ± 6.343	173.23 ± 5.61	.000
Weight (kg)	72.92 ± 13.76	72.68 ± 10.89	.879
SMI (g/m ²)	743.204 ± 46.5	988.53 ± 102.76	.000

SMI, skeletal muscle mass index. An independent *t*-test test was performed.

employed a stratified, clustered, multistage probability sampling design, involving a total of 35,737 individuals surveyed between 2008 and 2011. After excluding 33,339 participants outside the age range of 30 to 39 years, the dataset comprised 2,398 men. Subsequently, 1,004 subjects were further excluded due to the unavailability of data from their health surveys and the absence of dual X-ray absorptiometry (DEXA) measurements. Eventually, only data from 1,546 male participants between the ages of 30 and 39 years were considered for the final analysis. The inclusion criteria stipulated individuals within the age range of 30 to 39 years, while the exclusion criteria included individuals who did not undergo DEXA measurements and with incomplete health survey data. Among these participants, 49 individuals were classified into the sarcopenia group based on their skeletal muscle mass index score, and the remaining 1,497 individuals were assigned to the normal group. The study received approval from the institutional review board of the KDCA (approval numbers 2008-04EXP-01-C, 2009-01CON-03-2C, 2010-02CON-21-C, and 2011-02CON-06-C), and informed written consent was obtained from all participants. Table 1 describes the general characteristic of the participants in the study.

2. Diagnosis of Sarcopenia

Sarcopenia, a medical condition categorized under the ICD-10-CM code M62.84 by the World Health Organization (WHO) [32], is diagnosed through the measurement of

appendicular skeletal muscle mass. In this study, DEXA was employed to assess the appendicular skeletal muscle mass (ASM). The skeletal muscle mass index (SMI) was calculated as the ratio of ASM (in kg) to BMI (in kg/m²). The definition of sarcopenia in this study was based on the criteria established by the Foundation for the National Institutes of Health (FNIH) Sarcopenia Project wherein sarcopenia in men was defined as an SMI of less than 0.789 [33].

3. Variables

1) Somatometric Variables

The somatometric variables data were collected from all study participants. To ensure accurate measurements, individuals were requested to remove their shoes, socks, hats, and hairpins, and were instructed to wear light clothing. Height and weight were measured using calibrated automatic body measurement equipment, with precision recorded to the nearest .1 cm or kg. Subsequently, the body mass index (BMI) was calculated by dividing the weight (in kg) by the square of height (in m²). Additionally, waist circumference (WC) was measured to the nearest .1 cm in a horizontal plane at the midline between the last rib and the iliac crest, following a normal expiration [34].

2) Biochemical and Blood Pressure Variables

The biochemical variables including fasting glucose (FG), triglyceride, and total cholesterol (TC) levels were

analyzed using the LABOSPECT 008AS platform (Hitachi High-Tech Co., Tokyo, Japan). Blood samples were collected from the non-dominant arm after an overnight fast of at least eight hours. Immediately after collection, the blood was mixed with a coagulation promoter and centrifuged within a mobile examination vehicle. All tests were conducted within 24 hours of blood sample collection. A trained practitioner conducted the measurement of systolic blood pressure (SBP) and diastolic blood pressure (DBP) using a mercury sphygmomanometer. The blood pressure cuff was positioned at the heart level with participants in a seated position after a minimum of five minutes of rest [35,34].

3) Behavioral Habits

Data pertaining to behavioral habits including the smoking and drinking status of the participants were obtained through a survey. Based on their cigarette smoking and alcohol drinking habits, individuals were categorized into three groups: non-users, ex-users, or current users. These measured variables play a crucial role in evaluating various aspects of health and the potential contribution to diseases within the study population [34].

4. Statistical Analysis

This study conducted a comprehensive analysis of data, providing statistical values such as mean and standard deviation for each measurement. The complex sampling method involved a nationwide analysis incorporating individual weights provided by the KNHANES. The researchers used SPSS 22.0 Windows version (IBM Corporation, Armonk, NY, USA) for their statistical analysis. The data utilized a stratified, clustered, multistage probability sampling design. The researchers employed independent t-tests and Chi-square analyses to compare the variables between participants with and without sarcopenia. Additionally, multiple logistic regression was utilized to calculate the odds ratio of sarcopenia. The

researchers established a statistical significance level of $\alpha = .05$ for the study.

III. Results

1. Occurrence of Sarcopenia

The weighted occurrence of sarcopenia was reported as 3.24% (with a 95% confidence interval of 2.39% to 4.36%). These values were calculated by complex sampling analysis, after applying adjusted weights to individuals. The unweighted prevalence of sarcopenia was found to be 3.17% (Table 2).

2. Contributing Factors

1) Somatometric Factors

The contributing factors analyzed in this study included height, BMI, and WC and they exhibited statistically significant differences between the two groups ($p < .05$). However, there was no statistically significant difference ($p > .05$) in the weight between the groups. Table 3 provides a comprehensive overview of these contributing factors and their associations with the studied outcomes (Table 3).

2) Biochemical and Blood Pressure Variables

Triglyceride and TC levels exhibited statistically significant differences between the two groups ($p < .05$), indicating that these variables are important distinguishing factors between the studied groups. On the other hand, FG, SBP, and DBP did not show statistically significant

Table 2. Prevalence of sarcopenia

	Sarcopenia group (n = 49)	Normal group (n = 1,497)	Total
Un-weighted (%)	3.17	96.83	100
Weighted (%)	3.24 (2.39-4.36)	96.76 (95.64-97.61)	100

Weighted values present the 95% confidence interval.

Table 3. Analysis of somatometric factors in the study population

	Sarcopenia group (N = 49)	Normal group (n = 1,497)	p
Height (cm)	164.29 ± 6.343	173.23 ± 5.61	.000
Weight (kg)	72.92 ± 13.76	72.68 ± 10.89	.879
BMI (kg/m ²)	26.92 ± 4.24	24.18 ± 3.19	.000
WC (cm)	89.60 ± 10.20	83.56 ± 8.91	.000

BMI, body mass index; WC, waist circumference. An independent *t*-test was performed.

Table 4. Biochemical and blood pressure variables of study participants

	Sarcopenia group (N = 49)	Normal group (n = 1,497)	p
FG (mg/dL)	99.22 ± 17.74	94.35 ± 18.8	.077
Triglycerides (mg/dL)	236.56 ± 249.32	162.39 ± 131.84	.000
TC (mg/dL)	200.89 ± 28.84	189.68 ± 34.15	.025
SBP (mmHg)	117.75± 13.05	116.27 ± 12.65	.420
DBP (mmHg)	80.735 ± 10.72	79.45 ± 10.62	.407

FG, fasting glucose; TC, total cholesterol; SBP, systolic blood pressure; DBP, diastolic blood pressure. An independent *t*-test was performed.

Table 5. Behavioral habits of study participants

	Sarcopenia group (N = 49)	Normal group (n = 1,497)	p
Drinking status (%) (current-/ex-/non-drinker)	93.30 / 4.84 / 1.85	92.49 / 5.03 / 2.47	.957
Smoking status (%) (current-/ex-/non-smoker)	71.84 / 11.06 / 17.08	70.23 / 12.14 / 17.62	.971

A Chi-square test was performed.

differences ($p > .05$) between the two groups. These results highlight the relevance of triglyceride and TC levels in sarcopenia, while FG, SBP, and DBP did not demonstrate significant associations with the studied outcomes (Table 4).

3) Behavioral Habits

The drinking and smoking status had no statistical significance ($p > .05$) (Table 5).

3. Odd Ratios (ORs) for Contributing Factors

The variables of height, BMI, WC, SMI, triglyceride, and TC demonstrated statistically significant differences between the two groups ($p < .05$). The corresponding values were .00 (.00-.00), 7.66 (1.27-46.19), 1.78 (1.24-2.56), 42.68 (36.62-56.93), 1.00 (1.00-1.00), 1.08 (1.05-1.10) (Table 6).

Table 6. Odds ratio (OR) for male sarcopenia

	Odds ratio (95% CI)	p
Height	.00 (.00–.00)	.000
BMI	7.66 (1.27–46.19)	.000
WC	1.78 (1.24–2.56)	.002
SMI	42.68 (36.62–56.93)	.000
Triglycerides	1.00 (1.00–1.00)	.001
TC	1.08 (1.05–1.10)	.000

BMI, body mass index; WC, waist circumference; SMI, skeletal muscle mass index; TC, Total cholesterol.

Odds ratio values are present in the 95% confidence interval (CI).

Multiple logistic regression in complex sampling was performed.

IV. Discussion

This study aimed to explore the occurrence and contributing factors associated with sarcopenia among young men aged 30 to 39 residing in the community. The aging population in Korea and Asia is increasing rapidly, leading to a higher occurrence of sarcopenia, especially among males. Unfortunately, healthcare professionals, including physical and occupational therapists, often face challenges in accurately diagnosing the disease due to limited knowledge and diagnostic tools, resulting in missed diagnoses and potential complications. To address this issue, this study focused on easily accessible and cost-effective variables, such as somatometric, biochemical variables, and behavioral characteristics, which are useful for detecting potential sarcopenia patients. Early detection and prevention of sarcopenia are crucial for managing its potential negative consequences. The study identified several contributing factors associated with sarcopenia, including height, BMI, WC, and triglyceride and TC levels. Understanding these contributing factors provides valuable insights that can help healthcare professionals intervene at an early stage and implement effective measures to prevent sarcopenia in community-dwelling younger men.

The WC has been identified as a contributing factor for sarcopenia. Numerous studies have consistently shown

an association between WC and an increased incidence of sarcopenia in men [36,5,37]. For instance, Brown et al. conducted a study on a US cohort and reported an OR of 1.39 (95% CI [1.05,1.84]) for WC as a contributing factor for sarcopenia in men [5]. Similarly, a cohort study in Brazil among individuals with sarcopenia found a significant OR of 17.90 (95% CI [1.48, 201.16]) for waist circumference [37]. Another study in Japan, focusing on community-dwelling individuals, also revealed that those with sarcopenia had a larger WC compared to individuals without sarcopenia [36]. The underlying reason for the higher WC in adults with sarcopenia can be attributed to the intricate relationship between increased fat mass and decreased muscle mass [38]. Individuals with sarcopenia often experience difficulties in muscle power and function due to muscle loss, leading to reduced levels of physical activity, including challenges in activities such as sitting-to-stand and walking long distances indoors and outdoors [39]. This decreased physical activity is closely linked to reduced total daily energy expenditure and an increase in fat deposits, especially in the visceral and abdominal areas, thereby contributing to increased waist volume [39]. Conversely, higher fat volume, particularly visceral fat, produces pro-inflammatory cytokines such as interleukin (IL)-6 and C-reactive protein (CRP), which can hinder the anabolic response of muscle tissue [40].

Consequently, the relationship between decreased muscle mass and increased fat mass in individuals with sarcopenia is bidirectional and mutually reinforcing [41]. In summary, the WC has been consistently identified as a significant contributing factor for sarcopenia in men, with the interplay between decreased muscle mass and increased fat mass playing a pivotal role in this relationship.

Triglyceride levels were identified as another significant contributing factor for men, with an average level of 231.74 mg/dL in the sarcopenia group and 178.14 mg/dL in the normal group. These findings are consistent with previous research on sarcopenia [42-44]. For instance, Lu et al. [32] examined 600 elderly individuals in northern Taiwan and reported that the sarcopenic group had significantly higher triglyceride levels of 1.9 mmol/L compared to the normal group with 1.3 mmol/L [42]. Similarly, Buchmann et al. [43] investigated 1420 elderly individuals living in Berlin and concluded that participants with sarcopenia had higher triglyceride levels (108.7 mg/dL) compared to the non-sarcopenia group (92.1 mg/dL). Additionally, Du et al. [44] conducted a cross-sectional study on community-dwelling elderly people in East China and found that the male sarcopenia group had increased serum triglyceride levels. An underlying mechanism for the higher triglyceride and total cholesterol levels in sarcopenia may be related to insulin resistance [45] and elevated levels of inflammatory cytokines [52]. These factors could contribute to alterations in lipid metabolism, ultimately leading to higher triglyceride levels in individuals with sarcopenia. In short, the study highlights triglyceride levels as an important contributing factor for men with sarcopenia, consistent with previous studies. The association between increased triglyceride levels and sarcopenia could be attributed to insulin resistance and inflammation. Understanding these connections is crucial for managing and preventing the incidence of age-related low skeletal muscle mass in the population [46].

Research findings have consistently indicated that TC

levels act as a contributing factor for sarcopenia in men, which is in line with previous studies [29,33]. For instance, a study by Du et al. [33] revealed that men in the sarcopenic group had higher TC levels compared to those in the normal group. Similarly, in a study conducted by Sanada and colleagues involving 1488 Japanese individuals, it was observed that TC levels were significantly elevated in individuals with sarcopenia when compared to the normal group [29]. The potential underlying mechanism for the higher triglyceride and TC levels in sarcopenia could be associated with factors such as insulin resistance [34] and an increased presence of inflammatory cytokines [35]. These factors can disturb lipid metabolism, leading to elevated triglyceride and total cholesterol levels in individuals with sarcopenia. To sum up, research consistently highlights the significance of higher total cholesterol levels as a contributing factor for sarcopenia in men, corroborating findings from earlier studies. The connection between elevated triglyceride and total cholesterol levels in sarcopenia may be influenced by factors like insulin resistance and increased inflammatory cytokines. Gaining a deeper understanding of these associations is vital for effectively managing and preventing sarcopenia among men.

One crucial aspect of this study is the investigation of male-specific contributing factors in a representative population of Korean individuals in their 30s, a critical age when the loss of skeletal muscles typically begins. Unlike most studies that combine both sexes into a single group, this study focused specifically on men, providing valuable insights into the gender-specific factors contributing to sarcopenia [5,21,47].

However, it is essential to acknowledge a couple of limitations of this study that should be considered for undertaking future research. The cross-sectional design utilized in this study, despite incorporating a large sample size of participants representative of the entire population through statistical weighting, may present limitations in

establishing causal relationships for the identified contributing factors. Cross-sectional studies only capture data at a single point in time, making it challenging to infer causality between variables. Hence, to strengthen the findings, future studies should consider employing a longitudinal or randomized case-control study design, allowing for a better understanding of the temporal relationships and causative factors for sarcopenia among the male population. Secondly, the current study did not investigate the issue of sarcopenic obesity which presents with reduced muscle mass and being overweight. To enhance comprehension of the observed heightened levels of triglycerides and total cholesterol, it would have been advantageous to examine sarcopenic obesity. Third, at the time of selecting study subjects, patients with underlying diseases such as cardiovascular disease, chronic heart disease, and chronic respiratory disease were not excluded. The study could not rule out the possibility that these diseases could have affected muscle mass. Finally, the study did not include standardized measures of physical activity or exercise as part of the surveyed behavioral habits. The inclusion of such measures would have enhanced the robustness of the study's findings. It is recommended that forthcoming research should investigate deeper into physical activity or the exercise-related habits of the participants to bolster the comprehensiveness and reliability of the study results.

V. Conclusion

This paper presents a pioneering investigation focusing on the clinical contributing factors which raise the risk of sarcopenia in young Korean males in their 30s. The study's objective was to shed light on this condition by providing valuable insights into its prevalence and associated contributing factors. According to the results, the estimated prevalence of sarcopenia in this specific demographic population was 3.24% (95%CI [2.39,4.36])

The study identified several significant contributing factors related to sarcopenia. These factors included height, BMI, SMI, WC, triglyceride levels, and TC levels. By recognizing and considering these contributing factors, healthcare professionals can identify and detect potential sarcopenia patients, thereby promoting early intervention and treatment. Despite the valuable findings presented in this study, the relationship between the identified predictors and sarcopenia requires further exploration. Therefore, additional research is warranted to deepen our understanding of this condition. Specifically, longitudinal or randomized case-control study designs should be employed to strengthen the reliability and validity of the study's conclusions. Such efforts will contribute significantly to the advancement of knowledge in the field of sarcopenia and its management in males in their 30s.

References

- [1] Population Division, Department of Economic and Social Affairs. United Nations. World population aging. 2019. 4-11.
- [2] Wang C, Bai L. Sarcopenia in the elderly: Basic and clinical issues. *Geriatr Gerontol Int*. 2012;12(3):388-96.
- [3] Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age and ageing*. 2019;48(1):16-31.
- [4] Jung HW. Visualizing domains of comprehensive geriatric assessments to grasp frailty spectrum in older adults with a radar chart. *Ann Geriatr Med Res*. 2020;24(1):55-6.
- [5] Brown JC, Harhay MO, Harhay MN. Sarcopenia and mortality among a population-based sample of community-dwelling older adults. *J Cachexia Sarcopenia Muscle*. 2016;7(3):290-8.
- [6] Liu X, Hao Q, Yue J, et al. Sarcopenia, obesity and sarcopenia obesity in comparison: Prevalence, metabolic profile, and key differences: Results from WCHAT study. *J Nutr Health Aging*. 2020;24(4):429-37.

- [7] Hai S, Wang H, Cao L, et al. Association between sarcopenia with lifestyle and family function among community-dwelling Chinese aged 60 years and older. *BMC Geriatr.* 2017;17(1):187.
- [8] Chan R, Leung J, Woo J. A prospective cohort study to examine the association between dietary patterns and sarcopenia in chinese community-dwelling older people in Hong Kong. *J Am Med Dir Assoc.* 2016; 17(4):336-42.
- [9] Nishiguchi S, Yamada M, Fukutani N, et al. Differential association of frailty with cognitive decline and sarcopenia in community-dwelling older adults. *J Am Med Dir Assoc.* 2015;16(2):120-4.
- [10] Kang SY, Lim GE, Kim YK, et al. Association between sarcopenic obesity and metabolic syndrome in postmenopausal women: A cross-sectional study based on the Korean national health and nutritional examination surveys from 2008 to 2011. *J Bone Metab.* 2017;24(1):9-14.
- [11] Velazquez-Alva MC, Irigoyen Camacho ME, Lazarevich I, et al. Comparison of the prevalence of sarcopenia using skeletal muscle mass index and calf circumference applying the European consensus definition in elderly Mexican women. *Geriatr Gerontol Int.* 2017;17(1): 161-70.
- [12] Lee HN KB. Convergence factors affecting sarcopenia in middle-aged and older women in Korea: A cross sectional study by using 5th KNHANES. *J Korean Converg Soc.* 2020;11(11):405-16.
- [13] Mager D, Carroll M, Wine E, et al. Vitamin D status and risk for sarcopenia in youth with inflammatory bowel diseases. *Eur J Clin Nutr.* 2018;72(4):623-6.
- [14] Jung HN, Jung CH, Hwang Y-C. Sarcopenia in youth. *Metabolism.* 2023;144:155557.
- [15] Ramsey KA, Rojer AG, D'Andrea L, et al. The association of objectively measured physical activity and sedentary behavior with skeletal muscle strength and muscle power in older adults: A systematic review and meta-analysis. *Ageing research reviews.* 2021;67:101266.
- [16] Maggio M, Lauretani F, Ceda GP. Sex hormones and sarcopenia in older persons. *Curr Opin Clin Nutr Metab Care.* 2013;16(1):3-13.
- [17] Ascenzi F, Barberi L, Dobrowolny G, et al. Effects of IGF-1 isoforms on muscle growth and sarcopenia. *Aging Cell.* 2019;18(3):e12954.
- [18] Bugge AB. Lovin' it? food. *Culture & Society.* 2011;14(1):71-89.
- [19] Robinson S, Cooper C, Sayer AA. Nutrition and sarcopenia: a review of the evidence and implications for preventive strategies. *Clinical Nutrition and Aging.* 2017:3-18.
- [20] Stenholm S, Harris TB, Rantanen T, et al. Sarcopenic obesity-definition, etiology and consequences. *Curr Opin Clin Nutr Metab Care.* 2008;11(6):693.
- [21] Hashemi R, Shafiee G, Motlagh AD, et al. Sarcopenia and its associated factors in Iranian older individuals: Results of SARIR study. *Arch Gerontol Geriatr.* 2016; 66:18-22.
- [22] Santos VRd, Araujo MYC, Cardoso MR, et al. Association of insufficient physical activity with sarcopenia and sarcopenic obesity in individuals aged 50 years or more. *Revista de Nutrição.* 2017;30:175-84.
- [23] Huschtscha Z, Parr A, Porter J, et al. Sarcopenic characteristics of active older adults: a cross-sectional exploration. *Sports Med Open.* 2021;7(1):32.
- [24] Hwang J, Park S. Gender-specific risk factors and prevalence for sarcopenia among community-dwelling young-old adults. *Int J Environ Res Public Health.* 2022;19(12):7232.
- [25] Hwang J, Park S. Sex Differences of sarcopenia in an elderly asian population: the prevalence and risk factors. *Int J Environ Res Public Health.* 2022; 19(19):11980.
- [26] Hwang J. Prevalence, Anthropometric risk factors, and clinical risk factors in sarcopenic women in their 40s. *J Korean Soc Phys Med.* 2023;18(2):23-31.
- [27] Hwang J. Age-related loss of skeletal muscle and associated risk factors in middle-aged men: a comprehensive study. *J Korean Soc Phys Med.* 2023; 18(2):13-21.

- [28] Hwang J, Park S. Gender-specific prevalence and risk factors of sarcopenic obesity in the Korean elderly population: A nationwide cross-sectional study. *Int J Environ Res Public Health*. 2023;20(2):1140.
- [29] Janssen I, Heymsfield SB, Wang ZM, et al. Skeletal muscle mass and distribution in 468 men and women aged 18-88 yr. *J Appl Physiol (1985)*. 2000;89(1):81-8.
- [30] Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc*. 2002;50(5):889-96.
- [31] Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: Revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48(1):16-31.
- [32] World Health O. ICD-10: International statistical classification of diseases and related health problems: tenth revision. (2nd ed ed). Geneva. World Health Organization. 2004.
- [33] Studenski SA, Peters KW, Alley DE, et al. The FNIH sarcopenia project: rationale, study description, conference recommendations, and final estimates. *J Gerontol A Biol Sci Med Sci*. 2014;69(5):547-58.
- [34] Kweon S, Kim Y, Jang MJ, et al. Data resource profile: The Korea national health and nutrition examination survey (KNHANES). *Int J Epidemiol*. 2014;43(1):69-77.
- [35] Han K, Park YM, Kwon HS, et al. Sarcopenia as a determinant of blood pressure in older Koreans: Findings from the Korea National Health and Nutrition Examination Surveys (KNHANES) 2008-2010. *PLoS One*. 2014;9(1):e86902.
- [36] Sanada K, Miyachi M, Tanimoto M, et al. A cross-sectional study of sarcopenia in Japanese men and women: reference values and association with cardiovascular risk factors. *Eur J Appl Physiol*. 2010;110(1):57-65.
- [37] Confortin SC, Meneghini V, Ono LM, et al. Anthropometric indicators as a screening tool for sarcopenia in older adults from Florianopolis, Santa Catarina: EpiFloripa ageing study. *Revista De Nutricao-Brazilian J Nutr*. 2017;30(3):287-96.
- [38] Zamboni M, Mazzali G, Fantin F, et al. Sarcopenic obesity: A new category of obesity in the elderly. *Nutr Metab Cardiovasc Dis*. 2008;18(5):388-95.
- [39] Nair KS. Aging muscle. *Am J Clin Nutr*. 2005; 81(5):953-63.
- [40] Tilg H, Moschen AR. Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nat Rev Immunol*. 2006;6(10):772-83.
- [41] Cesari M, Kritchevsky SB, Baumgartner RN, et al. Sarcopenia, obesity, and inflammation—results from the Trial of Angiotensin Converting Enzyme Inhibition and Novel Cardiovascular Risk Factors study. *Am J Clin Nutr*. 2005;82(2):428-34.
- [42] Lu CW, Yang KC, Chang HH, et al. Sarcopenic obesity is closely associated with metabolic syndrome. *Obes Res Clin Pract*. 2013;7(4):e301-7.
- [43] Buchmann N, Nikolov J, Spira D, et al. Identifying sarcopenia in metabolic syndrome: Data from the Berlin aging study II. *J Gerontol A Biol Sci Med Sci*. 2016;71(2):265-72.
- [44] Du Y, Wang X, Xie H, et al. Sex differences in the prevalence and adverse outcomes of sarcopenia and sarcopenic obesity in community dwelling elderly in East China using the AWGS criteria. *BMC Endocr Disord*. 2019;19(1):109.
- [45] Cleasby ME, Jamieson PM, Atherton PJ. Insulin resistance and sarcopenia: Mechanistic links between common co-morbidities. *J Endocrinol*. 2016;229(2): R67-81.
- [46] Schragger MA, Metter EJ, Simonsick E, et al. Sarcopenic obesity and inflammation in the InCHIANTI study. *J Appl Physiol (1985)*. 2007;102(3):919-25.
- [47] Therakomen V, Petchlorlian A, Lakananurak N. Prevalence and risk factors of primary sarcopenia in community-dwelling outpatient elderly: A cross-sectional study. *Sci Rep*. 2020;10(1):19551.