



# Evaluation of schistosomula lung antigen preparation and soluble egg antigen vaccines on experimental schistosomiasis mansoni



Nagwa S. M. Aly<sup>1,2,\*</sup> , Hye-Sook Kim<sup>2</sup> , Maysa A. Eraky<sup>1</sup>, Asmaa A. El Kholy<sup>1</sup>, Basma T. Ali<sup>1</sup>, Shin-ichi Miyoshi<sup>3</sup>, Rabab E. Omar<sup>1</sup>

<sup>1</sup>Department of Parasitology, Benha Faculty of Medicine, Benha University, Egypt; <sup>2</sup>Division of International Infectious Diseases Control, Faculty of Pharmaceutical Sciences, Okayama University, Tsushima-Naka, Kita-Ku, Okayama 700-8530, Japan;

<sup>3</sup>Department of Sanitary Microbiology, Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, Okayama University, Okayama 700-8530, Japan

## Abstract

Received: 13 November 2022

Accepted: 20 July 2023

### \*Correspondence

(drnagwashaban@gmail.com)

### Citation

Aly NSM, Kim HS, Eraky MA, Kholy AAE, Ali BT, Miyoshi SI, Omar RE. Evaluation of schistosomula lung antigen preparation and soluble egg antigen vaccines on experimental schistosomiasis mansoni. Parasites Hosts Dis 2023;61(3):251-262.

Schistosomiasis causes significant morbidity and mortality worldwide. This study aimed to assess the effect of schistosomula lung antigen preparation (SLAP) and soluble egg antigen (SEA) on a murine schistosomiasis mansoni model. Ninety laboratory-bred male Swiss albino mice were divided into 6 groups. Two doses of the vaccine were given at 2-week intervals. All mice were subcutaneously infected with  $80 \pm 10$  *Schistosoma mansoni* cercariae 2 weeks after the last vaccination dose. They were sacrificed 7 weeks post-infection. Parasitological and histopathological studies were conducted to assess the effect of inoculated antigens (single or combined). The results showed that the combination of SLAP and SEA (combination group) led to a significant reduction in worm burden (65.56%), and liver and intestine egg count (59% and 60.59%, respectively). The oogram pattern revealed a reduction in immature and mature eggs ( $15 \pm 0.4$  and  $10 \pm 0.8$ , respectively) and an increased number of dead eggs in the combination group ( $P < 0.001$ ). In terms of histopathological changes, the combination group showed notably small compact fibrocellular egg granuloma and moderate fibrosis in the liver. A high percentage of destroyed ova was observed in the intestine of the combination group. This study demonstrates for the first time the prophylactic effect of combined SLAP and SEA vaccine. The vaccine induced a significant reduction in the parasitological and pathological impacts of schistosomiasis mansoni in hepatic and intestinal tissues, making it a promising vaccine candidate for controlling schistosomiasis.

**Keywords:** Schistosomiasis, vaccine, SLAP, SEA

## Introduction

Schistosomiasis is one of the most neglected tropical diseases associated with significant morbidity, mortality, and socioeconomic impact in many developing countries of Africa, Latin America, and Asia [1]. It is a public health concern in 51 endemic countries. Less than 50% of the 237 million people who required preventive chemotherapy in 2019 were treated [2]. In Africa, more than 112 and 54 million individuals were reported to have urogenital and intestinal schistosomiasis, respectively [3]. Praziquantel (PZQ), the only medication used to treat schistosomiasis at present, has been administered to large numbers of people to manage the disease over the past few decades. Because PZQ is inefficient against juvenile schistosomes and does not prevent re-infection, the prevalence and transmission

© 2023 The Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Author contributions**

Conceptualization: Aly N, Eraky MA, El kholy A, Omar RE  
 Data curation: Ali B  
 Formal analysis: Aly N, Eraky MA, El kholy A, Omar RE  
 Funding acquisition: Miyoshi SI  
 Investigation: Ali B  
 Methodology: Ali B, Omar RE  
 Resources: Ali B, Miyoshi SI  
 Software: Aly N, Kim HS  
 Supervision: Aly N, Eraky MA, El kholy A, Omar RE  
 Validation: Aly N, Kim HS, Eraky MA  
 Visualization: Ali B  
 Writing – original draft: Aly N, Omar RE  
 Writing – review & editing: Aly N, Kim HS, Eraky MA, El kholy A, Miyoshi SI

**Conflict of interest**

The authors have declared that no competing interests exist.

**ORCID**

Nagwa S. M. Aly  
 (<https://orcid.org/0000-0001-5550-479X>)  
 Hye-Sook Kim  
 (<https://orcid.org/0000-0001-6142-5983>)

of schistosomiasis have generally remained unabated despite the intense efforts of mass drug administration programs. Additionally, other efforts including controlling snail intermediate hosts and water sanitation and hygiene programs have had little to no effect. These problems show that the current control measures are completely ineffective at stopping transmission, necessitating the use of new control strategies [4]. Hence, novel vaccines and strategies for preventing schistosomiasis and reinfections are urgently needed [5]. Several studies have reported promising results of vaccines using either radiated cercariae or recombinant antigens [6]. A single-antigen vaccine is likely to elicit less protection against schistosomiasis than a multi-antigen vaccine [7]. The efficacy of a vaccine depends on the choice of parasite stage and use of an effective adjuvant [8]. Soluble egg antigen (SEA) acts as a potential anti-schistosome immunization candidate [9]. *Schistosoma* miracidium inside the ovum secretes glycoprotein antigens that pass through microscopic pores within the eggshell, hence the term SEA. These antigens elicit a vigorous immune response that encapsulates the ova in pre-granuloma collagen fibers and immune cells, predominantly eosinophils and macrophages. The granuloma formation presents a barrier to sequester egg toxicity and antigenicity [10].

The migrating schistosomulum stage, represented as schistosomula lung antigen preparation (SLAP), has been found to induce consistently high protective responses against *Schistosoma* infection in laboratory animals [11]. Although human vaccines against schistosomiasis are currently unavailable, strong evidence from human field studies and animal models supports the probability of developing vaccines for long-term protection [12].

No previous study used a combination of SLAP and SEA antigens as a vaccine candidate. Thus, this study aimed to assess the effect of SLAP and SEA on a murine model of schistosomiasis.

## Materials and Methods

### Parasite

*Schistosoma mansoni* cercariae were obtained from infected *Biomphalaria alexandrina* snails, which were purchased from the Schistosome Biological Supply Unit, Theodor Bilharz Research Institute, Giza, Egypt, and maintained in the laboratory according to the guidelines proposed by Becker and Lamprecht [13]. *Schistosoma mansoni* cercariae from the snails (after exposure to light for at least 4h) were used to infect the mice through the subcutaneous route.

### Experimental animals

Laboratory-bred male Swiss albino mice weighing nearly 20–30 g were used in this study. All animal-related procedures in this study were in accordance with the International Guiding Principles for Biomedical Research, as issued by the International Organizations of Medical Science [14]. The study was revised and approved by the Institutional Animal Care and Use Committee, Medical Ethics Committee of the Faculty of Medicine, Benha University (approval no. MD 1-8-2021).

### Antigen preparation

SLAP preparation: Schistosomula were extracted using the perfusion technique from pul-

monary tissue 7 days post-infection. The lungs were placed in a small beaker with 2 ml of RPMI 1640 medium (Sigma, St. Louis, MO, USA) with penicillin and streptomycin and finely chopped to allow the schistosomula to migrate out of the capillary beds. The suspension was placed in a 50-ml centrifuge tube, incubated for 3 h at 37°C, and then centrifuged at 100 × g at 4°C. Thereafter, the supernatant was removed. The last pellet was added to 2–3 ml of RPMI 1640/penicillin/streptomycin in a petri dish with a diameter of 60 and 15 mm. The number of schistosomula in the total dish was counted using a dissecting microscope. The schistosomula were gathered using a fine-tip Pasteur pipette and transferred into culture jars for testing [15].

SEA preparation: Eggs were collected from liver tissues and suspended in 4°C PBS at a concentration of 100,000 eggs/ml. The eggs were homogenized on ice using a pre-chilled homogenizer and tight pestle. The crude mixture was centrifuged at 4°C at 200 × g for 20 min. The supernatant was collected and centrifuged for 90 min at 100,000 × g at 4°C. The supernatant was sterilized by passing it through a 0.2-µm filter [16]. The protein content of prepared antigens was determined according to Bradford's method [17]. The antigens were added with Complete Freund's Adjuvant (CFA) (Sigma) and emulsified in phosphate-buffered saline at a ratio of 2:1.

### Experimental design

Six groups of Swiss albino mice (15 mice in each group) were classified as follows: group I, non-infected and non-immunized (healthy control group); group II, infected and non-immunized (infected control group); group III, received CFA (50 µl); group IV, immunized with SLAP (50 µl SLAP+50 µl CFA); group V, immunized with SEA (100 µl SEA+50 µl CFA); and group VI, immunized with SLAP and SEA (combination group; 50 µl CFA+25 µl SLAP +50 µl SEA).

Each mouse received a single subcutaneous injection of vaccine. Then, a booster dose of vaccine was given 2 weeks later. Two weeks after the last dose of vaccine, all mice were infected with 80 ± 10 *S. mansoni* cercariae, which were injected subcutaneously in the abdomen [18]. The mice were sacrificed 7 weeks post-infection through rapid decapitation, and parasitological and histopathological studies were conducted to assess the effect of antigens.

**Table 1.** Mean worm burden and egg count in different groups

Groups No.	Worm burden		Egg count per gram of liver		Egg count per gram of intestine	
	Mean ± SD	R (%)	Mean ± SD	R (%)	Mean ± SD	R (%)
I	-	-	-	-	-	-
II	17.2 ± 2.9	-	2,548.1 ± 225.1	-	2,780.0 ± 170.8	-
III	17.1 ± 2.6	0.6	2,525.2 ± 221.4	0.9	2,750.2 ± 140.9	1.1
IV	11.0 ± 1.3	28.6*	1,755.0 ± 164.6	45.2*	2,100.3 ± 68.0	35.0*
V	14.2 ± 1.4	17.6*	1,922.1 ± 124.2	40.3*	2,222.1 ± 73.6	21.9*
VI	6.8 ± 0.9	65.6*	1,044.7 ± 103.7	59.0*	1,262.7 ± 63.3	60.6*

SD, standard deviation; R, reduction rate.


\*P-value < 0.05.

Group I: non-infected non-immunized (healthy control group). Group II: infected non-immunized. Group III: received Complete Freund's adjuvant (CFA). Group IV: immunized with SLAP. Group V: immunized with SEA. Group VI: immunized with SLAP and SEA (combination group).


### Evaluation of the vaccine efficacy

#### Parasitological studies

1. Worm distribution: Perfusion of sacrificed mice to detect adult worm distribution. The principle of this technique is to isolate the region to be examined and perfuse the vessels in



**Fig. 1.** Mean reduction rate of worm burden. Group I, non-infected non-immunized (Healthy control group); Group II, infected non-immunized; Group III, received Complete Freund’s adjuvant (CFA); Group IV, immunized with SLAP; Group V, immunized with SEA; Group VI, immunized with SLAP and SEA (combination group).



**Fig. 2.** Mean reduction rate of egg count per gram of liver and intestine tissues. Experimental conditions are indicated in the legend of Fig. 1.

**Table 2.** Oogram pattern and granuloma parameters (number and diameter) in different groups

Group No.	Oogram pattern			Granuloma number		Granuloma diameter	
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	R (%)	Mean ± SD	R (%) Mean ± SD
I	-	-	-	-	-	-	-
II	45.2 ± 1.8	50.2 ± 1.66	4.7 ± 1.9	10.0 ± 2.0	-	184.0 ± 22.0	-
III	45.0 ± 1.7	49.1 ± 1.5	4.8 ± 2.1	9.0 ± 2.41	10.0	170.0 ± 56.16	7.6
IV	35.4 ± 1.2*	27.0 ± 1.1 <sup>a</sup>	30.2 ± 3.0 <sup>a</sup>	4.0 ± 0.7	60.0*	85.0 ± 19.6	53.8*
V	38.2 ± 1.4*	29.0 ± 1.0 <sup>a</sup>	20.0 ± 6.0*	6.1 ± 3.6	4.0*	100.0 ± 34.1	45.0*
VI	15.0 ± 0.4*	10.0 ± 0.8 <sup>a</sup>	50.5 ± 4.33 <sup>b</sup>	1.5 ± 6.7	85.0*	50.0 ± 6.5	72.8*


SD, standard deviation; R, reduction rate.

\*P-value < 0.05.

Experimental conditions are indicated in the ligand of Table1.

which the worms may be present. The needle was connected to the automatic pumping machine and inserted in the inferior vena cava. Saline was pumped from the machine through the inferior vena cava to the liver. The worms were washed out and came out of the cut portal vein. Then, they were collected and counted [19].

2. Tissue egg counts: Pieces of the liver and intestine were removed from each animal after perfusion to determine the number of eggs/gram of tissue. The gut was slightly dilated and cleaned with saline to remove any fecal waste that was in the lumen. Ten milliliters of 5% potassium hydroxide solution was used for each piece of liver or intestine after it had been dried on filter paper, weighed, and deposited separately. The tissues were hydrolyzed after 24 h of incubation at 37°C. Three samples (0.1 ml each) were pipetted from the digest and placed on a counting slide. After they had been agitated vigorously on a magnetic mixer for 1 min, they were analyzed. The number of ova in each tissue was counted, and the average number was counted [20].




**Fig. 3.** Histopathological study of the liver stained with hematoxylin and eosin stain ( $\times 200$ ). (A) Normal hepatic lobular architecture. Bar = 100  $\mu\text{m}$ . (B) Cellular egg granuloma with intact central ovum (black arrow). (C-E) Egg granulomas with degenerated central ova (black arrows). (F) Fibrocellular egg granuloma with regular outline and degenerated central ovum (black arrow).

3. Oogram pattern: The oogram pattern was used to evaluate the degree of ova maturity and viability, which may reflect the vaccine's effect on oviposition and maturation [21]. The oogram showed the developmental stages of ova (immature, mature, and dead) in the small intestine of experimental animals.

#### Histopathological studies


Sections were stained with hematoxylin and eosin [22] or Masson's trichrome stain to demonstrate collagen fibers and measure granuloma dimensions easily. The liver sections were histopathologically examined for the presence of *Schistosoma* ova or granuloma, and their diameter and number were counted in each group. Intestine sections were examined for the presence or absence of *S. mansoni* ova, granuloma, any villous abnormality, and intensity of inflammatory cellular reaction.



**Fig. 4.** Histopathological examination of the liver stained with Masson's trichrome stain ( $\times 200$ ). Bar = 100  $\mu\text{m}$ . (A) Deposition of fibrous tissue. (B) Mild fibrosis around the granuloma (+) (black arrow). (C) Marked fibrosis (+++) around the granuloma. (D-F) Moderate fibrosis (++) around the granuloma.

### Statistical analysis

Data were analyzed by SPSS program (version 20; SPSS Inc., Chicago, IL, USA). Quantitative data were analyzed using mean and standard deviation. Reduction % (R%) = (No. of worms in the infected non-treated control group - No. of worms in the infected treated group  $\times$  100) / (No. of worms in the infected non-treated control group) [23]. Differences among the studied groups were estimated by one-way analysis of variance, and the Tukey-Kramer test was used as the post hoc test. Statistical significance was considered at  $P \leq 0.05$ .



**Fig. 5.** Histopathological study of the intestine stained with hematoxylin and eosin ( $\times 400$ ). Experimental conditions are indicated in the legend of Fig. 1. (A) Group I mice showing normal villus (black arrow). Bar = 100  $\mu$ m. (B) Group II mice showing many *S. mansoni* ova within the mucosa (black arrow). Bar = 50  $\mu$ m. (C) Group III mice showing *S. mansoni* ovum surrounded by a large number of inflammatory cells with a high percentage of eosinophils (yellow arrows). Bar = 50  $\mu$ m. (D) Group IV mice showing many *S. mansoni* ova within the mucosa with some intact (red arrow) and degenerated (yellow arrow) miracidia. Bar = 50  $\mu$ m. (E) Group V mice showing many *S. mansoni* ova within the mucosa (yellow arrow). Bar = 50  $\mu$ m. (F) Group VI mice showing many *S. mansoni* ova within the mucosa with mostly destroyed ova (yellow arrows). Bar = 50  $\mu$ m.

## Results

### Parasitological assessment

The vaccine's effect on worm burden is shown in Table 1 and Fig. 1. Group VI (combination group) was the most affected group, in which the reduction rate of the worm burden was statistically significant (65.5%), followed by groups IV and V with significant reduction rates of 28.63% and 17.57%, respectively ( $P < 0.01$ ).

In terms of the mean egg count (the mean number of eggs/gram of tissue) in the liver and intestine, group VI showed a significant reduction rate (59.0% and 60.59%, respectively) followed by groups IV (45.2% and 35.03%, respectively) and V (40.3% and 21.9%, respectively) (Table 1; Fig. 2).

The oogram pattern showed that group VI presented a statistically significant reduction of the mean number of immature and mature ova ( $15 \pm 0.4$  and  $10 \pm 0.8$ , respectively) followed by groups IV ( $35.4 \pm 1.2$  and  $27 \pm 1.1$ , respectively) and V ( $38.2 \pm 1.43$  and  $29 \pm 1.0$ , respectively). Additionally, group VI showed a statistically significant increase in the mean number of dead ova ( $50.5 \pm 4.33$ ), followed by groups IV and V ( $30.2 \pm 3.0$  and  $20.0 \pm 6.0$ , respectively) (Table 2).

### Histopathological results

Table 2 and Fig. 3 show that group VI had a significant reduction rate of granuloma number and diameter (85% and 72.8%, respectively), followed by groups VI (60% and 53%, respectively) and V (40% and 45%, respectively). The histopathological study of the liver stained by Masson's trichrome stain is shown in Fig. 4.

Group I is showing deposition of fibrous tissue. Group II has mild fibrosis around the granuloma (+). Group III has marked fibrosis (+++) around the granuloma. Vaccinated groups (IV, V, and VI) have moderate fibrosis (++) around the granuloma. Regarding hematoxylin and eosin stains of the intestine (Fig. 5), the intestine sections of group I showed normal villous (left) and crypt (right) patterns. All infected groups showed viable ova with intact miracidium and acute inflammatory cellular reaction around *Schistosoma* ova, with a high percentage of eosinophils. Group VI showed mostly destroyed ova.

## Discussion

Identifying vaccine targets is difficult because of the complexity of the schistosome's life cycle. Many experimental trials to develop an effective human vaccine for schistosomiasis have been reported. More than 100 schistosome vaccine antigens have demonstrated some degree of protection in murine schistosomiasis models. A vaccine would lessen schistosomiasis morbidity by inducing immune responses that would decrease the parasite load and diminish egg production [24]. The *Schistosoma* parasite secretes and excretes various antigens into the host's bloodstream; these antigens are divided into cercarial, adult worm, and egg antigen categories depending on the state of the parasite's development [25]. Many laboratories have isolated different antigens from crude SEA and soluble worm antigen preparation (SWAP). A previous study has shown that antigens derived from various phases of the parasite's life can trigger a protective reaction in mice [26]. Studies on the preventive



and anti-fecundity effects of Sm-p80 have shown that combining different antigens can result in higher levels of vaccine-induced protection. For example, an adjuvant, synthetic hexaacetylated lipid A derivative (glucopyranosyl lipid A) formed in aluminum has been combined with recombinant Sm-p80 to increase the vaccine's efficacy. The biological functions of schistosomes such as schistosomula migration and fecundity represent key potential sites to target parasites for elimination through vaccination [27]. During the acute stage, the predominant immune response to schistosomula antigens and as the disease progresses, the body response is caused by egg antigens [28]. Therefore, this research assessed the prophylactic impact of SLAP combined with SEA as a new vaccine candidate on a murine model of schistosomiasis.

The results in the present study showed variable degrees of reduction in worm burden. A significant reduction in worm burden was observed in the combination group (65.56%). Many previous reports showed variable degrees of reduction in worm burden in vaccinated groups [29]. The variation between their data and our data may be a result of using different antigen doses and schistosomula age as schistosomula were extracted from the lungs 18 days post-infection.

Egg reduction is more influential in vaccine efficacy than the reduction of adult worms as a reduction in hepatic egg count reflects a decrease in eggs entrapped in the liver, leading to improvement in liver pathology, reduction in intestinal egg count, and decreased transmission rate and pathology of the parasite [30].

The combination of SLAP and SEA resulted in a significant reduction in egg count per gram of liver and intestine tissues (59% and 60.59%, respectively). Our results were in accordance with those of Etewa et al. [9], who observed a significant reduction of *S. mansoni* eggs in the liver. Lower values were obtained by El-Ahwany et al. [29], with 42.8% and 8.4% reduction of eggs in the liver and tissue, respectively.

The reduction of the immature and mature egg count in vaccinated groups goes in parallel with the significant reduction in the adult count recorded in the present study. These changes could be explained by the cessation of oviposition because of the death of adult worms or the reduction of female fecundity because of the decoupling effect [31].

In the present study, different groups had a reduction in granuloma count and granuloma diameter, with the combination group having the most significant reduction (85% and 72.8%, respectively). These results reflect that the reduction of granuloma number might be related to a decrease in tissue egg count rather than the inability of eggs to initiate granuloma formation. The granulomatous reaction may be attributed to SEA-specific stimulation of immune reaction against *Schistosoma* eggs in hepatic pathology. These results were consistent with those of other studies [32], indicating that the vaccine could induce a partial reduction in worm burden and decrease the granuloma number and diameter, which were attributed to an increase in CD8+ cells in immunized groups and a reduction of CD4+ cells.

Owing to the downregulation of immune response, newly formed granulomas are smaller and contain killed eggs; meanwhile, granulomas that are formed earlier gradually resolve to be replaced by fibrous tissue, leading to the amelioration of symptom severity [33]. Fibrosis is the excessive accumulation of fibrous connective tissue (e.g., collagen and fibronectin) in and around inflamed or damaged tissue [34]. This indicates a good protective

immune response to the combined vaccine in our study. In Etewa et al.'s study [9], the liver sections of the SEA-vaccinated group had fewer numbers and smaller sizes of granuloma; degenerated ova surrounded by empty spaces were enclosed in fibrocellular granulomas in addition to mild lobular inflammation.

In the present study, no statistical difference was observed between the group receiving CFA adjuvant (III) and the infected non-immunized group (II). In addition, the SLAP-immunized group (IV) obtained better results than the SEA-immunized group (V). The administration of SLAP and SEA (VI) prior to infection resulted in decreased worm load and hepatic and intestinal ova and changes in the oogram pattern. This could be due to the augmentation of immune response or a result of primary infection. According to Etewa et al. [9], SEA is preferable than SWAP. The ovum's *Schistosoma* miracidium (SEA) secretes glycoprotein antigens, which pass through tiny openings in the eggshell. A strong immune reaction to these antigens surrounds the ova in pre-granuloma collagen fibers and immune cells, primarily eosinophils and macrophages. Granuloma formation is a barrier to sequestering egg toxicity and antigenicity. Moreover, proteins were identified from *S. mansoni* SLAP, which can stimulate protective Th1 cell-mediated immune responses [35]. A purified SLAP preparation causes a significant decrease in worm burden, egg load, granuloma diameter, and collagen content and increase in the proportion of degenerated ova. Moreover, it can improve pathological alterations in pulmonary and hepatic tissues [29]. However, which of these immune responses is directly involved in worm-killing and protective immunity remains unknown. Further studies are needed to assess the effect of our vaccine on Th1 and Th2.

An effective anti-schistosome vaccine would greatly decrease the morbidity associated with schistosomiasis via protective immune responses, leading to reduced worm burden and decreased egg production and its pathology [24]. The combined vaccine apparently induced an improved immune response against antigenic components.

In conclusion, the obtained parasitological and histopathological findings highlighted the protective response of the combined vaccine (SLAP and SEA). The combination of SEA and SLAP antigens induced a marked decrease in worm burden, egg load, and granuloma count and diameter and an improvement of pathological changes in hepatic and intestine tissue.

## Acknowledgments

This study was partially supported by a grant from the Program of the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID, JP22wm0125004) from the Ministry of Education, Culture, Sports, Science and Technology in Japan (MEXT), and the Japan Agency for Medical Research and Development (AMED).

We are deeply grateful to Prof. Tarek Aboushousha, Department of Pathology, Theodor Bilharz Research Institute, for his important role in performing the histopathology of this study.

## References

1. Odhiambo GO, Musuva RM, Atuncha VO, Mutete ET, Odiere MR, et al. Low levels of awareness despite high prevalence of schistosomiasis among communities in Nyalenda informal settlement, Kisumu City, Western Kenya. *PLoS Negl Trop Dis* 2014;8(4):e2784. <https://doi.org/10.1371/journal.pntd.0002784>
2. World Health Organization. Schistosomiasis: Key Facts [Internet]. Available from: <https://www.who.int/news-room/factsheets/detail/schistosomiasis>
3. World Health Organization. Current estimated total number of individuals with morbidity and mortality due to *Schistosoma haematobium* and *S. mansoni* infection in Sub-Saharan Africa. Schistosomiasis; 2017. [Internet]. Available from: <https://www.who.int/teams/control-of-neglected-tropical-diseases/schistosomiasis/epidemiology>
4. Campbell SJ, Biritwum NK, Woods G, Velleman Y, Fleming E, et al. Tailoring water, sanitation, and hygiene (WASH) targets for soil-transmitted Helminthiasis and Schistosomiasis control. *Trends Parasitol* 2018; 34(1):53-63. <https://doi.org/10.1016/j.pt.2017.09.004>
5. Yepes E, Varela MRE, Lopez-Aban J, Roj-as-Caraballo A, Muro A. Inhibition of granulomatous inflammation and prophylactic treatment of schistosomiasis with a combination of Edelfosine and praziquantel. *PLoS Negl Trop Dis* 2015;9(7):30-50. <https://doi.org/10.1371/journal.pntd.0003893>
6. Molehin AJ, Rojo JU, Siddiqui SZ, Gray SA, Carter D, et al. Development of a schistosomiasis vaccine. *Expert Rev Vaccines* 2016;15(5):619-627. <https://doi.org/10.1586/14760584.2016.1131127>
7. Ewaisha RE, Bahey-El-Din M, Mossallam SF, Amer EL, Aboushleib HM, et al. Combination of the schistosomal antigens Sm14 and Sm29 elicits significant protection against experimental *Schistosoma mansoni* infection. *Exp Parasitol* 2014;145: 51-60. <https://doi.org/10.1186/s12879-015-0906-z>
8. Araujo JM, de Melo TT, de Sena IC, Alves CC, Araujo N, et al. *Schistosoma mansoni* schistosomula tegument (Smtcg) immunization in absence of adjuvant induce IL-10 production by CD4+ cells and failed to protect mice against challenge infection. *Acta Trop* 2012; 124:140-146. <https://doi.org/10.1371/journal.pone.0160118>
9. Etewa S, AL-Hoot AAA, Sharaf HM, Moawad HS, Mohamed SM, et al. Modelling approaches to predict and evaluate schistosomiasis immunization utilizing SEA loaded on chitosan nanoparticles via liver tissue differentiation and angiogenesis. *Parasitol United J* 2019; 12:197-208. <https://doi.org/10.21608/puj.2019.16532.1051>
10. Hams E, Aviello G, Fallon PG. The *Schistosoma* granuloma: friend or foe? *Front Immunol* 2013;4:1-8. <https://doi.org/10.3389/fimmu.2013.00089>
11. Al-Naseri A, Al-Absi S, El Ridi R. Comprehensive and critical overview of schistosomiasis vaccine candidates. *J Parasit Dis* 2021;45(2):557-580. <https://doi.org/10.1007/s12639-021-01387-w>
12. Fonseca CT, Oliveira SC, Alves CC. Eliminating Schistosomes through vaccination: what are the best immune weapons? *Front Immunol* 2015;6:95. <https://doi.org/10.3389/fimmu.2015.00095>
13. Becker W, Lamprecht I. Microcalorimetric investigations of the behavior and infectivity of miracidia of *Biomphalaria glabrata* and *Schistosoma mansoni*. *Z Parasitenkd* 1977;53(3):297-305 (in German). <https://doi.org/10.1007/BF00389947>
14. National Research Centre. Guide for the Care and Use of Laboratory Animals. 8th ed. National Academies Press. Washington DC, USA. 2011. Available from: <https://grants.nih.gov/grants/olaw/guide-for-the-care-and-use-of-laboratory>
15. Imohiosen AS. Early developmental changes of the schistosomula of *S. mansoni* in vitro and in mouse lung. *Parasitol* 1978; 76(3):317-326. <https://doi.org/10.1017/s0031182000048198>
16. Tucker MS, Karunaratne LB, Lewis FA, Freitas TC, Liang YS. Schistosomiasis. *Curr Protoc Immunol* 2013;103:19.1.1-19.1.58. <https://doi.org/10.1002/0471142735.im1901s103>
17. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal Biochem* 1976;72:248-254. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
18. Peters AP, Warren KS. A rapid method of infecting mice and other laboratory animals with *S. Mansoni* subcutaneous infection. *J Parasitol* 1969;55(3):558. <https://doi.org/10.2307/3277297>
19. Duvall RH, Dewitt WB. An improved perfusion technique for recovering adult schistosomes from laboratory animals. *Am J Trop Med Hyg* 1967;16(4):483-486. <https://doi.org/10.4269/ajtmh.1967.16.483>
20. Cheever AW. Conditions affecting the accuracy of potassium hydroxide digestion techniques for counting *Schistosoma mansoni* eggs in tissues. *Bull World Health Organ* 1968;39(2):328-331
21. Pellegrino J, Oliveira CA, Faria J, Cunha AS. New approach to the screening of drugs in experimental schistosomiasis mansoni. *Am J Trop Med Hyg* 1962;11:201-215. <https://doi.org/10.4269/ajtmh.1962.11.201>
22. Feldman AT, Wolfe D. Tissue processing and hematoxylin and eosin staining. *Methods Mol Biol* 2014;1180:31-43. [https://doi.org/10.1007/978-1-4939-1050-2\\_3](https://doi.org/10.1007/978-1-4939-1050-2_3)
23. Brown A. A new software for carrying out one-way ANOVA post hoc tests. *Comput Methods Programs Biomed* 2005;79(1): 89-95. <https://doi.org/10.1016/j.cmpb.2005.02.007>
24. Siddiqui AA, Bilal A, Siddiqui BA, Ganley-Leal L. Schistosomiasis vaccines. *Hum Vaccin* 2011;7(11):1192-1197. <https://doi.org/10.4161/hv.7.11.17017>
25. Van Lieshout L, Polderman AM, Deelder AM. Immunodiagnosis of schistosomiasis by determination of the circulating antigens CAA and CCA, in particular in individuals with recent or light infections. *Acta Trop* 2000;77(1):69-80. [https://doi.org/10.1016/S0167-2825\(99\)00089-9](https://doi.org/10.1016/S0167-2825(99)00089-9)

- doi.org/10.1016/s0001-706x(00)00115-7
26. Teixeira de Melo T, Michel de Araujo J, Do Valle Durães E, Caliari MV, Oliveira SC, et al. Immunization with newly transformed *Schistosoma mansoni* schistosomula tegument elicits tegument damage, reduction in egg and parasite burden. *Parasite Immunol* 2010;32(11-12): 749-759. <https://doi.org/10.1111/j.1365-3024.2010.01244.x>
  27. Gobert GN, You H, McManus DP. Gaining biological perspectives from schistosome genomes. *Mol Biochem Parasitol* 2014; 196(1):21-28. <https://doi.org/10.1016/j.molbiopara.2014.07.007>
  28. Costain A, Mac Donald A, Smits HH. Schistosome egg migration: mechanisms, pathogenesis and host immune responses. *Front Immunol* 2018;9:3042. <https://doi.org/10.3389/fimmu.2018.03042>
  29. El-Ahwany E, Bauomy IR, Nagy F, Zalat R, Mahmoud O, et al. Regulatory cell responses to immunization with a soluble egg antigen in *Schistosoma mansoni* infected mice. *Korean J Parasitol* 2012;50(1):29-35. <https://doi.org/10.3347/kjp.2012.50.1.29>
  30. Ranasinghe SL, Duke M, Harvie M, McManus DP. Kunitz-type protease inhibitor as a vaccine candidate against schistosomiasis *mansoni*. *Int J Infect Dis* 2018;66:26-32. <https://doi.org/10.1016/j.ijid.2017.10.024>
  31. Mati VL, Melo AL. Current applications of oogram methodology in experimental schistosomiasis; fecundity of female *Schistosoma mansoni* and egg release in the intestine of AKR/J mice following immunomodulatory treatment with pentoxifylline. *J Helminthol* 2013;87(1):115-24. <https://doi.org/10.1017/S0022149X12000144>
  32. Pinho JMR, Cardoso FC, Lopes DO, Pinheiro CS, Caliari MV. Immunization with SmIg, a novel tegument protein from *Schistosoma mansoni*, fails to induce protection in mice but reduces liver pathology. *Parasitology* 2010;137(7):1079-1088. <https://doi.org/10.1017/S0031182009991387>
  33. McManus DP, Dunne DW, Sacko M, Utzinger J, Vennervald BJ, et al. Schistosomiasis. *Nat Rev Dis Primers* 2018;4(1):13. <https://doi.org/10.1038/s41572-018-0013-8>
  34. Wynn TA. Integrating mechanisms of pulmonary fibrosis. *J Exp Med* 2011;208(7):1339-1350. <https://doi.org/10.1084/jem.20110551>
  35. Mounford AP, Harrop R. Vaccination against schistosomiasis: the case for lung-stage antigens. *Parasitol Today* 1998;14(3):109-114. [https://doi.org/10.1016/s0169-4758\(97\)01169-1](https://doi.org/10.1016/s0169-4758(97)01169-1)