



A case of hepatic anisakidosis caused by *Anisakis pegreffii* mimicking liver cancer



Minoru Yamada^{1,*} , Fumi Murakoshi^{1,2} , Hisashi Ikoma³ , Osamu Inamori⁴ , Akio Yanagisawa⁴ , Eiichi Konishi⁴

¹Department of Infectious Diseases, Kyoto Prefectural University of Medicine, Kyoto 602-8566, Japan; ²The Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Sendai 980-8578, Japan; ³Department of Gastroenterological Surgery, Kyoto Prefectural University of Medicine, Kyoto 602-8566, Japan; ⁴Department of Surgical Pathology, Kyoto Prefectural University of Medicine, Kyoto 602-8566, Japan

Abstract

Received: 9 May 2023
Accepted: 25 June 2023

*Correspondence
(minorori@koto.kpu-m.ac.jp)

Citation

Yamada M, Murakoshi F, Ikoma H, Inamori O, Yanagisawa A, Konishi E. A case of hepatic anisakidosis caused by *Anisakis pegreffii* mimicking liver cancer. Parasites Hosts Dis 2023;61(3):292-297.

Extra-gastrointestinal anisakidosis is rare. We herein report an *Anisakis pegreffii* infection in a patient with hepatic anisakidosis diagnosed based on its molecular identification. A 71-year-old male patient had a hepatic tumor presenting as a low-density area of 20 mm in diameter in segment 6 of the liver on abdominal ultrasonography, computed tomography, and magnetic resonance imaging. The surgically resected pathological specimen revealed a necrotizing eosinophilic granuloma containing nematode larvae, possibly an *Anisakis* larva. Molecular and phylogenetic analysis demonstrated *Anisakis* larvae belonging to *A. pegreffii*. The present results will help identify and characterize unknown *Anisakis* species in histological sections.

Keywords: *Anisakis pegreffii*, hepatic anisakidosis, granuloma

Introduction

Raw or undercooked marine fish consumption may cause infection with several helminths. Anisakidosis is a foodborne disease caused by the accidental ingestion of larval nematode (commonly *Anisakis* spp. or *Pseudoterranova* spp.) in raw marine fishes [1-3]. Anisakidosis incidence continues to increase globally due to the growing popularity of consuming raw or undercooked fish [4,5]. Annually, over 2,000 cases of anisakidosis are assumed to occur in most regions of Japan [3]. Approximately 95% of larvae parasitize the gastric, small intestinal, and colon mucosae, when *Anisakis* larvae orally infect the digestive tract lumen. However, they occasionally perforate the intestinal wall and resulting in ectopic anisakidosis cases. By the mid-1990s, 769 cases of anisakidosis caused by *Pseudoterranova decipiens* (this species is widely known in Japan) were reported in the northern part of Japan [3]. The ectopic form of anisakidosis is much less common than the gastric and intestinal forms, but ectopic anisakidosis caused by *Pseudoterranova* spp. has been reported in humans [6,7]. A biopsy sample obtained under endoscopy from patients with acute or subacute abdominal symptoms is used to generally diagnose *Pseudoterranova* infection. However, a previous study reported that the 4th-stage larva of *P. azarasi* was expectrated by strong coughing after an asymptomatic course [6]. To date, 3 sibling species of *Anisakis simplex* larvae have been reported: *A. simplex* sensu stricto (s.s.), *A. pegreffii*, and *A. simplex* C [8]. The most common species in Japan is *A. simplex* (s.s.) followed by *A. pegreffii*, and *A. simplex* (s.s.) caused 99% of all human anisakidosis [3,9]. The total number of anisakidosis cases

© 2023 The Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author contributions

Conceptualization: Yamada M, Murakoshi F, Ikoma H
 Investigation: Inamori O
 Methodology: Murakoshi F
 Supervision: Yanagisawa A, Konishi E
 Visualization: Ikoma H
 Writing – original draft: Yamada M, Murakoshi F
 Writing – review & editing: Yamada M, Murakoshi F

Conflict of interest

We have no conflict of interest related to this work.

ORCID

Minoru Yamada
<https://orcid.org/0000-0002-4941-6117>
 Fumi Murakoshi
<https://orcid.org/0000-0002-8784-717X>
 Hisashi Ikoma
<https://orcid.org/0000-0002-2647-6854>
 Osamu Inamori
<https://orcid.org/0000-0003-2366-7863>
 Akio Yanagisawa
<https://orcid.org/0009-0005-6063-3111>
 Eiichi Konishi
<https://orcid.org/0000-0002-7122-2271>

reported each year is 70 in the United States of America and 500 in Europe [1]. Herein, we report a 71-year-old male who presented with liver cancer and was diagnosed with extra-gastrointestinal anisakidosis caused by *Anisakis pegreffii*.

Case Records

A 71-year-old male patient was admitted to Kyoto Prefectural University of Medicine Hospital with a liver tumor impression. He had a previous history of hypertension, gout, and prostatomegaly. He underwent appendectomy at 30 years of age. He had a family history of lung cancer, which affected his mother. Abdominal ultrasonography performed in health check-up revealed a low-density area of 20 mm in diameter in segment 6 of the liver. He was referred to our hospital for a more detailed lesion examination after 1 month. Difficulties were associated in reaching a definitive diagnosis although computed tomography (CT)-ethoxybenzyl-diethylenetriamine pentaacetic acid-enhanced magnetic resonance imaging (EOB-MRI) images were investigated (Figs. 1, 2). Therefore, a liver biopsy was performed and tissue sections were examined. Preoperative diagnosis includes a liver tumor (S6), and a differential diagnosis from hepatocellular carcinoma, metastatic liver cell carcinoma, cholangiocarcinoma in the liver, and an inflammatory pseudotumor was required. Borderline-limited necrotizing granuloma comprising coagulation necrosis containing eosinophilic ghost cells and liquefactive necrosis with a desquamated cytoplasm was revealed in a pathological examination of a tissue section from a surgical sample obtained under laparoscopy. Necrotizing granuloma of the S6 lesion in the center of the tissue section was partially resected, revealing avascular necrosis in the cavity, and contained a necrotic larval nematode with a thick pellicle and amorphous content, indicating an anisakis infection (Figs. 3, 4). Necrotizing eosinophilic granulomatous connective tissues surrounding the larval nematode were visualized with hematoxylin and eosin staining, but were not clarified by Victorian-blue or elastica staining (Figs. 5, 6). DNA samples were extracted from parasite-positive and -negative regions in paraffin-embedded sections using the DEXPAT reagent (TaKaRa, Shiga, Japan) and the QIAmp DNA Mini Kit (Qiagen GmbH, Hilden, Germany) to




Fig. 1. Abdominal CT. (A) Early stage of arterial phase. Contrast CT after 30 sec. (B) Equilibrium stage of arterial phase. Contrast CT after 180 sec. Arrows show a low-density area of 20 mm in size in segment 6 of the liver. CT, computed tomography.




Fig. 2. (A) EOB-MRI. Left: Early stage of arterial phase. Right: Equilibrium stage of arterial phase. Arrows show a low-density area of 20 mm in size in segment 6 of the liver. (B) EOB-MRI. Liver cell phase of arterial phase. Arrow shows low-density signal regions that showed lighter color than those of the early and equilibrium stages. EOB-MRI, ethoxybenzyl-diethylenetriamine pentaacetic acid-enhanced magnetic resonance imaging.




Fig. 3. Necrotizing granuloma of the S6 lesions in the liver section partially resected.




Fig. 4. Low magnification of the tissue section infected with the worm in the S6 lesion of the liver. Bar = 1 cm.

molecularly identify the larvae in the nematode-positive lesion. A nuclear DNA region of internal transcribed spacer (ITS)1, 5.8S rRNA and ITS2, and a mitochondrial DNA region




Fig. 5. (A) Larval nematode, *Anisakis* sp. in the tissue section of the S6 lesion of the liver. Magnification ($\times 100$). Bar = 200 μm . (B) Higher magnification of the larval nematode, *Anisakis* sp. in the tissue section of the S6 lesion of the liver. Magnification ($\times 200$). Bar = 100 μm .




Fig. 6. Phylogenetic tree based on partial sequences of the COX2 sequence constructed using the maximum likelihood method with 468 nucleotides without gaps. Substitution model and optional parameters = GTR+ Γ . Only bootstrap values of > 50% from 1,000 pseudo-replicates are shown.

of COX2 are used to identify *Anisakis* larvae. The COX2 region was amplified by polymerase chain reaction (PCR) and sequenced in the present case. The primers used for COX2 were 5'-TCAGGATTTTGGTTTGATGTTT-3' and 5'-ATTCTCCATAAAACCTATACAC-3' [10]. A PCR reaction was performed under the following conditions: samples were denatured at 95°C for 3 min and then subjected to 40 cycles of 94°C for 30 sec, 48°C for 40 sec, and 72°C for 50 sec, with final extension at 72°C for 7 min on a thermocycler (GeneAmp PCR System 9700, Applied Bio-systems, Foster City, CA, USA). The PCR product (682 bp) was purified with a QIAquick PCR Purification Kit (Qiagen GmbH) and sequenced. Clustal X2 [11] aligned sequences and BioEdit 7.0.5.3 manually edited computed sequences [12].

All gaps were eliminated and *COX2* sequences were used for the phylogenetic analysis. A maximum likelihood (ML) analysis was performed with MEGA 6.0.6 [13]. The substitution model and optional parameter sets used were assessed with MEGA 6.0.6, and the most suitable sets were selected following the Akaike information criterion. The same datasets were used to construct 1,000 ML trees to calculate the bootstrap values. The final dataset contained 468 positions. Molecular and phylogenetic analysis of *COX2* (468 bp, partial sequence) sequences revealed *Anisakis* larvae belonging to *A. pegreffii* (Fig. 6).

Discussion

Anisakidosis is one of the most common fishborne helminthic diseases in Japan, which is contracted by ingesting the nematode *Anisakis* spp larvae, carried by marine fish [1]. The dominant species in fish caught offshore Japan included *A. simplex* (s.s.) and *A. pegreffii* [9]. A previous study revealed that significantly higher rates of *A. simplex* (s.s.) penetrated agar than *A. pegreffii*, indicating that its larvae can survive in acidic gastric juice to some extent and penetrate the stomach, small intestine, or colon of infected humans [14]. Furthermore, in vitro and in vivo studies revealed that *A. pegreffii* can pathogenically cause anisakidosis in humans when ingested, similar to *A. simplex* (s.s.) [15]. Recent cases in Italy indicated that *A. pegreffii* is a major cause for concern in countries in which it is dominant [16], similar to *A. simplex* (s.s.) in Japan [14]. The present case report is crucial because it describes a rare hepatic anisakidosis case caused by *A. pegreffii*, which is mostly encountered in gastric, intestinal, and extra-gastrointestinal (ectopic) cases mainly caused by *A. simplex* (s.s.) in Japan [14]. Difficulties were associated with differentiating between liver cancer and parasitic granuloma solely based on imaging modalities. An accurate diagnosis is possible using PCR based on *COX2* genes subjected to a mtDNA sequence analysis although ectopic anisakidosis is challenging to diagnose using pathological specimens. A molecular/genetic methodological approach allows for the easy and rapid *Anisakis* spp. identification in formalin-fixed and paraffin-embedded tissues obtained from gastric or intestinal human anisakidosis cases [16,17]. Laparoscopic partial hepatectomy for a pathological examination and the genetic anisakidosis identification by PCR were useful for achieving a final diagnosis even when degenerative granuloma surrounding larval bodies necrotizing in the resected specimen collapses with time. Only 8 hepatic anisakidosis cases have been reported from Japanese people who frequently consume undercooked fish [18,19]. In conclusion, the present results emphasize the rarity of ectopic anisakidosis, which may help identify and characterize unknown *Anisakis* species in biopsied histological sections misdiagnosed as liver cancer.

Acknowledgments

This study was performed in accordance with the ethical standards of the Institutional Review Board of the Kyoto Prefectural University of Medicine and with the Declaration of Helsinki as revised in 2013 and an informed consent for publication was obtained from the patient.

References

- Chai JY, Murrell KD, Lamberty AJ. Fish-borne parasitic zoonoses: status and issues. *Int J Parasitol* 2005;35(11-12):1233-1254. <https://doi.org/10.1016/j.ijpara.2005.07.013>
- Sohn WM, Chai JY. Anisakiosis (anisakidosis). In Palmer SR, Soulsby L, Torgerson PR, Brown WG, eds. *Oxford Textbook of Zoonoses*. 2nd ed. Oxford University Press. Oxford, UK. 2011, pp 774-786.
- Ishikura H. 5. Anisakiasis. (2) Clinical pathology and epidemiology. In Otsuru M, Kamegai S, Hayashi S. eds, *Progress of Medical Parasitology in Japan*. vol.8. Meguro Parasitological Museum, Tokyo, Japan. 2003, pp 451-473.
- Takahashi S, Ishikura H, Kikuchi K. Anisakidosis: global point of view. In Ishikura H ed, *Host Response to International Parasitic Zoonoses*. Springer-Verlag, New York, USA. 1998, pp 109-120.
- Takabe K, Ohki S, Kunihiro O, Sakashita T, Endo I, et al. Anisakidosis: a cause of intestinal obstruction from eating sushi. *Am J Gastroenterol* 1998;93(7):1172-1173. <https://doi.org/10.1111/j.1572-0241.1998.00356.x>
- Arizono N, Miura T, Yamada M, Tegoshi T, Onishi K. Human infection with *Pseudoterranova azarasi* roundworm. *Emerg Infect Dis* 2011;17(3):555-556. <https://doi.org/10.3201/eid1703.101350>
- Mitsuboshi A, Yamaguchi H, Ito Y, Mizuno T, Tokoro M, et al. Extra-gastrointestinal anisakidosis caused by *Pseudoterranova azarasi* manifesting as strangulated inguinal hernia. *Parasitol Int* 2017;66(6):810-812. <https://doi.org/10.1016/j.parint.2017.09.008>
- Mattiucci S, Nascetti G. Advances and trends in the molecular systematics of anisakid nematodes, with implications for their evolutionary ecology and host-parasite co-evolutionary processes. *Adv Parasitol* 2008;66:47-148. [https://doi.org/10.1016/S0065-308X\(08\)00202-9](https://doi.org/10.1016/S0065-308X(08)00202-9)
- Umehara A, Kawakami Y, Araki J, Uchida A. Molecular identification of the etiological agent of the human anisakiasis in Japan. *Parasitol Int* 2007;56(3):211-215. <https://doi.org/10.1016/j.parint.2007.02.005>
- Valentini A, Matticci S, Bondanelli P, Webb SC, Mignucci-Giannone AA, et al. Genetic relationships among *Anisakis* species (Nematoda: Anisakidae) inferred from mitochondrial *cox2* sequences and comparison with allozyme data. *J Parasitol* 2006;92(1):156-166. <https://doi.org/10.1645/GE-3504.1>
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettngen PA, et al. Clustal W and Clustal X version 2.0. *Bioinformatics* 2007;23(21):2947-2948. <https://doi.org/10.1093/bioinformatics/btm404>
- Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 1999;41:95-98.
- Tamura K, Stecher G, Peterson D, Filipiński A, Kumar S. MEGA6: molecular evolutionary genetics analysis Version 6.0. *Mol Biol Evol* 2013;30(12):2725-2729. <https://doi.org/10.1093/molbev/mst197>
- Arizono N, Yamada M, Tegoshi T, Yoshikawa M. *Anisakis simplex* sensu stricto and *Anisakis pegreffii*: biological characteristics and pathogenic potential in human anisakiasis. *Foodborne Pathog Dis* 2012;9(6):517-521. <https://doi.org/10.1089/fpd.2011.1076>
- Jeon CH, Kim JH. Pathogenic potential of the two sibling species, *Anisakis simplex* (s.s.) and *Anisakis pegreffii* (Nematoda: Anisakidae): *in vitro* and *in vivo* studies. *Biomed Res Int* 2015; 2015:983656. <https://doi.org/10.1155/2015/983656>
- Mattiucci S, Paoletti M, Borriani F, Palumbo M, Palmieri RM, et al. First molecular identification of the zoonotic parasite *Anisakis pegreffii* (Nematoda: Anisakidae) in a paraffin-embedded granuloma taken from a case of human intestinal anisakiasis in Italy. *BMC Infect Dis* 2011;11:82. <https://doi.org/10.1186/1471-2334-11-82>
- Mladineo I, Popović M, Drmić-Hofman I, Poljak V. A case report of *Anisakis pegreffii* (Nematoda, Anisakidae) identified from archival paraffin sections of a Croatian patient. *BMC Infect Dis* 2016;16:42. <https://doi.org/10.1186/s12879-016-1401-x>
- Kagei N, Orikasa H, Hori E, Sannomiya A, Yasumura Y. A case of hepatic anisakiasis with a literal survey for extra-gastrointestinal anisakiasis. *Jpn J Parasitol* 1995;44:346-351.
- Murata Y, Ando K, Usui M, Sugiyama H, Hayashi A, et al. A case of hepatic anisakiasis caused by *Pseudoterranova decipiens* mimicking metastatic liver cancer. *BMC Infect Dis* 2018; 18(1):619. <https://doi.org/10.1186/s12879-018-3540-8>