

# Effect of Synthetic CaM and NFAT Oligodeoxynucleotide on MPP<sup>+</sup>-Stimulated Mesencephalic Neurons

Jihyun Park<sup>1</sup>, Kyung Mi Jang<sup>2</sup>

<sup>1</sup>Department of Pathology, College of Medicine, Catholic University of Daegu, Daegu, Korea

<sup>2</sup>Department of Pediatrics, College of Medicine, Yeungnam University, Daegu, Korea

**Background:** Ca<sup>2+</sup> signaling plays a vital role in neuronal signaling and altered Ca<sup>2+</sup> homeostasis in Parkinson's disease (PD). Over-expression of αSYN significantly promote the Ca<sup>2+</sup>-Calmodulin (CaM) activity and subsequent nuclear translocation of nuclear factor of activated T cells (NFAT) transcription factor in dopaminergic neurons of midbrain. However, the exact role of Ca<sup>2+</sup>-CaM and NFAT in PD pathology is yet to be elucidated.

**Methods:** We designed the CaM-NFAT-oligodeoxynucleotide (ODN), a synthetic short DNA containing complementary sequence for NFAT transcription factor and CaM mRNA. Then, the effect of CaM-NFAT-ODN on 1-methyl-4-phenylpyridinium (MPP<sup>+</sup>)-mediated neurotoxicity was investigated in mimic PD model *in vitro*.

**Results:** First, the expression of αSYN and CaM was strongly increased in substantia nigra (SN) of PD and the expression of tyrosine hydroxylase (TH) was strongly increased in control SN. Additionally, the expression of apoptosis marker proteins was strongly increased in SN of PD. Transfection of CaM-NFAT-ODN repressed CaM and pNFAT, the target genes of this ODN in rat embryo primary mesencephalic neurons. It also reduced ERK phosphorylation, a downstream target of these genes. These results demonstrated that CaM-NFAT-ODN operated successfully in rat embryo primary mesencephalic neurons. Transfection of CaM-NFAT-ODN repressed TH reduction, αSYN accumulation, and apoptosis by MPP<sup>+</sup>-induced neurotoxicity response through Ca<sup>2+</sup> signaling and mitogen-activated protein kinases (MAPK) signaling.

**Conclusion:** Synthetic CaM-NFAT-ODN has substantial therapeutic feasibility for the treatment of neurodegenerative diseases.

**Key words:** Dopaminergic neurons, Oligodeoxynucleotide, Tyrosine hydroxylase, Apoptosis, Calmodulin, Nuclear factor of activated T cells

## ORIGINAL ARTICLE

**Received:** September 28, 2023

**Revised:** October 24, 2023

**Accepted:** October 26, 2023

**Correspondence to:** Kyung Mi Jang  
Department of Pediatrics, Yeungnam  
University School of Medicine, Yeungnam  
University Hospital, 170 Hyeonchung-ro,  
Nam-gu, Daegu 42415, Korea  
**Tel:** +82-53-620-3532  
**E-mail:** Fortune001j@ymc.yu.ac.kr

**ORCID**

<https://orcid.org/0000-0002-2226-9268>



Copyright © 2023, Interdisciplinary Society of Genetic & Genomic Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), provided the original work is properly cited.

## INTRODUCTION

Parkinson's disease (PD) is one of the most prevalent neurodegenerative disorders [1]. The pathology of the disease is the selective and progressive loss of the dopaminergic neurons, leading to the formation of Lewy bodies in substantia nigra (SN) pars compacta of midbrain [2]. Lewy bodies are formed by fibrils of alpha-synuclein (αSYN), and this cascade is an important step in the development of PD [3]. Aggregated αSYN regulates dopamine biosynthesis by triggering dopamine transporter (DAT) recruitment of dopamine and reducing the activity or the phosphorylation status of tyrosine hydroxylase (TH), thus leading to neuronal demise [4]. To date, no drugs have been developed to alleviate the loss of dopaminergic neurons in PD and the pathogenesis of PD remains controversial.

Ca<sup>2+</sup> homeostasis plays a key role in the degeneration of dopaminergic neurons [5]. Additionally, the increase in cytosolic Ca<sup>2+</sup> is dependent on Ca<sup>2+</sup> channel activity and αSYN expression, and these two pathogenic factors were reported to act in PD [4]. The degeneration of dopaminergic neurons is a major contributor to the pathogenesis of midbrain disorders [6]. Ca<sup>2+</sup> is important for many cellular processes, such as proliferation, growth, differentiation, development, and cell death [7]. Calcium signaling is essential for regulating dopamine concentration

levels in neurons [8]. Exposure to Ca<sup>2+</sup> overload has been reported to be an important determinant, and pharmacological treatments that inhibits Ca<sup>2+</sup> entry reduce neuronal damage in chemical models of PD [9]. Calmodulin (CaM), the best studied Ca<sup>2+</sup>-binding protein, is abundantly expressed in the brain [10]. CaM contains globular domains at its C- and N-terminus respectively [11]. Each globular domain of CaM has a pair of Ca<sup>2+</sup> binding sequences, and after binding Ca<sup>2+</sup>, a hydrophobic surface is exposed [12]. The expression of CaM is up regulated in response to the increase in intracellular Ca<sup>2+</sup> concentration and can thus lead to NFAT (nuclear factor of activated T-cells) dephosphorylation and nuclear translocation [9].

Recent reports have highlighted the important role of the NFAT transcription factor family in neurons involved in the regulation of synaptic plasticity, axonal growth and neuronal survival [13]. The NFAT regulatory domain is highly phosphorylated in the inactive state, covering the nuclear localization sequence and causing the NFAT protein to remain in the cytoplasm [14]. NFAT proteins directly regulate the expression of apoptosis-related genes and with the promotion of inflammatory process [15]. Both apoptosis and inflammation are processes widely involved in the development of PD [7,16].

The synthetic oligodeoxynucleotide (ODN) technique is a gene therapy strategy that DNA or RNA based molecular compounds that are utilized to disrupt gene transcription or translation [17]. Antisense ODN are used for selective impairment of protein synthesis based on the specific sequence of the target RNA [18,19]. Decoy ODN is employed to block transcription factor activity using synthetic double stranded ODN containing DNA binding sites, which works at the pre-transcription level [20,21]. Advances in gene therapy strategies have provided new technologies to inhibit dual-target gene expression based on short RNA and DNA strategies [22,23]. To improve a new therapeutic approach, in this study we used a combination of antisense ODN and decoy ODN to synthesize CaM-NFAT-ODN, which inhibits both CaM and NFAT. Although these ODN have proven beneficial in several disease models, it has not yet been demonstrated whether CaM-NFAT-ODN can attenuate the development of molecular mechanisms of neurotoxicity. There-

**Table 1.** Human postmortem tissues used for Figure 1A

KBBN ID	Age	Gender	Final diagnosis	Tissue
07004509	50–59	F	Control	SN
07004529	60–69	M	PD	SN

F, Female; M, Male; SN, Midbrain with substantia nigra; PD, Parkinson disease.

fore, we investigated the effect of CaM-NFAT-ODN on neuronal cytotoxicity in an *in vitro* model of Parkinson's disease.

## METHODS

### Synthesis of oligodeoxynucleotides

Synthetic ODNs were commissioned by MacroGen (Seoul, Korea). To generate the ODN of CaM, it was selected through sequential overlapping simulations of the secondary structure using the S-Fold program. The sequence of CaM antisense ODN is 5'-GAA TTC GGT ATC AAG AAC TAG ATA TAC C-3' and the sequence of NFAT decoy ODN is 5'-GAA TTC TGA CIT TCC TCA TGA AAA CAT GAG GAA AGT AC-3'. The underline is the target site as well as the agreed combining order. These ODNs were structured by annealing and each ODN was mixed with T4 ligase (Takara Bio, Otsu, Japan) to obtain a covalent ligation for ring-type ODN molecules.

### Human Brain Samples and rat embryo primary mesencephalic neurons culture

A total of eight postmortem brain samples were obtained (Tables 1 and 2); with clinically and neuropathologically diagnosed PD and controls. Brain sections and tissues were provided by the Korea Brain Bank Network (KBBN) operated through the National Brain Bank Project funded by the Ministry of Science and ICT. The project was approved by the Daegu Catholic University Institutional Review Board, DCU IRB (IRB number: CR-20-149).

Mesencephalic neuron cultures were prepared from the ven-

**Table 2.** List of human postmortem tissues used for Figure 1B

Sample No.	KBBN ID	Age	Gender	Final diagnosis	Tissue
1	07004507	50–59	F	Control	SN
2	07004508				
3	06012591	60–69	M		
4	06012592				
5	06012596	50–59	M		
6	06012597				
1	07004517	70–79	M	PD	SN
2	07004518				
3	07004527	60–69	M		
4	07004528				
5	05007340	70–79	F		
6	05007344	80–89	F		
7	05007348	70–79	M		

F, Female; M, Male; SN, Midbrain with substantia nigra; PD, Parkinson disease.

tral mesencephalic tissues of embryonic day 13–14 rats, as described previously [7,24]. All experimental protocols were approved by the Institutional Animal Care and Use Committee of the Daegu Catholic University Medical Center (EXP-IRB number: DCIAFCR-191112-07-Y) in accordance with criteria outlined in the Institutional Guidelines for Animal Research. Briefly, dissociated cells were seeded on poly-d-lysine and laminin-coated plates. Cells were cultured in a Dulbecco's modified Eagle's medium/Ham's F-12 medium (Gibco, NY, USA) containing ITS premix (Sigma-Aldrich, St Louis, MO, USA) and 1% penicillin-streptomycin (Gibco).

### Transfection and Morphology examination

Rat embryo primary mesencephalic neurons were transfected with synthetic ODN using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The transfection efficiency of synthetic ODN was observed using ODN labeled with fluorescein isothiocyanate (FITC) using the Label IT nucleic acid labeling kit (Mirus Bio, Madison, WI, USA). After transfection, the rat embryo primary mesencephalic neurons were cultured in 50  $\mu\text{M}$  of MPP<sup>+</sup> for 24 hr.

Nissl bodies in human brain slides were detected by a previously described Nissl staining method [25]. Nissl Staining Solution (Cresyl violet stain solution; Abcam, Cambridge, MA, USA) manufacturer's instructions. Tissues sections were washed in cold phosphate-buffered saline (PBS) for 15 min and block with universal blocking solution (0.3% Triton<sup>TM</sup> X-100, 1% BSA, 0.05% Tween 20, 0.1% cold fish gelatin and 0.05% sodium azide in PBS) for 1 hr at room temperature.

The morphology of rat embryo primary mesencephalic neurons was observed using an inverted phase contrast microscope (Olympus CKX41SF, Tokyo, Japan,  $\times 200$  or  $\times 400$  magnification).

### Immunoblot analysis

Protein samples were prepared from human brain tissue of the SN (Table 2) and rat embryo primary mesencephalic neuron with a protein extraction buffer (N-PER<sup>TM</sup>, Thermo Fisher Scientific, Waltham, MA, USA) according to the instruction manual. The protein samples were separated on precast gradient polyacrylamide gels (Bolt<sup>TM</sup> 4%–12% Bis-Tris Plus Gels; Thermo Fisher Scientific) and transferred to nitrocellulose membranes (GE Healthcare, Madison, WI, USA) by using Bolt<sup>TM</sup> Mini Blot Module and Mini Gel Tank (Thermo Fisher Scientific), according to the manufacturer's recommendations. The membrane blocked with 5% bovine serum albumin was probed with a primary antibody and horseradish peroxidase-conjugat-

ed secondary antibody. Following a repeat of the wash step, the membrane was kept in enhanced chemiluminescence detection reagents (Thermo Fisher Scientific). Signal intensity was measured with an image analyzer (ChemiDoc<sup>TM</sup> XRS+; Bio-Rad Laboratories). The protein expression values were normalized to GAPDH and  $\beta$ -actin expression values. The primary antibodies used were anti-TH (Cat no: 13106, Cell Signaling Technology, Danvers, MA, USA), anti- $\alpha$ SYN (Cat no: 2628, Cell Signaling Technology), anti-PARP (Cat no: 9542, Cell Signaling Technology), anti-CaM (Cat no: sc-137079, Santa Cruz Biotechnology, TX, USA), anti-pNFATc3 (Cat no: sc-8405, Santa Cruz Biotechnology), anti-NFATc3 (Cat no: sc-365883, Santa Cruz Biotechnology), anti-pERK (Cat no: 4377, Cell Signaling Technology), anti-ERK (Cat no: 4695, Cell Signaling Technology) and anti- $\beta$ -actin (Cat no: SAB3500350, Sigma-Aldrich), anti-GAPDH (Cat no: sc-47724, Santa Cruz Biotechnology).


### Immunofluorescent staining

Human brain tissues were deparaffinized in xylene and subjected to citrate antigen retrieval prior to immunohistochemistry [25]. Cells were incubated with primary antibodies for 1 hr at room temperature and they were incubated with the Alexa Flour 488 (excitation/emission = 495/519 nm, green, Invitrogen, CA, USA) and Alexa Flour 594 (excitation/emission = 590/617 nm, red, Invitrogen) for 30 min at room temperature. Cells were counterstained with DAPI (excitation/emission = 330–380 nm/460 nm, ImmunoChemistry, MN, USA). Slides were mounted using ProLong<sup>®</sup> Gold antifade reagent (Molecular Probes<sup>®</sup> by Life Technologies<sup>TM</sup>, CA, USA). Immunolabeling was examined using an Eclipse Ti-U and confocal microscope (Nikon, Tokyo, Japan).


## RESULTS

### Expression of TH, $\alpha$ SYN, PARP-1 and CaM in the SN of human Parkinson's disease brain

Increase in cytosolic Ca<sup>2+</sup> is dependent on Ca<sup>2+</sup> channel activity and  $\alpha$ SYN expression [4]. To determine whether these findings are relevant to PD, we examined human postmortem SN from control versus PD patients. Nissl-stained positive neurons in the SN were significantly reduced in the PD brain compared to human control brain (Table 1 and Fig. 1A).  $\alpha$ SYN and CaM expression was strongly increased in SN of PD, and expression of TH, a dopaminergic neuron marker, was strongly increased in control SN (Table 2 and Fig. 1B). Cleaved-PARP-1, an apoptosis marker protein, was also confirmed to be strongly increa-



**Fig. 1.** Expression of TH,  $\alpha$ SYN, PARP-1 and CaM in the substantia nigra of human Parkinson's disease brain. (A) Nissl staining, Magnification  $\times 200$ , (B) Immunoblot analysis of TH,  $\alpha$ SYN, PARP-1 and CaM in the SN of human control and Parkinson's disease brain. GAPDH was used to confirm equal sample loading.




**Fig. 2.** Effect of CaM-NFAT-ODN in rat embryo primary mesencephalic neurons. (A) Representative immunofluorescence images showing the fluorescence activity in both the cytoplasm and nucleus detected via FITC-labeled ODN deposition (green). Cells were counter-stained with DAPI (blue). Magnification  $\times 1,600$ . (B) Immunoblot analysis of CaM, pNFAT, NFAT, pERK and ERK in rat embryo primary mesencephalic neurons.  $\beta$ Actin was used to confirm equal sample loading.

sed in SN of PD (Fig. 1B). These results demonstrated that CaM is related to apoptosis of dopaminergic neurons and PD.

#### CaM-NFAT-ODN protects rat embryo primary mesencephalic neurons against MPP<sup>+</sup>-Induced Neurotoxicity

To confirm the efficacy and distribution of the CaM-NFAT-

ODN, we transfected rat embryo primary mesencephalic neurons with FITC-labeled ODN. As shown in Fig. 2A, FITC-CaM-NFAT-ODN was successfully transfected and delivered into the nucleus and cytoplasm of rat embryo primary mesencephalic neurons and observed for 24 hr. Moreover, immunoblot analysis showed that the CaM-NFAT-ODN decreased the protein levels of CaM, pNFAT, as well as the key downstream targets of



**Fig. 3.** The CaM-NFAT-ODN influenced the expression of TH,  $\alpha$ SYN and PARP-1 in MPP<sup>+</sup> treated rat embryo primary mesencephalic neurons. (A) The morphological changes, TH and  $\alpha$ SYN expression in dopaminergic neurons after exposure to MPP<sup>+</sup> in the presence or absence of APM. Immunofluorescence staining for TH (green) and  $\alpha$ SYN (red) localization. Cells were counterstained with DAPI (blue). Magnification  $\times 400$ . (B) CaM-NFAT-ODN strongly reduced expression of TH reduction, accumulation of  $\alpha$ SYN and cleaved PARP-1 expression in rat embryo primary mesencephalic neurons,  $\beta$ Actin was used to confirm equal sample loading.

ERK phosphorylation (Fig. 2B).

To determine whether Synthetic CaM-NFAT-ODN could regulate the expression of MPP<sup>+</sup>-induced TH and  $\alpha$ SYN in rat embryo primary mesencephalic neurons. The rat embryo primary mesencephalic neurons showed obvious neurites, and the neurons transfected with only CaM-NFAT-ODN were observed similar to normal neurons (Fig. 3A). When rat embryo primary mesencephalic neurons were exposed to the neurotoxin MPP<sup>+</sup>, neurites were reduced, and neuronal debris increased. Interestingly, it was rescued by CaM-NFAT ODN transfection.

Rat embryo primary mesencephalic neurons were confirmed to be healthy TH-positive neurons with extensive neurites through immunofluorescence staining using a TH antibody. However, exposure to neurotoxin for 24 hr decreased the number of TH-positive neurons and induced accumulation of  $\alpha$ SYN. When transfected with CaM-NFAT ODN, the number and morphology of TH-positive neurons were similar to normal neurons. Exposure of MPP<sup>+</sup> in CaM-NFAT-ODN transfected neurons appears to protect the cells from loss of TH-positive neurons and accumulation of  $\alpha$ SYN. Consistent with immunofluorescence staining, CaM-NFAT-ODN attenuated TH reduction and accumulation of  $\alpha$ SYN in rat embryo primary mesencephalic neurons (Fig. 3B). CaM-NFAT-ODN inhibited MPP<sup>+</sup>-induced cleaved PARP-1, apoptosis marker proteins, in rat embryo primary mesencephalic neurons.

## DISCUSSION

Appearance of PD symptoms is attributed to the loss of dopaminergic neurons in the striatum and substantia nigra [10].

Regarding PD, many CaM-regulated Ca<sup>2+</sup> channels seem to be implicated in this pathology [26]. Neuronal cell death may be due to increased Ca<sup>2+</sup>, which is commonly observed in neurons in various areas of PD brain [5,8]. Additionally, the increase in cytosolic Ca<sup>2+</sup> is not due to MPP<sup>+</sup>-induced oxidative stress but is dependent on Ca<sup>2+</sup> channel activity and  $\alpha$ SYN expression, and these two pathogenic factors have been reported to act in PD [27]. Our result showed that  $\alpha$ SYN and CaM expression was strongly increased in Parkinson's disease SN, and expression of TH, a dopaminergic neuron marker, was strongly increased in control SN.

A Study has reported on potential neuroprotective agents through attenuation of MPP<sup>+</sup> and Ca<sup>2+</sup>-overload-induced excitotoxicity in dopaminergic neurons [28]. Many reports suggest that Ca<sup>2+</sup> is involved in the pathogenesis of PD and that regulation of Ca<sup>2+</sup> may be a potential therapeutic target for neuroprotection in PD [29]. For an effective treatment that targets the causes of PD, it is necessary to systemically determine the therapeutic component based on the inhibition of initial molecule mechanism.

We have reported the anti-inflammatory and anti-fibrotic properties of synthetic targeted ODN in chronic diseases [17, 20,21,23]. Recently, we also reported the effects of a selective antagonist of the small-conductivity Ca<sup>2+</sup>-activated K<sup>+</sup> channels antagonist on neuroinflammation and neurotoxicity [7,30]. However, the molecular mechanisms of MPP<sup>+</sup>-induced neurotoxicity of ODN in PD models have not yet been elucidated.

In this study, we investigated the effects of CaM-NFAT-ODN, which simultaneously regulates the NFAT and CaM, on the neurotoxicity of rat embryo primary mesencephalic neurons.

In PC12 cells, MPP<sup>+</sup>-induced loss of cell viability was medi-

ated by increased intracellular  $\text{Ca}^{2+}$  levels and CaM action [31]. Overexpression of  $\alpha\text{SYN}$  activates the CaM-NFAT pathway in cell lines and dopaminergic neurons, whereas inhibition of this pathway protects dopaminergic neurons against  $\alpha\text{SYN}$ -mediated toxicity [32]. The  $\text{Ca}^{2+}$ -CaM-ERK mitogen-activated protein kinase signaling pathway induces neuronal cell death in rat brain [33]. Our result showed that CaM-NFAT-ODN inhibited ERK phosphorylation, reduction of TH,  $\alpha\text{SYN}$  expression and apoptosis in MPP<sup>+</sup>-induced rat embryo primary mesencephalic neurons. Thus, gene therapy targeted to suppress mRNA level of CaM and transcription activity of NFAT simultaneously might provide a new therapeutic strategy to prevent various neurological disorders.

## ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2021R111A1A01060030).

## CONFLICTS OF INTEREST

Not applicable.

## AUTHOR CONTRIBUTIONS

Investigation: Park J and Jang KM. Writing original draft preparation: Park J and Jang KM. Writing review: Jang KM. Funding acquisition: Park J. All authors have read and agreed to the published version of the manuscript.

## REFERENCES

- Zhao D, Feng PJ, Liu JH, Dong M, Shen XQ, Chen YX, et al. Electromagnetized-nanoparticle-modulated neural plasticity and recovery of degenerative dopaminergic neurons in the mid-brain. *Advanced Materials* 2020;32:2003800. doi: 10.1002/adma.202003800.
- Wu J, Xu X, Zheng L, Mo J, Jin X, Bao Y. Nilotinib inhibits microglia-mediated neuroinflammation to protect against dopaminergic neuronal death in Parkinson's disease models. *International Immunopharmacology* 2021;99:108025. doi: 10.1016/j.intimp.2021.108025.
- Wang X, Cao G, Ding D, Li F, Zhao X, Wang J, et al. Ferruginol prevents degeneration of dopaminergic neurons by enhancing clearance of  $\alpha$ -synuclein in neuronal cells. *Fitoterapia* 2022;156:105066. doi: 10.1016/j.fitote.2021.105066.
- Lieberman OJ, Choi SJ, Kanter E, Saverchenko A, Frier MD, Fiore GM, et al.  $\alpha$ -Synuclein-dependent calcium entry underlies differential sensitivity of cultured sn and vta dopaminergic neurons to a parkinsonian neurotoxin. *eNeuro* 2017;4. doi: 10.1523/eneuro.0167-17.2017.
- Sun Y, Zhang H, Selvaraj S, Sukumaran P, Lei S, Bimbaumer L, et al. Inhibition of L-type  $\text{Ca}^{2+}$  channels by TRPC1-STIM1 complex is essential for the protection of dopaminergic neurons. *Journal of Neuroscience* 2017;37:3364-77. doi: 10.1523/JNEUROSCI.3010-16.2017.
- Chen L, Huang Y, Yu X, Lu J, Jia W, Song J, et al. Corynoxine protects dopaminergic neurons through inducing autophagy and diminishing neuroinflammation in rotenone-induced animal models of parkinson's disease. *Front Pharmacol* 2021;12:642900. doi: 10.3389/fphar.2021.642900.
- Park J, Jang KM, Park KK. Effects of apamin on MPP(+)-induced calcium overload and neurotoxicity by targeting CaMKII/ERK/p65/STAT3 signaling pathways in dopaminergic neuronal cells. *Int J Mol Sci* 2022;23. doi: 10.3390/ijms232315255.
- Pawar A, Pardasani KR. Computational model of calcium dynamics-dependent dopamine regulation and dysregulation in a dopaminergic neuron cell. *The European Physical Journal Plus* 2023;138:30. doi: 10.1140/epjp/s13360-023-03691-1.
- Chen L, Song M, Yao C. Calcineurin in development and disease. *Genes & Diseases* 2022;9:915-27. doi: 10.1016/j.gendis.2021.03.002.
- Bohush A, Lesniak W, Weis S, Filipek A. Calmodulin and its binding proteins in Parkinson's disease. *Int J Mol Sci* 2021;22. doi: 10.3390/ijms22063016.
- Ren R, Guo J, Chen Y, Zhang Y, Chen L, Xiong W. The role of  $\text{Ca}^{2+}$ /Calcineurin/NFAT signalling pathway in osteoblastogenesis. *Cell Proliferation* 2021;54:e13122.
- An S. The emerging role of extracellular  $\text{Ca}^{2+}$  in osteo/odontogenic differentiation and the involvement of intracellular  $\text{Ca}^{2+}$  signaling: from osteoblastic cells to dental pulp cells and odontoblasts. *J Cell Physiol* 2019;234:2169-93. doi: 10.1002/jcp.27068.
- Sant'Anna R, Robbs BK, Freitas JAd, Santos PPd, König A, Outeiro TE, et al. Alpha-synuclein oligomers activate NFAT proteins modulating synaptic homeostasis and apoptosis. *bioRxiv* 2023:2023.02.21.529374. doi: 10.1101/2023.02.21.529374.
- Hogan PG, Chen L, Nardone J, Rao A. Transcriptional regulation by calcium, calcineurin, and NFAT. *Genes Dev* 2003;17:2205-32. doi: 10.1101/gad.1102703.
- Mognol GP, Carneiro FRG, Robbs BK, Faget DV, Viola JPdB. Cell cycle and apoptosis regulation by NFAT transcription factors: new roles for an old player. *Cell death & disease* 2016;7:e2199. doi: 10.1038/cddis.2016.97.
- Park J, Jang KM, Park KK. Apamin suppresses LPS-induced neuroinflammatory responses by regulating sk channels and TLR4-mediated signaling pathways. *Int J Mol Sci* 2020;21. doi: 10.3390/ijms21124319.
- Gu H, An HJ, Gwon MG, Bae S, Zouboulis CC, Park KK. The effects of synthetic SREBP-1 and PPAR-gamma decoy oligodeoxynucleotide on acne-like disease in vivo and in vitro via lipogenic

- regulation. *Biomolecules* 2022;12. doi: 10.3390/biom12121858.
18. Bennett CF, Swayze EE. RNA targeting therapeutics: molecular mechanisms of antisense oligonucleotides as a therapeutic platform. *Annual Review of Pharmacology and Toxicology* 2010; 50:259-93. doi: 10.1146/annurev.pharmtox.010909.105654.
  19. Mann MJ, Dzau VJ. Therapeutic applications of transcription factor decoy oligonucleotides. *The Journal of Clinical Investigation* 2000;106:1071-5. doi: 10.1172/JCI11459.
  20. Lee SJ, Kim YA, Park KK. Anti-Fibrotic effect of synthetic noncoding decoy ODNs for TFE3 in an animal model of chronic kidney disease. *International Journal of Molecular Sciences* 2022; 23:8138. doi: 10.3390/ijms23158138.
  21. Gwon MG, Leem J, An HJ, Gu H, Bae S, Kim JH, et al. The decoy oligodeoxynucleotide against HIF-1 $\alpha$  and STAT5 ameliorates atopic dermatitis-like mouse model. *Molecular Therapy-Nucleic Acids* 2023;34:102036. doi: 10.1016/j.omtn.2023.102036.
  22. Kim KH, Park KK. Small RNA-and DNA-based gene therapy for the treatment of liver cirrhosis, where we are? *World Journal of Gastroenterology: WJG* 2014;20:14696-705. doi: 10.3748/wjg.v20.i40.14696.
  23. Kim JY, An HJ, Kim WH, Gwon MG, Gu H, Park YY, et al. Anti-fibrotic effects of synthetic oligodeoxynucleotide for TGF- $\beta$ 1 and smad in an animal model of liver cirrhosis. *Mol Ther Nucleic Acids* 2017;8:250-63. doi: 10.1016/j.omtn.2017.06.022.
  24. Park JH, Seo YH, Jang JH, Jeong CH, Lee S, Park B. Asiatic acid attenuates methamphetamine-induced neuroinflammation and neurotoxicity through blocking of NF- $\kappa$ B/STAT3/ERK and mitochondria-mediated apoptosis pathway. *J Neuroinflammation* 2017;14:240. doi: 10.1186/s12974-017-1009-0.
  25. Nam JH, Park ES, Won SY, Lee YA, Kim KI, Jeong JY, et al. TRPV1 on astrocytes rescues nigral dopamine neurons in Parkinson's disease via CNTF. *Brain* 2015;138(Pt 12):3610-22. doi: 10.1093/brain/awv297.
  26. Bohush A, Leśniak W, Weis S, Filipek A. Calmodulin and its binding proteins in Parkinson's disease. *International Journal of Molecular Sciences* 2021;22:3016. doi: 10.3390/ijms22063016.
  27. Chen TS, Koutsilieris E, Rausch WD. MPP+ selectively affects calcium homeostasis in mesencephalic cell cultures from embryonal C57/B16 mice. *Journal of Neural Transmission/General Section JNT* 1995;100:153-63. doi: 10.1007/BF01271538.
  28. Egunlusi AO, Malan SF, Omoruyi SI, Ekpo OE, Palchykov VA, Joubert J. Open and rearranged norbornane derived polycyclic cage molecules as potential neuroprotective agents through attenuation of MPP(+)- and calcium overload-induced excitotoxicity in neuroblastoma SH-SY5Y cells. *Eur J Med Chem* 2020;204: 112617. doi: 10.1016/j.ejmech.2020.112617.
  29. Swart T, Hurley MJ. Calcium channel antagonists as disease-modifying therapy for Parkinson's disease: therapeutic rationale and current status. *Cns Drugs* 2016;30:1127-35. doi: 10.1007/s40263-016-0393-9.
  30. Park J, Jang KM, Park KK. Apamin suppresses LPS-induced neuroinflammatory responses by regulating SK channels and TLR4-mediated signaling pathways. *International Journal of Molecular Sciences* 2020;21:4319. doi: 10.3390/ijms21124319.
  31. Lee DH, Han YS, Han ES, Bang H, Lee CS. Differential involvement of intracellular Ca<sup>2+</sup> in 1-methyl-4-phenylpyridinium-or 6-hydroxydopamine-induced cell viability loss in PC12 cells. *Neurochemical Research* 2006;31:851-60. doi: 10.1007/s11064-006-9088-9.
  32. Sompol P, Furman JL, Pleiss MM, Kraner SD, Artiushin IA, Batten SR, et al. Calcineurin/NFAT signaling in activated astrocytes drives network hyperexcitability in A $\beta$ -bearing mice. *Journal of Neuroscience* 2017;37:6132-48. doi: 10.1523/JNEUROSCI.0877-17.2017.
  33. Sticozzi C, Belmonte G, Meini A, Carbotti P, Grasso G, Palmi M. IL-1 $\beta$  induces GFAP expression in vitro and in vivo and protects neurons from traumatic injury-associated apoptosis in rat brain striatum via NF $\kappa$ B/Ca<sup>2+</sup>-calmodulin/ERK mitogen-activated protein kinase signaling pathway. *Neuroscience* 2013;252:367-83. doi: 10.1016/j.neuroscience.2013.07.061.