

Nonlinear Functional Analysis and Applications

Vol. 29, No. 1 (2024), pp. 165-177

ISSN: 1229-1595(print), 2466-0973(online)

<https://doi.org/10.22771/nfaa.2024.29.01.11>

<http://nfaa.kyungnam.ac.kr/journal-nfaa>

Copyright © 2024 Kyungnam University Press



ANALYSIS OF EXISTENCE AND STABILITY RESULTS FOR FRACTIONAL IMPULSIVE \mathfrak{S} -HILFER FREDHOLM-VOLTERRA MODELS

Fawzi Muttar Ismaael

Ministry of Education, Open Education College, Department of Mathematics,

Salah Al-Din, Iraq

e-mail: fawzimuttar62@gmail.com

Abstract. In this paper, we investigate the suitable conditions for the existence results for a class of \mathfrak{S} -Hilfer fractional nonlinear Fredholm-Volterra models with new conditions. The findings are based on Banach contraction principle and Schauder's fixed point theorem. Also, the generalized Hyers-Ulam stability and generalized Hyers-Ulam-Rassias stability for solutions of the given problem are provided.

1. INTRODUCTION

Science and progress have advanced significantly in recent decades thanks to the fractional calculus theory, which comprises fractional equations. This is mostly because it has produced adequate models, especially for real-world issues. Furthermore, generalized differential equations are thought to include fractional differential equations (FDEs) [1, 6, 8, 14, 16, 17, 18, 20].

The integro-differential equation (IDE) is an operator that contains both integer-order derivatives and integer-order integrals as special cases, which is the reason why in the present, fractional calculus gains enormous popularity and many applications arise from the term in the field of mathematical

⁰Received July 12, 2023. Revised September 4, 2023. Accepted September 10, 2023.

⁰2020 Mathematics Subject Classification: 34A12, 26A33, 49K40.

⁰Keywords: \mathfrak{S} -Hilfer fractional derivative, Volterra-Fredholm equation, fixed point technique, generalized Ulam-Hyers-Rassias stability.

⁰Corresponding author: Fawzi Muttar Ismaael(fawzimuttar62@gmail.com).

physics where The integrals are essentially power law and have a convolution or logarithm type single kernels [3, 4, 5, 7, 8, 13, 23, 24].

The initial-boundary value difficulties posed by nonlinear parabolic differential and IDEs may be used to mimic many real phenomena; for examples, see [8, 15, 16, 17, 25, 26, 27, 28]. As a result, the Caputo fractional operator’s application and generalization in IDEs are justified.

Through the use of Caputo, Hilfer operators and their generalization, several scholars have recently examined the stability, uniqueness, and existence of various boundary value problems (BVPs). Non-instantaneous impulsive BVPs and the \mathfrak{S} -Caputo (or, more precisely, \mathfrak{S} -Liouville-Caputo) fractional derivative were explored by Asawasamrit et al. in [2]. The \mathfrak{S} -Hilfer fractional derivative including boundary conditions was explored by Ivaz et al. in [12].

The fractional Hilfer derivative under nonlocal boundary constraints was studied by Nuchpong et al. [19]

$$\begin{cases} {}^H\mathcal{D}^{p,q}X(\phi) = \Xi(\phi, X(\phi), \mathcal{I}^\delta X(\phi)), & \phi \in [\alpha_1, \alpha_2], \\ X(\alpha_1) = 0, \quad \nabla + \int_{\alpha_1}^{\alpha_2} X(l)dl = \sum_{k=1}^{\varrho-2} \varsigma_k X(\vartheta_k), \end{cases}$$

where they have used the the \mathcal{I}^δ -R-L and ${}^H\mathcal{D}^{p,q}$ -Hilfer fractional derivative, and Ξ is continuous function.

In [22], Salim et al. investigated the BVP for the implicit fractional-order generalised fractional impulses Hilfer-type of the following form:

$$\begin{cases} ({}^\varrho\mathcal{D}_{\tau^+}^{p,q}X)(\phi) = \Xi(\phi, X(\phi), ({}^\varrho\mathcal{D}_{\tau^+}^{p,q}X)(\phi)), & \phi \in \phi_c, \\ X(\phi) = \mathcal{B}_c(\phi, X(\phi)), & \phi \in (\phi_c, r_c], \quad c = 1, \dots, \varrho, \\ \varphi_1 ({}^\varrho\mathcal{I}_{\alpha_1^+}^{1-\epsilon})(\alpha_1) + \varphi_2 ({}^\varrho\mathcal{I}_{\tau^+}^{1-\epsilon})(\alpha_2) = \varphi_3, \end{cases}$$

where ${}^\varrho\mathcal{I}_{\alpha_1^+}^{1-\epsilon}$ and ${}^\varrho\mathcal{D}_{\tau^+}^{p,q}$ are the fractional generalized Hilfer-type integral and derivative and the function Ξ is continuous.

As started by Sousa et al. in [25], we examine the existence, and stability of fractional IDEs using the \mathfrak{S} -Hilfer fractional derivative in this work. A remark on the transformation may be found in [25]. Multiple fractional derivatives are combined into the fractional \mathfrak{S} -Hilfer derivative.

$$\begin{cases} D_{\alpha_+}^{\varphi,\beta;\mathfrak{S}}\xi(v) = \varpi(v, \xi(v), G\xi(v), K\xi(v)), & v \in J := (\alpha, \chi], \\ I_{\alpha_+}^{1-\gamma;\mathfrak{S}}\xi(\alpha) = \xi_\alpha, & \gamma = \varphi + \beta - \varphi\beta, \end{cases} \quad (1.1)$$

where $I_{\alpha_+}^{1-\gamma;\mathfrak{S}}$ is \mathfrak{S} -fractional integral and $D_{\alpha_+}^{\varphi,\beta;\mathfrak{S}}$ is fractional \mathfrak{S} -Hilfer derivative of order φ and type β , where $\varpi : J \times \mathbb{R}^3 \rightarrow \mathbb{R}$, $g, k : \Delta \times R \rightarrow \mathbb{R}$ are continuous,

with $\Delta = \{(v, \sigma) : \alpha \leq \sigma \leq v \leq \chi\}$, and

$$G\xi(v) = \int_{\alpha}^v g(v, \sigma, \xi(\sigma))d\sigma, \quad K\xi(v) = \int_{\alpha}^{\chi} k(v, \sigma, \xi(\sigma))d\sigma.$$

2. AN AUXILIARY RESULT

We review several definitions and outcomes from fractional calculus in this section. The observations that follow are drawn from [8, 11, 12, 16, 20]. Let $\xi : J \rightarrow \mathbb{R}$ in $\zeta[\alpha, \chi]$ with

$$\|\xi\| = \sup\{|\xi(v)| : v \in J\}.$$

The $\zeta_{\gamma, \mathfrak{S}}[\alpha, \chi]$ is weighted space, let ϖ on $(\alpha, \chi]$ is given by

$$\zeta_{\gamma, \mathfrak{S}}[\alpha, \chi] = \{\varpi : (\alpha, \chi] \rightarrow R : (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma} \varpi(v) \in \zeta[\alpha, \chi]\}, \quad 0 \leq \gamma < 1,$$

and the norm

$$\|\varpi\|_{\zeta_{\gamma, \mathfrak{S}}} = \|(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma} \varpi(v)\|_{\zeta[\alpha, \chi]} = \max_{v \in J} |(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma} \varpi(v)|.$$

The weighted space $\zeta_{\gamma, \mathfrak{S}}^n[\alpha, \chi]$ of ϖ on $(\alpha, \chi]$ is given by

$$\zeta_{\gamma, \mathfrak{S}}^n[\alpha, \chi] = \{\varpi : J \rightarrow R : \varpi(v) \in \zeta^{n-1}[\alpha, \chi]; \varpi(v) \in \zeta_{\gamma, \mathfrak{S}}[\alpha, \chi]\}, \quad 0 \leq \gamma < 1,$$

and the norm

$$\|\varpi\|_{\zeta_{\gamma, \mathfrak{S}}^n[\alpha, \chi]} = \sum_{k=0}^{n-1} \|\varpi^k\|_{\zeta[\alpha, \chi]} + \|\varpi^n\|_{\zeta_{\gamma, \mathfrak{S}}[\alpha, \chi]}.$$

For $n = 0$, we get $\zeta_{\gamma}^0[\alpha, \chi] = \zeta_{\gamma}[\alpha, \chi]$.

Definition 2.1. ([21]) The fractional integral left-sided of ϖ in relation to \mathfrak{S} on $[\alpha, \chi]$ is given by

$$\left(I_{\alpha^+}^{\varphi; \mathfrak{S}}\right) \varpi(v) = \frac{1}{\Gamma(\varphi)} \int_{\alpha}^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varphi-1} \varpi(\sigma) d\sigma, \quad v > \alpha. \quad (2.1)$$

Definition 2.2. ([21]) Let $\mathfrak{S}'(\xi) \neq 0 (-\infty < v < \chi < \infty)$. The fractional Liouville-Riemann derivative of ϖ in relation to \mathfrak{S} of order $\varphi > 0$ correspondent to the Liouville-Riemann, is given by

$$\left(D_{\alpha^+}^{\varphi; \mathfrak{S}}\right) \varpi(v) = \frac{1}{\Gamma(n-\varphi)} \left(\frac{1}{\mathfrak{S}'(v)} \frac{d}{dv}\right)^n \int_{\alpha}^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{n-\varphi-1} \varpi(\sigma) d\sigma, \quad (2.2)$$

where $n = [\varphi] + 1$, $n \in \mathbb{N}$.

Definition 2.3. ([21]) Let $\varphi > 0$, $I = [\alpha, \chi]$, $(-\infty < v < \chi < \infty)$, $\varpi, \mathfrak{S} \in \zeta^n([\alpha, \chi], \mathcal{R})$ be two functions, \mathfrak{S} be increasing and $\mathfrak{S}'(\xi) \neq 0$ for all $\xi \in I$. The left \mathfrak{S} -Caputo derivative of ϖ of order φ as

$$\left(D_{\alpha^+}^{\varphi; \mathfrak{S}} \varpi\right)(v) = I_{\alpha^+}^{n-\varphi; \mathfrak{S}} \left(\frac{1}{\mathfrak{S}'(v)} \frac{d}{dv}\right)^n \varpi(v), \quad (2.3)$$

where $n = [\varphi] + 1$, for $\varphi \notin N$ and $\varphi = n$ for $\varphi \in N$.

Definition 2.4. ([21]) According to function ϖ of order φ has the following fractional \mathfrak{S} -Hilfer derivative:

$$D_{\alpha^+}^{\varphi, \beta; \mathfrak{S}} \varpi(v) = I_{\alpha^+}^{\beta(1-\varphi); \mathfrak{S}} \left(\frac{1}{\mathfrak{S}'(v)} \frac{d}{dt}\right) I_{\alpha^+}^{(1-\beta)(1-\varphi); \mathfrak{S}} \varpi(v). \quad (2.4)$$

As previously stated, the \mathfrak{S} -Hilfer fractional derivative may be expressed as follows:

$$D_{\alpha^+}^{\varphi, \beta; \mathfrak{S}} \varpi(v) = I_{\alpha^+}^{\gamma-\varphi; \mathfrak{S}} D_{\alpha^+}^{\gamma; \mathfrak{S}} \varpi(v).$$

Lemma 2.5. ([8]) Assume $\varphi, \beta > 0$. Then we get

$$\left(I_{\alpha^+}^{\varphi; \mathfrak{S}} I_{\alpha^+}^{\beta; \mathfrak{S}} \varpi\right)(v) = \left(I_{\alpha^+}^{\varphi+\beta; \mathfrak{S}}\right)(v)$$

and

$$\left(D_{\alpha^+}^{\varphi; \mathfrak{S}} I_{\alpha^+}^{\varphi; \mathfrak{S}} \varpi\right)(v) = \varpi(v).$$

Lemma 2.6. ([8]) Let $\varphi, \beta > 0$.

(1) If $\varpi(\xi) = (\mathfrak{S}(v)\mathfrak{S}(\alpha))^{\beta-1}$, then

$$I_{\alpha^+}^{\varphi; \mathfrak{S}} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\beta-1}(v) = \frac{\Gamma(\beta)}{(\varphi + \beta)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\varphi+\beta-1}.$$

(2) If $g(\xi) = (\mathfrak{S}(v)\mathfrak{S}(\alpha))^{\varphi-1}$, then

$$D_{\alpha^+}^{\varphi; \mathfrak{S}} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\varphi-1}(v) = 0.$$

Lemma 2.7. ([22]) For $0 < \varphi < 1$, if $\varpi \in \zeta^n[\alpha, \chi]$, then

$$\left(I_{\alpha^+}^{\varphi; \mathfrak{S}} D_{\alpha^+}^{\varphi; \mathfrak{S}}\right)(v) = \varpi(v) - \frac{\left(I_{\alpha^+}^{1-\varphi; \mathfrak{S}} \varpi\right)(\alpha)}{\Gamma(\varphi)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\varphi-1}$$

for all $\xi \in (\alpha, \chi]$.

Lemma 2.8. ([11]) *Assume $n - 1 \leq \gamma < n$ and $\varpi \in \zeta_\gamma[\alpha, \chi]$. Then*

$$\left(I_{\alpha^+}^{\varphi; \mathfrak{S}} \varpi \right) (\alpha) = \lim_{v \rightarrow \alpha^+} \left(I_{\alpha^+}^{\varphi; \mathfrak{S}} \right) \varpi(v) = 0.$$

The weighted space is shown below as follows.

$$\zeta_{1-\gamma; \mathfrak{S}}^{\varphi, \beta}[\alpha, \chi] = \left\{ \varpi \in \zeta_{1-\gamma; \mathfrak{S}}[\alpha, \chi], D_{\alpha^+}^{\varphi, \beta; \mathfrak{S}} \varpi \in \zeta_{\gamma; \mathfrak{S}}[\alpha, \chi] \right\}$$

and

$$\zeta_{1-\gamma; \mathfrak{S}}^\gamma[\alpha, \chi] = \left\{ \varpi \in \zeta_{1-\gamma; \mathfrak{S}}[\alpha, \chi], D_{\alpha^+}^{\gamma; \mathfrak{S}} \varpi \in \zeta_{1-\gamma; \mathfrak{S}}[\alpha, \chi] \right\}.$$

There is no denying

$$\zeta_{1-\gamma; \mathfrak{S}}^\gamma[\alpha, \chi] \subset \zeta_{1-\gamma; \mathfrak{S}}^{\varphi, \beta}[\alpha, \chi].$$

Lemma 2.9. ([11]) *Let $0 \leq \beta \leq 1$, $0 < \varphi < 1$ and $\gamma = \varphi + \beta - \varphi\beta$. If $\zeta_{1-\gamma; \mathfrak{S}}^\gamma[\alpha, \chi]$, then*

$$I_{\alpha^+}^{\gamma; \mathfrak{S}} D_{\alpha^+}^{\gamma; \mathfrak{S}} \varpi = I_{\alpha^+}^{\varphi; \mathfrak{S}} D_{\alpha^+}^{\varphi, \beta; \mathfrak{S}} \varpi \tag{2.5}$$

and

$$D_{\alpha^+}^{\gamma; \mathfrak{S}} I_{\alpha^+}^{\varphi; \mathfrak{S}} \varpi = D_{\alpha^+}^{\beta(1-\varphi); \mathfrak{S}} \varpi. \tag{2.6}$$

Lemma 2.10. ([28]) *Let $\varpi \in L^1(\alpha, \chi)$. If $D_{\alpha^+}^{\beta(1-\varphi); \mathfrak{S}} \varpi$ exists on $L^1(\alpha, \chi)$. Then*

$$D_{\alpha^+}^{\varphi, \beta; \mathfrak{S}} I_{\alpha^+}^{\varphi; \mathfrak{S}} \varpi = I_{\alpha^+}^{\beta(1-\varphi); \mathfrak{S}} D_{\alpha^+}^{\beta(1-\varphi); \mathfrak{S}} \varpi.$$

Lemma 2.11. ([28]) *Suppose $\varpi \in \zeta^1[\alpha, \chi]$, $\varphi > 0$ and $0 \leq \beta \leq 1$. Then we get*

$$D_{\alpha^+}^{\varphi, \beta; \mathfrak{S}} I_{\alpha^+}^{\varphi; \mathfrak{S}} \varpi = \varpi.$$

Lemma 2.12. *Let $\varphi > 0$, $a(v)$ be an integrable on $a \leq v < \chi$ (some $\chi \leq \infty$), and let $g(v)$ be nondecreasing, continuous and nonnegative on $\alpha \leq v < \chi$ such that $g(v) \leq \Omega^*$ for some Ω . Let $\xi(v)$ be locally integrable on $\alpha \leq v < \chi$ with*

$$|\xi(v)| \leq a(v) + g(v) \int_{\alpha}^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varphi-1} \xi(\sigma) d\sigma, \quad v \in J$$

with some $\varphi > 0$. Then

$$|\xi(v)| \leq a(v) + \int_{\alpha}^v \left[\sum_{n=1}^{\infty} \frac{(g(\sigma) \Gamma(\varphi))^n}{\Gamma(n\varphi)} \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{n\varphi-1} \right] \xi(\sigma) d\sigma, \quad \alpha \leq v < \chi.$$

Proof. Theorem 1 in [28] is identical to the proof in this case. □

Lemma 2.13. ([28]) *Assume that $\gamma = \wp + \beta - \wp\beta$, where $0 \leq \beta \leq 1$ and $0 < \wp < 1$. If $\varpi : J \times \mathbb{R}^3 \rightarrow \mathbb{R}$ and $\varpi(\cdot, \xi(\cdot)) \in \zeta_{1-\gamma}[\alpha, \chi]$ for all $\xi \in \zeta_{1-\gamma}[\alpha, \chi]$. A function $\xi \in \zeta_{1-\gamma}[\alpha, \chi]$ is a solution of (1.1) if and only if ξ satisfies*

$$\begin{aligned} \xi(v) &= \frac{\xi_\alpha}{\Gamma(\gamma)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \\ &+ \frac{1}{\Gamma(\wp)} \int_\alpha^v (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\wp-1} \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma. \end{aligned} \quad (2.7)$$

3. EXISTENCE RESULTS

To back up our primary findings, we propose the following hypothesis.

(H1) Assume that $\varpi : J \times \mathbb{R} \rightarrow \mathbb{R}$ is a function and $\varpi(\cdot, \xi(\cdot)) \in \zeta_{1-\gamma, \mathfrak{S}}[J, \mathbb{R}]$ for any $\xi \in \zeta_{1-\gamma}[J, \mathbb{R}]$ for all $\xi, y \in \mathbb{R}$ and there exist $L > 0$ with

$$|\varpi(v, \xi) - \varpi(v, y)| \leq L|\xi - y|.$$

(H2) Assume that $g, k : \Delta \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous and there exist $G^*, K^* > 0$, and

$$\int_\alpha^v |g(v, \sigma, \xi) - g(v, \sigma, y)| ds \leq G^* |\xi - y|$$

and

$$\int_\alpha^\chi |k(v, \sigma, \xi) - k(v, \sigma, y)| ds \leq K^* |\xi - y|.$$

(H3) Assume that $\varpi : J \times \mathbb{R} \rightarrow \mathbb{R}$ is a function and there exists M, N with

$$|\varpi(v, \xi)| \leq M|\xi| + N, \quad \forall v \in J, \xi \in \mathbb{R}.$$

Theorem 3.1. *Assume that both (H1) and (H2) are met. Then, there is at least one solution to the Eq. (1.1).*

Proof. Incorporate the map $N : \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi] \rightarrow \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$. The operator-form representation of the Eq. (2.7)

$$\xi(v) = N\xi(v),$$

where

$$\begin{aligned} (\Lambda\xi)(v) &= \frac{\xi_\alpha}{\Gamma(\gamma)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \\ &+ \frac{1}{\Gamma(\wp)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\wp-1} \varpi(s, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma. \end{aligned} \quad (3.1)$$

Consider the ball

$$\mathbb{C}_r = \{\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi] : \|\xi\| \leq r\}.$$

The operator Λ is obviously well defined. It is obvious that the problem's solutions are the fixed points of the operator Λ . Then, for all $\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ and $v \in J$ we get,

$$\begin{aligned}
& |(\Lambda\xi)(v)(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma}| \\
&= \left| \frac{\xi_\alpha}{\Gamma(\gamma)} + \frac{1}{\Gamma(\wp)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma} \right. \\
&\quad \times \left. \int_\alpha^v \mathfrak{S}'(\sigma)(\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\wp-1} \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma \right| \\
&\leq \frac{\xi_\alpha}{\Gamma(\gamma)} + \frac{1}{\Gamma(\wp)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma} \\
&\quad \times \int_\alpha^v \mathfrak{S}'(\sigma)(\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\wp-1} |\varpi(\sigma, \xi(\sigma))| d\sigma \\
&\leq \frac{\xi_\alpha}{\Gamma(\gamma)} + \frac{1}{\Gamma(\wp)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma} \\
&\quad \times \int_\alpha^v \mathfrak{S}'(\sigma)(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\wp-1} (M|\xi(\sigma)| + \Lambda) d\sigma \\
&\leq \frac{\xi_\alpha}{\Gamma(\gamma)} + \frac{M}{\Gamma(\wp)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\wp+\gamma-1} B(\gamma, \wp) \|\xi\|_{\zeta_{1-\gamma, \mathfrak{S}}} \\
&\quad + \frac{\Lambda}{\Gamma(\wp+1)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^\wp \\
&\leq \frac{\xi_\alpha}{\Gamma(\gamma)} + \frac{1}{\Gamma(\wp)} (\mathfrak{S}(\chi) - \mathfrak{S}(\alpha))^\wp B(\gamma, \wp) \|\xi\|_{\zeta_{1-\gamma, \mathfrak{S}}} \\
&\quad + \frac{\Lambda}{\Gamma(\wp+1)} (\mathfrak{S}(\chi) - \mathfrak{S}(\alpha))^{\wp+1-\gamma}.
\end{aligned}$$

This demonstrates how the $\mathfrak{C}_r = \{\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi] : \|\xi\|_{\zeta_{1-\gamma, \mathfrak{S}}} \leq r\}$ is transformed by Λ into itself. The proof is broken down into different steps:

Step 1: Λ is continuous.

Let ξ_n and $\xi_n \rightarrow \xi$ in $\zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$. Then for all $v \in J$,

$$\begin{aligned}
& \left| ((\Lambda\xi_n)(v) - (\Lambda\xi)(v)) (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma} \right| \\
&\leq \left| \frac{(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{1-\gamma}}{\Gamma(\wp)} \right. \\
&\quad \times \int_\alpha^v \mathfrak{S}'(\sigma)(\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\wp-1} \varpi(\sigma, \xi_n(\sigma), G\xi_n(\sigma), K\xi_n(\sigma)) d\sigma \\
&\quad \left. - \frac{(\mathfrak{S}(v) - \mathfrak{S}(\alpha))}{\Gamma(\wp)} \int_\alpha^v \mathfrak{S}'(\sigma)(\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\wp-1} \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma \right|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{(\mathfrak{F}(v) - \mathfrak{F}(\alpha))^{1-\gamma}}{\Gamma(\varrho)} \int_{\alpha}^v \mathfrak{F}'(\sigma) (\mathfrak{F}(v) - \mathfrak{F}(\sigma))^{\varrho-1} \\
&\quad \times \left| \varpi(\sigma, \xi_n(\sigma), G\xi_n(\sigma), K\xi_n(\sigma)) - \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) \right| d\sigma \\
&\leq \frac{(\mathfrak{F}(v) - \mathfrak{F}(\alpha))^{1-\gamma}}{\Gamma(\varrho)} (\mathfrak{F}(v) - \mathfrak{F}(\alpha))^{\varrho+\gamma-1} B(\gamma, \varrho) \\
&\quad \times \left\| \varpi(\cdot, \xi_n(\cdot), G\xi_n(\cdot), H\xi_n(\cdot)) - \varpi(\cdot, \xi(\cdot), G\xi(\cdot), K\xi(\cdot)) \right\|_{\zeta_{1-\gamma, \mathfrak{S}}}
\end{aligned}$$

which implies

$$\begin{aligned}
\|\Lambda\xi_n - \Lambda\xi\|_{\zeta_{1-\gamma, \mathfrak{S}}} &\leq B(\gamma, \varrho) \frac{(\mathfrak{F}(\chi) - \mathfrak{F}(\alpha))^{\varrho}}{\Gamma(\varrho)} \\
&\quad \times \left\| \varpi(\cdot, \xi_n(\cdot), G\xi_n(\cdot), K\xi_n(\cdot)) - \varpi(\cdot, \xi(\cdot), G\xi(\cdot), K\xi(\cdot)) \right\|_{\zeta_{1-\gamma, \mathfrak{S}}}.
\end{aligned}$$

This means that Λ is continuous.

Step 2: $\Lambda(\mathfrak{C}_r)$ is uniformly bounded.

It is obvious that $\Lambda(\mathfrak{C}_r) \subset \mathfrak{C}_r$ is bounded

Step 3: $\Lambda(\mathfrak{C}_r)$ is relatively compact.

It makes sense when $\Lambda(\mathfrak{C}_r) \subset \mathfrak{C}_r$ that $\Lambda(\mathfrak{C}_r)$ is uniformly bounded, and to demonstrate that Λ is an equicontinuous operator. Assume that $v_1, v_2 \in J$, $v_1 < v_2$, \mathfrak{C}_r is bounded in $\zeta_{1-\gamma, \rho}[\alpha, \chi]$. Then,

$$\begin{aligned}
&|((\Lambda\xi)(v_1) - (\Lambda\xi)(v_2))| \\
&\leq \frac{\xi_{\alpha}}{\Gamma(\gamma)} \left| (\mathfrak{F}(v_1) - \mathfrak{F}(\alpha))^{\gamma-1} - (\mathfrak{F}(v_2) - \mathfrak{F}(\alpha))^{\gamma-1} \right| \\
&\quad + \frac{B(\gamma, \varrho)}{\Gamma(\varrho)} \left((\mathfrak{F}(v_1) - \mathfrak{F}(\alpha))^{\varrho+\gamma-1} - (\mathfrak{F}(v_2) - \mathfrak{F}(\alpha))^{\varrho+\gamma-1} \right) \|\varpi\|_{\zeta_{1-\gamma, \mathfrak{S}}}.
\end{aligned}$$

As $v_1 \rightarrow v_2$, The aforementioned inequality's right side goes to zero. We may infer that $\Lambda : \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi] \rightarrow \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ is continuous and totally continuous as a result of Steps 1 through 3 and the Arzela-Ascoli theorem. \square

4. GENERALIZED STABILITY

Let $\epsilon > 0$ and $\varphi : I \rightarrow \mathbb{R}^+$ be continuous. We consider the following inequalities:

$$\left| D_{\alpha_+}^{\varrho, \beta; \mathfrak{S}} \eta(v) - \varpi(v, \eta(v), G\eta(v), K\eta(v)) \right| \leq \epsilon, \quad v \in J, \quad (4.1)$$

$$\left| D_{\alpha_+}^{\varrho, \beta; \mathfrak{S}} \eta(v) - \varpi(v, \eta(v), G\eta(v), K\eta(v)) \right| \leq \epsilon \varphi(v), \quad v \in J, \quad (4.2)$$

$$\left| D_{\alpha_+}^{\varrho, \beta; \mathfrak{S}} \eta(v) - \varpi(v, \eta(v), G\eta(v), K\eta(v)) \right| \leq \varphi(v), \quad v \in J. \quad (4.3)$$

Definition 4.1. ([9]) The problem (1.1) is said to be Hyers-Ulam stable if for all $\epsilon > 0$, there exist $\zeta_{\varpi} > 0$ and for each solution $\eta \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ of the (4.1) there exists a solution $\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ of the problem (1.1) such that

$$|\eta(v) - \xi(v)| \leq \zeta_{\varpi} \epsilon, \quad v \in J.$$

Definition 4.2. ([10]) The problem (1.1) is said to be generalized Hyers-Ulam stable if there exists $\varphi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ with $\varphi_{\varpi}(0) = 0$ such that for all $\eta \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ solution of (4.1) there exists a solution $\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ of (1.1) with

$$|\eta(v) - \xi(v)| \leq \varphi_{\varpi} \epsilon, \quad v \in J.$$

Definition 4.3. ([10]) The problem (1.1) is said to be Hyers-Ulam-Rassias stable in relation to $\varphi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ if there exists a real number $\zeta_{\varpi, \varphi} > 0$ and for all $\epsilon > 0$ and $\eta \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ of the inequality (4.2) there exists a solution of (1.1), $\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ with

$$|\eta(v) - \xi(v)| \leq \zeta_{\varpi, \varphi} \varphi(v), \quad v \in J.$$

Definition 4.4. ([9]) The problem (1.1) is said to be generalized Hyers-Ulam-Rassias stable in relation to $\varphi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ if there exists $\zeta_{\varpi, \varphi} > 0$ such that for each solution $\eta \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ of (4.3) there exists $\xi \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ is a solution of (1.1) with

$$|\eta(v) - \xi(v)| \leq \zeta_{\varpi, \varphi} \varphi(v), \quad v \in J.$$

Remark 4.5. Clearly, we have the following implication:

- (1) The Hyers-Ulam stability is the generalized Hyers-Ulam stability.
- (2) The Hyers-Ulam-Rassias stability is generalized Hyers-Ulam-Rassias stability.
- (3) If the problem (1.1) is generalized Hyers-Ulam-Rassias stable with $\varphi(v) = 1$ then it is Hyers-Ulam stable.

Remark 4.6. Let a function $\eta \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$. Then η is a solution of the inequality (4.1),

$$\left| D_{\alpha_+}^{\varphi, \beta; \mathfrak{S}} \eta(v) - \varpi(v, \eta(v), G\eta(v), K\eta(v)) \right| \leq \epsilon, \quad v \in J$$

if and only if there exists $g \in \zeta_{1-\gamma, \mathfrak{S}}[\alpha, \chi]$ such that

- (i) $|g(v)| \leq \epsilon, v \in J$
- (ii) $D_{\alpha_+}^{\varphi, \beta; \mathfrak{S}} \eta(v) = \varpi(v, \eta(v), G\eta(v), K\eta(v)) + g(v), v \in J.$

Remark 4.7. If the inequality (4.1) has a solution of η , then the following inequality has a solution of η .

$$\left| \eta(v) - \frac{\eta_\alpha}{\Gamma(\gamma)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} - \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \right. \\ \left. \times (\varpi(\sigma, \eta(\sigma), G\eta(\sigma), K\eta(v))) d\sigma \right| \leq \epsilon \frac{(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^\varrho}{\Gamma(\varrho + 1)}.$$

Indeed, by Remark 4.6 we have that

$$D_{\alpha+}^{\varrho, \beta; \mathfrak{S}} \eta(v) = \varpi(v, \eta(v), G\eta(v), K\eta(v)) + g(v), \quad v \in J.$$

Then

$$\eta(v) = \frac{\eta_\alpha}{\Gamma(\gamma)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \\ + \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} (\varpi(\sigma, \eta(\sigma), G\eta(\sigma), K\eta(v)) + g(\sigma)) d\sigma.$$

From this it follows that

$$\left| \eta(v) - \frac{\eta_\alpha}{\Gamma(\gamma)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \right. \\ \left. - \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} (\varpi(\sigma, \eta(\sigma), G\eta(\sigma), K\eta(\sigma))) d\sigma \right| \\ = \left| \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} g(\sigma) d\sigma \right| \\ \leq \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} |g(\sigma)| d\sigma \\ \leq \epsilon \frac{(\mathfrak{S}(v) - \mathfrak{S}(\alpha))^\varrho}{\Gamma(\varrho + 1)}.$$

We make similar observations about inequality (4.2) and inequality (4.3).

The major findings of generalized Hyers-Ulam-Rassias stability are now presented in this section.

(H4) There are rising functions ($\lambda_\varphi > 0$) such that for each $v \in J$, $\varphi \in \zeta_{1-\gamma, \rho}[\alpha, \chi]$,

$$I_{\alpha+}^{\varrho; \mathfrak{S}} \varphi(v) \leq \lambda_\varphi \varphi(v).$$

Theorem 4.8. *If the hypotheses (H1) and (H4) are true, then the issue (1.1) is Hyers-Ulam-Rassias stable in general.*

Proof. Assuming that η is the solution to (4.3). Then ξ is the only possible solution to the

$$\begin{cases} D_{\alpha_+}^{\varrho, \beta; \mathfrak{S}} \xi(v) &= \varpi(v, \xi(v), G\eta(v), K\eta(v)), \quad v \in J, \\ I_{\alpha_+}^{1-\gamma; \mathfrak{S}} \xi(\alpha) &= I_{\alpha_+}^{1-\gamma; \mathfrak{S}} \eta(\alpha), \end{cases}$$

when

$$\frac{L(1 + G^* + K^*)}{\Gamma(\varrho)} B(\gamma \cdot \varrho) (\mathfrak{S}(\chi) - \mathfrak{S}(\alpha))^\varrho < 1.$$

Then we have

$$\begin{aligned} \xi(v) &= \frac{\eta_\alpha}{\Gamma(\varrho)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \\ &\quad + \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma. \end{aligned}$$

By differentiating inequality (4.3), we have

$$\begin{aligned} &\left| \eta(v) - \frac{\eta_\alpha}{\Gamma(\varrho)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \right. \\ &\quad \left. - \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \varpi(\sigma, \eta(\sigma), G\eta(\sigma), K\eta(\sigma)) d\sigma \right| \\ &\leq \lambda_\varphi \varphi(v). \end{aligned}$$

Hence it follows

$$\begin{aligned} &|\eta(v) - \xi(v)| \\ &\leq \left| \eta(v) - \frac{\eta_\alpha}{\Gamma(\varrho)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \right. \\ &\quad \left. - \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma \right| \\ &\leq \left| \eta(v) - \frac{\eta_\alpha}{\Gamma(\varrho)} (\mathfrak{S}(v) - \mathfrak{S}(\alpha))^{\gamma-1} \right. \\ &\quad \left. - \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \varpi(\sigma, \eta(\sigma), G\eta(\sigma), K\eta(\sigma)) d\sigma \right| \\ &\quad + \left| \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \varpi(\sigma, \eta(\sigma), G\eta(\sigma), K\eta(\sigma)) d\sigma \right. \\ &\quad \left. - \frac{1}{\Gamma(\varrho)} \int_\alpha^v \mathfrak{S}'(\sigma) (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} \varpi(\sigma, \xi(\sigma), G\xi(\sigma), K\xi(\sigma)) d\sigma \right| \\ &\leq \lambda_\varphi \varphi(v) + \frac{L(1 + H)}{\Gamma(\varrho)} \int_\alpha^v (\mathfrak{S}(v) - \mathfrak{S}(\sigma))^{\varrho-1} |\eta(\sigma) - \xi(\sigma)| d\sigma. \end{aligned}$$

According to Lemma 2.5, there exists a constant $M^* > 0$ that is not reliant on $\lambda_\varphi \varphi(v)$ such that

$$|\eta(v) - \xi(v)| \leq M^* \lambda_\varphi \varphi(v) := \zeta_{\varpi, \varphi} \varphi(v).$$

As a result, the problem (1.1) is Ulam-Hyers-Rassias stable in its generalized form. \square

Remark 4.9. Clearly, we have the followings:

- (1) We take into consideration the inequality (4.2) as well as the issue (1.1) under the premise of Theorem 4.8. The Ulam-Hyers-Rassias stability of the issue (1.1) may be confirmed by repeating the same procedure.
- (2) We take into consideration the inequality (4.1) and the issue (1.1) under the premise of Theorem 4.8. The same procedure may be used to confirm that the Ulam-Hyers stability of the issue (1.1) can be established.

REFERENCES

- [1] M. Alesemi, N. Iqbal and A.A. Hamoud, *The analysis of fractional-order proportional delay physical models via a novel transform*, Complexity, **2022** (2022), 1–13.
- [2] S. Asawasamrit, Y. Thadang, S.K. Ntouyas and J. Tariboon, *Non-instantaneous impulsive boundary value problems containing Caputo fractional derivative of a function with respect to another function and Riemann-Stieltjes fractional integral boundary conditions*, Axioms, **10** (2021), 130.
- [3] K. Balachandran, S. Kiruthika and J.J. Trujillo, *Existence results for fractional impulsive integrodifferential equations in Banach spaces*, Commu. Nonlinear Sci. Numer. Simu., **16** (2011), 1970–1977.
- [4] K. Balachandran and K. Uchiyama, *Existence of local solutions of quasilinear integrodifferential equations in Banach spaces*, Applicable Anal., **76**(1-2) (2000), 1–8.
- [5] K.M. Furati, M.D. Kassim and N.E. Tatar, *Existence and uniqueness for a problem involving Hilfer fractional derivative*, Comput. Math. Appl., **64** (2012), 1616–1626.
- [6] A. Hamoud, M.SH. Bani Issa and K. Ghadle, *Existence and uniqueness results for nonlinear Volterra-Fredholm integro-differential equations*, Nonlinear Funct. Anal. Appl., **23**(4) (2018), 797–805.
- [7] A. Hamoud, A. Khandagale, R. Shah and K. Ghadle, *Some new results on Hadamard neutral fractional nonlinear Volterra-Fredholm integro-differential equations*, Discontinuity, Nonlinearity, and Complexity, **12**(4) (2023) 893–903.
- [8] R. Hilfer, *Applications of fractional Calculus in Physics*, World scientific, Singapore, 1999.
- [9] R.W. Ibrahim, *Generalized Ulam-Hyers stability for fractional differential equations*, Int. J. Math., **23**(5) (2012), 1–14.
- [10] R.W. Ibrahim, *Stability of sequential fractional differential equation*, Appl. Comput. Math., **14**(2) (2015), 19–41.
- [11] A.G. Ibrahim and A.A. Elmandouh, *Existence and stability of solutions of \mathfrak{S} -Hilfer fractional functional differential inclusions with non-instantaneous impulses*, AIMS Mathematics, **6** (2021), 10802–10832.
- [12] K. Ivaz, I. Alasadi and A. Hamoud, *On the Hilfer fractional Volterra-Fredholm integro differential equations*, IAENG Int. J. Appl. Math., **52**(2) (2022), 426–431.
- [13] T. Jankowski, *Delay integro-differential equations of mixed type in Banach spaces*, Glasnik Math., **37**(57) (2002), 321–330.

- [14] M.D. Kassim and N.-E. Tatar, *Well-posedness and stability for a differential problem with Hilfer-Hadamard fractional derivative*, Abst. Appl. Anal., Article ID 605029 (2014), 1–7.
- [15] U.N. Katugampola, *Existence and uniqueness results for a class of generalized fractional differential equations*, Bull. Math. Anal. Appl., arXiv:1411.5229, v1 (2014).
- [16] A.A. Kilbas, H.M. Srivastava and J.J. Trujillo, *Theory and applications of fractional differential equations*, Amsterdam: Elsevier, 2006.
- [17] C. Kou, J. Liu and Y. Ye, *Existence and uniqueness of solutions for the Cauchy-Type problems of fractional differential equations*, Discrete Dyna. Nature and Soc., Article ID 142175 (2010), 15pages.
- [18] T. Li, A. Zada and S. Faisal, *Hyers-Ulam stability of n th order linear differential equations*, J. Nonlinear Sci. Appl., **9** (2016), 2070–2075.
- [19] C. Nuchpong, S.K. Ntouyas and J. Tariboon, *Boundary value problems of Hilfer-type fractional integro-differential equations and inclusions with nonlocal integro-multipoint boundary conditions*, Open Math., **18** (2020), 1879–1894.
- [20] D.S. Oliveira and E. Capelas de Oliveira, *Hilfer-Katugampola fractional derivative*, Mathematics, Cornell University, arxiv:1705.07733v1, (2017).
- [21] I. Podlubny, *Fractional differential equations*, Academic Press, San Diego, 1999.
- [22] A. Salim, M. Benchohra, J.R. Graef and J.E. Lazreg, *Boundary value problem for fractional order generalized Hilfer-type fractional derivative with non-instantaneous impulses*, Fractal Fract., **5** (2021), 1–21.
- [23] A. Sharif and A. Hamoud, *On ψ -Caputo fractional nonlinear Volterra-Fredholm integro-differential equations*, Discontinuity, Nonlinearity, and Complexity, **11**(1) (2022), 97–106.
- [24] A. Sharif, A. Hamoud and K.P. Ghadle, *On existence and uniqueness of solutions to a class of fractional Volterra-Fredholm initial value problems*, Discontinuity, Nonlinearity, and Complexity, **12**(4) (2023) 905–916.
- [25] J. Vanterler da C. Sousa and E. Capelas de Oliveira, *On the \mathfrak{S} -Hilfer fractional derivative*, Mathematics, Cornell University, arXiv:1704.08186, (2017).
- [26] J. Wang, L. Lv and Y. Zhou, *Ulam stability and data dependence for fractional differential equations with Caputo derivative*, Elect. J. Qualitative Theory Diff. Equ., **63** (2011), 1–10.
- [27] J. Wang and Y. Zhang, *Ulam-Hyers-Mittag-Leffler stability of fractional-order delay differential equations*, Optimization: A Journal of Mathematical Programming and Operations Research, **63**(8) (2014), 1181–1190.
- [28] H. Ye, J. Gao and Y. Ding, *A generalized Gronwall inequality and its application to a fractional differential equation*, J. Math. Anal. Appl., **328** (2007), 1075–1081.