



Research Note

Nutrient composition and taste properties of *Sargassum fusiforme* from the Wando region

Hae-In Lee¹, Cheong-Bin Kim², Mi-Kyung Lee^{1*}

¹Department of Food and Nutrition, Sunchon National University, Suncheon 57922, Korea

²Department of Physics Education, Sunchon National University, Suncheon 57922, Korea

Abstract This study examined the nutrient composition and taste properties of *Sargassum fusiforme* (SF) from two locations in the Wando region (Wando-eup: WD-A, Sinji-myeon: WD-B) and control region in Korea. The protein content of the SF in Wando regions was higher than that of the control, while the carbohydrate content was lower. The WD-A region had the lowest calorie content and the highest ash level when compared to the other regions. The potassium, iron, and zinc contents of the in WD-A were significantly higher than the other groups, but the sodium, phosphorus, and magnesium contents were lower. Free and constitutive amino acids, as well as the total and essential amino acid content, were higher in Wando regions than in the control. When analyzing using an electronic tongue, the SF from the WD-A region had the highest umami and sourness and the lowest bitterness and sweetness. Therefore, the results of this study can serve as fundamental data for seaweed research in the Wando region.

Keywords *Sargassum fusiforme*, Wando, nutrient composition, physicochemical properties, taste properties



OPEN ACCESS

Citation: Lee HI, Kim CB, Lee MK. Nutrient composition and taste properties of *Sargassum fusiforme* from the Wando region. Food Sci. Preserv., 31(2), 324-331 (2024)

Received: December 20, 2023
Revised: February 27, 2024
Accepted: February 28, 2024

***Corresponding author**
Mi-Kyung Lee
Tel: +82-61-750-3656
E-mail: leemk@snu.ac.kr

Copyright © 2024 The Korean Society of Food Preservation. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Marine algae, or seaweeds, are integral to Asian diets and traditional medicine, particularly in Korea. Their recognized health benefits have spurred global interest (Peñalver et al., 2020). Seaweeds are classified into brown, red, and green algae, each having distinct components (Lee et al., 2020a). *Sargassum fusiforme* (SF), a brown alga, is perennial and belongs to the Sargassaceae family (Lee et al., 2020b). Predominantly found along Korea's southwestern coast, SF has a unique flavor and texture, making it popular in Korea, Japan, and China (Park and Ryu, 2013). SF is a rich source of dietary fiber, minerals, vitamins, and functional compounds such as polyphenols and polysaccharides, including fucoidan, laminaran, and alginic acid (Shin et al., 2014). Its polysaccharide, alginate, consists of α -L-guluronic and β -D-mannuronic acid chains, which form the cell wall. The resistance of alginate to degradation *in vivo* and its unique structure mean that it is physiologically active (Kwon and Youn, 2017). SF also contains phlorotannins, secondary metabolites formed of phloroglucinol units (Ferrerres et al., 2012). Because of its high content of nutrients and bioactive compounds, SF has potential for disease prevention and treatment, for example, anti-inflammatory, anti-obesity, and anti-cancer effects (Sugiura et al., 2016; Wei et al., 2020). Ancient texts, such as the pharmacopoeia titled "Shen Nong's Canon of Materia Medica," document the therapeutic uses of SF, such as treating thyroid tumors and alleviating edema and reducing abdominal sounds (Zhang et al., 2020). In addition, the high glutamic and aspartic acid content, key flavor enhancers, of SF has led to the development of

seaweed-based processed foods (Dai et al., 2019).

The Jeollanam-do province produces 1.68 million tons of seaweed, accounting for 91% of Korea's overall seaweed yield, and the Wando region contributes 44% nationally (MOF, 2022; Wandogun, 2022). The ecological conditions of Wando, including a high ecological evaluation index, tidal currents, and species diversity, create an ideal seaweed habitat. This is attributed to the geography of the region, which ensure a steady influx of nutrient-rich coastal and open sea waters, including the cold bottom water of the Yellow Sea (Cho and Choi, 2005; Son, 2011). Notably, the nutrient profile and physicochemical properties of seaweed are strongly affected by the marine environment (temperature, tidal current, and inhabiting marine species), and this results in consumer preference for region-specific seaweeds (Lim and Choi, 2005).

In this study, we analyzed the nutritional composition and taste profiles of *Sargassum fusiforme* (SF), a seaweed vital to the economy of the Wando region and compared them to those of SF from another area in Jeollanam-do to establish a database for SF component analysis.

2. Materials and methods

2.1. Sample collection and preparation

SF samples were harvested from Wando (Wando-eup, WD-A; Sinji-myeon, WD-B) and a control region (Jodo-myeon, Jindo: control) in Dec 2022. After thorough washing under running water to remove impurities and salts, the SF was rinsed with distilled water, strained, and freeze-dried (Operon Co., Gimpo, Korea) for experiments.

2.2. Proximate composition analysis

The proximate composition of SF were determined following the AOAC (2019) method. Moisture was assessed using an air-oven at 105°C, crude fat was extracted by Soxhlet extraction, and crude protein was assessed using the Kjeldahl method with a nitrogen-protein analyzer (Kjeltec 8400 System, Foss, Hoganas, Sweden). The ash content was measured post-ashing in a muffle furnace (F6010, Thermo Fisher Scientific, Waltham, MA, USA) at 600°C. The carbohydrate content was calculated by subtracting the moisture, ash, protein, and fat content from the total weight of the sample, and the caloric value was calculated following

Korean Food Code standards.

2.3. Mineral content analysis

The samples (0.2-1.0 g) were mixed with nitric acid (7 mL) and hydrogen peroxide (1 mL) and heated for 30 minutes. After heating, the mixture was digested using a microwave digestion system (MARS 6, CEM Corporation, Matthews, NC, USA) and then cooled. Then, a sample was analyzed using inductively coupled plasma optical emission spectrometry (ICP-OES 8300, PerkinElmer Inc., Waltham, MA, USA).

2.4. Dietary fiber analysis

Total dietary fiber was quantified by the AOAC method (2019). In this process, 50 mM MES/Tris buffer (40 mL; pH 8.2), α -amylase (50 μ L), and sample (1 g) were mixed for 15 min at 95°C. After cooling to 60°C, the 100 μ L of protease solution was added, and allowed to react for 30 minutes before adjusting the pH to 4.5. Subsequently, aminoglucosidase solution was added, and hydrolysis was carried out for 30 min at 60°C. The precipitate was then filtered and dried to measure the fiber content.

2.5. Amino acid analysis

For constitutive amino acid analysis, a sample (0.1 g) underwent hydrolysis with 6 N HCl (10 mL) under nitrogen gas for 22 h at 110°C, followed by concentration in under reduced pressure. Free amino acids were extracted from the sample (0.5 g) with 70% ethanol (50 mL) for 30 min, followed by standing for 10 min; the sample was then centrifuged at 1,500 rpm for 15 min three times and concentrated in under reduced pressure. Pretreated samples were dissolved in 0.02 N HCl (20 mL), filtered, and analyzed with an automated amino acid analyzer (Hitachi L-8900, Hitachi, Tokyo, Japan) (Ham et al., 2021).

2.6. Taste pattern analysis by electronic sensor system

A sample (5 g) was added to distilled water (100 mL), stirred at 60°C for 1 h, and filtered (Whatman No. 2). The taste profile (sourness, sweetness, bitterness, saltiness, and umami) was assessed using an electronic sensor system (ASTREE II, Alpha MOS, Toulouse, France), and GPS and

SPS sensors for calibration. The samples introduced into the device were analyzed using seven sensors for 120 s, and the sensors were washed between sample analysis to prevent contamination.

2.7. Statistical analysis

Independent experiments were conducted in triplicate, and the results are presented as mean±standard deviation. One-way ANOVA (SPSS version 27.0, SPSS Inc., Chicago, IL, USA) was used to determine the statistical significance, and Duncan's multiple test post-hoc for group differences was set at $p < 0.05$. Discriminant function analysis (DFA) was used to compare the taste pattern obtained from electronic tongue analysis.

3. Results and discussion

3.1. Proximate composition and fiber content

Proximate composition analysis revealed that WD-A and WD-B SF had significantly higher crude protein contents than the control (Fig. 1). The ash content of WD-A was the significantly highest (40.67%), surpassing those of WD-B (37.20%) and the control (36.20%) (Fig. 1). The carbohydrate

levels varied: the control region had the highest value (48.73%), followed by those of WD-B and WD-A at 46.07% and 43.13%, respectively. The caloric value of WD-A was the lowest (242.70 kcal/100 g). Further, the dietary fiber and crude lipid were similar for all samples (Fig. 1). Consistent with the findings of Jung (2017) for SF from Wando, Goheung, and Jindo regions, in this study, the SF from Wando showed elevated protein and ash contents compared to the other samples but similar crude lipid levels. The high ash content is attributed to the quartz porphyry of Wando, suggesting the role of the environment on the proximate contents, but there is need for further correlation studies.

3.2. Mineral content

Seaweeds are valued for their mineral content, which can be similar to or exceed those of other foods (Lozano Muñoz and Díaz, 2020). WD-A SF contained significantly more potassium but less sodium, phosphorus, and magnesium than WD-B and the control (Fig. 2). Given the role of potassium in sodium excretion and the increase in sodium-related chronic diseases, potassium-rich foods such as SF are gaining attention. Na et al. (2022) linked a high sodium and low potassium intake to an increased prevalence of obesity and




Fig. 1. The proximate compositions of the freeze-dried *Sargassum fusiforme*. All values are mean±SD (n=3). Different superscript letters (a-c) on the bars indicate significant differences ($p < 0.05$) by Duncan's multiple range test. WD-A, *Sargassum fusiforme* in Wando-eup; WD-B, *Sargassum fusiforme* in Sinji-myeon of Wando; control, *Sargassum fusiforme* in Jindo.




Fig. 2. The mineral contents of the freeze-dried *Sargassum fusiforme*. All values are mean \pm SD (n=3). Different superscript letters (^{a-c}) on the bars indicate significant differences (p<0.05) by Duncan's multiple range test. WD-A, *Sargassum fusiforme* in Wando-eup; WD-B, *Sargassum fusiforme* in Sinji-myeon of Wando; control, *Sargassum fusiforme* in Jindo.

hypertension. As in our study, Yoon et al. (2022) found SF to have the highest potassium content among 11 seaweed types; specifically, a single serving (15 kcal) was found to provide 35% of the recommended potassium intake for adult males. Notably, the mineral profile of WD-A SF, which has high potassium and low sodium levels, stands out among the samples from different regions.

SF from WD-A contained 1.4- and 2.0-times greater iron levels than WD-B and the control, respectively, and this represents a significant difference (Fig. 2). Environmental factors such as ocean salinity, tidal currents, water temperature, and organic matter content influence the nutrient profiles of marine plants and benthic animals. Seaweeds, in particular, has been reported to be rich in minerals essential for growth and bioregulation. Because it selects and accumulates specific elements among the minerals in seawater (Yamamoto et al., 1984). The seawater of the Wando region is known for its high carbon absorption, and the iron concentration of the tidal flats, which is influenced by the quartz porphyry, is higher than other coastal areas, including the control region (data not shown). This environment likely contributes to the elevated iron levels in SF of WD-A. Additionally, SF of WD-A was found to have the highest zinc content (0.68

mg/100 g) of all samples (Fig. 2). Consistent with this finding, high zinc levels have been observed in samples of *Lajonkairia lajonkairii* from Wando, suggesting that both benthic organisms are affected by the unique marine conditions of the Wando region (Ham et al., 2021).

3.3. Amino acid composition

Amino acids are categorized into constitutive amino acids and free amino acids. Free amino acids enhance the taste and flavor of foods, thus affecting their palatability (Mustafa et al., 2007). The free amino acid analysis revealed that SF of WD-A had the highest aspartic acid levels, a key taste-contributing amino acid. In contrast, the seaweed from the control region was rich in glutamic acid and glycine, whereas SF of WD-B contained high levels of other amino acids (Table 1). Notably, taurine and tyrosine, known fatigue-reducing amino acids, were exclusively found in WD-A. Regarding the constitutive amino acids, WD-A had the highest histidine contents, but WD-B had higher levels than the control for all other amino acids (Table 2). Notably, histidine, which is an essential amino acid for infants but not adults, plays a role in hematopoiesis (Kim et al., 2017).

Table 1. The free amino acid composition of the freeze-dried *Sargassum fusiforme*

	WD-A	WD-B	control
Essential amino acids			
Valine	23.75±1.15 ^{1)(b2)}	25.73±0.65 ^b	16.30±0.35 ^a
Isoleucine	3.65±0.35 ^a	5.10±1.08 ^b	4.10±0.36 ^{ab}
Leucine	7.75±0.85 ^a	11.63±0.40 ^b	7.57±1.24 ^a
Threonine	14.40±0.47 ^a	21.90±0.79 ^b	13.97±0.47 ^a
Lysine	3.80±0.44 ^a	3.27±0.81 ^a	3.80±0.60 ^a
Total	53.57±0.75 ^b	67.63±3.55 ^c	45.73±1.97 ^a
Non-essential amino acids			
Taurine	32.10±0.23	– ³⁾	–
Tyrosine	2.20±2.43	–	–
Aspartic acid	326.55±5.97 ^c	313.87±5.50 ^b	225.50±6.09 ^a
Serine	12.75±0.40 ^a	16.07±0.40 ^b	12.23±0.40 ^a
Alanine	271.50±5.70 ^a	369.83±7.52 ^c	308.13±7.85 ^b
Cystathionine	9.75±0.55 ^a	14.80±0.61 ^c	12.30±0.46 ^b
Glutamic acid	114.15±2.57 ^b	99.93±3.59 ^a	146.40±5.27 ^c
Glycine	4.60±0.12 ^b	3.50±0.26 ^a	8.50±0.17 ^c
Total	836.27±16.14 ^b	885.63±20.88 ^c	758.80±21.97 ^a

¹⁾All values are mean±SD (n=3).

²⁾Means with different superscripts within each row are significantly different (p<0.05) by Duncan's multiple range test.

³⁾Not detected.

Essential amino acids, which must be obtained from the diet because they are not synthesized endogenously (An et al., 2020), were most abundant in WD-B, followed by WD-A, and both samples had significantly higher levels than the control. WD-A was particularly rich in valine, whereas WD-B contained all essential amino acids except lysine. Thus, SF from Wando is an excellent source of essential amino acids.

3.4. Taste profile analysis

Electronic tongue systems are capable of analyzing primary tastes such as umami (NMS), sweetness (PKS), bitterness (ANS), saltiness (CTS), and sourness (AHS) and have, thus, become prominent for food characteristic and taste profile analysis (Hong et al., 2020). Crucially, these systems provide objective taste measurements using non-destructive methods without pretreatment (Hong et al., 2021). Therefore, in this

study, we compared SF taste patterns from WD-A, WD-B, and the control region using an electronic tongue (Fig. 3). The NMS score of WD-A was 8.8, which is significantly higher than those of WD-B and the control (by 1.9- and 2.0-times, respectively). AHS was also ≥1.8-times higher in WD-A than in the WD-B and control samples. In contrast, the control region scored highest for PKS (7.6) and ANS (7.7); WD-B showed a similar trend but with lower values, whereas WD-A showed significantly lower PKS and ANS values. CTS was similar for all samples. Notably, humans can discern taste differences when primary taste values differ by ≥2.0, and greater differences indicate stronger taste perception (Jo et al., 2012). The high umami and sourness values of WD-A are consistent with its high glutamic and aspartic acid levels, key contributors to umami flavor. In contrast, WD-B had high sweetness and bitterness values. In the DFA to identify patterns based on these results, the discriminant function first score (DF1) was 97.68%, and the

Table 2. The constitutive amino acid composition of the freeze-dried *Sargassum fusiforme*

	WD-A	WD-B	control
Essential amino acids			
Valine	494.35±3.42 ^{1)bc2)}	512.77±5.09 ^c	448.60±4.38 ^a
Isoleucine	348.80±3.27 ^b	360.40±1.10 ^b	327.13±6.73 ^a
Leucine	671.15±8.28 ^b	730.27±9.49 ^c	638.50±12.03 ^a
Lysine	607.65±9.62 ^b	626.07±4.11 ^b	570.83±3.94 ^a
Threonine	502.15±4.52 ^b	547.50±16.67 ^c	460.20±5.17 ^a
Histidine	280.75±0.65 ^b	274.13±13.32 ^b	238.83±3.34 ^a
Methionine	253.75±1.24 ^a	280.73±6.52 ^b	241.90±3.35 ^a
Phenylalanine	383.35±7.08 ^a	414.10±3.45 ^b	379.20±6.55 ^a
Total	3,572±36 ^b	3,746±54 ^c	3,305±37 ^a
Non-essential amino acids			
Aspartic acid	1,232±22.37 ^b	1,246±32.49 ^b	1,051±8.49 ^a
Serine	446.25±7.05 ^b	482.47±19.45 ^c	419.90±6.25 ^a
Glutamic acid	1,301±29.83 ^b	1,435±25.38 ^c	1,176±19.01 ^a
Glycine	459.65±9.37 ^b	476.93±8.71 ^b	425.40±8.77 ^a
Alanine	441.75±3.65 ^b	443.03±13.31 ^b	391.07±4.12 ^a
Cysteine	739.60±11.10 ^a	847.93±20.77 ^b	722.07±7.09 ^a
Tyrosine	179.95±3.20 ^a	200.50±3.80 ^b	181.90±4.48 ^a
Arginine	191.30±14.64 ^a	224.33±14.13 ^b	185.03±2.73 ^a
Proline	242.35±25.51 ^{ab}	279.33±3.65 ^b	236.33±2.29 ^a
Total	8,869±165 ^b	9,381±159 ^c	8,094±110 ^a

¹⁾All values are mean±SD (n=3).

²⁾Means with different superscripts within each row are significantly different ($p < 0.05$) by Duncan's multiple range test.

discriminant function second score (DF2) was 2.32%. With DF1 values as the reference, WD-A clustered in the negative direction, while WD-B and control clustered in the positive direction, indicating a significant differentiation in taste patterns among the regions (Fig. 4). Overall these differences suggest that the marine origin, environment, and climate influence the taste of SF (Dong et al., 2017).

4. Conclusions

In this study, we assessed the nutritional property and taste characteristics of *Sargassum fusiforme* (SF) from two regions in Wando (Wando-eup, WD-A; Sinji-myeon, WD-B) and a control area (Jodo-myeon, Jindo: control), focusing on macronutrients, amino acids, and minerals. Additionally, taste

patterns were evaluated using an electronic tongue system. The crude protein content was significantly higher in the WD-A and WD-B SF compared to the control regions, and ash content was also highest in the WD-A regions. Conversely, WD-A had the lowest carbohydrate contents and caloric values. Mineral analysis indicated that WD-A SF had high potassium, iron, and zinc contents, whereas sodium, phosphorus, and magnesium were more abundant in the control. Amino acid analysis showed that both WD-A and WD-B SF had higher levels of constitutive, free, and essential amino acids, with WD-B leading, followed by WD-A and the control. Notably, taurine and tyrosine, which are associated with fatigue reduction, were exclusive to WD-A. Taste evaluation demonstrated that WD-A SF was characterized by pronounced umami and sourness, whereas WD-B and the control were




Fig. 3. The radar diagram of electronic tongue sensor response of the freeze-dried *Sargassum fusiforme*. The taste of the samples was measured using an electronic tongue with the following sensors: AHS sensor for sourness, PKS sensor for sweetness, CTS sensor for saltiness, NMS sensor for umami, and ANS sensor for bitterness. WD-A, *Sargassum fusiforme* in Wando-eup; WD-B, *Sargassum fusiforme* in Sinji-myeon of Wando; control, *Sargassum fusiforme* in Jindo.




Fig. 4. Discriminant function analysis plot of the electronic tongue sensor response data for the freeze-dried *Sargassum fusiforme*. WD-A, *Sargassum fusiforme* in Wando-eup; WD-B, *Sargassum fusiforme* in Sinji-myeon of Wando; control, *Sargassum fusiforme* in Jindo.

distinguished by heightened bitterness and sweetness. Therefore, the nutrient composition and taste profile of SF vary with regional marine conditions, and these findings provide valuable insights for future seaweed research in the Wando region.

Funding

The research was supported by Wando-gun.

Acknowledgements

None.

Conflict of interests

The authors declare no potential conflicts of interest.

Author contributions

Conceptualization: Kim CB, Lee MK. Data curation: Lee MK. Formal analysis: Lee HI. Validation: Lee MK. Writing - original draft: Lee HI. Writing - review & editing: Lee HI, Lee MK.

Ethics approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

ORCID

Hae-In Lee (First author)

<https://orcid.org/0000-0003-0221-2253>

Cheong-Bin Kim

<https://orcid.org/0009-0002-9663-7753>

Mi-Kyung Lee (Corresponding author)

<https://orcid.org/0000-0002-6064-5295>

References

- An GH, Han JG, Cho JH. Comparisons of biological activities and amino acid contents of edible mushrooms extracted using different solvents. *J Mushrooms*, 18, 53-62 (2020)
- AOAC. Official Methods of Analysis. 21th ed, Association of Official Analytical Chemists, Gaithersburg, USA (2019)
- Cho ES, Choi YK. The characteristics of marine environment and phytoplankton community around southwestern waters for ichthyotoxic dinoflagellate *Cochlodinium polykrikoides* monitoring programme. *Int J Environ Sci*, 14, 177-184 (2005)
- Dai YL, Jiang YF, Lee HG, Jeon YJ, Kang MC. Characterization and screening of anti-tumor activity of fucoidan from acid-processed hijiki (*Hizikia fusiforme*). *Int J Biol Macromol*, 139, 170-180 (2019)
- Dong H, Moon JY, Lee SH. Discrimination of geographical origins of raw ginseng using the electronic tongue. *Korean J Food Sci Technol*, 49, 349-354 (2017)
- Ferreres F, Lopes G, Gil-Izquierdo A, Andrade PB, Sousa C, Mouga T, Valentao P. Phlorotannin extracts from fucales

- characterized by HPLC-DAD-ESI-MSN: Approaches to hyaluronidase inhibitory capacity and antioxidant properties. *Mar Drugs*, 10, 2766-2781 (2012)
- Ham JR, Lee HI, Kim CB, Shin EC, Lee MK. Nutritional composition and taste properties of abalone and short-neck clam in Wando. *J Korean Soc Food Sci Nutr*, 50, 1010-1018 (2021)
- Hong SJ, Boo CG, Heo SU, Jo SM, Yoon S, Jeong H, Lee Y, Park SS, Shin EC. Physicochemical characteristics of wintering radish produced in Jeju Island by different processing methods. *J Korean Soc Food Sci Nutr*, 50, 748-755 (2021)
- Hong SJ, Cho JJ, Boo CG, Youn MY, Lee SM, Shin EC. Comparison of physicochemical and sensory properties of bean sprout and peanut sprout extracts, subsequent to roasting. *J Korean Soc Food Sci Nutr*, 49, 356-369 (2020)
- Jo HS, Kim KH, Kim MJ, Kim HJ, Im YJ, Kwon DH, Heu MS, Kim JS. Sensory characterization of domestic mottled skate *Raja pulchra* as affected by area caught, sex and fish weight. *Korean J Fish Aquat Sci*, 45, 619-626 (2012)
- Jung HJ. The minerals analysis and nutritional evaluation according to production area laver, Japanese kelp, sea mustard, hijiki in Korea (*Porphyra tenera*, *Saccharina japonicus*, *Undaria pinnatifida*, *Sargassum fusiforme*). MS Thesis, Pukyong National University, Korea, p 12-14 (2017)
- Kim KJ, Im SB, Yun KW, Je HS, Ban SE, Jin SW, Jeong SW, Koh YW, Cho IK, Seo KS. Content of proximate compositions, free sugars, amino acids, and minerals in five *Lentinula edodes* cultivars collected in Korea. *J Mushrooms*, 15, 216-222 (2017)
- Kwon YR, Youn KS. Antioxidant and physiological activities of *Hizikia fusiforme* by extraction methods. *Korean J Food Preserv*, 24, 631-637 (2017)
- Lee CH, Park YN, Lee SG. Analysis and comparison of bioactive compounds and total antioxidant capabilities of Korean brown algae. *Korean J Food Sci Technol*, 52, 54-59 (2020)
- Lee YJ, Jeon YJ, Kim YT. Comparison of antioxidant and physiological activities of various solvent extracts from *Hizikia fusiformis*. *Korean J Fish Aquat Sci*, 53, 886-893 (2020)
- Lim HS, Choi JW. Ecological impact of the dyke construction on the marine benthos community of the oligohaline Youngam lake. *J Kor Fish Soc*, 38, 172-183 (2005)
- Lozano Munoz I, Diaz NF. Minerals in edible seaweed: Health benefits and food safety issues. *Crit Rev Med Inform*, 62, 1592-1607 (2020)
- MOF. Statistical yearbook of oceans and fisheries. Available from: <https://www.mof.go.kr>. Accessed Dec. 30, 2022.
- Mustafa A, Aman P, Andersson R, Kamal-Eldin A. Analysis of free amino acids in cereal products. *Food Chem*, 105, 317-324 (2007)
- Na W, Choi S, Kim C, Sohn C. The relationship between dietary sodium-to-potassium ratio and the levels of energy intake from dish group and obesity in Korean adults of thirties and forties: Based on the 6th to 7th (2013-2018) national health and nutrition survey. *Korean J Human Ecol*, 31, 787-797 (2022)
- Park H, Ryu HS. Effect of *Hizikia fusiforme* water extracts on splenocyte proliferation and cytokine production in mice. *J Korean Soc Food Scid Nutr*, 42, 1924-1929 (2013)
- Penalver R, Lorenzo JM, Ros G, Amarowicz R, Pateiro M, Nirto G. Seaweeds as a functional ingredient for a healthy diet. *Mar Drugs*, 18, 301-328 (2020)
- Shin DB, Han EH, Park SS. Cytoprotective effects of Phaeophyta extracts from the coast of Jeju Island in HT-22 mouse neuronal cells. *J Korean Soc Food Sci Nutr*, 43, 224-230 (2014)
- Son B. Marine algal flora of the South-West Coast, Korea. Ph D Thesis, Chonnam National University, Korea, p 54-55 (2011)
- Sugiura Y, Kinoshita Y, Abe M, Murase N, Tanaka R, Matsushita T, Usui M, Hanaoka KI, Miyata M. Suppressive effects of the diethyl ether fraction from a brown alga *Sargassum fusiforme* on allergic and inflammatory reactions. *Fish Sci*, 82, 369-377 (2016)
- Wandogun. Statistical yearbook. Available from: <https://www.wandogun.go.kr>. Accessed Dec. 10, 2022.
- Wei B, Zhong QW, Ke SZ, Zhou TS, Xu QL, Wang SJ, Chen JW, Zhang HW, Jin WH, Wang H. *Sargassum fusiforme* polysaccharides prevent high-fat diet-induced early fasting hypoglycemia and regulate the gut microbiota composition. *Mar Drugs*, 18, 444-457 (2020)
- Yamamoto T, Otsuka K, Okamoto K. Character of each element on its distribution in seaweeds. *Hydrobiologia*, 116, 510-512 (1984)
- Yoon SH, Kim SY, Park ES, Choi MK. Nutritional assessment focusing on proximate composition and mineral content of Korean seaweeds. *J East Asian Soc Diet Life*, 32, 321-329 (2022)
- Zhang R, Zhang X, Tang Y, Mao J. Composition, isolation, purification and biological activities of *Sargassum fusiforme* polysaccharides: A review. *Carbohydr Polym*, 228, 115381 (2020)