




Metabolic syndrome awareness in the general Korean population: results from a nationwide survey

Hyun-Jin Kim¹, Mi-Seung Shin², Kyung-Hee Kim³, Mi-Hyang Jung⁴, Dong-Hyuk Cho⁵, Ju-Hee Lee⁶, and Kwang Kon Koh⁷

¹Division of Cardiology, Department of Internal Medicine, Hanyang University Guri Hospital, Guri; ²Division of Cardiology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon; ³Division of Cardiology, Incheon Sejong Hospital, Incheon; ⁴Division of Cardiology, Department of Internal Medicine, Seoul St. Mary's Hospital, Catholic Research Institute for Intractable Cardiovascular Disease, College of Medicine, The Catholic University of Korea, Seoul; ⁵Division of Cardiology, Department of Internal Medicine, Korea University Anam Hospital, Seoul; ⁶Division of Cardiology, Department of Internal Medicine, Chungbuk National University Hospital, Chungbuk National University College of Medicine, Cheongju; ⁷K-Heart Clinic & Lab, Incheon, Korea

Metabolic syndrome awareness in the general Korean population: results from a nationwide survey



Conclusion

The high-awareness group showed greater knowledge of the importance of lifestyle modifications and better health behaviors regarding MetS. The findings highlight the need for improved public education and awareness programs regarding MetS.

Background/Aims: Metabolic syndrome (MetS) raises the risk of cardiovascular disease and type 2 diabetes. An awareness of MetS is vital for early detection and proactive management, which can mitigate the risks associated with MetS. Therefore, our study aimed to investigate the level of awareness of MetS among the Korean population.

Methods: We conducted a nationwide survey between January and February 2023 among a representative sample of the Korean population using an online survey. Information regarding the awareness of MetS and its risk, the importance of lifestyle modification, and health behavior were collected. The question about the awareness of MetS was "How much do you think you know about MetS?" and there were five answers: 1) I know very well, 2) I know well, 3) I know a little, 4) I do not

know, and 5) I have no idea. The high-awareness group was defined as those who answered that they knew very well or well.

Results: Among 1,000 participants (mean age, 45.7 ± 13.2 yr), 29% were unaware of MetS, and only 20.8% had high awareness. The high-awareness group was significantly more knowledgeable about lifestyle modifications and demonstrated better health behaviors. After adjustment for possible confounding factors, younger age, low household income, and absence of comorbidity were independently associated with a lack of awareness regarding MetS.

Conclusions: The high-awareness group showed greater knowledge of the importance of lifestyle modifications and better health behaviors regarding MetS. The findings highlight the need for improved public education and awareness programs regarding MetS.

Keywords: Metabolic syndrome; Awareness; Health behavior; Cardiovascular disease

INTRODUCTION

Metabolic syndrome is recognized globally as a significant public health concern. It is characterized by a cluster of interconnected metabolic dysfunctions, including insulin resistance, hypertension, and dyslipidemia, that collectively increase the risk of cardiovascular disease (CVD) and type 2 diabetes [1,2]. Notably, its prevalence is increasing globally. In South Korea, this escalation in the prevalence rate of metabolic syndrome is attributed to rapid urbanization, an aging population, and lifestyle changes [3,4]. Despite this increasing prevalence, research by Scuteri et al. (2015) and others has revealed a general lack of public awareness of this syndrome [5]. This paucity of knowledge can have severe implications on health outcomes, as understanding the condition is crucial for encouraging individuals to engage in preventive measures such as dietary modifications and regular exercise [6]. Awareness is the initial step towards promoting behavioral changes and preventive actions, and risk-informed individuals are more likely to make lifestyle modifications to reduce their risk of developing metabolic syndrome [7]. Increased awareness can further facilitate the early detection and management of the syndrome, thereby reducing the potential for developing severe complications such as CVD and type 2 diabetes [8].

Despite this alarming situation in Korea, empirical evidence on the population's understanding of metabolic syndrome remains limited [4]. To address this gap, our study aimed to investigate the level of awareness about metabolic syndrome among the Korean population.

METHODS

Study design and data collection

This cross-sectional online survey was conducted between January and February 2023. Excluding the study design, the survey process was outsourced to MegaResearch (Seoul, Korea). The estimated target population (aged ≥ 20 yr) in December 2022 was 37,121,412. To represent Korean adults aged between 20 and 69 years, the survey population ($n = 1,000$) was selected using a stratified systematic sampling method, considering gender (men/women), age (20–29, 30–39, 40–49, 50–59, and 60–69 yr), and place of residence (eight metropolitan cities and eight non-metropolitan cities). The study was conducted through online interviews by sending web survey page links to randomly selected individuals from a panel of members of a subscription-based professional survey company. The respondents answered the web questionnaire using a mouse and keyboard, and the entered responses were automatically collected and managed on a data server. The web questionnaire was written using web programming languages that closely resemble offline survey procedures. Non-responses were systematically eliminated by the web page, which was programmed to not allow respondents to proceed to the subsequent page if a response was missed. Survey execution and management were conducted by the research company, and progress was monitored in real-time during the online survey.

Study population

The study population was derived from the target population, estimated by stratification based on gender, age, and

place of residence. Eligible participants were aged between 20 and 69 years, agreed to participate in the survey, and answered the questions after understanding them. Responses completed up to the last question were considered valid. We excluded the following participants: 1) those aged < 20 years or ≥ 70 years, 2) those with significantly low online accessibility, 3) those who did not agree to participate in the survey, 4) those who failed to respond to the last question, and 5) those of a specific gender and age group that exceeded the assigned survey ratio during the survey.

Questionnaire

Participant eligibility was determined on the basis of age, gender, and place of residence, which were the initial questions asked in the survey (Supplementary Material). The main body of the survey was divided into three sections:

1) The first section gathered demographic data, such as educational attainment; income; type of residence; marital status; and health status, including the presence of hypertension, dyslipidemia, diabetes, central obesity, CVD, cerebrovascular disease, social history of alcohol consumption and smoking habits, and family history of CVD.

2) The second section assessed participants' awareness of metabolic syndrome. Participants were asked, "How much do you think you know about metabolic syndrome?" with five potential responses: (1) I know very well, (2) I know well, (3) I know a little, (4) I do not know, and (5) I have no idea. These responses were based on subjective judgments of the participants.

3) The third and final section explored understanding the importance of lifestyle modifications and health behaviors in relation to metabolic syndrome. For each lifestyle modification category (calorie intake control, exercise, weight reduction, smoking cessation, and medical treatment [i.e., the need for medical consultation or intervention]), participants rated the importance on a 5-point scale from "very necessary" to "not necessary at all." For health behavior within the last three years, participants measured health parameters, including body weight, waist circumference, and blood pressure. Additionally, details on specific lifestyle modifications, such as regularly measuring blood pressure, getting adequate sleep, following a low-salt diet, and exercising regularly, were gathered.

Participants who responded that they knew about metabolic syndrome "very well" or "well" were categorized as the high awareness group, while those who responded with

"I know a little", "I do not know", or "I have no idea" were classified as the low awareness group.

Ethics

Personal information was not collected during the online

Table 1. Characteristics of the study population

Characteristic	Value (n = 1,000)
Age (yr)	45.7 ± 13.2
Men	509 (50.9)
Education attainment	
Middle school or less	9 (0.9)
High school	237 (23.7)
College or more	754 (75.4)
Household income (KRW 1,000/mo)	
≤ 1,000	45 (4.5)
1,000–2,000	80 (8.0)
2,000–3,000	187 (18.7)
3,000–4,000	209 (20.9)
4,000–5,000	164 (16.4)
> 5,000	315 (31.5)
Presence of underlying disease	
Hypertension	287 (28.7)
Dyslipidemia	289 (28.9)
Diabetes	122 (12.2)
Cardiovascular disease	44 (4.4)
Cerebrovascular disease	32 (3.2)
Central obesity	412 (41.2)
Current smoker	282 (28.2)
Socially alcohol drinker	231 (23.1)
Family history of cardiovascular disease	205 (20.5)
Marital status	
Single	340 (34.0)
Married	609 (60.9)
Divorced/widowed	49 (4.9)
Other	2 (0.2)
Type of residence	
Metropolitan area ^{a)}	446 (44.6)
Non-metropolitan area ^{b)}	554 (55.4)

Values are presented as mean ± standard deviation or number (%).

^{a)}Eight major metropolitan cities including Seoul, Busan, Dae-gu, Incheon, Gwangju, Daejeon, Ulsan, and Sejong.

^{b)}Regions beyond the eight major metropolitan cities including Gyeonggi, Gangwon-do, Chungbuk, Chungnam, Jeonbuk, Jeonnam, Gyeongbuk, Gyeongnam, and Jeju.

survey. The local Institutional Review Board of Gachon University Medical Center approved this study (GFIRB2023-084) and waived the requirement for written informed consent.

Statistical analysis

All categorical data are presented as frequencies and percentages. Pearson’s chi-squared test was used to compare categorical variables. A linear-by-linear association was employed to discern trends in clinical characteristics based on the responses. Additionally, a univariate analysis was conducted, followed by a multivariate logistic regression analysis to evaluate the risk of lack of awareness of metabolic

syndrome while accounting for individual risk factors. Variables that showed a significant predictive value ($p < 0.05$) in univariate analysis were subsequently incorporated into the regression analysis. The threshold for statistical significance was set at $p < 0.05$. All statistical analyses were performed using the SPSS software (version 22.0; IBM Corp., Armonk, NY, USA).

RESULTS

Characteristics of the study population

A total of 1,000 selected participants completed an online interview based on systematic stratification sampling. The study participants were evenly distributed according to age, gender, and place of residence. Table 1 presents the characteristics of the study population. The mean age of the study population was 45.7 ± 13.2 years (range, 20–69 yr), and approximately half of the participants were men (50.9%). The majority of participants had attained a college education or higher (75.4%), with a smaller percentage having a high school education (23.7%), and less than one percent (0.9%) having middle school education or less. In terms of income, 31.5% had a household income of more than 5,000 (KRW 1,000/mo), while 4.5% had an income of 1,000 (KRW 1,000/mo) or less. The remaining individuals fell within these income brackets. Regarding underlying health conditions, the most common underlying disease was central obesity (41.2%), followed by dyslipidemia (28.9%), hypertension (28.7%), and diabetes (12.2%); CVD and cerebrovascular disease were present in 4.4% and 3.2% of




Figure 1. The extent to which participants know about metabolic syndrome. Question: How much do you think you know about metabolic syndrome?




Figure 2. Awareness of metabolic syndrome and the awareness pathways. Question: Have you ever heard of metabolic syndrome?

participants, respectively. Of the participants, 28.2% were current smokers and 23.1% were social alcohol drinkers. Regarding marital status, 60.9% of the participants were married, 34.0% were single, and 4.9% were divorced or widowed.

Awareness regarding metabolic syndrome

Approximately 80% of the participants listed “I do not know,” “I have no idea” or “I know a little” about metabolic syndrome (Fig. 1). A mere 4.2% of participants, which corresponds to 42 individuals, answered that “I know very

Table 2. Metabolic syndrome awareness in the general population

Characteristic	Total	Answer					p value
		I know very well	I know well	I know a little	I do not know	I have no idea	
Total population		42 (4.2)	166 (16.6)	502 (50.2)	234 (23.4)	56 (5.6)	
Men		37 (3.7)	169 (16.9)	499 (49.9)	236 (23.6)	59 (5.9)	0.940
Age (yr)							< 0.001
20–29	175	8 (19.0)	23 (13.9)	71 (14.1)	53 (22.6)	20 (35.7)	
30–39	175	7 (16.7)	25 (15.1)	80 (15.9)	51 (21.8)	12 (21.4)	
40–49	216	6 (14.3)	29 (17.5)	111 (22.1)	60 (25.6)	10 (17.9)	
50–59	234	9 (21.4)	49 (29.5)	124 (24.7)	45 (19.2)	7 (12.5)	
60–69	200	12 (28.6)	40 (24.1)	116 (23.1)	25 (10.7)	7 (12.5)	
Education attainment							0.007
Middle school or less	9	0 (0.0)	0 (0.0)	7 (1.4)	2 (0.9)	0 (0.0)	
High school	237	8 (19.0)	31 (18.7)	117 (23.3)	60 (25.6)	21 (37.5)	
College or more	754	34 (81.0)	135 (81.3)	378 (75.3)	172 (73.5)	35 (62.5)	
Household income (KRW 1,000/mo)							< 0.001
≤ 1,000	45	1 (2.4)	3 (1.8)	18 (3.6)	16 (6.8)	7 (12.5)	
1,000–2,000	80	2 (4.8)	13 (7.8)	3 (6.6)	26 (11.1)	6 (10.7)	
2,000–3,000	187	7 (16.7)	29 (17.5)	92 (18.3)	48 (20.5)	11 (19.6)	
3,000–4,000	209	7 (16.7)	36 (21.7)	99 (19.7)	53 (22.6)	14 (25.0)	
4,000–5,000	164	7 (16.7)	20 (12.0)	93 (18.5)	36 (15.4)	8 (14.3)	
> 5,000	315	18 (42.9)	65 (39.2)	167 (33.3)	55 (23.5)	10 (17.9)	
Presence of underlying disease ^{a)}							<0.001
Yes	465	29 (69.0)	106 (63.9)	229 (45.6)	86 (36.8)	15 (26.8)	
No	535	132 (31.0)	60 (36.1)	273 (54.4)	148 (63.2)	41 (73.2)	
Marital status							<0.001
Single	340	9 (21.4)	41 (24.7)	153 (30.5)	105 (44.9)	32 (57.1)	
Married	609	30 (71.4)	115 (69.3)	320 (63.7)	122 (52.1)	22 (39.3)	
Divorced/widowed	49	3 (7.1)	10 (6.0)	27 (5.4)	7 (3.0)	2 (3.6)	
Other	2	0 (0.0)	0 (0.0)	2 (0.4)	0 (0.0)	0 (0.0)	
Type of residence							0.054
Metropolitan area ^{b)}		20 (47.6)	86 (51.8)	221 (44.0)	96 (41.0)	23 (41.1)	
Non-metropolitan area ^{c)}		22 (52.4)	80 (48.2)	281 (56.0)	138 (59.0)	33 (58.9)	

Values are presented as number only or number (%).

^{a)}Presence of hypertension, dyslipidemia, diabetes, cardiovascular disease, or cerebrovascular disease.

^{b)}Eight major metropolitan cities including Seoul, Busan, Daegu, Incheon, Gwangju, Daejeon, Ulsan, and Sejong.

^{c)}Regions beyond the eight major metropolitan cities including Gyeonggi, Gangwon-do, Chungbuk, Chungnam, Jeonbuk, Jeonnam, Gyeongbuk, Gyeongnam, and Jeju.

well.” Further, an additional 16.6% of the population, equivalent to 166 participants, reported that “I know well.” Three-quarters (78.8%) of the participants answered that they had heard of metabolic syndrome in the previous year (Fig. 2). The most common sources of information were television and radio (40.4%) followed by the internet or YouTube (28.9%); other sources included friends, relatives, hospital care staff, newspapers, and magazines.

Notably, awareness of metabolic syndrome varied significantly across different demographic parameters, such as in age group, education level, household income, presence of underlying disease, and marital status (Table 2). Among the different age groups, the highest awareness levels were observed in individuals aged 60–69 years and 50–59 years (percentage of those who answered “I know very well” or “I know well”), while the youngest age group (20–29 yr)

Table 3. Differences in awareness of the importance of lifestyle modifications for metabolic syndrome in the general population according to awareness

Lifestyle modification	All (n =1000)	High awareness group (n = 208)	Low awareness group (n = 792)	<i>p</i> value
Calorie intake control				< 0.001
Very necessary	363 (36.3)	123 (59.1)	240 (30.3)	
Somewhat necessary	306 (30.6)	62 (29.8)	244 (30.8)	
Slightly necessary	178 (17.8)	16 (7.7)	162 (20.5)	
Not sure	145 (14.5)	7 (3.4)	138 (17.4)	
Not necessary at all	8 (0.8)	0 (0.0)	8 (1.04)	
Exercise				< 0.001
Very necessary	497 (49.7)	144 (69.2)	353 (44.6)	
Somewhat necessary	265 (26.5)	49 (23.6)	216 (27.3)	
Slightly necessary	137 (13.7)	13 (6.3)	124 (15.7)	
Not sure	99 (9.9)	2 (1.0)	97 (12.2)	
Not necessary at all	2 (0.2)	0 (0.0)	2 (0.3)	
Weight reduction				< 0.001
Very necessary	535 (53.5)	132 (63.5)	403 (50.9)	
Somewhat necessary	252 (25.2)	51 (24.5)	201 (25.4)	
Slightly necessary	142 (14.2)	19 (9.1)	123 (15.5)	
Not sure	66 (6.6)	6 (2.9)	60 (7.6)	
Not necessary at all	5 (0.5)	0 (0.0)	5 (0.6)	
Smoking cessation				< 0.001
Very necessary	479 (47.9)	120 (57.7)	359 (45.3)	
Somewhat necessary	261 (26.1)	62 (29.8)	199 (25.1)	
Slightly necessary	136 (13.6)	15 (7.2)	121 (15.3)	
Not sure	117 (11.7)	9 (4.3)	108 (13.6)	
Not necessary at all	7 (0.7)	2 (1.0)	5 (0.6)	
Medical treatment				< 0.001
Very necessary	339 (33.9)	105 (50.5)	234 (29.5)	
Somewhat necessary	387 (38.7)	81 (38.9)	306 (38.6)	
Slightly necessary	167 (16.7)	15 (7.2)	152 (19.2)	
Not sure	104 (10.4)	7 (3.4)	97 (12.2)	
Not necessary at all	3 (0.3)	0 (0.0)	3 (0.4)	

Values are presented as number (%).

showed the lowest awareness levels (percentage of those who answered “I have no idea” or “I do not know”). Additionally, awareness increased with an increase in the level of education and household income. Participants with comorbid conditions demonstrated a higher level of awareness of metabolic syndrome. Married individuals exhibited a higher level of awareness, whereas those living alone exhibited a lower level. Additionally, respondents with a high level of metabolic syndrome awareness tended to live in the eight major metropolitan cities.

Understanding the importance of lifestyle modifications and health behaviors in metabolic syndrome

Table 3 and Supplementary Figure 1 present differences in the awareness of the general population regarding the importance of lifestyle modifications in relation to metabolic syndrome. Participants were divided into two groups accord-

ing to their level of awareness: a high-awareness group (n = 208) and a low-awareness group (n = 792). Significantly different distributions between the high- and low-awareness groups were observed in each category, as indicated by *p* values < 0.001. For calorie intake control, 59.1% of the high awareness group viewed it as “very necessary,” compared to only 30.3% in the low awareness group. Similar trends were observed in the exercise, weight reduction, smoking cessation, and medical treatment categories, with higher percentages of the high awareness group rating these interventions as “very necessary.” Interestingly, the highest levels of “very necessary” responses across both groups were observed for weight reduction, with 63.5% of the high awareness group and 50.9% of the low awareness group agreeing. Even in the high awareness group, only 50.5% perceived medical treatment as “very necessary,” highlighting a lack of understanding of the importance of medical treatment among individuals with metabolic syndrome.

Table 4. Differences in health behaviors about metabolic syndrome in the general population according to awareness

	All (n = 1,000)	High awareness group (n = 208)	Low awareness group (n = 792)	<i>p</i> value
Body weight measurement in the last 3 years				< 0.001
Often	391 (39.1)	104 (50.0)	287 (36.2)	
Sometimes	369 (36.9)	76 (36.5)	293 (37.0)	
Rarely	197 (19.7)	24 (11.5)	173 (21.8)	
Never	43 (4.3)	4 (1.9)	39 (4.9)	
Waist circumference measurement in the last 3 years				< 0.001
Often	67 (6.7)	34 (16.3)	33 (4.2)	
Sometimes	362 (36.2)	108 (51.9)	254 (32.1)	
Rarely	363 (36.3)	51 (24.5)	193 (24.4)	
Never	208 (20.8)	15 (7.2)	193 (24.4)	
Blood pressure measurement in the last 3 years				< 0.001
Often	200 (20.0)	75 (36.1)	125 (15.8)	
Sometimes	437 (43.7)	96 (46.2)	341 (43.1)	
Rarely	298 (29.8)	32 (15.4)	266 (33.6)	
Never	65 (6.5)	5 (2.4)	60 (7.6)	
Lifestyle modification				
Measure blood pressure regularly	339 (33.9)	94 (45.2)	245 (30.9)	< 0.001
Have deep sleep	335 (33.5)	70 (33.7)	265 (33.5)	0.958
Low-salt diet	331 (33.1)	90 (43.3)	241 (30.4)	< 0.001
Regular exercise	538 (53.8)	127 (61.1)	411 (51.9)	0.018

Values are presented as number (%).

Table 4 and Supplementary Figure 2 detail differences in health behaviors of the general population related to metabolic syndrome between the two groups. In the high awareness group, a higher proportion of individuals “often” measured their body weight (50.0%) compared to the low awareness group (36.2%). Similarly, the high awareness group more frequently measured their waist circumference and blood pressure, with 16.3% and 36.1% doing so “often” compared to 4.2% and 15.8% in the low awareness group, respectively. These differences are statistically significant, as indicated by p values < 0.001 . In terms of lifestyle modifications, more participants in the high awareness group regularly measured blood pressure (45.2% vs. 30.9%), followed a low-salt diet (43.3% vs. 30.4%), and exercised regularly (61.1% vs. 51.9%) compared to the low awareness group. These differences were also statistically significant ($p < 0.05$), excluding the practice of ensuring a restful sleep, where no significant difference was observed between the two groups ($p = 0.958$).

Risk factors for lack of awareness of metabolic syndrome

The univariate analysis in Table 5 showed that the following variables were associated with a lack of awareness about metabolic syndrome (“I do not know” and “I have no idea”) in the general population: increasing age (from 20s to 60s), a below college level of education, a household income $< 2,000$ (in 1,000 KRW/mo), absence of comorbidity, and single-person households.

After adjusting for possible confounding factors in the multivariate analysis, each decade increase in age from the

20s to the 60s was associated with a 25% decreased hazard (odds ratio of 0.75) for lack of awareness of metabolic syndrome. This indicates that awareness increases with age. Lower household income ($< 2,000$ in 1,000 KRW/mo) was associated with a 2.13-fold increased risk of lack of awareness, suggesting that lower income significantly contributes to decreased awareness. The absence of comorbidities was also independently associated with a lack of awareness, increasing the risk by 56% (odds ratio, 1.56). However, having a below college level of education and single-person households had no statistically significant associations toward lower awareness.

DISCUSSION

Our recent cross-sectional web-based survey offers pivotal findings regarding the awareness level of metabolic syndrome within the adult Korean population, specifically among those aged 20–69 years. Although a stratified systematic sampling approach was employed to ensure a wide representation of the demographic groups, the data indicated a substantial awareness gap regarding metabolic syndrome among the respondents. Additionally, this study underscored the correlation between level of awareness and participant recognition of the importance of lifestyle alterations to enhance health. Individuals with higher awareness levels demonstrated superior comprehension and involvement in behaviors that promote better health. Interestingly, specific risk factors such as younger age, lower household income, and lack of comorbid conditions were found to be

Table 5. Factors associated with lack of awareness (I do not know or I have no idea) about metabolic syndrome in the general population

Factor	Univariate		Multivariate	
	OR (95% CI)	p value	OR (95% CI)	p value
Age (every decade from the 20s to the 60s)	0.72 (0.652–0.801)	< 0.001	0.75 (0.661–0.854)	< 0.001
Educational attainment $<$ college ^{a)}	1.35 (0.988–1.833)	0.060	1.29 (0.928–1.803)	0.129
Household income $<$ 2,000 (1,000 KRW/mo) ^{b)}	2.14 (1.458–3.140)	< 0.001	2.13 (1.388–3.275)	0.001
Absence of comorbidity ^{c)}	1.97 (1.484–2.612)	< 0.001	1.56 (1.149–2.115)	0.004
Single person house hold ^{d)}	1.95 (1.476–2.572)	< 0.001	1.02 (0.716–1.447)	0.922

CI, confidence interval; OR, odds ratio.

^{a)}Educational attainment $<$ college vs. college or higher educational attainment.

^{b)}Household income $<$ 2,000 (1,000 KRW/mo) vs. household income \geq 2,000 (1,000 KRW/mo).

^{c)}Including hypertension, dyslipidemia, diabetes, cardiovascular disease, or cerebrovascular disease.

^{d)}Married vs. single person house hold (single, or divorced/widowed).

associated with reduced awareness. This knowledge could help shape targeted educational strategies to boost awareness and understanding of metabolic syndrome.

Metabolic syndrome, which includes a cluster of interconnected factors, such as insulin resistance, hypertension, and dyslipidemia, is recognized as a major worldwide health concern [9,10]. Globally, the prevalence of metabolic syndrome is increasing, with significant increases occurring in certain regions and specific ethnic groups. In the US, the rate of metabolic syndrome is escalating faster in Asian and Hispanic adults than in other racial groups [11]. In the Asia-Pacific region, the prevalence of metabolic syndrome varies from 11.9 to 49.0% in the adult population, with a notable increase in East Asia [12]. Notably, the metabolic dysfunction associated with metabolic syndrome poses a significant risk of developing CVD [13]. Moreover, each additional metabolic syndrome component has been found to increase the risk of a significant cardiovascular events by 22% [14].

Building on this global perspective, understanding regional differences and the impact of lifestyle and socioeconomic changes on the prevalence of metabolic syndrome is of paramount importance. A range of factors, including obesity, diet, physical activity, and socioeconomic influences, contribute to the development of metabolic syndrome [1]. Countries experiencing swift economic changes often see drastic lifestyle changes and subsequent shifts in the incidence and prevalence of chronic diseases [15]. Economic growth was paralleled by a significant nutritional transition, as consumption of processed foods and animal products increased, intake of plant foods decreased, and a shift occurred in the primary cause of death from infectious to CVDs [16]. Although previous studies have investigated the progression of metabolic syndrome in South Korea, certain aspects remain unaddressed. The prevalence of the syndrome was found to have increased from 24.9% in 1998 to 30.52% in 2013, but alterations in lifestyle and socioeconomic factors were not considered, and the time period investigated was relatively short [17,18]. In contrast, a more recent study examined changes in the prevalence of metabolic syndrome from 2001 to 2020, accounting for alterations in the components of metabolic syndrome and related lifestyle factors, using data from the Korean National Health and Nutrition Examination Survey (KNHANES) [19].

In our study, as we delved into socioeconomic influences and changing lifestyle patterns, it became clear that enhanced awareness and recognition of metabolic syndrome

are crucial. The increasing global prevalence of metabolic syndrome underscores the importance of recognizing its severity and the need for effective management and prevention. Despite South Korea's status as a developed country, there remains an urgent need to enhance awareness of this prevalent condition [20]. The situation in South Korea is pressing, with studies revealing a marked lack of awareness of metabolic syndrome and its connection with lifestyle and socioeconomic factors [17]. Cultural influences can significantly affect awareness, as established in previous research, thereby stressing the need for tailored educational initiatives [21]. In China, a considerable number of adults remain unaware of their metabolic risk factors [22], further emphasizing the need for easily understandable measures like the "A Body Shape Index" and the "Waist to Height Ratio" [23]. Moreover, studies have highlighted that certain lifestyle risk factors, such as with sedentary and low activity jobs, pose significant risks for developing total and central obesity, particularly in middle-aged male workers [24]. This knowledge substantiates the importance of raising awareness of these factors as part of an overall strategy to combat metabolic syndrome. Our study illustrates that awareness directly influences an individual's commitment to lifestyle modifications to manage metabolic risks. Those with higher awareness perceived these changes as more vital and engaged more frequently in health-promoting behaviors than their lower awareness counterparts.

Having discussed the role of awareness in managing metabolic syndrome, our study examined the specific factors contributing to this lack of awareness. Remarkably, our study found an association between young age, low household income, absence of comorbidities, and lack of awareness of metabolic syndrome. There are several possible reasons for this observation. Young individuals may perceive themselves as invincible to health conditions such as metabolic syndrome, which is often associated with older age groups. Consequently, they may not actively seek information about health risks. People with lower incomes may face more pressing immediate needs or lack access to resources that facilitate health education, thereby limiting their exposure to vital health information. Finally, those without preexisting health conditions might not interact as frequently with the healthcare system, which is a significant source of health risk education. These findings underline the need to target health education initiatives towards at-risk demographics in order to combat the rising prevalence of

metabolic syndrome.

The present study has certain limitations. First, there may be some selection bias. Health conscious individuals with an inherent interest in wellness, or with a higher monthly household income may be more inclined to participate in our survey. Their knowledge about metabolic syndrome, and better access to health information due to their socioeconomic status might have influenced their responses, resulting in an overestimation of the overall awareness level in the general population. Therefore, the awareness of metabolic syndrome could be lower than that seen in our results. Second, the generalizability of our findings is limited because of the geographic scope of our study, which was restricted to Korean adults aged 20–69 years. Therefore, the results may not be applicable to other ethnicities or age groups. Third, the evaluation of metabolic syndrome awareness was based on subjective responses rather than on a standardized scoring system. This subjective approach might have limited accuracy in assessing the participants' actual knowledge of metabolic syndrome. However, future studies might benefit from incorporating more objective measures to further validate self-reported awareness levels. Despite these potential limitations, our study aimed to bridge the gap between subjective self-perception and actual knowledge of metabolic syndrome in the Korean population. The results of awareness, as self-reported by the participants, can be considered meaningful in shedding light on this crucial public health issue.

Our study offers valuable insights into awareness of metabolic syndrome and its influencing factors in the Korean adult population. This study unveils a substantial gap in the awareness of metabolic syndrome among Korean adults, despite the varied demographic representation ensured by stratified systematic sampling. Our findings underscore the crucial role of awareness in recognizing the importance of lifestyle modifications and actively engaging in health-promoting behaviors. Furthermore, the identification of certain risk factors, including younger age, lower household income, and absence of comorbid conditions, contributes to the deficit in awareness and provides a roadmap for designing targeted interventions. As we move forward, it is imperative to deploy concerted efforts towards comprehensive educational strategies that are personalized according to these risk factors to bolster the understanding and management of metabolic syndrome. Such an approach will not only help individuals become active participants in their own

health management, but also enable us to navigate towards a healthier society.

In conclusion, it is clear that our study highlights a significant awareness gap regarding metabolic syndrome among Korean adults, revealing that age, income, and absence of comorbid conditions can affect this awareness. These findings emphasize the need for personalized educational strategies, promotion of better self-management of health, and steering society towards improving overall health.

KEY MESSAGE

1. There was a significant gap in the awareness of metabolic syndrome among Korean adults.
2. The high-awareness group showed more knowledge of the importance of lifestyle modifications and better health behaviors regarding metabolic syndrome.
3. Certain risk factors, including younger age, lower household income, and absence of comorbid conditions, contribute to the deficit in awareness of metabolic syndrome.
4. The findings highlight the need for improved public education and awareness programs regarding metabolic syndrome.

REFERENCES

1. Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640-1645.
2. Jang YN, Lee JH, Moon JS, et al. Metabolic syndrome severity score for predicting cardiovascular events: a nationwide population-based study from Korea. *Diabetes Metab J* 2021;45: 569-577.
3. Huh JH, Kang DR, Kim JY, Koh KK. Metabolic syndrome fact sheet 2021: executive report. *Cardiometab Syndr J* 2021;1: 125-134.
4. Ko SH, Kim DJ, Park JH, et al.; Task Force Team for Diabetes Fact Sheet of the Korean Diabetes Association. Trends of anti-diabetic drug use in adult type 2 diabetes in Korea in 2002-

- 2013: nationwide population-based cohort study. *Medicine (Baltimore)* 2016;95:e4018.
5. Scuteri A, Laurent S, Cucca F, et al.; Metabolic Syndrome and Arteries Research (MARE) Consortium. Metabolic syndrome across Europe: different clusters of risk factors. *Eur J Prev Cardiol* 2015;22:486-491.
 6. Jahangiry L, Farhangy MA, Rezaei F. Framingham risk score for estimation of 10-years of cardiovascular diseases risk in patients with metabolic syndrome. *J Health Popul Nutr* 2017;36:36.
 7. Moore SE, Harris C, Wimberly Y. Perception of weight and threat to health. *J Natl Med Assoc* 2010;102:119-124.
 8. Saklayen MG. The Global Epidemic of the Metabolic Syndrome. *Curr Hypertens Rep* 2018;20:12.
 9. Lobene AJ. The increasing prevalence of metabolic syndrome in Korea: a multifarious disease with a multifactorial etiology. *JACC Asia* 2023;3:503-505.
 10. Lee SH, Park SY, Choi CS. Insulin resistance: from mechanisms to therapeutic strategies. *Diabetes Metab J* 2022;46:15-37.
 11. Hirode G, Wong RJ. Trends in the prevalence of metabolic syndrome in the United States, 2011-2016. *JAMA* 2020;323:2526-2528.
 12. Ranasinghe P, Mathangasinghe Y, Jayawardena R, Hills AP, Misra A. Prevalence and trends of metabolic syndrome among adults in the asia-pacific region: a systematic review. *BMC Public Health* 2017;17:101.
 13. Isomaa B, Almgren P, Tuomi T, et al. Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care* 2001;24:683-689.
 14. Guembe MJ, Fernandez-Lazaro CI, Sayon-Orea C, Toledo E, Moreno-Iribas C; RIVANA Study Investigators. Risk for cardiovascular disease associated with metabolic syndrome and its components: a 13-year prospective study in the RIVANA cohort. *Cardiovasc Diabetol* 2020;19:195.
 15. Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* 2012;70:3-21.
 16. Kim S, Moon S, Popkin BM. The nutrition transition in South Korea. *Am J Clin Nutr* 2000;71:44-53.
 17. Lim S, Shin H, Song JH, et al. Increasing prevalence of metabolic syndrome in Korea: the Korean National Health and Nutrition Examination Survey for 1998-2007. *Diabetes Care* 2011;34:1323-1328.
 18. Lee SE, Han K, Kang YM, et al.; Taskforce Team of Diabetes Fact Sheet of the Korean Diabetes Association. Trends in the prevalence of metabolic syndrome and its components in South Korea: Findings from the Korean National Health Insurance Service Database (2009-2013). *PLoS One* 2018;13:e0194490.
 19. Park D, Shin MJ, Després JP, Eckel RH, Tuomilehto J, Lim S. 20-year trends in metabolic syndrome among Korean adults from 2001 to 2020. *JACC Asia* 2023;3:491-502.
 20. Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 2002;287:356-359.
 21. Aekplakorn W, Abbott-Klafter J, Premgamone A, et al. Prevalence and management of diabetes and associated risk factors by regions of Thailand: third National Health Examination Survey 2004. *Diabetes Care* 2007;30:2007-2012.
 22. Yu S, Guo X, Yang H, Zheng L, Sun Y. An update on the prevalence of metabolic syndrome and its associated factors in rural northeast China. *BMC Public Health* 2014;14:877.s
 23. Krakauer NY, Krakauer JC. A new body shape index predicts mortality hazard independently of body mass index. *PLoS One* 2012;7:e39504.
 24. Choi B, Schnall PL, Yang H, et al. Sedentary work, low physical job demand, and obesity in US workers. *Am J Ind Med* 2010;53:1088-1101.

Received : August 27, 2023

Revised : October 3, 2023

Accepted : October 5, 2023

Correspondence to

Mi-Seung Shin, M.D., Ph.D.

Division of Cardiology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, 21 Namdong-daero 774beon-gil, Namdong-gu, Incheon 21565, Korea

Tel: +82-32-460-3663, Fax: +82-32-469-1906

E-mail: msshin@gilhospital.com

<https://orcid.org/0000-0002-0273-0109>

Acknowledgments

The authors thank the Korean Society of CardioMetabolic Syndrome for providing the funds for this online survey.

CRedit authorship contributions

Hyun-Jin Kim: methodology, data curation, writing - original draft; Mi-Seung Shin: conceptualization, methodology, writing - review & editing, supervision, project administration; Kyung-Hee Kim: methodology; Mi-Hyang Jung: methodology; Dong-Hyuk Cho: methodology; Ju-Hee Lee: methodology; Kwang Kon Koh: writing - review & editing, supervision

Conflicts of interest

The authors disclose no conflicts.

Funding

None