



CERTAIN NEW RESULTS ON RATIONAL FUNCTIONS WITH PRESCRIBED POLES

R. Mohammad¹, Mridula Purohit² and Ab. Liman³

¹Department of Mathematics, Vivekananda Global University, India
e-mail: shafatruqia@gmail.com

²Department of Mathematics, Vivekananda Global University, India
e-mail: mirdula_purohit@vgu.ac.in

³Department of Mathematics, National Institute of Technology Srinagar-190006, India
e-mail: abliman22@yahoo.com

Abstract. Let R_n be the space of rational functions with prescribed poles. If $r \in R_n$, does not vanish in $|z| < k$, then for $k = 1$

$$|r'(z)| \leq \frac{|B'(z)|}{2} \sup_{z \in T} |r(z)|,$$

where $B(z)$ is the Blaschke product.

In this paper, we consider a more general class of rational functions $rof \in R_{m^*n}$, defined by

$$(rof)(z) = r(f(z)),$$

where $f(z)$ is a polynomial of degree m^* and prove a more general result of the above inequality for $k > 1$.

We also prove that

$$\sup_{z \in T} \left[\left| \frac{r^{*'}(f(z))}{B'(z)} \right| + \left| \frac{r'(f(z))}{B'(z)} \right| \right] = \sup_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|,$$

and as a consequence of this result, we present a generalization of a theorem of O'Hara and Rodriguez for self-inverse polynomials. Finally, we establish a similar result when supremum is replaced by infimum for a rational function which has all its zeros in the unit circle.

⁰Received February 6, 2023. Revised July 7, 2023. Accepted October 3, 2023.

⁰2020 Mathematics Subject Classification: 30A10, 30C15, 30D15.

⁰Keywords: Rational functions, polynomials, inequalities, poles, zeros.

⁰Corresponding author: R. Mohammad(shafatruqia@gmail.com).

1. INTRODUCTION

Let \mathbb{P}_n be the space of complex polynomials of degree at most n and \mathbb{C} be the complex plane. Let $T = \{z \in \mathbb{C} : |z| = 1\}$ and D^- denotes the region inside of T and D^+ denotes the region outside of T .

Let $p \in \mathbb{P}_n$. Then concerning the estimate of p' on T , we have by a famous result due to Bernstein [4]:

$$\sup_{z \in T} |p'(z)| \leq n \sup_{z \in T} |p(z)|. \quad (1.1)$$

This result is sharp and equality holds for the polynomials having all zeros at origin.

As a refinement of result of Bernstein [4], we mention the following result due to Aziz [2];

Theorem 1.1. ([2]) *If $p \in P_n$ and $p^*(z) = z^n \overline{p(\frac{1}{\bar{z}})}$, then*

$$\sup_{z \in T} [|(p^*(z))'| + |p(z)|] = n \sup_{z \in T} |p(z)|. \quad (1.2)$$

The next result was conjectured by Erdős and later verified by Lax [6]: If $p \in \mathbb{P}_n$, and all the zeros of $p(z)$ lie in TUD^+ , then for $z \in T$, we have

$$\sup_{z \in T} |p'(z)| \leq \frac{n}{2} \sup_{z \in T} |p(z)|.$$

Equality holds for $p(z) = \lambda z^n + \mu$, $|\lambda| = |\mu| = 1$.

Let a_1, a_2, \dots, a_n be n given points in D^+ . We consider the following space of rational functions R_n with prescribed poles:

$$R_n := R_n(a_1, a_2, \dots, a_n) = \left[\frac{p(z)}{w(z)} : p \in P_n \right],$$

where

$$w(z) = (z - a_1)(z - a_2) \dots (z - a_n).$$

Let

$$B(z) := \frac{z^n \overline{w(1/\bar{z})}}{w(z)} = \prod_{k=1}^n \frac{1 - \bar{a}_k z}{z - a_k},$$

where $B(z) \in R_n$ is called Blaschke product. Note that $|B(z)| = 1$, when $z \in T$.

Li, Mohapatra and Rodriguez [8] extended Bernstein inequality to rational functions $r \in R_n$ with prescribed poles a_1, a_2, \dots, a_n and replaced z^n by Blaschke product $B(z)$ and proved:

Theorem 1.2. ([8]) *If $z \in T$, then for any $r \in R_n$,*

$$|r'(z)| \leq |B'(z)| \sup_{z \in T} |r(z)|.$$

Further more, the inequality is sharp and equality holds for $r(z) = \alpha B(z)$ with $|\alpha| = 1$.

The next result is due to Borwein and Erdelyi [4]:

Theorem 1.3. ([4]) *If $r \in R_n$ and all the zeros of $r(z)$ lie in TUD^+ , then for $z \in T$*

$$|r'(z)| \leq \frac{|B'(z)|}{2} \sup_{z \in T} |r(z)|.$$

Equality holds for $r(z) = \lambda B(z) + \mu$, $|\lambda| = |\mu| = 1$.

Now, we consider a class of rational functions $r(f(z))$ defined by

$$(rof)(z) = r(f(z)) = \frac{p(f(z))}{w(f(z))},$$

where $f(z)$ is a polynomial of degree m^* and $r(z)$ is a rational function of degree n , so that $rof \in R_{m^*n}$, and

$$w(f(z)) = \prod_{j=1}^{m^*n} (z - a_j).$$

Hence, in case of Balaschke product $B(z)$ is given by

$$B(z) := \frac{w^*(f(z))}{w(f(z))} = \frac{z^{m^*n} \overline{w(f(\frac{1}{\bar{z}}))}}{w(f(z))} = \prod_{j=1}^{m^*n} \left(\frac{1 - \bar{a}_j z}{z - a_j} \right).$$

Now onwards, we shall always assume that all poles $a_1, a_2, \dots, a_{m^*n}$ of $r(f(z))$ lie in D^+ . For the case when all poles are in D^- , we can obtain analogous results with suitable transformations.

2. LEMMAS

The first two lemmas are due to Li, Mohapatra and Rodriguez [8].

Lemma 2.1. ([8]) *Suppose $\lambda \in T$. Then the equation $B(z) = \lambda$ has exactly n simple roots, say t_1, t_2, \dots, t_n and all lie on the unit circle T . Moreover*

$$\frac{t_k B'(t_k)}{\lambda} = \sum_{j=1}^n \frac{|a_j|^2 - 1}{|t_k - a_j|^2} \quad \text{for } k = 1, 2, 3, \dots, n. \tag{2.1}$$

Lemma 2.2. ([8]) *If $|x| = |y| = 1$, then*

$$(x - y)^2 = -xy|x - y|^2. \quad (2.2)$$

Qasim [5] have proved:

Lemma 2.3. ([5]) *If $rof \in R_{m^*n}$ and $z \in T$, then*

$$B'(z)r(f(z))-f'(z)r'(f(z))[B(z)-\lambda] = \frac{B(z)}{z} \sum_{k=1}^{m^*n} c_k r(f(t_k)) \left| \frac{B(z) - \lambda}{z - t_k} \right|^2, \quad (2.3)$$

where $c_k = c_k(\lambda)$ is defined for $k = 1, 2, 3, \dots, m^*n$ by

$$c_k^{-1} = \sum_{j=1}^{m^*n} \frac{|a_j|^2 - 1}{|t_k - a_j|^2}. \quad (2.4)$$

Furthermore, for $z \in T$

$$\frac{zB'(z)}{B(z)} = \sum_{k=1}^{m^*n} c_k \left| \frac{B(z) - \lambda}{z - t_k} \right|^2 \quad (2.5)$$

and also

$$|B'(z)| = \frac{zB'(z)}{B(z)} = \sum_{k=1}^{m^*n} \frac{|a_k|^2 - 1}{|z - a_k|^2}, \quad (2.6)$$

where $t_k, k = 1, 2, 3, \dots, m^*n$ are defined in Lemma 2.1.

Aziz and Dawood [1] proved the following lemma :

Lemma 2.4. ([1]) *Let $p(z)$ be a polynomial of degree n , having all zeros in $T \cup D^-$. Then*

$$\inf_{z \in T} p'(z) \geq n \inf_{z \in T} p(z).$$

Next Lemma has been proved by Idrees Qasim [5]:

Lemma 2.5. ([5]) *Let $r(f(z)) \in R_{m^*n}$. Then for $z \in T$,*

$$|r'(f(z))| + |r^{*'}(f(z))| \leq \frac{|B'(z)|}{m^*m} \sup_{z \in T} |r(f(z))|, \quad (2.7)$$

where $r^*(f(z)) = \overline{B(z)r(f(\frac{1}{\bar{z}}))}$. The result is sharp and equality holds for $r(f(z)) = aB(z)$ with $a \in T$, where $f(z) = z^{m^*}$.

Now, we shall prove the following lemma:

Lemma 2.6. *Let $r(f(z)) \in R_{m^*n}$. Then for $z \in T$,*

$$\operatorname{Re} \frac{z((wof)(z))'}{(wof)(z)} = \frac{n - |B'(z)|}{2}$$

and

$$\operatorname{Re} \frac{z((wof)^*(z))'}{(wof)^*(z)} = \frac{n + |B'(z)|}{2}.$$

Proof. We have

$$(wof)^*(z) = z^n \overline{(wof)(1/\bar{z})}.$$

Also

$$z(wof)^*(z)' = nz^n \overline{(wof)(1/\bar{z})} - z^{n-1} f'(1/\bar{z}) \overline{(w'(f(1/\bar{z})))}.$$

Now for $z \in T$, we have

$$\frac{z((wof)^*(z))'}{(wof)^*(z)} = n - \frac{\overline{(w'(f(z)))f'(z)}}{(wof)(z)}.$$

This gives

$$\operatorname{Re} \frac{z((wof)^*(z))'}{(wof)^*(z)} + \operatorname{Re} \frac{w'(f(z))f'(z)}{(wof)(z)} = n. \tag{2.8}$$

Also we know that

$$B(z) := \frac{w^*(f(z))}{w(f(z))} = \frac{z^{m^*n} \overline{w(f(\frac{1}{\bar{z}}))}}{w(f(z))} = \prod_{j=1}^{m^*n} \left(\frac{1 - \bar{a}_j z}{z - a_j} \right).$$

Since $\frac{zB'(z)}{B(z)}$ is a positive real number, this gives

$$\frac{zB'(z)}{B(z)} = \sum_{k=1}^{m^*n} z \left[\frac{-\bar{a}_j}{1 - \bar{a}_j z} - \frac{1}{z - a_j} \right] = \sum_{k=1}^{m^*n} \frac{(|a_j|^2 - 1)}{(\bar{z} - \bar{a}_j)(z - a_j)}$$

and

$$|B'(z)| = \frac{zB'(z)}{B(z)} = \sum_{k=1}^{m^*n} \frac{|a_k|^2 - 1}{|z - a_k|^2}.$$

Again

$$\frac{zB'(z)}{B(z)} := \frac{z \left(\frac{(wof)^*(z)}{(wof)(z)} \right)'}{\left(\frac{(wof)^*(z)}{(wof)(z)} \right)} = z \frac{((wof)^*(z))'}{(wof)^*(z)} - \frac{((wof)(z))'}{(wof)(z)}.$$

Equivalently

$$\frac{zB'(z)}{B(z)} = z \frac{((wof)^*(z))'}{(wof)^*(z)} - \frac{(w'(f(z)))f'(z)}{(wof)(z)}.$$

Therefore, for $z \in T$, we get

$$|B'(z)| = \operatorname{Re} \left[z \frac{((wof)^*(z))'}{(wof)^*(z)} \right] - \operatorname{Re} \left[\frac{(w'(f(z)))f'(z)}{(wof)(z)} \right]. \quad (2.9)$$

From (2.8) and (2.9), we obtain for $z \in T$,

$$\operatorname{Re} \frac{z((wof)(z))'}{(wof)(z)} = \frac{n - |B'(z)|}{2}$$

and

$$\operatorname{Re} \frac{z((wof)^*(z))'}{(wof)^*(z)} = \frac{n + |B'(z)|}{2}.$$

This completes the proof. \square

3. MAIN RESULTS

We first prove the following generalization of Theorem 1.3:

Theorem 3.1. *Let $rof \in R_{m^*n}$ and all zeros of $r(f(z))$ lie in $T_k \cup D_k^+$, where $k \geq 1$, T_k is circle of radius k (centre origin) and D_k^+ represents exterior of T_k . Then for $z \in T$,*

$$|r'(f(z))| \leq \frac{1}{2} \left[\frac{|B'(z)|}{mm^*} + \frac{n|rof|^2}{(\sup_{z \in T} |r(f(z))|)^2 |f'(z)|^2} \left(\frac{2m^*}{k+1} - 1 \right) mm^* \right] \\ \times \sup_{z \in T} |r(f(z))|,$$

where $m = \inf_{z \in T} |f(z)|$.

Proof. Let $(rof)(z) = \frac{(pof)(z)}{(wof)(z)} \in R_{m^*n}$. If $b_1, b_2, \dots, b_{m^*n^*}$ are all zeros of $(pof)(z)$, then $m^*n^* \leq m^*n$, $|b_j| \geq k > 1$, $j = 1, 2, 3, \dots, m^*n^*$ and we have

$$z \left(\frac{(rof)(z)}{(rof)(z)} \right)' = z \frac{\left(\frac{(pof)(z)}{(wof)(z)} \right)'}{\frac{(pof)(z)}{(wof)(z)}} \\ = z \left[\frac{p'(f(z))f'(z)}{(pof)(z)} - \frac{zw'(f(z))f'(z)}{(wof)(z)} \right] \\ = \sum_{k=1}^{m^*n^*} \frac{z}{z - b_j} - \frac{zw'(f(z))f'(z)}{(wof)(z)}.$$

For $z \in T$, this gives, with the help of Lemma 2.6, that

$$\operatorname{Re} \left[z \left(\frac{(rof)(z)}{(rof)(z)} \right)' \right] = \operatorname{Re} \sum_{k=1}^{m^*n^*} \frac{z}{z - b_j} - \operatorname{Re} \frac{zw'(f(z))f'(z)}{(wof)(z)}.$$

$$Re \left[z \frac{((rof)(z))'}{(rof)(z)} \right] = Re \sum_{k=1}^{m^*n^*} \frac{z}{z - b_j} - \left[\frac{n - |B'(z)|}{2} \right]. \tag{3.1}$$

Now it can be easily verified that for $z \in T$, $|b| \geq k > 1$, since $Re(z) \leq |z|$

$$Re \left(\frac{z}{z - b} \right) \leq \frac{1}{k + 1}.$$

Using this in (3.1), we get for $z \in T$,

$$Re \left[z \frac{((rof)(z))'}{(rof)(z)} \right] \leq \sum_{k=1}^{m^*n^*} \frac{1}{k + 1} - \left[\frac{n - |B'(z)|}{2} \right]$$

and

$$Re \left[z \frac{((rof)(z))'}{(rof)(z)} \right] \leq \frac{m^*n^*}{k + 1} - \left[\frac{n - |B'(z)|}{2} \right].$$

Since $m^*n^* \leq m^*n$, this gives

$$Re \left[z \frac{((rof)(z))'}{(rof)(z)} \right] \leq \frac{|B'(z)|}{2} + \frac{m^*n}{k + 1} - \frac{n}{2}$$

or

$$Re \left[z \frac{((rof)(z))'}{(rof)(z)} \right] \leq \frac{|B'(z)|}{2} + n \left[\frac{m^*}{k + 1} - \frac{1}{2} \right].$$

Since

$$\left((rof)^*(z) \right)' = \left[B'(z) \overline{\left((rof)\left(\frac{1}{\bar{z}}\right) \right)} \right]$$

and

$$z \frac{\left((rof)^*(z) \right)'}{(rof)(z)} = \left[|B'(z)| - \frac{r'(f(z))f'(z)}{(rof)(z)} \right] B(z),$$

we have

$$\begin{aligned}
 \left| z \frac{\left((rof)^*(z) \right)'}{(rof)(z)} \right|^2 &= \left| |B'(z)| - \frac{r'(f(z))f'(z)}{(rof)(z)} \right|^2 \\
 &= |B'(z)|^2 + \left| \frac{r'(f(z))f'(z)}{(rof)(z)} \right|^2 \\
 &\quad - 2|B'(z)| \operatorname{Re} \left[\frac{zr'(f(z))f'(z)}{(rof)(z)} \right] \\
 &\geq |B'(z)|^2 + \left| \frac{r'(f(z))f'(z)}{(rof)(z)} \right|^2 \\
 &\quad - |B'(z)| \left[|B'(z)| + n \left(\frac{2m^*}{k+1} - 1 \right) \right] \\
 &\geq \left| \frac{r'(f(z))f'(z)}{(rof)(z)} \right|^2 - n \left(\frac{2m^*}{k+1} - 1 \right) |B'(z)|.
 \end{aligned}$$

This implies, for $z \in T$,

$$\left| \left((rof)^*(z) \right)' \right| \geq \left[|r'(f(z))f'(z)|^2 - n|(rof)(z)|^2 \left(\frac{2m}{k+1} - 1 \right) |B'(z)| \right]^{\frac{1}{2}}$$

and

$$\begin{aligned}
 &\left[|r'(f(z))f'(z)|^2 + n|(rof)(z)|^2 \left(\frac{2m^*}{k+1} - 1 \right) |B'(z)| \right]^{\frac{1}{2}} + |r'(f(z))f'(z)| \\
 &\leq |r^*(f(z))f'(z)| + |r'(f(z))f'(z)|.
 \end{aligned}$$

Combining this with Lemma 2.5, we get

$$\begin{aligned}
 &\left[|r'(f(z))f'(z)|^2 - n|(rof)(z)|^2 \left(\frac{2m^*}{k+1} - 1 \right) |B'(z)| \right]^{\frac{1}{2}} + |r'(f(z))f'(z)| \\
 &\leq |f'(z)| \frac{|B'(z)|}{mm^*} \sup_{z \in T} |r(f(z))|,
 \end{aligned}$$

where m^* is degree of the polynomial $f(z)$ and $m = \inf_{z \in T} |f(z)|$. This gives

$$\begin{aligned}
 &|r'(f(z))f'(z)|^2 + n|(rof)(z)|^2 \left(\frac{2m^*}{k+1} - 1 \right) |B'(z)| \\
 &\leq \left[|f'(z)| \frac{|B'(z)|}{mm^*} \sup_{z \in T} |r(f(z))| - |r'(f(z))f'(z)| \right]^2.
 \end{aligned}$$

Equivalently

$$|r'(f(z))f'(z)|^2 - n|(rof)(z)|^2 \left(\frac{2m^*}{k+1} - 1 \right) |B'(z)|$$

$$\leq \left(\left| \frac{|f'(z)||B'(z)|}{mm^*} \sup_{z \in T} |r(f(z))| \right)^2 + |r'(f(z))f'(z)|^2 - 2|f'(z)| \frac{|B'(z)|}{mm^*} \sup_{z \in T} |r(f(z))||r'(f(z))f'(z)|. \right.$$

Which after simplification yields for $z \in T$,

$$|r'(f(z))f'(z)| \leq \frac{1}{2} \left[\frac{|B'(z)|}{mm^*} + \frac{n|rof|^2}{(\sup_{z \in T} |r(f(z))|)^2 |(f'(z))^2|} \left(\frac{2m^*}{k+1} - 1 \right) mm^* \right] \times \sup_{z \in T} |r(f)|.$$

□

For $k = 1$, we get the following:

Corollary 3.2. *Let $rof \in R_{m^*n}$ and all zeros of $r(f(z))$ lie in $T \cup D^+$. Then for $z \in T$,*

$$|r'(f(z))| \leq \frac{1}{2} \left[\frac{|B'(z)|}{mm^*} + \frac{n|rof|^2}{(\sup_{z \in T} |r(f(z))|)^2 |(f'(z))^2|} (m^* - 1) mm^* \right] \sup_{z \in T} |r(f(z))|.$$

Set $f(z) = z$, Theorem 3.1 yields the following:

Corollary 3.3. *Let $r \in R_n$ and all zeros of r lie in $T_k \cup D_k^+$, where $k \geq 1$. Then for $z \in T$,*

$$|r'(z)| \leq \frac{1}{2} \left[|B'(z)| - \frac{n(k-1)}{(k+1)} \frac{|r(z)|^2}{(\sup_{z \in T} |r(z)|)^2} \right] \sup_{z \in T} |r(z)|.$$

Equality holds for

$$r(z) = \left[\frac{(z+k)}{(z-a)} \right]^2,$$

where $a > 1, k \geq 1$ and $B(z) = \left[\frac{(1-az)}{(z-a)} \right]^2$ and $z = 1$.

Remark 3.4. For $k = 1$, Corollary 3.2 reduces to Theorem 1.3.

Next, we shall prove the following result:

Theorem 3.5. *Let $rof \in R_{m^*n}$ and $r^*(f(z)) = B(z)\overline{r(f(\frac{1}{z}))}$. Then*

$$\sup_{z \in T} \left[\left| \frac{r^{*'}(f(z))}{B'(z)} \right| + \left| \frac{r'(f(z))}{B'(z)} \right| \right] = \sup_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|.$$

Furthermore, in above equation suprema on both sides is attained at the same point $z_0 \in T$.

Proof. Let $rof \in R_{m*n}$. Then we have

$$(rof)^*(z) = B(z) \overline{r\left(f\left(\frac{1}{z}\right)\right)}.$$

Therefore,

$$\frac{((rof)^*(z))'}{(rof)^*(z)} = \frac{zB'(z)}{B(z)} - \frac{\overline{r'(f(\frac{1}{z}))f'(\frac{1}{z})}}{z(rof)(\frac{1}{z})}.$$

Since

$$|B'(z)| = \frac{zB'(z)}{B(z)},$$

for $z \in T$

$$\frac{((rof)^*(z))'}{(rof)^*(z)} = |B'(z)| - \frac{\overline{r'(f(\frac{1}{z}))f'(\frac{1}{z})}}{z(rof)(\frac{1}{z})}.$$

This gives

$$\begin{aligned} |B'(z)| &= \left| \frac{((rof)^*(z))'}{(rof)^*(z)} + \frac{\overline{r'(f(\frac{1}{z}))f'(\frac{1}{z})}}{z(rof)(\frac{1}{z})} \right| \\ &\leq \left| \frac{((rof)^*(z))'}{(rof)^*(z)} \right| + \left| \frac{\overline{zr'(f(\frac{1}{z}))f'(\frac{1}{z})}}{(rof)(\frac{1}{z})} \right|. \end{aligned} \quad (3.2)$$

Since for $z \in T$, $|(rof)^*(z)| = |(rof)(z)|$, it follows from (2.2) that

$$|B'(z)||rof)(z)| \leq |((rof)^*(z))'| + |r'(f(z))f'(z)|. \quad (3.3)$$

Equivalently

$$\sup_{z \in T} \left[\left| \frac{r^{*'}(f(z))}{B'(z)} \right| + \left| \frac{r'(f(z))}{B'(z)} \right| \right] \geq \sup_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|. \quad (3.4)$$

From Lemma 2.5, we have

$$\sup_{z \in T} \left[\left| \frac{r^{*'}(f(z))}{B'(z)} \right| + \left| \frac{r'(f(z))}{B'(z)} \right| \right] \leq \sup_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|. \quad (3.5)$$

From (3.4) and (3.5), we get

$$\sup_{z \in T} \left[\left| \frac{r^{*'}(f(z))}{B'(z)} \right| + \left| \frac{r'(f(z))}{B'(z)} \right| \right] = \sup_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|.$$

This completes the proof. \square

Set $f(z) = z$ in Theorem 3.5, we get the following result:

Corollary 3.6. *Let $r \in R_n$ and $r^*(z) = B(z)\overline{r(\frac{1}{\bar{z}})}$. Then*

$$\sup_{z \in T} \left[\left| \frac{r^{*'}(z)}{B'(z)} \right| + \left| \frac{r'(z)}{B'(z)} \right| \right] = \sup_{z \in T} \left| \frac{r(z)}{B(z)} \right|.$$

Furthermore, in above equation suprema on both sides is attained at the same point $z_0 \in T$.

The rational function $r \in R_n$ is self-inversive if $r^*(z) = s\overline{r(\frac{1}{\bar{z}})}$, for $s \in T$. The following result, which is a generalization of Theorem 1 of [7] for self-inversive polynomials, follows from Theorem 3.5.

Corollary 3.7. *If $r \in R_n$ is self-inversive function, then*

$$2 \sup_{z \in T} \left| \frac{r'(z)}{B'(z)} \right| = \sup_{z \in T} \left| \frac{r(z)}{B(z)} \right|.$$

Theorem 3.8. *Suppose $rof \in R_{m^*n}$ has m^*n zeros and all the zeros of rof lie in TUD^- . If*

$$(rof)^*(z) = B(z)\overline{r(f(\frac{1}{\bar{z}}))},$$

then for $z \in T$,

$$\inf_{z \in T} \left[\left| \frac{r'(f(z))}{B'(z)} \right| - \left| \frac{r^{*'}(f(z))}{B'(z)} \right| \right] = \inf_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|.$$

Moreover, in above equation infima on both sides is attained at the same point $z_0 \in T$.

Proof. Suppose all zeros m^*n of $rof \in R_{m^*n}$ lie in TUD^- and let $m_* = \inf_{z \in T} |(rof)(z)|$, then we have $m_* \leq |(rof)(z)|$, for $z \in T$. We show for any complex number α with $|\alpha| < 1$, then rational function $F(z) = c(z) + \alpha m_*$ has all its zeros in TUD^- . This is obvious if $m_* = 0$ i.e, if $(rof)(z)$ has a zero on T . So we suppose all the zeros of $(rof)(z)$ lie in D^- so that $m_* \neq 0$ and we have for $z \in T$

$$|\alpha m_*| < |m_*| \leq |(rof)(z)|.$$

Apply Rouché's theorem, it follows that

$$F(z) = c(z) + \alpha m_*$$

has all its zeros in D^- . Hence in any case $F(z)$ has all its zeros in TUD^- for every $\alpha, |\alpha| < 1$. Let

$$\begin{aligned} F^*(z) &= B(z)\overline{f(\frac{1}{\bar{z}})} \\ &= B(z)\overline{(rof)(\frac{1}{\bar{z}})} + \bar{\alpha}m_*B(z) \\ &= (rof)^*(z) + \bar{\alpha}m_*B(z). \end{aligned}$$

Then all the zeros of $F^*(z)$ lie in TUD^+ . Now, it follows from Theorem 3.1 with $k = 1$, $z \in T$ and $(rof)^*$ replaced by F^* ,

$$|(F^*(z))'| \leq |(F(z))'|.$$

This implies

$$|((rof)^*(z))' + \bar{\alpha}mB'(z)| \leq |((rof)(z))'|.$$

Choosing argument of α suitably, we get for $z \in T$

$$|((rof)^*(z))'| + m|\alpha||B'(z)| \leq |((rof)(z))'|.$$

Letting $|\alpha| \rightarrow 1$, we get

$$m|B'(z)| \leq |((rof)^*(z))'| - |((rof)(z))'|.$$

This implies

$$\inf_{z \in T} \left[\left| \frac{r'(f(z))}{B'(z)} \right| - \left| \frac{r^{*'}(f(z))}{B'(z)} \right| \right] \geq \inf_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|. \quad (3.6)$$

Again

$$(rof)^*(z) = B(z)r \left(\overline{f\left(\frac{1}{\bar{z}}\right)} \right).$$

Therefore,

$$\begin{aligned} |(r^*(f(z)))'| &= \left| B'(z)r \left(\overline{f\left(\frac{1}{\bar{z}}\right)} \right) - \frac{1}{z^2}B(z)r' \left(\overline{f\left(\frac{1}{\bar{z}}\right)} \right) f' \left(\frac{1}{\bar{z}} \right) \right| \\ &\geq \left| B'(z)r \left(\overline{f\left(\frac{1}{\bar{z}}\right)} \right) \right| - \left| B(z)r' \left(\overline{f\left(\frac{1}{\bar{z}}\right)} \right) f' \left(\frac{1}{\bar{z}} \right) \right|. \end{aligned}$$

This implies

$$\inf_{z \in T} \left[\left| \frac{r'(f(z))}{B'(z)} \right| - \left| \frac{r^{*'}(f(z))}{B'(z)} \right| \right] \leq \inf_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|. \quad (3.7)$$

From (3.6) and (3.7), we get

$$\inf_{z \in T} \left[\left| \frac{r'(f(z))}{B'(z)} \right| - \left| \frac{r^{*'}(f(z))}{B'(z)} \right| \right] = \inf_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|.$$

We now show that the infimum of both sides are attained at the same point $z_0 \in T$. Let

$$\inf_{z \in T} |(rof)(z)| = |(rof)(z_0)|.$$

Since

$$|(rof)(z)||B'(z)| \geq |((rof)(z))'| - |((rof)^*(z))'|, \quad (3.8)$$

we have

$$|(rof)(z_0)| \geq \left| \frac{((rof)(z_0))'}{B'(z_0)} \right| - \left| \frac{((rof)^*(z_0))'}{B'(z_0)} \right|. \quad (3.9)$$

From (3.8) and (3.9), we get

$$\inf_{z \in T} \left[\left| \frac{r^{*'}(f(z_0))}{B'(z_0)} \right| + \left| \frac{r'(f(z_0))}{B'(z_0)} \right| \right] = \inf_{z \in T} \left| \frac{(rof)(z_0)}{f'(z_0)} \right|.$$

This completes the proof. □

From Theorem 3.3 we can deduce the following:

Corollary 3.9. *Suppose $rof \in R_{m^*n}$ has m^*n zeros and all the zeros of rof lie in TUD^- . If*

$$(rof)^*(z) = \overline{B(z)r\left(f\left(\frac{1}{\bar{z}}\right)\right)},$$

then for $z \in T$,

$$\inf_{z \in T} \left[\left| \frac{r'(f(z))}{B'(z)} \right| \right] \geq \inf_{z \in T} \left| \frac{(rof)(z)}{f'(z)} \right|.$$

Set $f(z) = z$, we get the following result:

Corollary 3.10. *If $r \in R_n$ has n zeros and all the zeros of r lie in TUD^- . If*

$$(r)^*(z) = \overline{B(z)r\left(\frac{1}{\bar{z}}\right)},$$

then for $z \in T$,

$$\inf_{z \in T} \left[\left| \frac{r'(z)}{B'(z)} \right| \right] \geq \inf_{z \in T} |r(z)|.$$

REFERENCES

- [1] A. Aziz and Q.M. Dawood, *Inequalities for a polynomial and its derivative*, J. Approx. Theory, **53** (1988), 155-162.
- [2] A. Aziz and Q.G. Mohammad, *Simple proof of a Theorem of Erdős and Lax*, Proc. Amer. Math. Soc., **80** (1980), 119-122.
- [3] P. Borwein and T. Erdelyi, *Polynomial Inequalities*, Springer-Verlag, New York, 1995.
- [4] P. Borwein and T. Erdelyi, *Sharp extensions to rational spaces*, Mathematika, **43** (1996), 413-423.
- [5] Idrees Qasim and A. Liman, *Berstein Type Inequalities for rational functions*, Indian J. Pure Appl. Math., **46**(3) (2015), 337-348,
- [6] P.D. Lax, *Proof of a conjecture of P. Erdős on the derivative of a polynomia*, Bull. Amer. Math. Soc., **50** (1944), 503-513.
- [7] P.J. O'Hara and R.S. Rodriguez, *Some properties of self-inversive polynomials*, Proc. Amer. Math. Soc., **44** (1974), 331-335.
- [8] Xin Li, R.N. Mohapatra and R.S. Rodriguez, *Berstein Type Inequalities for rational functions with prescribed poles*, J. London, Math. Soc., **51** (1995), 523-531.