

ARTICLE

Effect of Adding Cultured Meat Tissue on Physicochemical and Taste Characteristics of Hybrid Cultured Meat Manufactured Using Wet-Spinning

 So-Hee Kim^{1,†}, Swati Kumari^{1,†}, Chan-Jin Kim¹, Eun-Yeong Lee¹, AMM Nurul Alam¹, Yong-Sik Chung², Young-Hwa Hwang^{3,4}, and Seon-Tea Joo^{1,3,4,*}
¹Division of Applied Life Science (BK21 Four), Gyeongsang National University, Jinju 52828, Korea

²Department of Organic Materials and Textile Engineering, Jeonbuk National University, Jeonju 54896, Korea

³Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea

⁴Orange CAU Co., Ltd., Jinju 52839, Korea


OPEN ACCESS
Received October 6, 2024

Revised October 15, 2024

Accepted October 15, 2024

***Corresponding author** : Seon-Tea Joo
 Division of Applied Life Science
 (BK21 Four), Gyeongsang National
 University, Jinju 52828, Korea
 Tel: +82-55-772-1943
 Fax: +82-55-772-1949
 E-mail: stjoo@gnu.ac.kr

***ORCID**
 So-Hee Kim
<https://orcid.org/0000-0003-3966-6160>
 Swati Kumari
<https://orcid.org/0009-0001-3330-7821>
 Chan-Jin Kim
<https://orcid.org/0000-0001-5020-6873>
 Eun-Yeong Lee
<https://orcid.org/0000-0002-3467-7349>
 AMM Nurul Alam
<https://orcid.org/0000-0003-3153-3718>
 Yong-Sik Chung
<https://orcid.org/0009-0009-9693-2825>
 Young-Hwa Hwang
<https://orcid.org/0000-0003-3687-3535>
 Seon-Tea Joo
<https://orcid.org/0000-0002-5483-2828>

[†] These authors contributed equally to this work.

Abstract This study investigated effect of adding cultured meat tissue (CMT; 10%, 20%, and 30%) to plant protein on quality of imitation muscle fiber (IMF) and hybrid cultured chicken meat (HCCM) manufactured using wet-spinning. The composite plant-based protein (CPP) solution consisted of pea protein, wheat protein, and sodium alginate. Adding 10%, 20% and 30% of CMT to CPP significantly reduced pH and Warner-Bratzler shear force of IMF ($p < 0.05$). However, texture profile analysis revealed that hardness, gumminess, and cohesiveness of the CMT 30% sample were significantly higher while springiness was lower in CPP without adding CMT ($p < 0.05$). Chewiness of CMT 20% was the highest among HCCM samples ($p < 0.05$). As the amount of CMT added increased, sourness decreased significantly, while bitterness and richness increased significantly (all $p < 0.05$). As CMT addition level increased, essential amino acid levels also increased comprehensively except phenylalanine, leading to improved nutritional quality of HCCM. These results imply that adding CMT could compensate for amino acids that are absent or lacking in CPP and enhance the taste of HCCM.

Keywords wet spinning, hybrid meat, cultured meat, fabrication, meat alternatives

Introduction

Plant-based meat alternative has secured a significant market share around the world (Ismail et al., 2020; Kumari et al., 2023). However, consumers are not ready to compromise on the taste or quality parameter (McClements and Grossmann, 2021). Thus, researchers and industrialists are trying to find a way in between (Hoek et al., 2013), including replacing a portion of meat with more sustainable protein source, either plant based or

cultured meat (Alam et al., 2024; Molfetta et al., 2022). These products are known as hybrid meat products (Grasso and Jaworska, 2020). Various types of hybrid meat products have been developed around the world. The production of hybrid meat is expected to solve food related problems due to the increasing population and consumer perception for balanced diets (Grasso, 2024). Products conventionally available in the market are manufactured by combining different sources with real meat such as chicken and beef (Annoh-Quarshie, 2018; He et al., 2021; Zahari et al., 2021). This combining strategy creates a flow of simple processed products such as hamburger patties and chicken nuggets. These processes require the use of a high temperature. Most of the hybrid meat products are produced using a high-temperature processing which causes loss of nutrients with deteriorating effects on cooking parameters of final products (Chandler and McSweeney, 2022; Grasso, 2024). In other words, the manufacturing process of hybrid meat has a limitation (Alam et al., 2024). To overcome problems related to high-temperature processing, a wet spinning approach can be considered.

The wet-spinning technique has been traditionally used in the textile industry. This particular technique is based on a bottom-up approach that requires complicated facilities (Dekkers et al., 2018; Kyriakopoulou et al., 2019). However, it is easier to control characteristics of fibers. In recent years, researchers have added a small amount of protein into the spinning solution to improve physicochemical properties of fibers (Cui et al., 2022). Based on this notion, a preliminary study has been conducted to utilize different protein sources including pea protein and wheat protein (WP; Kumari et al., 2024). It was found that imitation fiber from a combination of two kinds of plant protein had the potential to mimic conventional meat. Additionally, incorporating cultured meat with plant protein has not been reported yet. This creates a gap in this field of hybrid meat product. To leverage advantages of wet spinning and fulling this gap, plant protein and cultured meat tissue (CMT) were utilized in this study. Plants-based protein can control production price increases due to relatively expensive CMT. Nutrients that cannot be provided by only plant proteins could be supplemented with CMT. Therefore, in this study, differences in quality characteristics between manufactured hybrid cultured chicken breasts were examined by adding CMT at different concentrations to plant-based protein using wet-spinning.

Materials and Methods

Cell culture and harvest

Chicken satellite cells (CSC) were isolated from hindlimb muscles as previously published study (Kim et al., 2022; Kim et al., 2023). Isolated CSC was suspended in growth media (GM) containing 20% fetal bovine serum (FBS; S1-004, Welgene, Gyeongsan, Korea), 1% GlutaMAXTM supplement (35050061, Gibco, London, UK), 1% antibiotic-antimycotic (15240062, Gibco), and 5 ng/mL basic fibroblast growth factor (233-FB-025, R&D Systems, Minneapolis, MN, USA) in DMEM. The cells were initially cultured in 175T flasks at a density of 3,000 cells/cm² in 41°C and 5% CO₂ incubator for scale-up. When the cell confluency came over 70%, the supernatant was aspirated, and the cells were dissociated using 0.25% trypsin-EDTA (LS015-10, Welgene). The cell suspensions were centrifuged at 800×g for 5 min to harvest the cells for further 3D culture. Cytodex 1 (Cytiva, Marlborough, MA, USA) microcarriers were sterilized by autoclaving at 121°C for 20 minutes, and subsequently hydrated in GM for 1 hour prior to use. The cells were seeded into the spinner flasks with Cytodex 1 microcarriers and cultured at a density of 3,000 cells/cm² in 41°C and 5% CO₂ incubator with stirring at 50 rpm. When the cells reached 100% confluency, the GM was aspirated, the cells were rinsed three times with DPBS. The cells were then dissociated using 0.25% trypsin-EDTA for 5 mins and the cell suspensions were passed through a 100 µm sieve to remove Cytodex 1 microcarriers. The cell suspension was centrifuged at 800×g for 5 min to harvest CMT. The harvested CMT was

lyophilized using freeze-dryer (OPERON OPR-FDB-5503 FREEZE DRY SYSTEM, OPERON, Gimpo, Korea) and then kept in a -70°C deep freezer until a sufficient amount of CMT was collected for the next experiments.

Materials for wet-spinning solution

Pea protein isolate (PPI) and WP were purchased from an online platform. Sodium alginate (SA) with high viscosity was obtained from online market (ESfood, Gunpo, Korea). Calcium chloride was purchased from Qingdao Soda Ash Industrial Development (Qingdao, China). All materials used for experiments were of food grade.

Sample preparation

Plant protein solution was prepared by dissolving 4% (w/v) WP and PPI in distilled water (DW) respectively. SA solution was formulated by dispersing SA in DW at a concentration of 2% (w/v). All solutions were kept at 4°C overnight to achieve complete hydration and a stable state. The plant protein solution was prepared by mixing WP and PPI solutions in equal amounts. Then, for making composite plant-based protein (CPP) solution, SA solution was mixed with plant protein solution in equal ratio. Hybrid cultured chicken meat (HCCM) contains 4% PPI, 4% WP, and CMT at concentrations of 10%, 20%, and 30%, which were mixed in equal volumes. SA solution was also added to mixture for HCCM production. All the solutions were uniformly mixed for 20 min and degassed for 20 min at room temperature with at 20 kHz using ultrasonicator (VCX 750, SONICS, Newtown, CT, USA).

Manufacturing of imitation muscle fiber and muscle

Imitated muscle fiber (IMF) was manufactured using wet-spinning according to the method of Kumari et al. (2024). In a coagulation bath, CPP solution or CPP solution containing CMT were extruded through a needle of 0.13 mm in diameter into a 3% calcium chloride (w/w) at room temperature (20°C – 25°C). IMFs were washed in the washing bath containing DW to remove the excess or remaining calcium chloride from surfaces of IMFs. After collecting IMFs, a cellulose membrane produced by electrospinning technique was used for warping each IMF. The process was repeated a number of times to make several muscle bundles which were then surrounded by a secondary membrane to provide a mimicking effect like conventional meat. CPP samples added with 10%, 20%, and 30% concentrations of CMT were designated as CMT 10%, CMT 20%, and CMT 30% respectively. Fig. 1 is a diagram showing the structure of artificial imitation meat that mimics the




Fig. 1. A diagram showing the structure of artificial imitation meat (A) that mimics the structure of meat (B).

structure of conventional meat.

Cellulose membrane by electrospinning

Cellulose acetate (CA), glacial acetic acid (AA), and citric acid anhydrous (CAA) of food grade were purchased from an online platform. CA solution stock solution was prepared by blending into 20% (w/v) of CA dissolved in 85% (V/V) AA. The process was carried out at 45°C with continuous stirring at 750 rpm for 12 hours until the solution became fully homogenized. For crosslinking, CAA was added to the stock solution of CA (30%). This solution was mixed with a magnetic stirrer for 30 minutes at 25°C with shaking at 1,500 rpm until a homogenous solution was obtained. Prepared solutions were then loaded into a 10 mL syringe with a 23 G needle and put into an electrospinning device (Electrospinning System, Nano NC, Seoul, Korea). Based on preliminary examinations, optimized electrospinning parameters were: a voltage of 18 kV, a needle-to-collector distance of 12 cm, a flow rate of 0.4 mL/h, and a collector rotating speed of 500 rpm.

Fabrication and appearance characterization of hybrid cultured chicken breast

HCCM was manufactured by introducing CMT into CPP solution at different levels (10%, 20%, and 30%). Fig. 2 illustrates the manufacturing process of HCCMs. Wet-spinning and electro-spinning techniques were used to produce IMF and artificial muscle membrane, respectively. The CPP solution without CMT was designated as CPP, while HCCMs produced by adding CMT to CPP were designated as CMT 10%, CMT 20%, and CMT 30% according to CMT content. Each IMF produced through coagulation using wet-spinning was fabricated by wrapping IMF with an artificial muscle membrane in order to replicate the structure of traditional muscle.

Measurements of imitated muscle fiber quality

Color

Color values of IMFs (CPP, CMT 10%, CMT 20% and CMT 30%) were measured with a Chroma Meter (CR-300, Konica Minolta, Osaka, Japan). Color values (CIE L*, CIE a*, and CIE b*) are presented as average values obtained from five measurements for each sample. Results are expressed as mean±SD.

pH

IMF was homogenized with DW at a ratio of 1:9. pH was measured triplicate using a digital pH meter (A211 pH Meter, Thermo Fisher Scientific, Waltham, MA, USA).

Warner-Bratzler shear force

IMF samples were cut into pieces with dimensions of 1 cm×1 cm×1 cm (length×width×height). Warner-Bratzler shear force (WBSF) was measured with a texture analyzer (AMETEK, Berwyn, PA, USA) and a V-shaped shear blade on its shear mode. The analysis was performed at a speed of 100 mm/min with a force of 50 kg. Data were processed and expressed as mean and SD of values measured five times.

Texture profile analysis

Texture profile analysis (TPA) was conducted using a double compression test, involving compression of the sample under




Fig. 2. A schematic diagram of processing flow for hybrid cultured chicken meat (HCCM) by wet-spinning.

fixed conditions. TPA of IMF was performed with a texture analyzer (AMETEK). All samples were shaped into 1 cm×1 cm×1 cm cubes. Compression and decompression were conducted twice at a fixed speed of 100 mm/min and a maximum load of 180 kg on a measuring cell. TPA parameters included hardness, springiness, gumminess, chewiness, and cohesiveness of each sample. Data were processed and expressed as mean and SEM for values measured five times.

Total amino acid contents

IMF (500 mg) was hydrolyzed with 6 N HCl in a dry oven at 110°C for 16 h. The hydrolysate was filtered with a Whatman filter no.1 paper and diluted with DW to a concentration of 0.1 N. Sample vials were then prepared by filter through a 0.22 µm PTFE syringe filter. Total amino acid content was analyzed using an OPA derivatization protocol provided by Agilent.

Sensory evaluation with electronic tongue

An electronic tongue system (ETS; INSERT SA402B Electric Sensing System, Insent, Tokyo, Japan) was used for measuring relative sensory characteristics of each sample with the technique exemplified by Ismail et al. (2020). The ETS can distinguish five different flavors with five different taste sensors (CA0, C00, and AAE) to analyze relative intensities of sourness, bitterness, umami, and richness, respectively. All membranes in sensors were stabilized in a standard meat taste (SMT) solution containing 0.01% lactic acid (DAEJUNG, Busan, Korea), 0.25% monosodium glutamate (DAEJUNG, Korea), and 0.0005% quinine hydrochloride (TCI, Tokyo, Japan). A solution obtained by mixing IMF with DW at 100°C was used for sensory analysis at ratio of 1:4 for 30 min. The agitated solution was centrifuged at 1,000×g for 15 min. The supernatant was collected and stored at -70°C for further analysis.

Statistical analysis

Statistical analyses were conducted using GraphPad Prism 10 software 10.0.2 (GraphPad, San Diego, CA, USA). All data are presented as mean and SD or SEM. The WBSF and TPA were measured in quintuplicate, while all other experiments were conducted in triplicate. Results were subjected to one-way analysis of variance (ANOVA) with Tukey's multiple comparison test. Principal component analysis (PCA) was conducted to assess the variation in overall qualities among the treatment groups (CPP, CMT 10%, CMT 20%, and CMT 30%). The variables used in PCA included the data of physicochemical analysis, amino acid analysis, and sensory analysis, respectively. A score plot was illustrated for the differences in distribution among groups. Statistical significance was considered when p-value was less than 0.05.

Results and Discussion

Effect of addition of cultured meat tissue on pH of solution for wet-spinning

The solution pH for wet-spinning decreased significantly ($p < 0.05$) after adding CMT to CPP (Table 1). The pH of the CPP solution (6.83) was mild neutral due to the presence of pea protein and WP along with SA. This trend could be due to increased amounts of glutamic acid and aspartic acid as total amino acid content was increased with each addition (Ferreira and Teixeira, 2003). Additionally, with increasing concentration of CMT, the buffering capacity of plant proteins may have been insufficient to neutralize additional acidic by-products from CMT, causing the overall pH to drop (Ebert et al., 2021).

This change in pH due to the addition of CMT is likely to interfere with homogeneous distribution of particles in the solution, leading to difficulties in dissolving materials. Moreover, in a previous study (Kumari et al., 2024) using wet-spinning, the highest water-holding capacity (82.66%) was observed at pH 6.44. This indicates that pH is directly related to water-holding capacity, highlighting the importance of pH control during the preparation of a solution for wet-spinning.

Changes in color and appearance of hybrid cultured chicken meat with addition of cultured meat tissue

Results of color measurements including CIE L*, CIE a* and CIE b* are shown in Table 1. Overall color values showed a

Table 1. Effects of cultured meat tissue (CMT) addition to a composite plant-based protein (CPP) on pH, moisture, and color of hybrid cultured chicken meat (HCCM)

Measurements	CPP	CMT 10%	CMT 20%	CMT 30%
pH	6.83±0.002 ^A	6.70±0.001 ^B	6.60±0.001 ^C	6.49±0.004 ^D
Color				
CIE L*	70.98±0.28 ^D	73.21±0.17 ^C	77.84±0.24 ^B	80.75±0.47 ^A
CIE a*	-1.91±0.03 ^D	-1.58±0.02 ^C	-1.38±0.03 ^B	-0.38±0.01 ^A
CIE b*	7.06±0.14 ^D	8.06±0.19 ^C	9.19±0.001 ^B	10.15±0.14 ^A

The pH value are presented as mean±SD (n=3).

The color value are presented as mean±SD (n=5).

^{A-D} Different superscripts in the same row indicate significant difference (p<0.05).

significant increase compared to CPP as the concentration of CMT in HCCM increased (p<0.05). This indicates that the addition of CMT makes the color brighter, redder, and more yellow. This color change was caused by a brighter and more yellowish color of the CMT than other plant-based protein relatively, indicating that the color of HCCM produced through wet-spinning could be greatly influenced by the color of materials used in its production (Cui et al., 2014; Fraeye et al., 2020). It has been suggested that color attributes can enhance visual attractiveness of hybrid cultured chicken breast, potentially increasing consumer acceptance by adding CMT (Lee et al., 2020). In addition, the cross-sectional view of HCCM produced using wet-spinning showed a fibrous structure more similar to that of muscle fibers in conventional meat when CMT was added (Fig. 2). This suggests that the addition of CMT during the manufacture of HCCM using wet-spinning techniques has the potential to achieve a more similar appearance to conventional meat.

Changes in shear force and texture of hybrid cultured chicken meat with addition of cultured meat tissue

Table 2 shows changes in tenderness and texture of HCCM with the addition of CMT. WBSF of CPP was significantly higher than those of CMT 20% and 30% HCCM. The WBSF decreased significantly (p<0.05) when CMT content was increased. This decrease of WBSF implies that addition of CMT can make tender HCCM. Adding CMT could have created a softer texture by interfering with the rigid cross-linking structure of plant protein with SA, resulting in a reduction in WBSF for CMT 20% and CMT 30% HCCMs. For CMT 10% containing, it was thought that plant-based proteins interacted with

Table 2. Effects of cultured meat tissue (CMT) addition to a composite plant-based protein (CPP) on tenderness and texture of hybrid cultured chicken meat (HCCM)

Measurements	CPP	CMT 10%	CMT 20%	CMT 30%	SEM	p-value
WBSF (kg/cm ²)	3.247 ^A	3.322 ^A	2.210 ^B	2.019 ^C	0.136	<0.0001
Hardness (N)	10.600 ^B	10.900 ^B	12.580 ^A	12.930 ^A	0.260	<0.0001
Springiness	0.895 ^A	0.895 ^A	0.891 ^A	0.869 ^B	0.003	0.0019
Gumminess (N)	4.339 ^C	3.973 ^D	4.845 ^B	5.207 ^A	0.115	<0.0001
Chewiness (N)	3.955 ^C	3.497 ^D	4.761 ^A	4.529 ^B	0.115	<0.0001
Cohesiveness	0.389 ^B	0.324 ^D	0.364 ^C	0.406 ^A	0.007	<0.0001

All values are presented as means and SEM (n=5).

^{A-D} Different superscripts in the same row indicate significant difference (p<0.05).

WBSF, Warner-Braztler shear force.

CMT to form strong gels with increased structural integrity of the structure, resulting in a stiffness of HCCM. In a study by Bakhsh et al. (2021) and Caine et al. (2003), the WBSF of plant-based meat analog (PBMA) patty was approximately 2.74 kgf/cm². This value fell between values for conventional beef and pork patties. Similarly, in the present study, HCCM produced by wet-spinning had WBSF values generally ranging from 2.0 to a maximum of 3.2, close to the WBSF value of conventional meat. This result suggests that wet-spinning techniques can control tenderness more easily by adjusting the composition of IMFs than the high-temperature extrusion method for manufacturing textured vegetable protein (TVP).

On the other hand, the addition of CMT resulted in a significant change in the texture of HCCM. Similar to WBSF results, hardness values of CMT 20% and CMT 30% were significantly higher than that of CPP ($p < 0.05$). However, CMT 10% and CPP showed no significant difference in hardness ($p > 0.05$). This trend was observed similarly in a previous study (Kumari et al., 2024). The springiness displayed no significant difference until CMT was introduced into CPP at 20% ($p > 0.05$). However, springiness showed a significant decline in CMT 30% ($p < 0.05$). The decrease of springiness in hybrid meat containing 30% of CMT could be due to interactions of different protein types. These proteins might have affect cross-linking with SA during the process (Nagamine et al., 2023). Gumminess, chewiness, and cohesiveness were significantly lower in CMT 10% among HCCMs. This could be due to an antagonist effect of the plant-based protein and a low concentration of the CMT protein (Ježek et al., 2020). With increased concentration of CMT, the overall cultured meat protein content might have cross-linked with each other, creating a firm and more cohesive structure (Kumari et al., 2024; Younis et al., 2023).

Changes in amino acid compositions with addition of cultured meat tissue

Amino acid analysis was conducted to determine how much CMT should be incorporated into CPP to have an effect on amino acid compositions and contents of HCCM. Results of amino acid analysis are shown in Table 3. Essential amino acid levels were increased comprehensively except for phenylalanine, improving the nutritional quality of the HCCM. These significant changes in essential amino acid level were attributed to the incorporation of CMT into CPP which resulted in increased total amino acid content. Results of amino acid composition analysis showed a distinctive decrease in glutamic acid due to wheat gluten in CPP solution because gluten could produce glutamic acid by hydrolysis (Manning, 1950). Glutamic acid sees a large increase, reflecting a high protein content in cultured meat, making it a key contributor to the blend (Qi et al., 2017). Lysine showed a dramatic rise displaying that the addition of CMT has compensated for its lower levels in plant proteins, along with a significant increase in amounts of leucine, isoleucine, and valine due to their abundance in animal-derived proteins, highlighting the impact of cultured meat on enhancing the solution's nutritional value. On the other hand, proline and tyrosine showed smaller increases, with significant changes emerging at higher cultured meat levels (CMT 20% and 30%). Additionally, the other amino acids such as phenylalanine, aspartic acid, and arginine also increased notably with each addition of cultured meat, further enriching the overall amino acid profile.

The increase in amino acid content with the addition of CMT could be due to its rich protein profile, which complements plant proteins in pea and wheat (Treich, 2021). Therefore, this study confirms that adding CMT could compensate for amino acids that are lower or missing in plant proteins, such as lysine, proline, and branched-chain amino acids (leucine, isoleucine, valine), leading to significant improvements in the overall nutritional quality (Wu, 2021).

Effect of adding cultured meat tissue on taste characteristics of hybrid cultured chicken meat

Changes in taste characteristics of HCCM evaluated by electronic tongue with addition of CMT to CPP are shown in Fig. 3. Overall, taste profiles including sourness, bitterness, and richness were significantly increased except for Umami ($p < 0.05$).

Table 3. Effects of cultured meat tissue (CMT) addition to a composite plant-based protein (CPP) on amino acid compositions of hybrid cultured chicken meat

Amino acids	CPP	CMT 10%	CMT 20%	CMT 30%	SEM	p-value
Asp	11.59 ^C	11.46 ^B	13.94 ^A	14.03 ^A	0.337	<0.0001
Glu	34.32 ^A	33.85 ^B	28.01 ^B	25.31 ^C	0.916	<0.0001
Ser	5.09 ^A	5.08 ^B	4.81 ^B	4.75 ^B	0.049	0.001
His	0.65 ^A	0.71 ^A	0.80 ^A	1.00 ^A	0.046	0.8463
Gly	1.45 ^A	1.50 ^A	1.58 ^A	1.82 ^A	0.038	0.0537
Thr	3.47 ^C	3.58 ^{BC}	4.48 ^{AB}	5.05 ^A	0.149	0.0014
Arg	4.45 ^B	4.66 ^B	4.47 ^B	5.47 ^A	0.143	0.0011
Ala	3.02 ^B	3.04 ^A	4.51 ^A	4.86 ^A	0.183	0.005
Tyr	3.18 ^B	3.23 ^B	3.24 ^B	3.45 ^A	0.032	0.0002
Val	4.86 ^A	4.76 ^A	5.05 ^A	5.03 ^A	0.031	0.9266
Met	0.86 ^B	0.91 ^B	1.27 ^{AB}	1.53 ^A	0.094	0.0056
Phe	6.20 ^A	6.17 ^{AB}	5.64 ^{BC}	5.28 ^C	0.123	0.0014
Ile	4.76 ^B	4.74 ^{AB}	5.26 ^{AB}	5.23 ^A	0.049	0.0254
Leu	7.94 ^C	7.97 ^{BC}	8.41 ^B	8.82 ^A	0.086	0.0002
Lys	2.96 ^C	3.03 ^B	4.32 ^B	5.79 ^A	0.283	<0.0001
Pro	5.13 ^A	5.24 ^B	4.14 ^B	2.52 ^C	0.246	0.0001

All values are presented as means and SEM (n=3).

^{A-C} Different superscripts in the same row indicate significant difference (p<0.05).

In general, considering that a decrease in the sourness of HCCM improves the overall taste, it is presumed that the addition of CMT can positively enhance the taste of HCCM. However, there was no change in umami level with an increase in the amount of CMT. The reason for the unchanged umami level can be due to the antagonist effect resulting from an increase of aspartic acid and a decrease of glutamic acid (Table 3). Although the richness of HCCM increased significantly, indicating that the overall mouthfeel may have increased due to increasing overall amino acid profile and the protein interaction (p<0.05; Paradowska et al., 2021; Xu and Falsafi, 2023). Meanwhile, the CMT addition increased the bitterness, especially in CP30 could be due to additional peptides and amino acids (e.g., histidine, arginine isoleucine, leucine; Tagliamonte et al., 2024).

Principal component analysis of quality characteristics of hybrid cultured chicken meat

PCA was conducted to analyze variations in quality among the four treatment groups (CPP, CMT 10%, CMT 20%, and CMT 30%). The first principal component (PC1) accounted for 66.19% of the total variance and the second principal component (PC2) explained an additional 12.29% of the total variance (Fig. 4). These two components explained approximately 78.48% of the total variance, providing most of the differences among treatment groups.

CPP and CMT 10% groups were mainly positioned on the negative side of PC1. This indicates that CPP and CMT 10% have similar characteristics. The result indicated that although CMT was added to the CMT 10% group, the effect on quality was minimal, leading to a closely clustered grouping with CPP in the PCA plot.

CMT 20% and CMT 30% groups showed positive values in PC1 compared to the other two groups. In particular, CMT




Fig. 3. Effects of cultured meat tissue (CMT) addition to a composite plant-based protein (CPP) on taste characteristics of hybrid cultured chicken meat (HCCM) evaluated by electronic tongue. ^{A-D} Different superscripts indicate significant difference ($p < 0.05$).

30% was located on the far right of the PC1 plot, which was clearly separated from the other three groups. When comparing CPP and CMT 30% groups, these two groups were most distinctly separated by PC1 in the PC plot, indicating significant differences in their quality. The addition of CMT to hybrid cultured chicken breast has a significant impact on the quality of HCCM, indicating that selecting the optimal combination of plant and animal proteins in hybrid meat production can considerably enhance both its nutritional value and overall quality.

Conclusion

The addition of CMT to CPP significantly improved the quality of IMF and HCCM produced via wet spinning. Specifically, CMT incorporation reduced pH and WBSF but enhanced essential amino acid levels, thus improving nutritional




Fig. 4. Principal component analysis of quality characteristics of hybrid cultured chicken meat (HCCM). CPP, composite plant-based protein; CMT 10%, cultured meat tissue 10%; CMT 20%, cultured meat tissue 20%; CMT 30%, cultured meat tissue 30%.

quality. Texture and sensory properties also improved from CMT addition, with higher content increasing the hardness, chewiness, and flavor richness. Overall, CMT can effectively compensate for deficiencies in plant proteins, enhancing both nutritional and sensory qualities of HCCM.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by the National Research Foundation of Korea (NRF) grantfunded by the Korean government (MSIT) (No. 2020R1I1A2069379 and 2023R1A2C1004867).

Author Contributions

Conceptualization: Chung YS, Joo ST. Data curation: Kim SH, Kumari S. Formal analysis: Kim SH, Kumari S, Joo ST. Methodology: Kumari S, Kim CJ, Lee EY. Software: Kim SH, Kim CJ, Hwang YH. Validation: Kim SH, Alam AN. Investigation: Kumari S, Chung YS, Hwang YH, Joo ST. Writing - original draft: Kim SH, Kumari S, Joo ST. Writing - review & editing: Kim SH, Kumari S, Kim CJ, Lee EY, Alam AN, Chung YS, Hwang YH, Joo ST.

Ethics Approval

The procedure for animal use and treatment was approved by the Institutional Animal Care and Use Committee (IACUC) of Gyeongsang National University (approval no. GNU-231017-C0196). All experimental processes were conducted in accordance with the IACUC standard procedure.

References

- Alam AN, Kim CJ, Kim SH, Kumari S, Lee SY, Hwang YH, Joo ST. 2024. Trends in hybrid cultured meat manufacturing technology to improve sensory characteristics. *Food Sci Anim Resour* 44:39-50.
- Annoh-Quarshie J. 2018. Development and comparison of processes for the extraction of dietary protein from yellow peas. Stellenbosch University, Stellenbosch, South Africa.
- Bakhsh A, Lee SJ, Lee EY, Hwang YH, Joo ST. 2021. Evaluation of rheological and sensory characteristics of plant-based meat analog with comparison to beef and pork. *Food Sci Anim Resour* 41:983-996.
- Caine WR, Aalhus JL, Best DR, Dugan MER, Jeremiah LE. 2003. Relationship of texture profile analysis and Warner-Bratzler shear force with sensory characteristics of beef rib steaks. *Meat Sci* 64:333-339.
- Chandler SL, McSweeney MB. 2022. Characterizing the properties of hybrid meat burgers made with pulses and chicken. *Int J Gastron Food Sci* 27:100492.
- Cui B, Liang H, Li J, Zhou B, Chen W, Liu J, Li B. 2022. Development and characterization of edible plant-based fibers using a wet-spinning technique. *Food Hydrocoll* 133:107965.
- Cui M, Fang L, Zhou H, Yang H. 2014. Effects of amino acids on the physiochemical properties of potato starch. *Food Chem* 151:162-167.
- Dekkers BL, Boom RM, van der Goot AJ. 2018. Structuring processes for meat analogues. *Trends Food Sci Technol* 81:25-36.
- Ebert S, Baune MC, Broucke K, Van Royen G, Terjung N, Gibis M, Weiss J. 2021. Buffering capacity of wet texturized plant proteins in comparison to pork meat. *Food Res Int* 150:110803.
- Ferreira RMB, Teixeira ARN. 2003. Amino acids | metabolism. In *Encyclopedia of food sciences and nutrition*. 2nd ed. Caballero B (ed). Academic Press, Cambridge, MA, USA. pp 197-206.
- Fraeye I, Kratka M, Vandeburgh H, Thorrez L. 2020. Sensorial and nutritional aspects of cultured meat in comparison to traditional meat: Much to be inferred. *Front Nutr* 7:35.
- Grasso S. 2024. Opportunities and challenges of hybrid meat products: A viewpoint article. *Int J Food Sci Technol* 59:8693-8696.
- Grasso S, Jaworska S. 2020. Part meat and part plant: Are hybrid meat products fad or future? *Foods* 9:1888.
- He J, Zhao Y, Jin X, Zhu X, Fang Y. 2021. Material perspective on the structural design of artificial meat. *Adv Sustain Syst* 5:2100017.
- Hoek AC, Elzerman JE, Hageman R, Kok FJ, Luning PA, de Graaf C. 2013. Are meat substitutes liked better over time? A repeated in-home use test with meat substitutes or meat in meals. *Food Qual Prefer* 28:253-263.
- Ismail I, Hwang YH, Joo ST. 2020. Meat analog as future food: A review. *J Anim Sci Technol* 62:111-120.
- Ježek F, Kameník J, Macharáčková B, Bogdanovičová K, Bednář J. 2020. Cooking of meat: Effect on texture, cooking loss and microbiological quality: A review. *Acta Vet Brno* 88:487-496.
- Kim CJ, Kim SH, Lee EY, Son YM, Bakhsh A, Hwang YH, Joo ST. 2023. Optimal temperature for culturing chicken satellite

- cells to enhance production yield and umami intensity of cultured meat. *Food Chem Adv* 2:100307.
- Kim SH, Kim CJ, Lee EY, Son YM, Hwang YH, Joo ST. 2022. Optimal pre-plating method of chicken satellite cells for cultured meat production. *Food Sci Anim Resour* 42:942-952.
- Kumari S, Alam AN, Hossain MJ, Lee EY, Hwang YH, Joo ST. 2023. Sensory evaluation of plant-based meat: Bridging the gap with animal meat, challenges and future prospects. *Foods* 13:108.
- Kumari S, Kim SH, Kim CJ, Chung YS, Hwang YH, Joo ST. 2024. Development and comparative evaluation of imitated fiber from different protein sources using wet-spinning. *Food Sci Anim Resour* 44:1156-1166.
- Kyriakopoulou K, Dekkers B, van der Goot AJ. 2019. Plant-based meat analogues. In *Sustainable meat production and processing*. Galanakis CM (ed). Academic Press, Cambridge, MA, USA. pp 103-126.
- Lee CC, Tomas M, Jafari SM. 2020. Optical analysis of nanoencapsulated food ingredients by color measurement. In *Characterization of nanoencapsulated food ingredients*. Jafari SM (ed). Elsevier, Amsterdam, The Netherlands. pp 505-528.
- Manning PD. 1950. US Patent No. 2,505,129. Patent and Trademark Office, Washington, DC, USA.
- McClements DJ, Grossmann L. 2021. The science of plant-based foods: Constructing next-generation meat, fish, milk, and egg analogs. *Compr Rev Food Sci Food Saf* 20:4049-4100.
- Molfetta M, Morais EG, Barreira L, Bruno GL, Porcelli F, Dugat-Bony E, Bonnarme P, Minervini F. 2022. Protein sources alternative to meat: State of the art and involvement of fermentation. *Foods* 11:2065.
- Nagamine S, Akagi M, Nakagawa K, Kobayashi T. 2023. Effect of coagulant concentration on soy protein-based fiber for meat substitute by wet spinning combined with ionic cross-linking of alginate. *J Chem Eng Japan* 56:2256377.
- Paradowska-Stolarz A, Wieckiewicz M, Owczarek A, Wezgowiec J. 2021. Natural polymers for the maintenance of oral health: Review of recent advances and perspectives. *Int J Mol Sci* 22:10337.
- Qi J, Wang H, Zhou G, Xu X, Li X, Bai Y, Yu X. 2017. Evaluation of the taste-active and volatile compounds in stewed meat from the Chinese yellow-feather chicken breed. *Int J Food Prop* 20:S2579-S2595.
- Tagliamonte S, Oliviero V, Vitaglione P. 2024. Food bioactive peptides: Functionality beyond bitterness. *Nutr Rev* (in press). doi: 10.1093/nutrit/nuae008
- Treich N. 2021. Cultured meat: Promises and challenges. *Environ Resour Econ* 79:33-61.
- Wu G. 2021. *Amino acids: Biochemistry and nutrition*. CRC Press, Boca Raton, FL, USA.
- Xu S, Falsafi SR. 2023. Juiciness of meat, meat products, and meat analogues: Definition, evaluation methods, and influencing factors. *Food Rev Int* 40:2344-2377.
- Younis K, Ashfaq A, Ahmad A, Anjum Z, Yousuf O. 2023. A critical review focusing the effect of ingredients on the textural properties of plant-based meat products. *J Texture Stud* 54:365-382.
- Zahari I, Ferawati F, Purhagen JK, Rayner M, Ahlström C, Helstad A, Östbring K. 2021. Development and characterization of extrudates based on rapeseed and pea protein blends using high-moisture extrusion cooking. *Foods* 10:2397.