

**SHORT COMMUNICATION**

# Assessment of Heat Processing Effects on Cortisol Concentration in Dairy Milk Products

 Mohammad Ataallahi, Geun-Woo Park, Eska Nugrahaeningtyas,  
 and Kyu-Hyun Park\*

 Department of Animal Industry Convergence, Kangwon National University,  
 Chuncheon 24341, Korea

 OPEN ACCESS

**Received** February 21, 2024

**Revised** April 29, 2024

**Accepted** May 1, 2024

\***Corresponding author** : Kyu-Hyun Park  
 Department of Animal Industry Convergence,  
 Kangwon National University, Chuncheon  
 24341, Korea  
 Tel: +82-33-250-8621  
 E-mail: [kpark74@kangwon.ac.kr](mailto:kpark74@kangwon.ac.kr)

\***ORCID**  
 Mohammad Ataallahi  
<https://orcid.org/0000-0003-0234-8863>  
 Geun-Woo Park  
<https://orcid.org/0000-0003-0336-4080>  
 Eska Nugrahaeningtyas  
<https://orcid.org/0000-0002-4931-7952>  
 Kyu-Hyun Park  
<https://orcid.org/0000-0002-6390-5478>

**Abstract** The presence of cortisol residue in processed dairy milk may be a good parameter for assessing the quality of dairy milk products and an alternative indicator of the overall welfare of dairy cattle. Thus, this study investigated the impact of heat processing on milk cortisol concentration (MCC). In total, 36 milk samples (50 mL) were collected from three Holstein dairy cattle at a research farm over two consecutive days. The samples were divided into experimental groups: unheated, heated at 65°C for 30 min, and heated at 121°C for 5 min. Additionally, 11 commercial dairy milk products were purchased under three heating conditions: low temperature, low time (LTLT), ultra-short time (UST), and ultra-high temperature (UHT). MCC was analyzed using an enzyme immunoassay. The average farm MCC (ng/mL) for the unheated milk, milk heated at 65°C, and milk heated at 121°C were 0.88±0.16, 0.86±0.19, and 0.80±0.15, respectively. MCC was not significantly affected by the heating process. The average market MCC (ng/mL) in LTLT, UST, and UHT were 0.16±0.07, 0.15±0.08, and 0.15±0.07, respectively. Overall, cortisol levels in fresh farm milk were unaffected by the heating process. Monitoring cortisol levels in processed milk could offer a valuable alternative indicator for assessing product quality and animal welfare, particularly when access to raw milk is limited.

**Keywords** animal stress indicator, dairy product safety, milk cortisol analysis, thermal milk processing

## Introduction

Dairy farming is vital to the agricultural industry as a supplier of essential products such as milk and cheese. However, recent environmental changes have also affected cattle health and milk production (Bokharaeian et al., 2023; Guzmán-Luna et al., 2022). Therefore, ensuring the quality and safety of dairy products is of paramount importance to consumer health. It is also important for animal welfare and behavior (Castellini et al., 2023).

Stress has been identified as a factor that can negatively affect the health and welfare of animals, leading to lower milk production and quality (Bokharaeian et al., 2023; Park

et al., 2022). Heat stress can alter the biological composition of cattle by affecting hormone secretion, immune responses, and milk composition (Ma et al., 2019). In animal studies, cortisol levels are commonly monitored to detect stress responses and potential welfare issues (Ito et al., 2017). Cortisol can be detected in common biomatrices, such as blood, saliva, urine, hair, and milk, involving both stressful and non-stressful procedures to evaluate whether an animal's condition is normal or abnormal (Ataallahi et al., 2022; Ghassemi Nejad et al., 2022). Changes in cortisol levels in the bloodstream can affect milk cortisol concentration (MCC). The measurement of MCC is useful because abnormal levels have been linked to cattle stress or health conditions during milk production (Ataallahi et al., 2023; Fazio et al., 2015; Sgorlon et al., 2015).

Heat treatment is the most common processing method used to inactivate harmful components of dairy milk to ensure food safety and extend the shelf life of dairy milk products (Čurlej et al., 2022; Franzoi et al., 2022). Heat processing changes the structure of milk components such as proteins (Genene et al., 2019; Moro et al., 2019), vitamins, and volatile flavors (Coutinho et al., 2018) depending on the heat exposure of the milk during processing (Kilic-Akyilmaz et al., 2022; Lykholat et al., 2016). Monitoring the MCC in commercially processed milk products may be an alternative and noninvasive method for assessing the health and welfare of dairy cattle at the time of milk production. In addition, it may help ensure the safety and quality of processed dairy products. Therefore, this study aimed to explore the effects of various thermal processing methods, including low temperature, long duration, and high temperature, short duration on the cortisol concentrations in milk obtained from farms and markets.

## Materials and Methods

### Experimental animal

Three healthy Holstein dairy cattle (30 to 70 months old) with a milk yield between 19 and 24 liters, and a milk composition of 4.0% fat and 2.8% protein were used in the current study. They were housed in open pen with a winch curtain. The dairy cattle were fed with concentrate (12 kg/head/day), ray grasses (*Lolium*) and Sudan grasses (*Sorghum × drummondii*). All cattle were administered water *ad-libitum*. They belonged to a livestock research farm located at Kangwon National University in Chuncheon, Korea.


### Farm milk collection and preparation

Farm milk samples (n=36) were obtained within two days. Each sample was placed in a conical tube (50 mL) during regular milking in the morning (08:00) and afternoon (16:00). The sample tubes were placed in a cooled transport bag and transported to the laboratory and refrigerated at 4°C.

The whole milk tubes were vortexed, then centrifuged at 2,000×g for 20 min at 4°C and milk fat was removed from the surface (Fukasawa et al., 2008). Then, the skim milk was divided into three portions in new conical tubes. One part as a raw skim milk (unheated), the other two parts of skim milk were heated at 65°C for 30 min using a laboratory water bath (Wesolowska et al., 2019), and heated at 121°C for 5 min using an autoclave (Papapanagiotou et al., 2005). Milk samples were thoroughly mixed. The first aliquot of skim milk sample was placed in a water bath heated to 65°C for 30 min. The samples were then left to cool inside the water bath to 4°C. The same procedure was used for the second aliquot with the autoclave heated to 121°C at 5 min (Fig. 1).

### Commercial market milk collection and preparation

Eleven commercial dairy milk products were purchased for analysis. All market milk products were made from 100% raw



**Fig. 1.** Flow diagram for processing and analysis of farm and market milk.

milk and they were heated by either low temperature low time (LTLT) at 63°C for 30 min, ultra-short time (UST) at range 120°C to 130°C for 2 to 3 seconds, or ultra-high temperature (UHT) at range 135°C to 150°C for 1 to 4 seconds. For each milk product, 50 mL was sampled in a conical tube for subsequent analysis. No additional preparation steps such as centrifugation were required (Fig. 1).

### Milk cortisol analysis

MCC was analyzed using the procedures described by Ataallahi et al. (2023) and Fukasawa et al. (2008). For cortisol extraction, the milk samples were thawed in a 37°C water bath and 0.1 mL of milk was mixed with 0.9 mL diethyl ether under a fume hood. The mixture was vortexed for 1 min and two organic (ether phase) and inorganic (aqueous phase) layers were formed. The organic layer containing cortisol was transferred into a separate 2 mL microcentrifuge tube and evaporated under a laboratory fume hood for 2 h. The extracted residues were stored at -20°C for further analysis. It should be noted that multiple extractions (two or three times) were sufficient for the full extraction of cortisol, and the amount of evaporated organic layers should be measured for the final calculation. The dried evaporated residue was re-dissolved in 0.25 mL of assay buffer supplied with a colorimetric competitive enzyme immunoassay (ELISA) kit (Enzo Life Science, Farmingdale, NY, USA) at room temperature, vortexed for 1 min and 0.1 mL of the reconstituted sample was assayed in duplicate in a 96-well plate (Fig. 1). The MCC of morning and afternoon milk samples were measured individually. The sensitivity of the cortisol ELISA kit was 56.72 pg/mL. The OD of the samples was measured at a wavelength of 405 nm using a microplate reader (SpectraMax Absorbance Reader, Molecular Devices, San Jose, CA, USA). MCC was reported in ng/mL. The average coefficient of variation (CV%) calculated for farm MCC was 9.2% for the intra-assay and 14.4% for the inter-assay. The average CV% calculated for the commercial MCC was 15.4% for the intra-assay and 19.8% for the inter-assay.

## Statistical data analysis


Statistical analysis was performed using SAS software (version 9.4; SAS Institute, Cary, NC, USA). To compare the farm MCC before and after heat processing, as well as the MCC in commercial milk products, an analysis of variance (ANOVA) was used. Data were considered statistically significant at  $p < 0.05$ . All results are reported as mean  $\pm$  SD.

## Results and Discussion

### Effect of heat treatment on the milk cortisol concentration from farm milk

The results revealed that the average MCC (ng/mL) in unheated milk, heated milk at 65°C, and heated milk at 121°C, were  $0.88 \pm 0.16$ ,  $0.86 \pm 0.19$ , and  $0.80 \pm 0.15$  respectively (Fig. 2). Accordingly, there was no significant difference ( $p > 0.05$ ) in MCC between unheated and heat-treated milk samples or between the heat treatment methods.

Cortisol, the main stress hormone in animals, is found at high levels in the blood and can be transferred to the mammary glands of dairy animals (Hechler et al., 2018). The range of MCC from 0.5 to 11.7 ng/mL was observed in previous studies (Gellrich et al., 2015; Verkerk et al., 1998). In addition, Malekinejad and Rezabakhsh (2015), reported the range of glucocorticoid hormone concentrations to be between 0.46 to 18 ng/mL in dairy milk, 20 to 136 ng/mL in human milk, and 144 ng/mL in rat milk. In our study, the average farm MCC was  $0.83 \pm 0.17$  ng/mL, falling within the range of glucocorticoid concentrations (0.7 to 1.4 ng/mL) previously reported by Jouan et al. (2006) in milk products. We observed no significant differences in the MCC before and after heat processing. This is consistent with previous studies indicating that steroid hormones are unaffected by heat because cholesterol, a key substrate for steroid biosynthesis, comprises stable carbon-based ring structures, making the molecule more stable below its melting point of 148.5°C (Derewiaka and Molińska, 2015; Malekinejad and Rezabakhsh, 2015; Van Der Voorn et al., 2017). In the dairy products industry, thermal processing is typically performed at temperatures ranging from 63°C to 100°C (Franzoi et al., 2022; Snoj et al., 2018). We found that that temperatures between 65°C for 30 min and 121°C for 5 min, unaffected the level of cortisol.




**Fig. 2.** Boxplot displays the effect of heat treatments (low and high) on farm milk cortisol concentrations (MCC). Error bars represent the range of minimum and maximum cortisol values observed and the symbol  $\times$  represents the mean of data. The MCC was not statistically different ( $p > 0.05$ ).

### Measurement of milk cortisol concentration from market milk

The results revealed that the average MCC (ng/mL) of commercially milk products processed under LTLT, UST, and UHT were  $0.16 \pm 0.07$ ,  $0.15 \pm 0.08$ , and  $0.15 \pm 0.07$ , respectively (Fig. 3). No significant differences ( $p > 0.05$ ) were observed in the MCC between commercial milk products subjected to different heat processing methods.

Heat treatment is known to influence the composition of milk depending on the intensity, duration, and temperature of the storage conditions (Krishna et al., 2021). Previous studies have indicated that protein-based hormones, including insulin-like growth factors, growth hormones, leptin, adiponectin, and insulin, are more susceptible to denaturation during heating than lipid-based hormones such as cortisol (Arslanoglu et al., 2013; Van Der Voorn et al., 2017; van Lieshout et al., 2020). In this study, despite employing different heat treatments (LTLT, UST, and UHT), no significant differences were observed in the MCC, suggesting that the heat processing methods did not significantly affect the MCC. This finding suggests that cortisol residue in milk remains relatively stable despite variations in the heat treatment processes commonly used in commercial dairy products (Ataallahi et al., 2023; Malekinejad and Rezabakhsh, 2015). Cortisol is a 21 carbon steroid with the characteristic 3-keto-4-ene structure of active steroids (Honour, 2022). The structural stability of cortisol is because of presence of strong covalent bounds within the molecule, particularly the bounds between carbon atoms in the rings and the hydrogen atoms attached to them. The composition of milk is a complex mixture of proteins, fats, carbohydrates, and other ingredients. The stability of cortisol in milk can be affected by interactions with these components. Cortisol may form complexes with proteins present in milk, which can affect its degradation by enzymes.

Under normal conditions, 80% to 90% of circulating cortisol is bound to corticosteroid-binding globulin and 10% to 15%, is bound to albumin, and remaining 5% to 10% circulate as free and active hormone (Mormède et al., 2007). The basic molecular structure of cortisol remains the same whether its in the bloodstream or present in milk. But, when cortisol is found in milk, it may be in lower levels compared to its levels in the bloodstream, and it might also be found to carrier proteins specific to milk. Moreover, cortisol in milk might undergo some metabolic processes or interactions with other milk



**Fig. 3.** Boxplot displays the milk cortisol concentrations (MCC) in commercially milk products processed under three heat conditions. Low temperature low time (LTLT) at 63°C for 30 min, ultra-short time (UST) at range 120°C to 130°C for 2 to 3 seconds, or ultra-high temperature (UHT) at range 135°C to 150°C for 1 to 4 seconds. Error bars represent the range of minimum and maximum cortisol values observed. The outliers are shown as circles and the symbol × represents the mean of data. The MCC was not statistically different ( $p > 0.05$ ).

components, although these alterations typically do not change the fundamental structure of cortisol molecule itself. The exact extent of structural changes to cortisol during heat treatment can vary depending on factors such as temperature degree, duration of heating, and presence of other milk ingredients. High temperatures can lead to the denaturation of proteins, and it may affect cortisol indirectly by these processes.

Therefore, the stress levels in farm animals can be a determining factor affecting milk quality. The presence of cortisol in milk, a steroid hormone produced by the adrenal glands, has been linked to environmental stressors and poor welfare conditions in cattle (Jouan et al., 2006; Qu et al., 2018). However, only a small amount of glucocorticoid is transferred from the bloodstream to milk (Jouan et al., 2006; Qu et al., 2018). Poor welfare conditions not only affect milk production, leading to lower fat and protein levels (Castellini et al., 2023; Kawonga et al., 2012) but also affect the hormonal composition of milk. Despite the potential influence of poor welfare conditions on milk composition, our results suggest that the heat treatments commonly employed in the dairy industry do not significantly alter cortisol levels.

It is essential to note that milk quality is affected by various factors such as thermal processing (pasteurization and sterilization). Although effective at eliminating harmful components and bacteria, these processes may degrade heat-sensitive vitamins and enzymes, thereby reducing the nutritional value of milk (Macdonald et al., 2011; Vass et al., 2020). Furthermore, they may not effectively eliminate hormonal residues such as cortisol and estrogen. Understanding the stability of MCC under different heat treatment conditions has important implications for the dairy industry and product quality. This information will help improve the safety and quality of dairy products, while considering the welfare of dairy cattle. In addition, it will be explained whether the level of cortisol in milk and even processed milk can be used as an indicator of the overall good welfare of dairy farms. Monitoring hormonal residues, such as cortisol, in commercial processed milk may be a good parameter to evaluate the quality of milk and dairy products in the market.

## Conclusion

Cortisol in farm milk was unaffected by the thermal processes tested. Therefore, the cortisol concentrations in commercial milk are expected to be similar to those in fresh milk products. This suggests that the stress indicator cortisol, remains stable during common heat processes in dairy milk products. Moreover, monitoring cortisol residues in processed milk products could be an alternative indicator to improve the quality of milk and mitigate cattle stress if raw milk is unavailable.

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, and Forestry (IPET) and Korea Smart Farm R&D Foundation (KosFarm) through Smart Farm Innovation Technology Development Program, funded by Ministry of Agriculture, Food, and Rural Affairs (MAFRA) and Ministry of Science and ICT (MSIT), Rural Development Administration (RDA) (Grant number: 421013-03). The authors would like to special thanks to colleagues and staff at the livestock research farm of Kangwon National University for their kind assistance with sample collection.

## Author Contributions

Conceptualization: Park KH. Data curation: Ataallahi M, Park GW, Nugrahaeningtyas E. Formal analysis: Ataallahi M, Park GW, Nugrahaeningtyas E. Methodology: Ataallahi M, Park GW, Park KH. Software: Ataallahi M, Park GW, Nugrahaeningtyas E. Validation: Ataallahi M, Park GW. Investigation: Ataallahi M, Park GW, Park KH. Writing - original draft: Ataallahi M. Writing - review & editing: Ataallahi M, Park GW, Nugrahaeningtyas E, Park KH.

## Ethics Approval

The experimental procedures and methods were approved by the Institutional Animal Care and Use Committee (IACUC) of Kangwon National University, Chuncheon, Korea (KW-200520-1).

## References

- Arslanoglu S, Corpeleijn W, Moro G, Braegger C, Campoy C, Colomb V, Decsi T, Domellöf M, Fewtrell M, Hojsak I, Mihatsch W. 2013. Donor human milk for preterm infants: Current evidence and research directions. *J Pediatr Gastroenterol Nutr* 57:535-542.
- Ataallahi M, Cheon SN, Park GW, Nugrahaeningtyas E, Jeon JH, Park KH. 2023. Assessment of stress levels in lactating cattle: Analyzing cortisol residues in commercial milk products in relation to the temperature-humidity index. *Animals* 13:2407.
- Ataallahi M, Ghassemi Nejad J, Park KH. 2022. Selection of appropriate biomatrices for studies of chronic stress in animals: A review. *J Anim Sci Technol* 64:621-639.
- Bokharaeian M, Toghdory A, Ghoorchi T, Ghassemi Nejad J, Esfahani IJ. 2023. Quantitative associations between season, month, and temperature-humidity index with milk yield, composition, somatic cell counts, and microbial load: A comprehensive study across ten dairy farms over an annual cycle. *Animals* 13:3205.
- Castellini G, Barello S, Bosio AC. 2023. Milk quality conceptualization: A systematic review of consumers', farmers', and processing experts' views. *Foods* 12:3215.
- Coutinho NM, Silveira MR, Rocha RS, Moraes J, Ferreira MVS, Pimentel TC, Freitas MQ, Silva MC, Raices RSL, Senaka Ranadheera C, Borges FO, Mathias SP, Fernandes FAN, Rodrigues S, Cruz AG. 2018. Cold plasma processing of milk and dairy products. *Trends Food Sci Technol* 74:56-68.
- Čurlej J, Zajác P, Čapla J, Golian J, Benešová L, Partika A, Fehér A, Jakabová S. 2022. The effect of heat treatment on cow's milk protein profiles. *Foods* 11:1023.
- Derewiaka D, Molińska E. 2015. Cholesterol transformations during heat treatment. *Food Chem* 171:233-240.
- Fazio E, Medica P, Cravana C, Ferlazzo A. 2015. Release of  $\beta$ -endorphin, adrenocorticotrophic hormone and cortisol in response to machine milking of dairy cows. *Vet World* 8:284-289.
- Franzoi M, Costa A, Vigolo V, Penasa M, De Marchi M. 2022. Effect of pasteurization on coagulation properties of bovine milk and the role of major composition traits and protein fractions. *J Food Compos Anal* 114:104808.
- Fukasawa M, Tsukada H, Kosako T, Yamada A. 2008. Effect of lactation stage, season and parity on milk cortisol concentration in Holstein cows. *Livest Sci* 113:280-284.
- Gellrich K, Sigl T, Meyer HHD, Wiedemann S. 2015. Cortisol levels in skimmed milk during the first 22 weeks of lactation

- and response to short-term metabolic stress and lameness in dairy cows. *J Anim Sci Biotechnol* 6:1-7.
- Genee A, Hansen EB, Eshetu M, Hailu Y, Ipsen R. 2019. Effect of heat treatment on denaturation of whey protein and resultant rennetability of camel milk. *LWT-Food Sci Technol* 101:404-409.
- Ghassemi Nejad J, Ghaffari MH, Ataallahi M, Jo JH, Lee HG. 2022. Stress concepts and applications in various matrices with a focus on hair cortisol and analytical methods. *Animals* 12:3096.
- Guzmán-Luna P, Mauricio-Iglesias M, Flysjö A, Hospido A. 2022. Analysing the interaction between the dairy sector and climate change from a life cycle perspective: A review. *Trends Food Sci Technol* 126:168-179.
- Hechler C, Beijers R, Marianne Riksen-Walraven J, de Weerth C. 2018. Are cortisol concentrations in human breast milk associated with infant crying? *Dev Psychobiol* 60:639-650.
- Honour JW. 2022. *Steroids in the laboratory and clinical practice*. Elsevier, Amsterdam, The Netherlands. pp 3-32.
- Ito T, Aoki N, Tsuchiya A, Kaneko S, Akiyama K, Uetake K, Suzuki K. 2017. Detection of stress hormone in the milk for animal welfare using QCM method. *J Sens* 2017:6486891.
- Jouan PN, Pouliot Y, Gauthier SF, Laforest JP. 2006. Hormones in bovine milk and milk products: A survey. *Int Dairy J* 16:1408-1414.
- Kawonga BS, Chagunda MGG, Gondwe TN, Gondwe SR, Banda JW. 2012. Characterisation of smallholder dairy production systems using animal welfare and milk quality. *Trop Anim Health Prod* 44:1429-1435.
- Kilic-Akyilmaz M, Ozer B, Bulat T, Topcu A. 2022. Effect of heat treatment on micronutrients, fatty acids and some bioactive components of milk. *Int Dairy J* 126:105231.
- Krishna TC, Najda A, Bains A, Tosif MM, Papliński R, Kaplan M, Chawla P. 2021. Influence of ultra-heat treatment on properties of milk proteins. *Polymers* 13:3164.
- Lykholat OA, Grigoryuk IP, Lykholat TY. 2016. Metabolic effects of alimentary estrogen in different age animals. *Ann Agrar Sci* 14:335-339.
- Ma L, Yang Y, Zhao X, Wang F, Gao S, Bu D. 2019. Heat stress induces proteomic changes in the liver and mammary tissue of dairy cows independent of feed intake: An iTRAQ study. *PLOS ONE* 14:e0209182.
- Macdonald LE, Brett J, Kelton D, Majowicz SE, Snedeker K, Sargeant JM. 2011. A systematic review and meta-analysis of the effects of pasteurization on milk vitamins, and evidence for raw milk consumption and other health-related outcomes. *J Food Prot* 74:1814-1832.
- Malekinejad H, Rezabakhsh A. 2015. Hormones in dairy foods and their impact on public health: A narrative review article. *Iran J Public Health* 44:742-758.
- Mormède P, Andanson S, Aupérin B, Beerda B, Guémené D, Malmkvist J, Manteca X, Manteuffel G, Prunet P, van Reenen CG, Richard S. 2007. Exploration of the hypothalamic–pituitary–adrenal function as a tool to evaluate animal welfare. *Physiol Behav* 92:317-339.
- Moro GE, Billeaud C, Rachel B, Calvo J, Cavallarin L, Christen L, Escuder-Vieco D, Gaya A, Lembo D, Wesolowska A, Arslanoglu S, Barnett D, Bertino E, Boquien CY, Gebauer C, Grovslie A, Weaver GA, Picaud JC. 2019. Processing of donor human milk: Update and recommendations from the European Milk Bank Association (EMBA). *Front Pediatr* 7:49.
- Papapanagiotou EP, Fletouris DJ, Psomas EI. 2005. Effect of various heat treatments and cold storage on sulphamethazine residues stability in incurred piglet muscle and cow milk samples. *Anal Chim Acta* 529:305-309.
- Park GW, Ataallahi M, Ham SY, Oh SJ, Kim KY, Park KH. 2022. Estimating milk production losses by heat stress and its



- impacts on greenhouse gas emissions in Korean dairy farms. *J Anim Sci Technol* 64:770-781.
- Qu X, Su C, Zheng N, Li S, Meng L, Wang J. 2018. A survey of naturally-occurring steroid hormones in raw milk and the associated health risks in Tangshan City, Hebei Province, China. *Int J Environ Res Public Health* 15:38.
- Sgorlon S, Fanzago M, Guiatti D, Gabai G, Stradaoli G, Stefanon B. 2015. Factors affecting milk cortisol in mid lactating dairy cows. *BMC Vet Res* 11:1-8.
- Snoj T, Zuzek MC, Cebulj-Kadunc N, Majdic G. 2018. *Short communication*: Heat treatment and souring do not affect milk estrone and  $17\beta$ -estradiol concentrations. *J Dairy Sci* 101:61-65.
- Van Der Voorn B, De Waard M, Dijkstra LR, Heijboer AC, Rotteveel J, Van Goudoever JB, Finken MJ. 2017. Stability of cortisol and cortisone in human breast milk during holder pasteurization. *J Pediatr Gastroenterol Nutr* 65:658-660.
- van Lieshout GAA, Lambers TT, Bragt MCE, Hettinga KA. 2020. How processing may affect milk protein digestion and overall physiological outcomes: A systematic review. *Crit Rev Food Sci Nutr* 60:2422-2445.
- Vass RA, Bell EF, Colaizy TT, Schmelzel ML, Johnson KJ, Walker JR, Ertl T, Roghair RD. 2020. Hormone levels in preterm and donor human milk before and after Holder pasteurization. *Pediatr Res* 88:612-617.
- Verkerk GA, Phipps Am, Carragher JF, Matthews LR, Stelwagen K. 1998. Characterization of milk cortisol concentrations as a measure of short-term stress responses in lactating dairy cows. *Anim Welfare* 7:77-86.
- Wesolowska A, Sinkiewicz-Darol E, Barbarska O, Bernatowicz-Lojko U, Borszewska-Kornacka MK, van Goudoever JB. 2019. Innovative techniques of processing human milk to preserve key components. *Nutrients* 11:1169.